Defining subpopulations of differential drug response to reveal novel target populations

Nirmal Keshava [1,#], Tzen S. Toh [2,3,#], Haobin Yuan[4], Bingxun Yang[4], Michael P. Menden [5,6,7,*], Dennis Wang [3,4,8*]

1. Constellation Analytics, LLC., Needham MA, USA
2. The Medical School, University of Sheffield, Sheffield, S10 2RX, UK
3. Sheffield Institute for Translational Neuroscience, University of Sheffield, Sheffield, S10 2HQ, UK
4. Department of Computer Science, University of Sheffield, Sheffield, S1 4DP, UK
5. Institute of Computational Biology, Helmholtz Zentrum München - German Research Center for Environmental Health, 85764 Neuherberg, Germany
6. Department of Biology, Ludwig-Maximilians University Munich, 82152 Martinsried, Germany
7. German Centre for Diabetes Research (DZD e.V.), 85764 Neuherberg, Germany
8. NIHR Sheffield Biomedical Research Centre, Sheffield, S10 2HQ, UK , UK

[#] These authors contributed equally to this work.
* These authors are corresponding authors.
Contacts: Michael P. Menden (michael.menden@helmholtz-muenchen.de) or Dennis Wang (dennis.wang@sheffield.ac.uk)

Abstract

Personalised medicine has predominantly focused on genetically-altered cancer genes that stratify drug responses, but there is a need to objectively evaluate differential pharmacology patterns at a subpopulation level. Here, we introduce an approach based on unsupervised machine learning to compare the pharmacological response relationships between 327 pairs of cancer therapies. This approach integrated multiple measures of response to identify subpopulations that react differently to inhibitors of the same or different targets to understand mechanisms of resistance and pathway cross-talk. MEK, BRAF, and PI3K inhibitors were shown to be effective as combination therapies for particular *BRAF* mutant subpopulations. A systematic analysis of preclinical data for a failed phase III trial of selumetinib combined with docetaxel in lung cancer suggests potential indications in urogenital and colorectal cancers with *KRAS* mutation. This data-informed study exemplifies a method for stratified medicine to identify novel cancer subpopulations, their genetic biomarkers, and effective drug combinations.

Keywords

patient stratification, machine learning, biomarker discovery, precision medicine, cancer genomics, pharmacogenomics

Introduction

Drug developers face a conundrum in predicting the efficacy of their investigational
compound compared to existing drugs used as the standard of care treatment. Systematic
screening of drug compounds across a variety of genomic backgrounds in cancer cell lines
has improved clinical trial design and personalized treatments [1]. Following the pioneering
NCI-60 screen comprised of 59 unique cell lines [2], modern high-throughput screens such as
the Genomics of Drug Sensitivity in Cancer (GDSC) [3,4], the Cancer Cell Line Encyclopedia
(CCLE) [5] and the Cancer Therapeutics Response Portal (CTRP) [6–8] have characterised
>1,000 cancer cell lines with the goal of establishing the genetic landscape of cancer. The
deep molecular characterisation of these large cell line panels is complemented with high-
throughput drug screens, which enables the discovery of drug response biomarkers. For
example, analysis of the generic BRAF inhibitors PLX4720, SB590885 and CI-1040
reproduced drug sensitivity association with *BRAF* mutation in melanoma, or afatinib
sensitivity with *ERBB2* amplifications in breast cancer [3,4,9]. These associations between
genetic variants and treatment response have helped identify specific patient subpopulations
who are most likely to benefit from treatment. In Phase III clinical trials, however, for new
drugs to be successful, they must demonstrate a significant improvement over the existing
standard of care. Accurately defining in which subpopulations a new drug demonstrates
improved differential efficacy over other drugs targeting the same disease could lead to both
better clinical outcomes as well as new targeted therapies.

While several methods have been proposed to identify drug response biomarkers in cell
lines for precision medicine and drug repositioning [4,5,10,11], there is a need for more objective
and unsupervised approaches for identifying subpopulations with differences in drug
response (differential drug response), and consequently systematically gain mechanistic
insights from biomarkers. Most approaches capable of comparing multiple drugs measure
the overall similarity (or correlation) based on a single response summary metric [7,12], which
permits drug repositioning based on subpopulations with similar behavior, but neglects ones
that behave differently (**Figure S1A**). Here, we used an unsupervised technique to identify
the perimeters of differentially sensitive or resistant subpopulations and which may be
generalized to stratify the pharmacology response for any pair (or n-tuple) of targets using
any number of drug response summary metrics (e.g. IC_{50} or AUC). Segmentation of the
overall population occurs top-down and along globally-optimal contours that are derived
explicitly and maximize the differences between the two resulting subpopulations. The
segmentation continues recursively and is modulated by multiple user-defined criteria such
as the size or separability of the resulting subpopulations. Higher threshold values for both
result in less granular subpopulations but increase certainty that the subpopulations and the
quantities estimated from them are both distinct and accurate.

We present results from our platform, SEABED (SEgmentation And Biomarker Enrichment
of Differential treatment response), to demonstrate how unsupervised machine learning can
discover intrinsic partitions in the drug response measurements of two or more drugs that
directly correspond to distinct pharmacological patterns of response with therapeutic
biomarkers. Addressing the challenges in comparing the response of two drugs, SEABED
initially assesses two gold standards with established clinical biomarkers, namely the
differential response of a BRAF inhibitor and MEK inhibitor with anticipated *BRAF* and *KRAS*
mutations [13–16], and an EGFR inhibitor and MEK inhibitor with expected biomarkers of
EGFR, *ERBB2* and *KRAS* mutations [17–20]. Next, we systematically compare how different
drugs targeting the MAPK and PI3K-AKT pathway yield different patterns of response within
subpopulations. We show how differential drug response may indicate benefit for drug

combinations explained through independent action rather than probable synergy by
examining subpopulations uniquely sensitive to a single drug [21], which may be precisely
targeted by identified biomarkers. Finally, we demonstrate how the analysis of differential
response can guide the design of clinical trials by revealing specific indications where an
investigational therapy may be more effective than the standard treatment.

103 Results

We applied our technique to discover subpopulations of cell lines in which two or more
compounds, possibly addressing the same disease state or even targeting the same genetic
alteration, have a common pharmacological pattern of response. By further associating
enriched genetic alterations in subpopulations with specific patterns of response, we shed
light into molecular mechanisms responsible for patient subpopulations that respond
differently to two drugs.

Identifying subpopulations of differential drug response
We first considered the specific circumstance in which two drugs engage different targets
within the same signalling pathway, namely agents targeting MAPK signaling. SEABED used
nearly 1,000 cancer cells derived from the GDSC database, and we evaluated two
established drug response measures: the drug concentration required to reduce cell viability
by half (IC_{50}) and the area under the dose-response curve (AUC; **Figure 1A**). SEABED
employed a multivariate similarity measure to compare the vector patterns of response for
each distinct pair of cell lines without requiring *a priori* assumptions on the number or
distribution of the subpopulations. The result is a diverse cell line population segmented into
distinct subpopulations having homogeneous patterns of drug response (**Figure 1B**). Here
exemplified, we show that the drug response of 802 cell lines treated with either SB590885
(BRAF inhibitor) or CI-1040 (MEK inhibitor) could be segmented into 7 distinct
subpopulations with a median size of 40 cell lines by integrating the two metrics of drug
response, AUC and IC_{50} (**Figure 1C;** see **Figures S1B and S1C** for individual cell lines
segmented by IC_{50} and AUC respectively). We comprehensively evaluate pan-cancer
somatic events to nominate biomarkers (**see Methods**) [4], and found that the subpopulation
sensitive to both inhibitors was significantly enriched for *BRAF* mutants (P=3.87e-14,
hypergeometric test), while another subpopulation was exclusively sensitive to the MEK
inhibitor and significantly enriched for *KRAS* mutations (P=0.00589, hypergeometric test).

In another example we examined a case where one inhibitor might overcome resistance to
another inhibitor targeting the same pathway; AZD6244/ARRY-142886 selumetinib (MEK
inhibitor) with afatinib (EGFR and ERBB2 dual inhibitor) across 839 cell lines (**Figure 1D**).
Strong markers of sensitivity for selumetinib are subpopulations carrying known associated
KRAS, *NRAS* and *BRAF* mutations (**Figures 1D and 1E**). A less anticipated association is
APC loss-of-function sensitivity to selumetinib, albeit this was also found with trametinib
(another MEK inhibitor) in APC deficient mice [22]. We reproduced the well-established
associations of afatinib with either *EGFR* and *ERBB2* amplifications [4,23], and surprisingly our
unsupervised segmentation returned two subpopulations enriched for *EGFR* amplifications.
The more sensitive subpopulation is solely enriched for *EGFR* amplifications, whilst the less
sensitive subpopulation additionally includes activating *PIK3CA* mutations. In concordance
with recent literature, PI3K-AKT signaling drives acquired drug resistance to EGFR inhibitors
in lung cancer [24].

Drug response segmentation resulted in 14 subpopulations with a median size of 37 (**Figure
1D**). The subpopulation enriched for *EGFR*, *ERBB2* and *PI3KCA* variants, has an average
$\log(IC_{50})$ of 1.01µM for selumetinib and -0.672µM for afatinib. In contrast, the *BRAF* mutation
was enriched in a subpopulation where the average $\log(IC_{50})$ for selumetinib was -1.097µM

and 0.625µM for afatinib. The difference in response between afatinib and selumetinib was
significantly greater (t-test P<0.01) between the subpopulations identified and the total
population of *PIK3CA* or *BRAF* mutant cell lines (**Figures 1F and 1G**).
153
Cross-comparison of multiple drugs redefines best-in-class drugs for specific
subpopulations
Although there is a larger portfolio of clinical drugs with identical putative targets, their
responses may differ substantially in subpopulations as a consequence of multiple factors,
for example mode-of-action, different off-target effects and binding properties. The ability to
discover cell line subpopulations with distinct pharmacological patterns of response
characterised by genetic mutations re-defines best-in-class drugs by their differential
response to other drugs in a specific subpopulation, rather than their absolute response
across an entire population.
163
In order to demonstrate this approach for drug discovery, we applied SEABED to 745 cell
lines across cancer types to evaluate the differential response in those cell lines to five
inhibitors (CI-1040, PD0325901, RDEA119, selumetinib, and trametinib) which all target the
MEK protein (**Figure 2A**). The segmentation of cell lines revealed 13 subpopulations with
different patterns of response and three having enriched biomarkers (**Figure S2A**). Two
subpopulations were sensitive to all MEK inhibitors, with trametinib achieving the greatest
sensitivity. In one subpopulation the *KRAS* mutation was enriched (Fisher exact p-value =
1.12e-4 and 40.8% of the cell lines) while another had the *BRAF* mutation enriched (Fisher
exact p-value = 1.39e-7 and 50% of the cell lines). In contrast, another subpopulation was
enriched with the *RB1* mutation (Fisher exact p-value = 3.84e-2 and 21.6% of cell lines),
within which the cell lines were almost uniformly resistant to all MEK inhibitors.
175
Distribution of subpopulations highlight distinct pharmacological relationships
between PI3K-AKT and MAPK signaling
Next, we used SEABED to investigate the cross-talk between two frequently active cancer
pathways, MAPK and PI3K-AKT signalling, by systematically comparing pairs of drugs
targeting different genes of each pathway (**Figure 2A**, **B**). In total, SEABED performed 324
pairwise comparisons of 18 PI3K-AKT and 18 MAPK pathway inhibitors. Each drug pair was
classified into five categories based on the distribution of subpopulation responses: (i) no
preferential response, (ii) sensitive to both MAPK and PI3K-AKT pathway inhibitors (**Figure**
S2B), (iii) greater sensitivity to MAPK pathway inhibitors (**Figure S2C**), (iv) greater sensitivity
to PI3K-AKT pathway inhibitors (**Figure S2D**), (v) sensitive to either a MAPK pathway or a
PI3K-AKT pathway inhibitor, i.e. divergent response (**Figure S2E**). While this approach
classified the distribution of subpopulations based on somewhat arbitrary thresholds for
sensitivity, we also measured the weighted average Pearson correlation across
subpopulations to identify interesting drug pairs (**Figure S3**).
190
We found 20 drug pairs subpopulations with sensitivity to both PI3K-AKT and MAPK
pathway inhibition. This association between subpopulation size and sensitive response was
significant when comparing a CRAF inhibitor (TL-2-105) to PI3K-AKT signaling inhibitors
(hypergeometric test P=2.358e-5). The same trend was observed for inhibiting ERK (FR-
180204) or RSK (FMK) compared to inhibiting any PI3K-AKT signaling gene (P=0.0185 and
P=2.358e-5, respectively; hypergeometric test), but interestingly there was no mutual
sensitivity when comparing to BRAF inhibitors and only two cases for MEK inhibitors.
198
There were 20 drug pairs with a significantly high proportion of subpopulations (P < 0.05)
exhibiting greater sensitivity to MAPK pathway inhibition. This phenotype is strongly
pronounced in pairs with BRAF inhibitors (hypergeometric test P=0.0166). 20 drug pairs
were found with significantly high proportions of preferential PI3K-AKT pathway inhibition.
This is observed when comparing MEK inhibitors (CI-1040 and selumetinib-1) to PI3K-AKT
signaling inhibitors (P=0.0185 and P=0.00273 respectively; hypergeometric test).

205
In 22 cases, we observed drug pairs with sensitivity to either a MAPK pathway or a PI3K-
AKT pathway inhibitor, i.e. divergent response. This response type was enriched for pairs of
any PI3K-AKT pathway inhibitors and BRAF (PLX4720-2) or MEK inhibitors (P=0.026 and
P=0.0135, respectively; hypergeometric test), and even significant for PI3K inhibitors in
comparison with either the EGFR (erlotinib) or MEK inhibitors (P=0.0315 and P=0.0173;
hypergeometric test). Response patterns for all drug pairs can be explored in our portal
(**Website S1**; https://szen95.github.io/SEABED).
213

Subpopulations of differential response identifies drug combination efficacy
Previous studies have hypothesised that the efficacy of many approved drug combinations
can be explained by the independent action of single agents on different patient
subpopulations with cancers driven by multiple pathways [21]. We hypothesised that SEABED
comparisons of drug pairs would highlight subpopulations of differential response that would
exhibit synergistic or independent action effects when the drugs are tested in combination.
Additionally, our method enables to explore putative biomarkers of such populations.
Systematic comparison of responses between two drugs highlighted subpopulations of cell
lines in which there was sensitivity to either drug but not both (divergent response). We
observed this phenomenon in 22 drug pair comparisons, including a MEK inhibitor
(RDEA119-2) which showed divergent responses to four PI3K inhibitors (PI-103,
GSK2126458, ZSTK474, and CAL-101; **Figure 3A**; **Figures S4A-D** respectively). Drug pairs
with divergent response were also observed in cell lines treated with PLX4720-2 (BRAF
inhibitor) and two PI3K inhibitors (PI-103 and ZSTK474; **Figures 3B and S4E; Figure S4F**).
Two subpopulations with a high proportion of *BRAF* mutations were identified with greater
sensitivity to the BRAF inhibitor (**Figure 3B**). These two subpopulations had a combined
total of 79 cell lines (**Figure 3C**). The subpopulation with an average log(IC_{50}) of -0.0232µM
for PLX4720-2 and 0.771µM for PI-103 had 21 cell lines with *BRAF* mutation (87.5%;
P=9.75e-17). In contrast, the subpopulation with an average log(IC_{50}) of 0.808µM for
PLX4720-2 and 0.562µM for PI-103 had 22 cell lines with *BRAF* mutation (40.4%; P=5.12e-
7) (**Figure 3B**). We observed 17 individual cell lines with *BRAF* mutation that are resistant to
both drugs (**Figure S4G**).
236

We next examined the drug pairs as combination therapies in cell lines [25] and patient-
derived tumor xenograft models (PDXs) [26] to investigate whether the drug pairs with
divergent response and subpopulations with preferential sensitivity to one drug would be
associated with efficacy of their combination treatment (**Figure 3D**). SEABED first compared
the single drug responses of BRAF, MEK and PI3K inhibitors as before to identify BRAF
mutant subpopulations with differential response. When the drugs were tested as
combinations in BRAF mutant cell lines, the MEK/PI3K inhibitor combination had a surprising
similar level of synergy as BRAF/MEK combinations, which was recently a clinically
approved combination [27,28]. Also surprising, these two combinations had significantly higher
synergistic effect when used on BRAF mutant cell lines compared to all cell lines (t-test
P=0.0204), and compared to all drug combinations tested (t-test P=1.46e-5; **Figure 3E**;
Figure S4H). In terms of overall efficacy in PDXs, we observed a similar level of inhibition to
tumour volume for the BRAF/PI3K inhibitor combination on BRAF mutant cells when
compared to the clinically approved BRAF/MEK combination and a significantly greater (t-
test P=0.0418) inhibition of tumour growth compared to all combinations (**Figure 3F**; **Figure
S4I**). Notably, previous work suggested *in vivo* efficacy of drug combinations is mostly driven
by the monotherapy agents targeting independent mechanisms [26]; however, they did not
exclude the possibility that such drug combinations may also be synergistic. Here, we
highlight a drug combination example, where *in vivo* efficacy is driven by targeting
independent mechanisms, and complementary being synergistic. This example highlights
that both concepts, synergy and targeting independent mechanism may contribute to
combination efficacy in patients.
259

Lack of subpopulations of differential response may explain clinical failure

Sometimes, despite strong preclinical evidence, some drugs do not succeed in clinical trials [29]. One such trial was SELECT-1 (**Table S1**) which compared the efficacy of combining selumetinib and docetaxel to docetaxel alone in patients with advanced *KRAS*-mutant non–small cell lung cancer (NSCLC) [30]. Although there were *KRAS* mutant cell lines sensitive to selumetinib in preclinical testing [31], we re-examined the pharmacological data with SEABED to assess whether there were distinct subpopulations that justified the patient selection criteria for *KRAS* mutation.

In this analysis, instead of only inspecting the subpopulation identified by SEABED when the segmentation algorithm terminated, we thoroughly examined all possible subpopulations. SEABED identified a total of 61 possible subpopulations from 840 cell lines across tissue types tested with selumetinib and docetaxel (**Figure 4A**). 10 subpopulations were more sensitive to selumetinib than docetaxel (**Figure 4B**), and 5 of those subpopulations were enriched for *KRAS* mutation. However, those subpopulations enriched for NSCLC *KRAS* mutants were small in size and mostly exhibited less sensitivity to selumetinib compared to docetaxel (**Figures S5A and S5B**). The distribution of different *KRAS* mutations (p.G12C vs p.G12V) was also no different in selumetinib sensitive subpopulations compared to resistant subpopulations (**Figures S5C and S5D**). Independent of mutation status, only 8.9% of NSCLC cell lines were found in subpopulations more sensitive to selumetinib, whereas 14.8% cell lines originating from aerodigestive cancer types (eg. esophageal) were found in these subpopulations (**Figures S5E and S5F**).

Next, we focused on *subpopulation_60*, which had the greatest difference in sensitivity (IC_{50} and AUC) to selumetinib compared to docetaxel (**Figure 4C**). This subpopulation of 122 cell lines was enriched in *KRAS* mutations (28.8%, P=3.061e-4) found across multiple tissue types. NSCLC cell lines accounted for only 8% of this subpopulation, with 50% of those cell lines being *KRAS* mutants. Colorectal and pancreatic cell lines accounted for 15% and 8% respectively of the subpopulation, and they both had a higher proportion of *KRAS* mutations (56% and 100% respectively; **Figure 4D**).

Discussion

The ability to identify distinct subpopulations based on multiple measures of drug response (eg. IC_{50} and AUC) and extract their biomarkers is the basis for personalised therapeutics, which may ultimately increase the likelihood of successful clinical trials [32,33]. Using a network-based segmentation algorithm coupled with biomarker detection (SEABED), we investigated well-established pharmacological targets and clinical biomarkers by comparing the response patterns for BRAF (SB590885) and MEK (CI-1040) inhibition, which expectedly reproduced subpopulations sensitive to both enriched for *BRAF* mutants [34–36]. In another example, SEABED compared EGFR/ERBB2 (afatinib) and MEK (selumetinib) inhibition to reveal expected biomarkers such as *BRAF*, *KRAS* and *NRAS* mutations for selumetinib [13–16], and afatinib associated with *EGFR* and *ERBB2* amplifications [37,38]. Interestingly, the more afatinib-resistant subpopulation was enriched for *PI3KCA*-activating mutation, which may cause acquired resistance [24]. When we systematically compared inhibitors of the MAPK and PI3K-AKT signaling pathways, we observed subpopulations sensitive to both CRAF, ERK or RSK targeted drugs and other drugs targeting the PI3K-AKT pathway, however, there were few instances of these subpopulations for inhibitors targeting other genes in the MAPK signalling [39]. We found many more subpopulations that were more sensitive to BRAF inhibitors than other PI3K-AKT inhibitors, and as expected, many contained *BRAF* mutations [34]. In contrast, there were not significantly more subpopulations sensitive to MEK inhibition compared to inhibition of PI3K-AKT signalling targets, but *BRAF* mutant subpopulations may have greater differential response [14]. Divergent response was observed when comparing EGFR, BRAF and MEK inhibitors to drugs targeting the PI3K-AKT pathway. Our results

314　comparing the MAPK and PI3K-AKT pathways based on drug response profiles highlights
315　how intertwined those two pathways are in pharmacology space [39].
316
Arguably, the divergent response type is the most exciting for personalised treatment, since
it may identify cases where independent drug action and synergy may guide effective drug
combinations [21]. Here exemplified, we showed that PI3K inhibitors combined with either
BRAF or MEK inhibitors increase *in vitro* synergy and reduce tumour volume of *in-vivo*
models. Furthermore, we were able to show that synergistic and overall effect can be further
enhanced by the correct biomarker indication, in this instance, *BRAF* mutant subpopulations
[40,41]. The *BRAF* mutant subpopulation with high efficacy for the BRAF inhibitor and not the
other inhibitor could be cases where independent drug action explains drug combination
efficacy, whereas, the subpopulation with lower efficacy for single treatments of either drug
may be cases for synergistic effects when the drugs are combined.
327
In examining the preclinical evidence for trial testing combination treatment of NSCLC in
which the *KRAS* mutation was the biomarker [42], SEABED revealed a high proportion of
NSCLC subpopulations having the *KRAS* mutation that are resistant to both selumetinib and
docetaxel, suggesting a smaller likelihood of efficacy for the drug combination. Alternately,
we identified a subpopulation with differential response to selumetinib for a small proportion
of *KRAS* NSCLC cell lines, but this subpopulation contained a higher proportion of colorectal
and pancreatic cancer cells with *KRAS* mutations. Previous studies have shown the
plausibility in treating colorectal cancer using MEK inhibitor combinations [43,44]. With
consideration of *KRAS* mutations in subpopulations having greater sensitivity to selumetinib,
SEABED suggests that while the correct biomarker was used for the clinical trial, there may
be other potential indications for selumetinib. Although response in cell lines may not always
correspond to response clinically, the use of data-informed approaches to examine large
populations of cells may reveal clinically relevant drug response patterns. Future studies
may need to account for differences between *in vitro* and *in vivo* responses.
342
While iterative and hierarchical clustering techniques have been used widely to attribute
molecular markers to differences in subpopulation drug response and outcomes [45,46], we use
an approach that does not require an explicit estimate of the number of subpopulations and
is not greedy, i.e., each incremental step is optimal but the overall algorithm is not. In
biomedical data processing, there has been substantial concern, particularly regarding
applications to molecular data, that rival unsupervised machine learning optimize different
criteria and consequently yield diverging answers [47,48]. In our effort, we are concerned with
discovering subpopulations having high homogeneity and statistical separability, while
avoiding subpopulations that are so small that extracting statistically significant biomarkers is
unlikely. We demonstrated the utility of SEABED over conventional approaches of K-Means
and hierarchical clustering for drug response comparisons (see **Supplementary Materials,**
Table S2, Table S3, and Figure S6). Further, the ability to segment non-convex regions,
which can arise to describe different disease states, is advantageous. Consequently,
because of their success in other industries [49,50] as well as fast, efficient implementations,
spectral methods based on network models are powerful methods for discovering distinct
subpopulations [51–53].
359
SEABED builds upon previous work using network models in biomedical contexts [51,54,55] that
explicitly partition a population of cell lines described by multiple variables into distinct
subpopulations, using a "top-down" approach of recursively identifying optimal cuts for graph
bisection. Similarly, while our segmentation capitalizes on past progress made in spectral
clustering [56,57], our effort distinguishes itself from past attempts by integrating all variables
into a single network model using a multivariate similarity and local and global network
statistics. Deeper interpretations of embedded matrix subspaces in network models may
provide further insight into the linkage between subpopulations of cancer cell lines and
drugs.

369
As a whole, this study demonstrates several important insights about the pharmacological
pattern of response for different cancer drugs by applying an unsupervised machine learning
platform to segment a large pan-cancer *in vitro* pharmacology data set. By organizing cell
lines along similar pharmacological patterns of response, we identified distinct, intrinsic
subpopulations sensitive to one drug but resistant to others, and in some cases identified
genetic alterations that can be used as biomarkers for those subpopulations. In the context
of analytical frameworks for increasing drug R&D productivity by sharpening the focus of
drugs [58], our work demonstrates the value of advanced analytical approaches in translational
medicine to enable decision making that is more data-informed and less ambiguous.
Moreover, by analyzing different pharmacological responses and interpreting its outputs in
the context of the underlying genetics and molecular pathways, we have created a multi-
faceted landscape for developing and assessing new drug therapies.
382
383

Methods

385
386 **CONTACT FOR REAGENT AND RESOURCE SHARING**
387
388 All code for the pipeline is open source and available at: https://github.com/szen95/SEABED.
389 Further information and requests should be directed to and will be fulfilled by the co-
390 corresponding authors, Michael P. Menden (michael.menden@helmholtz-muenchen.de) or
391 Dennis Wang (dennis.wang@sheffield.ac.uk).
392
393 **METHOD DETAILS**
394
Pharmacology data
The discovery pharmacology dataset was extracted from the The Genomics of Drug
Sensitivity in Cancer (GDSC) database [3,4], while leads from the analysis were validated with
the Cancer Cell Line Encyclopedia (CCLE) [5] and the Cancer Therapeutics Response Portal
(CTRP) [6–8]. Furthermore, suggested drug combinations were validated with cell line
responses from the AstraZeneca-DREAM challenge dataset [25] and patient derived xenograft
(PDX) models from Gao *et al.* [26].
402
403 For a given cell line in GDSC, the drug response was fitted with a sigmoid curve [59] and
404 consecutively quantified as area under the curve (AUC) or the concentration required to
405 reduce cell viability by half (IC_{50}). GDSC contains 265 compounds tested in 1074 cell lines,
406 whilst we focus on a subset of 38 drugs targeting either the PI3K-AKT or MAPK signalling,
407 which leads to 327 experiments considered for evaluation.
408
Deep molecular characterisation of the cancer cell lines
The GDSC project [4] provides the characterisation of >1,000 cell lines including whole exome
sequencing, targeted PCR sequencing/split probe FISH analysis and SNP6.0 arrays, which
enabled to quantify somatic mutations, gene fusions and copy number variations (CNVs),
respectively. In our analysis, we focus on somatic mutational state of 300 cancer genes and
10 gene fusions. Additionally, we considered 425 recurrent CNVs, split into 117
amplifications and 308 deletions. In total, we consider 735 cancer functional events, which is
summarized in the binary event matrix (BEM) from Iorio *et al.* [4].
417
Pre-processing cell line and pharmacological data
For every pair of drugs that was computationally analyzed, the subset of GDSC cell lines
having valid IC_{50} and AUC values for both drugs was retained. Typically there were roughly
700 cell lines across all cancer types that met these criteria in each experiment. Cancer

types and subtypes were stored along with the pharmacological data in a table for each drug
for subsequent recall and analysis.
424
425 **Processing drug response measures (AUC/IC$_{50}$ values)**
426 We build network models for a set of N cell lines, $C = \{C_1, .., C_N\}$, that are separately
427 exposed to two distinct drugs, D_1 and D_2 , which results in two sets of M measurement
428 variables, $X_i = [x_1, .., x_M], i = 1,2$, describing the response to each compound:

$$X_{i,j} = D_i(C_j), \quad i = 1,2; \; j = 1, \ldots, N \quad \text{(Equation 1)}$$

We use a network model that is an undirected graph, G, consisting of N vertices, $V_i, i =$
$1, \ldots, N$, (one for each cell line in C) with weighted edges, $W_{i,j}(V_i, V_j), i, j = 1, \ldots, N, i \neq j$,
between every distinct pair of vertices. Our approach
uses a single multivariate similarity measure (Equation 2), to construct one network model,
with the advantage that the subspace properties of the resulting adjacency and Laplacian
matrices are fully embedded with the complete characteristics of C. The weight is the
similarity, $w_{i,j}$, between i-th and j-th composite $2M \; x \; 1$ dose response profile (DRP), $X_i =$
$[X_{1,i}, X_{2,i}]$, for C_i and C_j.
440
We characterize drug response by two important continuous-valued measurements
extrapolated from the cell line pharmacology screens: the IC$_{50}$ and the AUC values of the
dose-response curve (**Table S4**) observed when one compound is applied *in vitro* to a single
cell line sample at successively greater concentrations. Since every cell line possesses a
length-4 DRP for a given pair of drugs, the similarity, w, between any two cell lines resides
on $(0,1)$ and is calculated by a multivariate quasi-Gaussian comparison that differences the
elements of the DRPs but also weighs the differences by a combination of local and global
network statistics. Similarity between the response vectors, X_i and X_j, is given by:

$$w(X_i, X_j) = e^{-d(X_i, X_j)} \quad \text{(Equation 2)}$$

$$d(X_i, X_j) = (\Delta X_{i,j}^T \, \Sigma_{i,j}^{-1} \Delta X_{i,j})^\beta \quad \text{(Equation 3)}$$

454 The similarity between two cell lines equals one when both have identical covariate values,
455 and approaches zero as their covariates increasingly differ. Additionally, $w(X_i, X_j) = w(X_j, X_i)$.
456 $\Delta X_{i,j}$ is a $4 \; x \; 1$ vector whose entries are the difference of the DRP values in X_i and X_j and β
457 modulates the similarity between two patients. We selected $\beta = 0.5$ for our experiments
458 based on experimentation and the observations of previous efforts.
459
460 $\Sigma_{i,j}$ is a 4 x 4 covariance-like matrix that is estimated for every distinct (i,j)-pair and captures
461 the variability of individual variables as well as their inter-relationships. While $\Sigma_{i,j}$ is an
462 explicit function of the two patients being compared, it also captures network-wide
463 characteristics. For diagonal elements, $\Sigma_{i,j}(a,a), a = 1, ..4$, the entries are:

$$\Sigma_{i,j}(a,a) = \frac{\left(\Delta X_{i,j}(a) + \overline{Ngbd(\Delta X_I(a))} + + \overline{Ngbd(\Delta X_J(a))}\right)^2}{9}, \; a = 1,2,3,4 \quad \text{(Equation 4)}$$

467 where $Ngbd\,(\Delta X_I(a))$ corresponds to all edges neighboring vertex-i, and the overbar is the
468 averaging operator. The off-diagonal elements, $\Sigma_{i,j}(a,b), a, b = 1, ..4, a \neq b$ are:

$$\Sigma_{i,j}(a,b) = \frac{\left(\Delta X_{i,j}(a) + \overline{Ngbd(\Delta X_I(a))} + + \overline{Ngbd(\Delta X_J(a))}\right)\left(\Delta X_{i,j}(b) + \overline{Ngbd(\Delta X_I(b))} + + \overline{Ngbd(\Delta X_J(b))}\right)}{9}, \; a =$$

471 1,2,3,4 (Equation 5)
472

473　The Moore-Penrose pseudo-inverse was used to avoid problems with low-rank during matrix
474　inversion. The framework is generalizable to include more variables and different measures
475　of similarity. The symmeric, positive semi-definite, $N \times N$ weighted adjacency matrix, W,
476　holds the pairwise similarities.
477
Segmentation
The set of cell lines, C, is segmented recursively into distinct subpopulations using the
Fiedler eigenvector derived from the eigendecomposition of W [60]. Each parent subpopulation
of cell lines is successively segmented into two offspring subpopulations until at least one of
3 constraints is satisfied:
● The size of the parent subpopulation falls below a user-defined threshold;
● The size of either offspring subpopulation is below a user-defined threshold;
● The offspring subpopulations are not sufficiently dissimilar, where we measure class
separability using the silhouette metric [61], a common non-parametric method in which
values range between -1 (highly similar) and 1 (highly dissimilar) and values near 0
indicate the two subpopulations are just barely overlapping. Higher values result in
fewer, less granular, subpopulations.
In our experiments, we required the size of the parent subpopulation to be at least 40 in
order for segmentation to be performed, and required both offspring subpopulations to have
20 or more members in order to be retained. The silhouette metric threshold was set to 0.25.
Generally, criteria and thresholds can be modified and adapted to emphasize relevant
factors in a particular problem. The whole segmentation process yields K mutually exclusive
subpopulations $P_k, k = 1, \ldots, K$, where $C = \cup_{k=1} P_k$. Successive segmentation results in sub-
populations with increasingly homogeneous DRPs.
497
498
499　**QUANTIFICATION AND STATISTICAL ANALYSIS**
500
Enrichment of features to nominate biomarkers
Because genetic alterations in each cell line are known, each subpopulation can be
evaluated by non-parametric statistical tests to identify enriched alterations that may be
attributed to patterns of sensitivity or resistance in the DRP across both drugs. For each
subpopulation, we measured the number of cell lines in the subpopulation with a particular
gene mutation, and the number of cell lines outside of the subpopulation with the mutation. A
2X2 contingency table was generated from the cell line counts of with/without mutation and
inside/outside of subpopulation. Significance of observed enrichment of mutations within
subpopulations were calculated using the Fisher's exact test. The resulting p-values were
corrected for multiple testing using the Benjamini and Hochberg (BH) procedure (**Table S5**).
511
Classification of drug pairs based on the distribution of subpopulations
We made 324 pairwise comparisons of drugs targeting the MAPK and PI3K-AKT pathways.
Based on the distribution of log(IC_{50}) values across all cell lines tested with both drugs, we
determined the 20th-percentile of log(IC_{50}) values for each drug. The 20th percentile cutoffs
P_{20} for drugs A and B was used to categorise each subpopulation *i* into four categories
based on their average log(IC_{50}) *y*:
$y_i < P_{20, A}$ and $y_i < P_{20, B}$ = *sensitive to drugs A and B*
$y_i < P_{20, A}$ and $y_i \geq P_{20, B}$ = *more sensitive to drug A*
$y_i \geq P_{20, A}$ and $y_i < P_{20, B}$ = *more sensitive to drug B*
$y_i \geq P_{20, A}$ and $y_i \geq P_{20, B}$ = *resistant to drugs A and B*
The number of subpopulations in each category were recorded in a 2X2 contingency matrix
and using a binomial test compared this to the proportion of cells expected in each category
if SEABED segmentation was not performed.
525

After classification of pairwise drug responses, we assessed whether a drug was significantly enriched for one category in comparison with all other drugs. Testing was carried out using the hypergeometric test (*phyper* R package).

2-D visualization of drug response profiles
To visualize DRPs across cell lines and drug comparisons, we calculated the average $\log(IC_{50})$ values for each drug in subpopulations generated based on their response to the tested drug pairs (**Table S5**). We then plotted the mean $\log(IC_{50})$ values as circles on a 2-D scatter plot using the Matplotlib Python library. Dashed lines indicative of local 20th percentile of $\log(IC_{50})$ values for each drug were also plotted on the scatter plot unless stated otherwise. The radii of the circles is proportional to the subpopulation size. Due to potential misidentification, cell lines named 'NA' found in the output files after segmentation with SEABED, were not used for the calculation of the mean $\log(IC_{50})$ values of each subpopulation and hence, not used for the 2-D visualization.

Tree visualization of subpopulations
We utilized tree diagrams to visualize the data generated. The tree diagrams illustrate how the cancer cell lines are segmented into different subpopulations, based on whether they are sensitive or resistant to the drugs that are being tested. We did not visualize further down the tree in the figures when previously observed significant enrichment of genetic biomarkers is no longer observed in all current subpopulations. The tree diagrams were generated through an open-source Python library called Graphviz. The style of each component of the tree diagram was first initialized through a class. This included the colours, shapes, and fonts of the edges and nodes of the tree diagram. A method to create tree diagrams was developed to accept the number of vertices and leaves, the labels for the leaves, and the tree diagram filename. The tree diagram is finally generated and saved by calling the method.

DATA AND SOFTWARE AVAILABILITY

All code for the pipeline is open source and available at: https://github.com/szen95/SEABED. All data used in the paper are published previously and publicly available at the GDSC, CCLE, and CTRP databases. Datasets used are listed in Table S4, Table S5, and the Key Resources Table.

ADDITIONAL RESOURCES

Response patterns of 324 pair-wise comparisons of 18 PI3K-AKT and 18 MAPK pathway inhibitors: https://szen95.github.io/SEABED/

References

1. Weinstein, J. N. Cell lines battle cancer. *Nature* **483**, 544–545 (2012).

2. Shoemaker, R. H. The NCI60 human tumour cell line anticancer drug screen. *Nat. Rev. Cancer* **6**, 813–823 (2006).

3. Garnett, M. J. *et al.* Systematic identification of genomic markers of drug sensitivity in cancer cells. *Nature* **483**, 570–575 (2012).

4.   Iorio, F. *et al.* A Landscape of Pharmacogenomic Interactions in Cancer. *Cell* **166**, 740–

754 (2016).

5.   Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling of

anticancer drug sensitivity. *Nature* **483**, 603–607 (2012).

6.   Rees, M. G. *et al.* Correlating chemical sensitivity and basal gene expression reveals

mechanism of action. *Nat. Chem. Biol.* **12**, 109–116 (2016).

7.   Seashore-Ludlow, B. *et al.* Harnessing Connectivity in a Large-Scale Small-Molecule

Sensitivity Dataset. *Cancer Discov.* **5**, 1210–1223 (2015).

8.   Basu, A. *et al.* An interactive resource to identify cancer genetic and lineage

dependencies targeted by small molecules. *Cell* **154**, 1151–1161 (2013).

9.   Yang, W. *et al.* Genomics of Drug Sensitivity in Cancer (GDSC): a resource for

therapeutic biomarker discovery in cancer cells. *Nucleic Acids Res.* **41**, D955–61

(2013).

10. Silverbush, D. *et al.* Cell-Specific Computational Modeling of the PIM Pathway in Acute

Myeloid Leukemia. *Cancer Res.* **77**, 827–838 (2017).

11. Costello, J. C. *et al.* A community effort to assess and improve drug sensitivity

prediction algorithms. *Nat. Biotechnol.* **32**, 1202–1212 (2014).

12. Ravikumar, B., Alam, Z., Peddinti, G. & Aittokallio, T. C-SPADE: a web-tool for

interactive analysis and visualization of drug screening experiments through compound-

specific bioactivity dendrograms. *Nucleic Acids Res.* **45**, W495–W500 (2017).

13. Patel, S. P. *et al.* Clinical responses to selumetinib (AZD6244; ARRY-142886)-based

combination therapy stratified by gene mutations in patients with metastatic melanoma.

Cancer **119**, 799–805 (2013).

14. Catalanotti, F. *et al.* Phase II trial of MEK inhibitor selumetinib (AZD6244, ARRY-

142886) in patients with BRAFV600E/K-mutated melanoma. *Clin. Cancer Res.* **19**,

2257–2264 (2013).

15. Metro, G. *et al.* Selumetinib: a promising pharmacologic approach for KRAS-mutant

advanced non-small-cell lung cancer. *Future Oncol.* **9**, 167–177 (2013).

16. Kim, D. W. & Patel, S. P. Profile of selumetinib and its potential in the treatment of

melanoma. *Onco. Targets. Ther.* **7**, 1631–1639 (2014).

17. Tricker, E. M. *et al.* Combined EGFR/MEK Inhibition Prevents the Emergence of

Resistance in EGFR-Mutant Lung Cancer. *Cancer Discov.* **5**, 960–971 (2015).

18. Sequist, L. V. *et al.* Phase III Study of Afatinib or Cisplatin Plus Pemetrexed in Patients

With Metastatic Lung Adenocarcinoma WithEGFRMutations. *J. Clin. Oncol.* **31**, 3327–

3334 (2013).

19. Misale, S. *et al.* Emergence of KRAS mutations and acquired resistance to anti-EGFR

therapy in colorectal cancer. *Nature* **486**, 532–536 (2012).

20. Geyer, C. E. *et al.* Lapatinib plus capecitabine for HER2-positive advanced breast

cancer. *N. Engl. J. Med.* **355**, 2733–2743 (2006).

21. Palmer, A. C. & Sorger, P. K. Combination Cancer Therapy Can Confer Benefit via

Patient-to-Patient Variability without Drug Additivity or Synergy. *Cell* **171**, 1678–

1691.e13 (2017).

22. Fujishita, T., Kajino-Sakamoto, R., Kojima, Y., Taketo, M. M. & Aoki, M. Antitumor

activity of the MEK inhibitor trametinib on intestinal polyp formation in Apc(Δ716) mice

involves stromal COX-2. *Cancer Sci.* **106**, 692–699 (2015).

23. De Grève, J. *et al.* Clinical activity of afatinib (BIBW 2992) in patients with lung

adenocarcinoma with mutations in the kinase domain of HER2/neu. *Lung Cancer* **76**,

123–127 (2012).

24. Jacobsen, K. *et al.* Convergent Akt activation drives acquired EGFR inhibitor resistance

in lung cancer. *Nat. Commun.* **8**, 410 (2017).

25. Menden, M. P. *et al.* A cancer pharmacogenomic screen powering crowd-sourced

advancement of drug combination prediction. (2017). doi:10.1101/200451

26. Gao, H. *et al.* High-throughput screening using patient-derived tumor xenografts to

predict clinical trial drug response. *Nat. Med.* **21**, 1318–1325 (2015).

27. Martin-Liberal, J. Encorafenib plus binimetinib: an embarrassment of riches. *Lancet*

Oncol. **19**, 1263–1264 (2018).

28. Dummer, R. *et al.* Overall survival in patients with BRAF-mutant melanoma receiving

encorafenib plus binimetinib versus vemurafenib or encorafenib (COLUMBUS): a

multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* **19**, 1315–1327 (2018).

29. Paller, C. J. *et al.* Design of Phase I Combination Trials: Recommendations of the

Clinical Trial Design Task Force of the NCI Investigational Drug Steering Committee.

Clin. Cancer Res. **20**, 4210–4217 (2014).

30. Jänne, P. A. *et al.* Selumetinib Plus Docetaxel Compared With Docetaxel Alone and

Progression-Free Survival in Patients With KRAS-Mutant Advanced Non–Small Cell

Lung Cancer. *JAMA* **317**, 1844 (2017).

31. Garon, E. B. *et al.* Identification of common predictive markers of in vitro response to the

Mek inhibitor selumetinib (AZD6244; ARRY-142886) in human breast cancer and non-

small cell lung cancer cell lines. *Mol. Cancer Ther.* **9**, 1985–1994 (2010).

32. Long, G. V. *et al.* Combined BRAF and MEK inhibition versus BRAF inhibition alone in

melanoma. *N. Engl. J. Med.* **371**, 1877–1888 (2014).

33. Van Allen, E. M. *et al.* The genetic landscape of clinical resistance to RAF inhibition in

metastatic melanoma. *Cancer Discov.* **4**, 94–109 (2014).

34. Holderfield, M., Deuker, M. M., McCormick, F. & McMahon, M. Targeting RAF kinases

for cancer therapy: BRAF-mutated melanoma and beyond. *Nat. Rev. Cancer* **14**, 455–

467 (2014).

35. Hall, R. D. & Kudchadkar, R. R. BRAF mutations: signaling, epidemiology, and clinical

experience in multiple malignancies. *Cancer Control* **21**, 221–230 (2014).

36. Abdel-Wahab, O. *et al.* Efficacy of intermittent combined RAF and MEK inhibition in a

patient with concurrent BRAF- and NRAS-mutant malignancies. *Cancer Discov.* **4**, 538–

545 (2014).

37. Paz-Ares, L. *et al.* Afatinib versus gefitinib in patients with EGFR mutation-positive

advanced non-small-cell lung cancer: overall survival data from the phase IIb LUX-Lung

7 trial. *Ann. Oncol.* **28**, 270–277 (2017).

38. Takezawa, K. *et al.* HER2 amplification: a potential mechanism of acquired resistance to

EGFR inhibition in EGFR-mutant lung cancers that lack the second-site EGFRT790M

mutation. *Cancer Discov.* **2**, 922–933 (2012).

39. Mendoza, M. C., Er, E. E. & Blenis, J. The Ras-ERK and PI3K-mTOR pathways: cross-

talk and compensation. *Trends Biochem. Sci.* **36**, 320–328 (2011).

40. Temraz, S., Mukherji, D. & Shamseddine, A. Dual Inhibition of MEK and PI3K Pathway

in KRAS and BRAF Mutated Colorectal Cancers. *Int. J. Mol. Sci.* **16**, 22976–22988

(2015).

41. Villanueva, J., Vultur, A. & Herlyn, M. Resistance to BRAF inhibitors: unraveling

mechanisms and future treatment options. *Cancer Res.* **71**, 7137–7140 (2011).

42. Jänne, P. A. *et al.* Selumetinib Plus Docetaxel Compared With Docetaxel Alone and

Progression-Free Survival in Patients With KRAS-Mutant Advanced Non–Small Cell

Lung Cancer. *JAMA* **317**, 1844 (2017).

43. Ebert, P. J. R. *et al.* MAP Kinase Inhibition Promotes T Cell and Anti-tumor Activity in

Combination with PD-L1 Checkpoint Blockade. *Immunity* **44**, 609–621 (2016).

44. Bendell, J. *et al.* LBA-01Safety and efficacy of cobimetinib (cobi) and atezolizumab

(atezo) in a Phase 1b study of metastatic colorectal cancer (mCRC). *Ann. Oncol.* **27**,

ii140–ii140 (2016).

45. Gholami, A. M. *et al.* Global proteome analysis of the NCI-60 cell line panel. *Cell Rep.* **4**,

609–620 (2013).

46. Eisen, M. B., Spellman, P. T., Brown, P. O. & Botstein, D. Cluster analysis and display

of genome-wide expression patterns. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 14863–14868

(1998).

47. Wiwie, C., Baumbach, J. & Röttger, R. Comparing the performance of biomedical

clustering methods. *Nat. Methods* **12**, 1033–1038 (2015).

48. Monti, S., Tamayo, P., Mesirov, J. & Golub, T. Consensus Clustering: A Resampling-

Based Method for Class Discovery and Visualization of Gene Expression Microarray

Data. *Mach. Learn.* **52**, 91–118 (2003).

49. Sarwar, B., Karypis, G., Konstan, J. & Reidl, J. Item-based collaborative filtering

recommendation algorithms. in *Proceedings of the tenth international conference on*

World Wide Web - WWW '01 (2001). doi:10.1145/371920.372071

50. Scott, J. *Social Network Analysis*. (SAGE, 2017).

51. Wang, B. *et al.* Similarity network fusion for aggregating data types on a genomic scale.

Nat. Methods **11**, 333–337 (2014).

52. Li, L. *et al.* Identification of type 2 diabetes subgroups through topological analysis of

patient similarity. *Sci. Transl. Med.* **7**, 311ra174 (2015).

53. Nicolau, M., Levine, A. J. & Carlsson, G. Topology based data analysis identifies a

subgroup of breast cancers with a unique mutational profile and excellent survival. *Proc.*

Natl. Acad. Sci. U. S. A. **108**, 7265–7270 (2011).

54. Kim, D. *et al.* Knowledge boosting: a graph-based integration approach with multi-omics

data and genomic knowledge for cancer clinical outcome prediction. *Journal of the*

American Medical Informatics Association (2014). doi:10.1136/amiajnl-2013-002481

55. Li, L. *et al.* Identification of type 2 diabetes subgroups through topological analysis of

patient similarity. *Sci. Transl. Med.* **7**, 311ra174 (2015).

56. Ng, A. Y., Jordan, M. I. & Weiss, Y. On spectral clustering: Analysis and an algorithm. in

Advances in neural information processing systems (2002).

57. Shi, J. & Malik, J. Normalized cuts and image segmentation. *IEEE Transactions on*

Pattern Analysis and Machine Intelligence 888–905 (2000).

58. Cook, D. *et al.* Lessons learned from the fate of AstraZeneca's drug pipeline: a five-

dimensional framework. *Nat. Rev. Drug Discov.* **13**, 419–431 (2014).

59. Vis, D. J. *et al.* Multilevel models improve precision and speed of IC50 estimates.

Pharmacogenomics **17**, 691–700 (2016).

60. Fiedler, M. A property of eigenvectors of nonnegative symmetric matrices and its

application to graph theory. *Czechoslovak Math. J.* **25**, 619–633 (1975).

61. Rousseeuw, P. J. Silhouettes: A graphical aid to the interpretation and validation of

cluster analysis. *Journal of Computational and Applied Mathematics* **20**, 53–65 (1987).

Supplemental Information

1. Supplementary Materials.pdf
   Document S1. Figures S1–S6 and Tables S1-3.

2. Table S4.xlsx
   Table S4. Input Data for Segmentation, Related to Figures 1-4, S1-S6, and Website S1
   Input data: $\log(IC_{50})$ and AUC values for 1074 cancer cell lines treated with 265 anti-cancer drugs.

3. Table S5.xlsx
   Table S5. Output Data and Enriched Biomarkers After Segmentation, Related to Figures 1-4, S2-S6, and Website S1
   Output data: IC_{50} 20% cutoff, minimum and maximum IC_{50} concentration, and average IC_{50} response of each subpopulation towards each drug (327 cancer drug pairs) tested on the cancer cell lines. The subpopulation number and the number of cell lines in each subpopulation are recorded. Each individual cell line in every subpopulation together with individual cell line tissue types are also shown. **Enriched biomarkers:** Biomarkers found within subpopulations (adjusted p-value and/or p-value < 0.05), together with the subpopulation number, the number of cell lines in each subpopulation, percentage of the biomarkers found within each subpopulation, the number of cell lines in the subpopulation with the biomarker, the p-value, and adjusted p-value.

Acknowledgements

We would like to thank Ben Sidders (AstraZeneca plc.), Jonathan Dry (AstraZeneca plc.), Francesco Iorio, (Sanger Institute), Michael Schubert (EMBL-EBI), Mi Yang (RWTH Aachen) and Winston Hide (Harvard University) for useful discussions. This work is supported by funding from the NIHR Sheffield Biomedical Research Centre, Rosetree Trust (ref: A2501), and the Academy of Medical Sciences Springboard (REF: SBF004\1052).

Author contributions

NK, TST, MM, and DW contributed to the conceptualization of the project. NK, TST, and DW were responsible for data curation. Formal analysis was performed by NK, TST, HY, BY, MM, and DW. MM and DW were responsible for funding acquisition. The methodology was developed by NK, TST, MM, and DW. TST and DW were responsible for administration of the project. Resources for the experiments were prepared by DW and NK. Various software for the project was developed and implemented by NK, TST, HY and DW. MM, and DW oversaw supervision of the project. NK, TST, MM, and DW wrote the manuscript.

Declaration of Interests

NK is an employee of Constellation Analytics, LLC.

760 # Figure Legends

763
Figure 1: Segmentation of a population based on pharmacological patterns of
response discovers subpopulations with differential response. (A) Dose
response curves of two or more drugs are measured across a population of up to
1,000 cancer cell lines. **(B)** The population is segmented into distinct and
homogeneous subpopulations based on their response to multiple drugs. When
comparing two drugs, subpopulations can be categorised based on their mean
log(IC$_{50}$s): sensitive to both drugs (orange), sensitive to drug A but not drug B
(green), sensitive to drug B but not drug A (blue), resistant to both drugs (grey). **(C)**
Segmentation results for a BRAF inhibitor (SB590885) and a MEK inhibitor (CI-
1040). Tree nodes contain the number of cell lines and are colored based upon their
category of response. Significance testing of 735 cancer functional events reveals
subpopulations enriched for *BRAF* and *KRAS* mutations. **(D)** Scatter plot showing
derived subpopulations based on their pharmacological responses for afatinib and
selumetinib. Dashed lines indicate 20th percentile of log(IC$_{50}$) values for each drug.
PIK3CA, EGFR, ERBB2, KRAS, NRAS, BRAF, APC, TCF4 and *RB1* mutations were
found enriched in the associated subpopulations. **(E)** OncoPrint visualizing the
percentage of mutations of selected genes in cell line panel treated with either
afatinib (EGFR inhibitor) or selumetinib (MEK inhibitor). The waterfall plots comparing
the response of the cell lines to afatinib and to selumetinib. **(F)** Boxplot of difference
in log(IC$_{50}$) values between afatinib and selumetinib response for wild-type cell lines,
all cell lines with *PI3KCA* mutation and cell lines in derived subpopulations with
enriched *PI3KCA* mutation. **(G)** Same as panel **(F)**, but for *BRAF* mutations.
786

787
788

789
790   **Figure 2: Distinct drug response types after unsupervised segmentation of**
791   **pharmacological response pattern for targeting MAPK or PI3K-AKT signaling.**
792   **(A)** MAPK and PI3K-AKT pathways illustrating drugs in purple boxes which were

assessed, and their different gene targets in the pathway. Genes in the green boxes
are involved in the PI3K-AKT pathway while genes in the blue boxes are involved in
the MAPK pathway. **(B)** Heatmap illustrating 324 pairwise comparison of responses
for 18 different inhibitors targeting the MAPK pathway and 18 different inhibitors
targeting the PI3K-AKT pathway. The 20th-percentile of log(IC_{50}) values for each
drug was determined based on the distribution of log(IC_{50}) values across all cell lines
after SEABED segmentation. Based on these 20th percentile cutoffs for sensitivity,
we assessed whether there was enrichment of subpopulations in each quadrant and
categorised each drug pair into five categories: (i) no preferential response (grey), (ii)
subpopulations sensitive to both MAPK and PI3K-AKT pathway inhibitors (pink), (iii)
greater MAPK pathway sensitivity (yellow), (iv) greater PI3K-AKT pathway sensitivity
(green) or (v) sensitive to either inhibitor but not both (blue).

Figure 3: Divergent response exemplified with PI3K inhibition in comparison to
MEK and BRAF inhibitors. (A) Scatter plot showing subpopulations that exhibit
divergent pharmacological response to RDEA119-2 (MEK inhibitor) and
GSK2126458 (PI3K inhibitor). Dashed lines indicate 20th percentile of log(IC$_{50}$)
values for each drug. *BRAF* and *KRAS* mutations were found enriched in the
associated subpopulations. **(B)** Subpopulations from comparison of PLX4720-2
(BRAF inhibitor) and PI-103 (PI3K inhibitor) responses. **(C)** Individual cell line
responses from PLX4720-2 and PI-103 coloured by the subpopulations they were
grouped in. **(D)** Workflow illustrating cell lines being tested with individual inhibitors
and their joint pharmacological patterns of response. Drug pairs with divergent
response and *BRAF* mutant subpopulations suggest a target for drug combination
therapies, which are validated in cell lines [25] and patient-derived tumor xenograft
(PDX) models [26]. **(E)** *In vitro* synergistic effect of combining MEK inhibitors with PI3K
or BRAF inhibitors in all cell lines and just those with *BRAF* mutations. This also
compared to measured synergy for all combinations tested in cell lines. **(F)** *In vivo*
effect of combining BRAF inhibitors with PI3K or MEK inhibitors. Measured response
is the change in tumor volume following treatment. BRAF and MEK inhibitor
combinations have also been shown to be effective in the patients [28].

827
828

829
830 **Figure 4: Subpopulations of cell lines indicating differential response between**
831 **docetaxel and selumetinib. (A)** Tree diagram illustrating the segmentation process
832 of 840 cell lines across cancer types into subpopulations based on their response to
833 docetaxel and selumetinib. Branch colours distinguish subsets of subpopulations with
834 darker colours indicating increasing number of cell lines in the subpopulation. **(B)**

Scatter plot of subpopulations discovered through progressive segmentation based on average $\log(IC_{50})$ values. Dashed lines indicate 20th percentile of $\log(IC_{50})$ values for each drug. The *KRAS* mutation is enriched in 5 subpopulations exhibiting sensitivity to selumetinib and resistance to docetaxel, including subpopulation 60. The colours of the subpopulations correspond to the location of the subpopulation in the tree diagram. **(C)** Bar plot illustrating the difference in AUC values for each cell line. The orange bars highlight the cell lines within subpopulation 60. **(D)** Heatmap showing the percentage of each cancer type enriched within the cell lines in subpopulation 60. The bar plot illustrates the *KRAS* mutations within the cell lines that are highlighted in green.

A

B

Pan-cancer cell lines

resistant

Selumetinib (MEKi)
log10(IC50) [uM]

sensitive

KRAS

KRAS

KRAS

KRAS

subpop. 60

sensitive   Docetaxel (Microtubules)   resistant
log10(IC50) [uM]

Number of cell lines
in subpopulation
● 100 cells
● 50 cells
· 10 cells

C

Difference in AUC
(Selumetinib − Docetaxel)

■ subpop. 60
■ other subpops.

D

breast   <1%
lymphoma   <1%
nervous system   <1%
soft tissue   <1%
smal cell lung   2%
myeloma   2%
thyroid   2%
neuroblastoma   3%
gastric   4%
kidney   4%
aerodigestive tract   5%
leukemia   8%
non-small cell lung   8%
pancreatic   8%
urogenital   8%
colorectal   15%
melanoma   27%

Difference in AUC
(Selumetinib − Docetaxel)

■ *KRAS* mutations