1   **Title:**

2   A hierarchical watershed model of fluid intelligence in childhood and adolescence

3
4   **Running Title:**

5   Fluid intelligence in childhood and adolescence

6
7   **Authors:**

8   Fuhrmann, D.[1*], Simpson-Kent, I. L.[1], Bathelt, J.[1], the CALM team [1] & Kievit, R. A.[1]

9   [1] MRC Cognition and Brain Sciences Unit, University of Cambridge, Cambridge, CB2 7EF, UK

10   [*] corresponding author: Delia Fuhrmann, MRC CBU, 15 Chaucer Road, Cambridge, CB2 7EF,

11   UK, email: delia.fuhrmann@mrc-cbu.cam.ac.uk

12   Abstract

Fluid intelligence is the capacity to solve novel problems in the absence of task-specific

knowledge, and is highly predictive of outcomes like educational attainment and

psychopathology. Here, we modelled the neurocognitive architecture of fluid intelligence in

two cohorts: CALM (N = 551, aged 5 - 17 years) and NKI-RS (N = 335, aged 6 - 17 years). We

used multivariate Structural Equation Modelling to test a preregistered watershed model of

fluid intelligence. This model predicts that white matter contributes to intermediate cognitive

phenotypes, like working memory and processing speed, which, in turn, contribute to fluid

intelligence. We found that this model performed well for both samples and explained large

amounts of variance in fluid intelligence (R^2_{CALM} = 51.2%, R^2_{NKI-RS} = 78.3%). The relationship

between cognitive abilities and white matter differed with age, showing a dip in strength

around ages 7 - 12 years. This age-effect may reflect a reorganization of the neurocognitive

architecture around pre- and early puberty. Overall, these findings highlight that intelligence

is part of a complex hierarchical system of partially independent effects.

26   Keywords

Fluid intelligence (g_f) is a core part of human cognition and refers to the capacity to solve

novel problems in the absence of task-specific knowledge. It is highly predictive of a number

of important life span outcomes, including educational attainment (Primi et al. 2010; Roth et

al. 2015) and psychopathology (Gale et al. 2010). Despite years of investigation, however, our

understanding of the neurocognitive architecture of g_f remains limited. Longstanding debates

have considered, for instance, how g_f relates to more fundamental cognitive functions such

as working memory and processing speed, and how all of these cognitive functions relate to

brain structure and function (Kyllonen and Christal 1990; Fry and Hale 2000; Chuderski 2013;

Ferrer et al. 2013).

Working memory is the ability to hold and manipulate information in the mind short-term. It

has been suggested that working memory is a key determinant of g_f by limiting mental

information processing capacity (Fukuda et al. 2010; Chuderski 2013). Proponents of this

working memory account of g_f cite high correlations between the two domains ranging from

0.5 to 0.9 in meta-analyses (Ackerman et al. 2005; Oberauer et al. 2005). Such high

correlations have led some to suggest that g_f and working memory are, in fact, isomorphic

(Kyllonen and Christal 1990). However, more recent work has highlighted that this

isomorphism only arises under conditions of high time constraints for g_f tasks (Chuderski

2013). This suggests that g_f and working memory are, in fact, separable constructs and

underlines the importance of processing speed for g_f.

Processing speed, the speed of mental computations, is thought to be rate-limiting to g_f and

is therefore sometimes proposed to be a particularly good predictor of g_f (Kail and Salthouse

1994; Salthouse 1996; Ferrer et al. 2013; Kail et al. 2015; Schubert et al. 2017). Proponents of

the processing speed account of g_f cite moderate but robust correlations between g_f and

processing speed of 0.2 in meta-analyses (Sheppard and Vernon 2008) as well as longitudinal

evidence (Finkel et al. 2005; Coyle et al. 2011; Kail et al. 2015). Salthouse (1996) argued in the

context of cognitive aging, that processing speed determines high-level cognitive

performance because slow processing means that relevant sub-operations cannot be

completed in a set amount of time or are not available for successful integration. A

complementary explanation of individual differences in g_f proposes that processing speed

may be a direct reflection of fundamental neuroarchitectonic properties of the brain, such as

myelination or white matter microstructure (Lu et al. 2011; Chevalier et al. 2015).

White matter shows protracted development throughout childhood and adolescence, and

into the third decade of life (Mills et al. 2016). White matter tracts can be characterised *in*

vivo using diffusion-tensor imaging (DTI), which is sensitive, but not necessarily specific, to

white matter microstructural properties such as myelination or axonal density (Jones et al.

2013; Wandell 2016). Fractional anisotropy (FA) is the most commonly investigated DTI

measure and quantifies the directionality of water diffusion in different white matter tracts

(Pfefferbaum et al. 2000; Wandell 2016). Working memory, processing speed and g_f have

each been linked to individual differences in FA (Vestergaard et al. 2011; Kievit, Davis,

Griffiths, Correia, CamCAN, et al. 2016; Bathelt et al. 2018). While some studies, using

Principal Component Analysis, have posited that FA in different tracts can be summarized by

sizable single components (Penke et al. 2010; Cox et al. 2016), formal investigations using

confirmatory factor analysis have demonstrated that single-factor models of FA generally

show poor fit and do not adequately capture individual differences in white matter

microstructure (Lövdén et al. 2013; Kievit, Davis, Griffiths, Correia, Cam-CAN, et al. 2016). In

a similar vein, there is a growing body of literature showing specific associations between

white matter tracts and cognitive abilities, with those connecting frontoparietal regions

usually showing largest contributions to complex cognitive functions like g_f (Vestergaard et al.

2011; Kievit et al. 2016; Bathelt et al. 2018).

We here seek to address several critical outstanding issues in the field: First, there is limited

systematic evidence on the concurrent relationships between g_f, working memory,

processing speed and white matter. This leaves the relative contributions of processing speed

and working memory to g_f unclear, which, in turn, poses challenges for the design of effective

cognitive training interventions. Second, studies usually use a single task as a proxy for

complex and abstract constructs such as processing speed, working memory, and g_f. This

raises questions about the generalizability of findings (Noack et al. 2014). Third, our

understanding of how the relationships between relevant cognitive domains and between

brain and cognition change with age remains limited, raising the possibility that brain-

behaviour relationships may change with age (Garrett 1946; Johnson 2000; Tamnes et al.

2017).

To address these issues, we here used structural equation modelling (SEM) to model the

associations between g_f, working memory, processing speed, and white matter

microstructure and age in two large, independent samples: the Centre for Attention, Leaning

and Memory sample (CALM, N = 551, aged 5 - 17 years), which consists of children and

adolescents referred to a clinic for having problems with attention, learning and memory

(Holmes et al. 2018), and the Enhanced Nathan Kline Institute – Rockland Sample (NKI-RS, N =

335, aged 6 - 17 years), a community-ascertained sample (Nooner et al. 2012).

To investigate the neurocognitive architecture of g_f in a principled way, we used a watershed

model of individual differences. Based on the metaphor of a watershed, the model predicts a

hierarchical many-to-one mapping of partially independent effects such that upstream

tributaries (e.g. brain structure) contribute to intermediate cognitive phenomena (cognitive

endophenotypes, e.g. working memory and processing speed), which then contribute to

downstream, complex cognitive phenomena such as g_f (Cannon and Keller 2006; Kievit, Davis,

Griffiths, Correia, CamCAN, et al. 2016). See Figure 1 for a representation of the model.

103
Figure 1. The Watershed Model. Schematic representation of the watershed model
developed by (Cannon and Keller 2006) and adapted for the present study. Fluid ability is
hypothesized to be the downstream product of working memory and processing speed,
which are, in turn, the product of white matter contributions. Figure adapted from Kievit et
al. (2016).

SEM, as a statistical technique, is uniquely suited to modeling the kinds of complex

multivariate brain-behavior associations posited by the watershed model (Kievit et al. 2011;

Kline 2015). SEM combines factor analysis and path analysis (a variant of regression analysis).

It can model abstract cognitive constructs like g_f, by estimating latent variables from

observed task scores (i.e. manifest variables). This feature of SEM allowed us to model g_f,

working memory, and processing speed in two independent samples, and thereby provided a

direct test of the generalizability of our findings. Second, SEM can test the simultaneous

relations between multiple cognitive and neural variables, allowing us to address the relative

contributions of different white matter tracts and different cognitive endophenotypes to g_f.

Finally, using SEM Trees (Brandmaier et al. 2013), a novel, decision-tree-based extension of

SEM, we investigated whether the associations in the watershed model change with age.

Based on the watershed model we made the following preregistered predictions

(http://aspredicted.org/blind.php?x=u5pf6z):

1.   Working memory, g_f and processing speed are separable constructs.

2.   Individual differences in g_f are predicted by working memory and processing speed.

3.   White matter microstructure is a multi-dimensional construct.

4.   There is a hierarchical relationship between white matter microstructure, cognitive

endophenotypes (working memory and processing speed) and g_f, such that white

matter contributes to working memory and processing speed, which, in turn

contribute to g_f.

5.   The contribution of working memory and processing speed to g_f changes with age.

Materials and Methods

Samples

We analysed data from the CALM and NKI-RS sample, as described in detail by (Holmes et al.

2018) and (Nooner et al. 2012) respectively. See also Simpson-Kent et al. (2019). We had also

preregistered to analyse data from the ABCD cohort (Volkow et al. 2018). The latter cohort

contains only data for 9 - and 10 - year olds at present, however, which limits comparability

to CALM and NKI-RS, and makes it unsuitable for investigations of developmental differences.

We therefore opted to not analyse ABCD data here and instead recommend a replication of

the analyses presented here in ABCD once longitudinal data is available. The CALM sample

consists of children and adolescents referred by health and educational professionals as

having difficulties in attention, learning and/or memory. The NKI-RS is a community-

ascertained, lifespan sample, and representative of the general population of Rockland, New

York, and the United States as a whole, in terms of ethnicity, socioeconomic status etc. For

NKI-RS, we included data for participants under the age of 18 only to match the age range of

CALM and excluded data that were completed more than half a year after enrolment. The

latter criterion was implemented to ensure that age at assessment did not differ

substantively between cognitive measures. The final samples included 551 participants from

CALM (30.85% female, aged 5.17 - 17.92 years, $N_{\text{Neuroimaging}}$ = 165) and 335 participants from

NKI-RS (43.48% female, aged 6.06 - 17.92 years, $N_{\text{Neuroimaging}}$ = 67). See Table 1 for prevalence

of relevant disorders and learning difficulties in the samples.

Table 1. Prevalence of Relevant Disorders and Learning Difficulties in the CALM and NKI-RS
cohorts

Variable	Percentage CALM	Percentage NKI-RS
ADHD	31.94	17.01
Dyslexia	5.81	5.67
Autism	6.72	0.60
Mood disorder	0.54	0.90
Anxiety disorder	2.36	18.21
Medicated[1]	10.53	17.01
Speech/language problems	38.11	19.40

152   *Note.* [1] unspecified medication for NKI-RS, ADHD-medication for CALM

Cognitive Tasks

154   We included cognitive tasks measuring the domains of g_f, working memory or processing

155   speed for CALM and NKI-RS. See Table 2 for the complete list of tasks used, and the

156   Supplementary Methods for task descriptions. Supplementary Figure 1 and 2 show raw

157   scores on all tasks. The tasks modelled here were preregistered for CALM but not NKI-RS.

158   *Table 2. Cognitive Tasks Modelled*

	CALM	NKI-RS
Working memory	AWMA Digit Recall (forward digit span)	WISC-R Forward Digit Span
	AWMA Backward Digit Span	WISC-R Backward Digit Span
	AWMA Dot Matrix	-
	AWMA Mr X	-
	-	CNB N-back task
g_f	WASI-II Matrix Reasoning	WASI-II Matrix Reasoning
	-	WASI-II Block Design
	-	WASI-II Similarities
	-	CNB Verbal Reasoning
Processing speed	DKEFS Trail-Making	DKEFS Trail-Making
	PhAB Rapid Naming	-
	TEA-Ch RBBS	-
	-	CNB Motor Speed
	-	CNB Sensory Motor Speed

Note. See the Supplementary Methods for task descriptions. Abbreviations: AWMA -
Automated Working Memory Assessment (Alloway 2007), CNB - Computerized
Neurocognitive Battery (Gur et al. 2001), DKEF - Delis-Kaplan Executive Functioning System
(Delis et al. 2004), PhAB – Phonological Assessment Battery (Gallagher and Frederickson
1995), TEA-Ch RBBS - Test of Everyday Attention for Children, Red & Blues, Bags & Shoes
subscale (Manly et al. 2001), WASI - Wechsler Abbreviated Scale of Intelligence - Second
Edition (Wechsler 2011), WISC-R - Wechsler Intelligence Scale for Children – Revised
(Kaufman 1975).

White Matter Microstructure

We modelled mean FA for all ten tracts of the Johns Hopkins University (JHU) white matter

tractography atlas (Hua et al. 2008) averaged over the hemispheres (Figure 2). See

Supplementary Methods for details of the MRI acquisition and processing and Supplementary

Figure 3 and 4 for raw FA values in all tracts.

Forceps Minor
Anterior Thalamic Radiation
Cingulate Gyrus
Superior Longitudinal Fasciculus
Inferior Longitudinal Fasciculus

Corticospinal Tract
Forceps Major
Uncinate Fasciculus
Cingulum Hippocampus
Inferior Fronto-Occipital Fasciculus

173   *Figure 2. White Matter Tracts Modelled in the Analyses.*

Analysis Methods and Structural Equation Modelling

Covariance matrices and scripts replicating key analyses can be obtained from:

https://github.com/df1234/gf_development. Supplementary Figure 5 and 6 show correlation

matrices of all tasks and white matter tracts modelled. We modelled raw scores for g_f and

working memory tasks, as preregistered. Raw scores on processing speed tasks were

transformed. This step was not preregistered, but found necessary to achieve model

convergence to ensure interpretability of scores. First, we inverted response time scores

(using the formula y = 1/x) to obtain more intuitive measures of 'speed' for all but the CNB

Motor Speed task, for which raw scores were already a measure of speed. Afterwards, we

applied a log-transformation to reaction time tasks to increase normality and aid estimation.

For the CNB Motor Speed task only, we additionally removed values ± 2 *SD* of the mean (*N* =

6) because the presence of these outliers had caused convergence problems.

We modelled the associations between cognition and white matter microstructure using SEM

in R (R core team 2015) using the package lavaan (Rosseel 2012). All models were fit using

maximum likelihood estimation with robust Huber-White standard errors and a scaled test

statistic. Missing data was addressed using full information maximum likelihood estimation.

We used SEM Trees to investigate whether the associations among cognitive and neural

measures differed with age. SEM Trees use decision tree methods to hierarchically split a

dataset into subgroups if parameter estimates differ significantly based on a covariate of

interest - in this case age (Brandmaier et al. 2013). We first ran the watershed model in

OpenMx (Boker et al. 2011) and then passed this model object to semtree to compute the

SEM Trees. We ran one SEM Tree for each parameter of interest (e.g. the covariance

between working memory and processing speed). All other parameters in each semtree

object were set to be invariant across groups to ensure that splits were specific to the

parameter of interest. We used a 10 - fold cross-validation estimation method as recommend

by (Brandmaier et al. 2013). For the path from the cingulate to working memory only we

used 5 - fold cross-validation because the model did not converge using 10 - fold cross-

validation. Minimum sample size in age group was set to N = 50 to ensure reliable estimation

of standard errors. Note that this choice effectively limited search space for potential splits to

ages 6.58 - 12.42 years for CALM and 8.08 - 15.49 years for NKI-RS.

Results

To evaluate the hypotheses generated by the watershed model, we built up the watershed

model in steps and carried our comprehensive tests of model fit at each step. First, we

assessed the overall fit of our models to the data using the chi-square test, root mean square

error of approximation (RMSEA), comparative fit index (CFI) and standardized root mean

square residual (SRMR). Good absolute fit was defined as RMSEA < 0.05, CFI > 0.97 and SRMR

< 0.05; acceptable fit as RMSEA = 0.08 - 0.05, CFI = 0. 95 - 0.97, SRMR = 0.05 - 0.10

(Schermelleh-Engel et al. 2003). Second, we assessed specific predictions from our models by

comparing them to alternative models. Comparative model fit for nested models was

assessed using the chi-square difference test. Non-nested models were compared using the

Akaike (AIC) weights, which indicates the probability of a model being the data-generating

model compared to all other models tested (Wagenmakers and Farrell 2004). Lastly, we

evaluated the significance and strength of relationships between specific variables in our

models by inspecting the Wald test for individual parameters, noting the joint R^2 where

relevant and reporting standardized parameter estimates. Absolute standardized parameter

estimates above 0.10 were defined as small effects, 0.20 as typical and 0.30 as large (Gignac

and Szodorai 2016).

The Measurement Model of Cognition

To examine the neurocognitive architecture of g_f, we started by modelling the cognitive

components of the watershed model: g_f, working memory and processing speed. Specifically,

we fit a three-factor model of cognition (Figure 3) and compared it to alternative

measurement models. This approach allowed us to test Hypothesis 1: namely that g_f, working

memory and processing speed form three separable, albeit likely correlated cognitive factors.

The Three-Factor Model (Figure 3) showed excellent *absolute fit* for both the CALM and NKI-

RS sample (Table 3), indicating that overall, the data was compatible with a model of g_f,

working memory and processing speed as three separate factors.

Figure 3. *Different Measurement Models of Cognition.* Abbreviations: WM: working memory, PS: processing speed

The Three-Factor Model also showed very good *comparative fit* for NKI-RS, with a 96.60% probability of being the data-generating model compared to all alternative models tested, as indicated by its AIC weight (Figure 3). The evidence was more mixed for CALM, for which the Three-Factor Model showed a 27.15% probability of being the data-generating model, while Two-Factor Model B (Figure 3, treating working memory and g_f as a unitary factor) showed a 72.85% probability of being the data-generating model, highlighting a close relationship between g_f and working memory for this sample. The Single-Factor Model and Two-Factor Model A (Figure 3, treating speed and g_f as a unitary factor) showed a very low (approximately 0%) probability of being the data-generating model, indicating that speed and g_f were clearly separable in both samples.

243 *Table 3. Model Fit of Competing Measurement Models*

Single-Factor Model	Two-Factor Model A	Two-Factor Model B	Three-Factor Model
CALM			
$\chi^2(20) = 70.28$, $p < .001$	$\chi^2(19) = 67.99$, $p < .001$	$\chi^2(19) = 41.66$, $p = .002$	$\chi^2(18) = 41.74$, $p = .001$
RMSEA = .068	RMSEA = .068	RMSEA = .047	RMSEA = .049
[.051-.085]	[.052 - .086]	[.027 - .066]	[.030 - .068]
CFI = .963	CFI = .964	CFI = .983	CFI = .983
SRMR = .047	SRMR = .043	SRMR = .032	SRMR = .032
AIC = 9697.18	AIC = 9696.44	AIC = 9668.58	AIC = 9670.55
BIC = 9800.66	BIC = 9804.24	BIC = 9776.37	BIC = 9782.66
AIC$_{weight}$ = 0%	AIC$_{weight}$ = 0%	AIC$_{weight}$ = 72.85%	AIC$_{weight}$ = 27.15%
NKI-R			
$\chi^2(35) = 109.96$, $p < .001$	$\chi^2(34) = 108.15$, $p < .001$	$\chi^2(34) = 64.85$, $p = .001$	$\chi^2(32) = 54.15$, $p = .009$
RMSEA = .080	RMSEA = .081	RMSEA = .052	RMSEA = .045
[.064 - .097]	[.064 - .098]	[.033 - .071]	[.024 - .065]
CFI = .936	CFI = .936	CFI = .974	CFI = .981
SRMR = .045	SRMR = .044	SRMR = .035	SRMR = .030
AIC = 7155.64	AIC = 7155.74	AIC = 7109.43	AIC = 7102.74
BIC = 7270.07	BIC = 7273.98	BIC = 7227.67	BIC = 7228.60
AIC$_{weight}$ = 0%	AIC$_{weight}$ = 0%	AIC$_{weight}$ = 3.40%	AIC$_{weight}$ = 96.60%

244
Note. See Figure 3 for the configuration of different models. Abbreviations: Akaike Information Criterion (AIC), Bayesian Information Criterion (BIC), Akaike weight (AIC$_{weight}$)

Overall, these result provide mixed evidence for Hypothesis 1: Even though working memory,

processing speed and g_f were highly correlated in both samples (Table 4), processing speed

formed a clearly separable factor from working memory and g_f in both samples. Working and

g_f, however, were clearly separable only in NKI-RS, but not CALM, suggesting greater

similarity between g_f and working memory in the CALM sample. To facilitate comparison

across samples and in accordance with our preregistered analysis plan we nonetheless used

the three-factor measurement model (Table 4, Supplementary Table 1) in all subsequent

analyses.

Table 4. Covariance between Cognitive Measures in the Three-Factor Model

Sample	Path	Standardized Estimate
CALM	g_f <-> memory	0.71, $z = 28.42$, $p < .001$
	g_f <-> speed	0.55, $z = 12.20$, $p < .001$
	memory <-> speed	0.79, $z = 19.35$, $p < .001$
NKI-RS	g_f <-> memory	0.91, $z = 19.51$, $p < .001$
	g_f <-> speed	0.81, $z = 24.73$, $p < .001$
	memory <-> speed	0.87, $z = 17.43$, $p < .001$

Note. See Supplementary Table 1 for factor loadings.

The Relationship between Working Memory, Processing Speed and g_f

We next examined the relationships between working memory, processing speed and g_f in more detail. Specifically, we fit a SEM including regression paths between working memory and g_f, as well as speed and g_f, to test Hypothesis 2, that working memory and processing speed each predict individual differences in g_f. We found that this model showed good absolute fit for both samples (CALM: $\chi^2(18) = 41.74$, $p = .001$; RMSEA = .049 [.030 - .068]; CFI = .983; SRMR = .032, NKI-RS: $\chi^2(32) = 54.15$, $p = .009$; RMSEA = .045 [.024 - .065]; CFI = .981; SRMR = .030), indicating that, overall, the data was compatible with our model.

To further scrutinize the relationship between g_f, working memory and speed, we compared our freely-estimated model to a set of alternative models with different constraints imposed upon the regression paths. First, to test whether working memory and speed each made different contributions, we tested an alternative model in which the paths from processing speed and working memory to g_f were constrained to be equal. In CALM ($\Delta\chi^2(1) = 15.53$, $p < .001$), but not NKI-RS ($\Delta\chi^2(1) = 3.25$, $p = .072$), the freely-estimated model fit better than the equality-constrained model, indicating that working memory and speed each made different contributions in CALM but not NKI-RS. Next, we tested whether the freely estimated model fit better than a model in which the path between g_f and working memory was constrained

to zero. We found that that the freely estimated model fit better for both samples (CALM:

$\Delta\chi^2(1) = 20.77$, $p < .001$; NKI-RS: $\Delta\chi^2(1) = 12.97$, $p < .001$). In line with our hypothesis, this

result indicates that working memory makes a significant incremental contribution to g_f.

Finally, we tested a model in which the path between g_f and processing speed was

constrained to zero. This model showed no difference in fit to the freely estimated model for

CALM ($\Delta\chi^2(1) = 0.02$, $p = .875$) or NKI-RS ($\Delta\chi^2(1) = 0.04$, $p = .849$). Contrary to our hypothesis,

this indicates that there was no clear incremental contribution of processing speed to g_f.

Finally, we inspected standardized path estimates of the freely estimated model to assess the

effect seizes of working memory and processing speed. Parameter estimates showed that

working memory showed a greater effect on g_f than processing speed, particularly in CALM

(Table 5) even though raw correlations between g_f and speed were high in both samples

(Table 4).

Table 5. Regression Path Estimates.

Sample	Path	Standardized Estimate
CALM	speed -> g_f	-0.01, $z = -0.16$, $p = .876$
	memory -> g_f	0.72, $z = 7.65$, $p < .001$
NKI-RS	speed -> g_f	0.06, $z = 0.21$, $p = .208$
	memory -> g_f	0.86, $z = 1.81$, $p = .070$

Overall these results provide mixed evidence for Hypothesis 2: There was good evidence that

working memory and speed made a significant joint contribution to g_f, and that working

memory made an incremental contribution to g_f in CALM. Contrary to our hypothesis, and

the watershed model, however, processing speed showed no significant incremental

contribution to g_f, above and beyond working memory. We explore likely explanations for this

finding in the Discussion.

The Measurement Model of White Matter

We next examined the measurement model of white matter to test Hypothesis 3, namely that white matter microstructure is a multi-dimensional construct. Specifically, we tested whether white matter microstructure could be adequately captured by a single factor by examining absolute model fit. As expected, the single-factor model of white matter microstructure did not fit the data well (CALM: $\chi^2(35) = 124.63$, $p < .001$; RMSEA = .125 [.103 - .147]; CFI = .933; SRMR = .039; NKI-RS: $\chi^2(35) = 132.33$, $p < .001$; RMSEA = .204 [.167 - .242]; CFI = .885; SRMR = .023). This indicates that white matter microstructure could not be reduced to a single 'global FA' dimension in our samples, in line with (Lövdén et al. 2013; Kievit, Davis, Griffiths, Correia, CamCAN, et al. 2016) and supporting Hypothesis 3. We therefore modelled each of the ten white matter tracts separately in all subsequent models.

The Watershed Model: Relationships between Cognition and White Matter

Next, we fit the full watershed model including white matter, working memory, processing speed and g_f. Following our general analysis procedure, we investigated overall model fit, alternative models and individual path estimates to gain a comprehensive understanding of the relationships in the watershed model and to test Hypothesis 4 - that white matter contributes to working memory capacity and processing speed, which, in turn, contribute to g_f.

We found largely converging results across samples. The watershed model showed good absolute fit in CALM ($\chi^2(78) = 107.78$, $p = .014$; RMSEA = .026 [.012 - .038]; CFI = .981; SRMR = .043) and acceptable fit in NKI-RS ($\chi^2(112) = 219.22$, $p < .001$; RMSEA = .053 [.043 - .064]; CFI = .928; SRMR = .088). White matter explained large amounts of variance in working memory ($R^2_{CALM} = 32.3\%$; $R^2_{NKI-RS} = 46.1\%$) and processing speed ($R^2_{CALM} = 38.2\%$; $R^2_{NKI-RS} =$

54.4%), which, in turn, explained even more variance in g_f (R^2_{CALM} = 51.2%; R^2_{NKI-RS} = 78.3%).

In line with Hypothesis 4, this indicates that the watershed model fit the data overall.

Comparing the freely estimated watershed model to alternative, constrained, models

showed that white matter contributed significantly to memory and processing speed.

Specifically, a model in which paths from white matter to processing speed were constrained

to zero fit worse than the freely-estimated model (CALM: $\Delta\chi^2$(10) = 50.26, p < .001; NKI-RS:

$\Delta\chi^2$(10) = 27.19, p = .002), as did a model in which paths from white matter to working

memory were constrained to zero (CALM: $\Delta\chi^2$(10) = 52.26, p < .001; NKI-RS: $\Delta\chi^2$(10) = 25.85,

p = .004). As hypothesised, white matter therefore contributed to both processing speed and

working memory.

We next inspected that relationship between individual white matter tracts and working

memory and speed in more detail. A model in which paths from white matter to working

memory and speed were constrained to be equal, fit worse than the freely-estimated

watershed model for CALM ($\Delta\chi^2$(18) = 47.76, p < .001) and NKI-RS ($\Delta\chi^2$(18) = 30.42, p = .034),

indicating that the role of white matter microstructure in supporting working memory and

processing speed differed across tracts. This supports the notion that there is a many-to-one

mapping between white matter and cognition - a core tenet of the watershed model.

Investigating individual standardised parameter estimates of the different white matter tracts

showed that for CALM, only the anterior thalamic radiation contributed significantly to

processing speed, whereas the superior longitudinal fasciculus, forceps major and cingulum

were significantly, independently and positively related to working memory (Figure 4). For

NKI-RS, the superior longitudinal fasciculus was significantly and positively related to

processing speed and working memory (Figure 5). Two tracts showed an unexpected,

strongly negative (< -1), relationship: the forceps minor for CALM and the inferior fronto-

occipital fasciculus for NKI-RS. We found that these negative estimates occurred only when

all other brain to cognition pathways were also estimated: When estimated on their own,

path estimates were positive (forceps minor to working memory: standardized estimate =

0.36, *z* = 4.05, *p* < .001; inferior fronto-occipital fasciculus to working memory: standardized

estimate = 0.14, *z* = 0.859, *p* = .390; inferior fronto-occipital fasciculus to processing speed:

standardized estimate = 0.26, *z* = 1.41, *p* = .158). This sign-flip suggests that the negative

pathways were potentially due to modelling several, highly-correlated paths at the same time

(Jöreskog 1999). Overall, these results further support the watershed prediction that multiple

white matter tracts map onto working memory and processing speed.

349
Figure 4. The Watershed Model in CALM. See Supplementary Table 2 for regression
estimates. Residual covariances between white matter tracts were allowed but are not
shown for simplicity. Abbreviations: uncinate fasciculus (UF), superior longitudinal fasciculus
(SLF), inferior fronto-occipital fasciculus (IFOF), anterior thalamic radiations (ATR),
cerebrospinal tract (CST), forceps major (FMaj), forceps minor (FMin), dorsal cingulate gyrus
(CG), ventral cingulate gyrus (CH), inferior longitudinal fasciculus (ILF).

Figure 5. The Watershed Model in NKI-RS. See Supplementary Table 3 for regression estimates. Residual covariances between white matter tracts were allowed but are not shown for simplicity.

Finally, we probed the watershed model in more detail by testing a set of alternative expressions of the watershed model still compatible with the core tenants of the watershed model – as well as a set of alterative models incompatible with the watershed model. We compared all alternatives (see Figure 6 for graphical representations) to the original watershed model by inspecting their relative probability of being the data-generating model as indicated by their AIC weights (Wagenmakers and Farrell 2004). We found that the original watershed model showed a very high probability (98.58%) of being the data-generating model for CALM but only a 0.10% probability for NKI-RS. For NKI-RS, a different expression of the watershed model, such that g_f was regressed on working memory, which was regressed on processing speed, which was then regressed on white matter (Alternative A, Figure 6) showed a 95.04% probability of being the data-generating model. This model only showed a

0.37% probability for CALM. Another expression of the watershed model, in which all tasks

were modelled separately as manifest, rather than latent, variables (Alternative B, Figure 6),

showed no advantage over the watershed model for CALM (0.00% probability) or NKI-RS

(0.00% probability). We next tested two alternative models incompatible with the tenants of

the watershed model. We found that a model in which the hierarchy between cognitive

endophenotypes and g_f was inverted (Alternative C, Figure 6) showed comparatively low

probability of being the data-generating model for both CALM (0.00%) and NKI-RS (2.86%).

Similarly, a model in which g_f was directly regressed on white matter, working memory and

processing speed (Alternative D, Figure 6), showed no clear advantage over the watershed

model for CALM (1.05% probability) or NKI-RS (0.00% probability). Overall these model

comparisons highlight that while the watershed model fit the data for both samples and had

large explanatory power (as indicated by R^2s), the precise configuration of the watershed

model may differ somewhat between cohorts.

In summary, we found that the watershed model performed well overall for both cohorts. As

hypothesised, white matter contributed to working memory and processing speed, which, in

turn, contributed to g_f, and explained large amounts of variance therein. Also as predicted by

the watershed model, there was a many-to-one mapping between white matter tracts and

cognition. The exact configuration of the watershed model, however, may differ slightly

between cohorts. These differences may be a function of cohort differences in sample size,

average levels of cognitive ability and/or the specific tasks used – a topic we will return to in

the Discussion.

Figure 6. Configuration of Alternative Models. Alternatives A and B are watershed-compatible, while C and D are watershed-incompatible. The best-fitting model for CALM is highlighted in blue; the best-fitting model for NKI-RS is highlighted in green. Regression paths only are shown for simplicity. Square shapes denote manifest variables and oval shapes latent variables.

Testing for potential confounds

We carried out a series of supplementary and non-preregistered analyses to examine whether possible confounders influenced our models. These analyses showed that our findings were robust to the inclusion of covariates such as scanner motion or socio-economic status. They were also robust across genders and participants taking or not taking medication. There were no differences in the structure of the model between participants with and without diagnosed disorders for CALM. Potential small differences cannot be ruled out for NKI-RS, likely due to the low number of diagnosed participants of $N = 106$ (Supplementary Material).

Age-Related Differences in the Neurocognitive Architecture of g_f

Finally, we tested Hypothesis 5 - that that the contribution of working memory and processing speed to g_f varied with age. We first inspected cross-sectional differences in g_f, working memory and processing speed, and then used SEM trees to investigate potential age-differences in the relationships between these factors. In additional, non-preregistered, analyses we also used SEM Trees to investigate potential age-differences in the relationship between white matter and cognitive endophenotypes by inspecting paths that were significant in the watershed model (Figure 4 and 5).

SEM trees combine SEMs with decision tree methods, separating a dataset into subgroups (in this case age groups) if SEM parameter estimates of interest differ sufficiently (Brandmaier et al. 2013). SEM trees allowed us to investigate age as a potential moderator without imposing a-priori categorical age splits. We initially allowed for no more than two age groups. This yielded inconsistent results for CALM and NKI-RS (Supplementary Table 4). To test whether these inconsistencies were an artefact of allowing for only two groups, we repeated our analysis and allowed for up to four age groups. This analysis yielded consistent results between CALM and NKI-RS (Table 6). This pattern of results indicates that the initial parameters of our analysis caused us to miss relevant age differences.

As shown in Figure 7, g_f, working memory and processing speed factor scores increased with age for all three cognitive phenotypes. In line with our hypothesis, SEM trees showed that there were pronounced age-related differences in brain-behaviour in childhood and adolescence (Table 6). For both samples and all but one path, there was an initially strong relationship between components of the watershed model, then a dip around ages 7 - 9 years for CALM and age 8 for NKI-RS, followed by an increase in path strength around ages 11

431 – 12 (see Supplementary Figure 7 for a graphical representation of these results).

432 Speculatively, this pattern of results is consistent with an interpretation of a reorganization of

433 neurocognitive faculties in late childhood, followed by a consolidation of neurocognitive

434 pathways around the onset of adolescence (Johnson 2000, 2011).

435 *Table 6. SEM Tree Results for the Watershed Model.*

Path	Est. Before	Age Split 1	Est. Betw.	Age Split 2	Est. Betw.	Age Split 3	Est. After
CALM							
memory <–> speed	0.85	**8.46**	0.97	**9.46**	0.74	-	-
memory –> g_f	0.83	**9.38**	0.42	**10.04**	1.14	**10.88**	0.94
speed –> g_f	0.04	**6.88**	-0.19	**11.21**	0.17	-	-
SLF –> memory	0.67	**7.21**	0.18	**11.21**	0.76	-	-
FMaj –> memory	0.59	**7.71**	0.14	**9.29**	0.33	**11.13**	0.74
CG –> memory [1]	0.64	**6.96**	0.09	**11.04**	0.70	-	-
ATR –> speed	0.96	**7.13**	0.68	**7.96**	0.17	**11.96**	0.65
NKI-RS							
memory <–> speed	0.90	**9.82**	0.48	**14.72**	1.11	-	-
memory –> g_f	1.10	**8.59**	0.59	**12.67**	1.03	-	-
speed –> g_f	0.53	**8.59**	-0.12	**12.96**	0.52	-	-
SLF –> memory	2.15	**8.30**	1.47	**12.15**	1.93	-	-
SLF –> speed	3.12	**8.63**	1.83	**15.09**	2.31	-	-

Note. The table shows differences in parameter estimates for paths of interest (as shown in
Figure 4 and 5) depending on participants' age in years. Our analyses allowed for a maximum
of three age splits (and thus four age groups). An absence of a third age split (denoted by '-'
in the table), indicates that the SEM tree split only twice, suggesting no further changes in
parameter strength after the second split. See Supplementary Figure 7 for a graphical
representation of these results.

442
443   *Figure 7. Cognitive Factor Scores by Age.*

444   Discussion

We here used multivariate statistical techniques to investigate the neurocognitive

architecture of g_f in two large (N_{CALM} = 551, N_{NKI-RS} = 335) developmental cohorts and, for the

first time, investigated how the neurocognitive architecture of g_f changes dynamically with

age. We tested a preregistered watershed model of g_f, which predicts a hierarchy of partially

independent effects. As might be expected from a multi-cohort study, there were some

differences between the community-ascertained cohort (NKI-RS) and the cohort of children

and adolescents with learning difficulties (CALM) in specific path estimates. Overall however,

we found strikingly convergent results across these two heterogeneous samples. The

watershed model performed well for both CALM and NKI-RS: White matter contributed to

working memory and processing speed, which, in turn, contributed to g_f and explained 51%

of variance therein for the CALM sample and 78% of variance for NKI-RS. Models were robust

across genders, participants taking or not taking medication and when controlling for socio-

economic status and scanner motion. Investigations of age effects showed that the

relationship between cognitive abilities and white matter dipped in strength around ages 7-

12 years. Speculatively, this age-effect may reflect a reorganization of the neurocognitive

architecture during pre-puberty and early puberty (Byrne et al. 2017). These findings have

implications for understanding and targeting cognitive impairments in populations with

learning difficulties.

The watershed model tested here consists of three levels: g_f forms the most down-stream

point, with working memory and processing speed as intermediate tributaries, and white

matter microstructural tracts as upstream sources. Previous studies suggested that matter

microstructure is best characterised by a single, 'global FA' factor (Penke et al. 2010) while

others have contended that association patterns among different white matter tracts are

more complex (Lövdén et al. 2013; Kievit, Davis, Griffiths, Correia, CamCAN, et al. 2016). Here

we found strong evidence for a multifactorial view of white matter tracts – for both samples,

a unidimensional model of white matter fit poorly, and for CALM, multiple tracts also showed

partially independent contributions to distal cognitive outcomes. This is in line with the

watershed model. There were some differences between cohorts as to which tracts

contributed most to working memory and processing speed: In line with previous research

(Kievit, Davis, Griffiths, Correia, CamCAN, et al. 2016; MacPherson et al. 2017; Bathelt et al.

2018), we found that the anterior thalamic radiation was related to processing speed, as

were the forceps major, forceps minor and the cingulum to working memory for CALM.

However, these tracts were not significant for NKI-RS. These differences between samples

may reflect differences in brain-behaviour mapping between more atypical and typical

cohorts (Bathelt et al. 2018), or simply sampling variance across two independent cohorts

where one cohort (NKI-RS) has a more modest number of participants with imaging data. Of

note, however, the superior longitudinal fasciculus was consistently associated with working

memory in both samples. For NKI-RS, the superior longitudinal fasciculus was also associated

with processing speed. The superior longitudinal fasciculus is a large, bilateral association

fibre connecting temporal, occipital, parietal and frontal regions (Kamali et al. 2014). It is

therefore well-situated for supporting cognitive processes such as g_f, which rely on

integrative multiple-demand systems (Jung and Haier 2007; Fedorenko et al. 2013; Parlatini

et al. 2017).

The cognitive levels of the watershed model highlighted a close relationship between

working memory and g_f. Previous studies had variably suggested that g_f and working memory

(Kyllonen and Christal 1990; Fukuda et al. 2010), or g_f and processing speed (Kail and

Salthouse 1994; Salthouse 1996; Coyle et al. 2011; Ferrer et al. 2013) may be most closely

related. We found that all three cognitive factors were highly correlated for both samples.

Nonetheless, processing speed formed a cognitive factor clearly separable from working

memory and g_f. Working memory and g_f, in turn, were separable in the community-

ascertained NKI-RS but not in CALM, the cohort of children and adolescents with learning

difficulties. This close relationship between g_f and working memory was also evident in other

models of CALM where processing speed and working memory were used as joint predictors

of g_f. Contrary to our hypotheses, processing speed became non-significant after controlling

for working memory here. There are several possible, and not mutually exclusive,

explanations for this finding. First, and in line with previous work showing that time-

constraints increase isomorphism of g_f and working memory (Chuderski 2013), even standard

implementations of g_f tasks may place considerable time-pressure on struggling learners,

whereby increasing g_f - working memory covariance. Second, a broader set of speed tasks

(which might be captured by several latent variables for clerical speed tasks, choice reaction

time tasks and variability indices) might show higher predictive power than the single latent

variable for speed, which could be modelled here. Third, the watershed model might be

configured somewhat differently for some populations, such that speed forms an

intermittent level in the hierarchy between white matter and working memory (Alternative A,

Figure 6). There was some evidence for this in NKI-RS, indicating that the hierarchy of the

watershed model might be differentiated more in cohorts of older ages and/or higher ability

levels. We note that all of these explanations would still be compatible with the notions of

the watershed model, and remain to be teased apart by future research. For now, we suggest

that our findings support the notion that mental information processing capacity, as

measured by working memory, is a key determinant of g_f (Kyllonen and Christal 1990; Fukuda

et al. 2010).

The associations in the watershed model differed between ages in a complex, non-monotonic

fashion. Previous research had suggested either a decrease in covariance among cognitive

domains with age (age differentiation) (Garrett 1946), an increase in covariance with age (age

de-differentiation) (Blum and Holling 2017), or no changes with age (Tucker-Drob 2009; de

Mooij et al. 2018). These investigations have traditionally focussed on relations between

cognitive domains, however, not on relationships between brain and cognition - although

see de Mooij et al. (2018). Possible linear and non-linear changes in brain-behaviour mapping

with age have remained mostly unexplored (Tamnes et al. 2017). Using structural equation

modelling trees, a novel decision-tree-based technique, we here found evidence of complex

developmental differences that were consistent across samples and relationships in the

watershed model: Initially strong path estimates showed a pronounced decrease in strength

around ages 7 - 9 years, followed by a renewed increase in the strength, even surpassing

initial levels, around ages 10 - 15.

There are at least two possible explanations for this developmental dip in brain-cognition

relationships. First, there may be a true decrease in relationship strength during this time of

life. Possibly, other cognitive skills such as verbal reasoning temporarily support g_f, resulting

in weaker relationships between g_f and working memory, for instance. Alternatively, the

configuration of the watershed model may change temporarily during this time, which could

also manifest in an apparently weaker covariance structure. In this case, the true relationship

between g_f, memory, speed and white matter may still be strong, just configured differently

from the watershed model. Both explanations are compatible with the interactive

specialization theory (Johnson 2000, 2011), which predicts as remapping of the relationships

between brain substrates and cognitive abilities during development.

On a physiological level, this age effect may be driven by neuroendocrine changes during pre-

and early puberty. Puberty is driven by a complex and only partially understood set of

hormonal events including gonadarche and andrenarche (Sisk and Zehr 2005). Gonadarche

begins with the secretion of gonadotropin-releasing hormone from the hypothalamus around

ages 10-11 years and closely tracks the overt bodily changes of puberty (Dorn 2006).

Andrenarche, beginning with the maturation of the andrenal gland, starts as early as six years

of age, and is increasingly recognized as a complimentary driver of puberty and brain

development (Byrne et al. 2017). It is possible that the hormonal changes of andrenarche and

early gonadarche may lead to a level of neural reorganization, which may initially appear as

weaker relationships in the watershed model. The sweeping bodily, social and cognitive

changes happening in early adolescence may then drive a consolidation of the neurocognitive

architecture of g_f.

On a more general level, this age effect suggests the existence of potential non-linear

changes in brain-behaviour mapping during childhood and adolescence and underlines the

value of modern statistical approaches, such as SEM Trees, for the study of age-related

differences. Nonetheless, it is worth noting that these findings, based on an inherently

exploratory technique, will need to be replicated in future confirmatory studies with fine-

grained data on puberty and larger sample sizes. The latter will also allow for detailed

investigations of potential gender differences.

Testing our model in two different samples allowed us to address several critical questions:

First, participants from both samples completed a small set of common and a larger set of

different cognitive tasks. Therefore, the results obtained here are likely not task-specific, but

rather can be expected to generalize to the domains of working memory, processing speed

and g_f (Noack et al. 2014). For instance, we show that even though one cohort (NKI-RS)

performed mainly clerical speed tasks and the other (CALM) mainly choice reaction time

tasks, our finding that processing speed did not significantly contribute to g_f after controlling

for working memory, was robust across these samples and tasks. Second, by comparing a

community-ascertained sample (NKI-RS) and a sample of children and adolescents with

learning difficulties (CALM), we demonstrated that the watershed model performed well for

very different populations. We are not able to make general claims about potential

differences between more typical and atypical populations, however. CALM and NKI-RS were

collected in countries with somewhat different socio-economic conditions (United Kingdom

and United States of America), the samples were of different sample sizes, and participants'

ages were distributed more evenly in one cohort (NKI-RS) than the other (CALM). While DTI

images were processed with the same pipeline across sites, the scanner and MRI acquisition

protocol were different. Although previous work suggests that FA is relatively robust measure

in multi-site comparisons (Vollmar et al. 2010), we cannot rule out site differences as a

potential confound. It will therefore be necessary to replicate these findings in large typical

and atypical cohorts collected in the same setting.

Our study illustrates some of the advantages and challenges of preregistered secondary data

analyses. We agree with others in the field that secondary data analysis need not be and

should not be confounded with purely exploratory research (Mills and Tamnes 2014; Orben

and Przybylski 2019; Scott and Kline 2019). Preregistrations, as well as dedicated multivariate

methods such as SEM, can help to reduce the scope for analytic flexibility and increase

scientific rigour when using secondary datasets, which are often rich and multivariate in

nature. Preregistrations also do not preclude the use of exploratory methods (such as SEM

trees used here) or the ability to ask exploratory questions (such as looking at age differences

in the relationships between brain and cognition reported here) - they merely facilitate the

distinction between exploratory and confirmatory research (Wagenmakers et al. 2012). There

are, however, some unique challenges to preregistering secondary data analyses worth

noting. First, information on the precise measures collected is not always easily available

prior to data access, which can limit the level of detail in which an analysis can be

preregistered. Second, data quality and the level of data-processing (the latter is particularly

relevant for MRI data) is not always clear a priori (e.g. see Kievit et al. 2018), which can

necessitate changes to analyses plans after data inspection. Third, convergence issues are

fairly common when using complex multivariate methods such as SEM. We found it

necessary to transform some of our speed variables, for instance, to achieve model

convergence. Such post-hoc modifications, not guided by the palatability of the results, but

rather by unforeseen, and sometimes unforeseeable, practical considerations, mean that

preregistration can sometimes fall short of full compliance. Nevertheless, we believe that

even imperfect preregistrations, alongside shared code, data and the transparent

presentation of results, can help the reader distinguish between confirmatory and

exploratory results, and adjust their level of confidence in conclusions accordingly. For

guidance on maximizing transparency in preregistration of secondary data, see Weston et al.

(2018).

A key limitation of our study is that our samples were cross-sectional, and not longitudinal.

Although the relatively narrow age range makes large cohort effects unlikely, it may still be

that there were differences in recruitment and selection that varied across the age range.

Moreover, while we were able to investigate *individual differences* in g_f, we could not assess

intra-individual changes during childhood and adolescence. As such, the cross-sectional

nature of our samples limits our ability to make inferences about developmental dynamics.

Longitudinal data will also be necessary to scrutinize the causal flow of effects in the

watershed model.

The findings from our study have implications understanding and targeting cognitive

impairments in populations with learning difficulties. First, the close relationship between

working memory and g_f found here and in other studies (Fukuda et al. 2010; Chuderski 2013),

indicates that children and adolescents struggling with working memory are likely to also

struggle in terms of complex reasoning tasks. Either reducing working memory load,

decreasing time constraints, or training working memory and fluid ability capacity in such

populations may therefore be promising lines of inquiry for intervention studies. It is worth

highlighting, however, that cognitive training studies have so far shown little evidence of (far)

transfer: Training abstract reasoning, a common measure of g_f, has not resulted in robust

increases in working memory (Knoll et al. 2016) and working memory training has not been

shown to transfer to reasoning skills or school performance (Dunning et al. 2013;

Schwaighofer et al. 2015). Similarly, transfer from processing speed to reasoning seems to be

limited (Mackey et al. 2011). The results obtained here suggest that interventions may

increase their chance of success by implementing programs of sufficient complexity to affect

the entire neurocognitive architecture of effects (see also Kievit et al. 2016). The level of

intensity required to produce sustained benefits may need to be as demanding and

consistent as education itself, which shows robust effects in increasing general cognitive

abilities over time (Ritchie and Tucker-Drob 2018). This work and work by others (Noack et al.

2014) also highlights the value of assessing, modeling, and potentially intervening on,

multiple tasks, rather than relying on a single task to capture complex cognitive domains such

as g_f. Finally, the age-related differences in the relationships of the watershed model

observed using SEM-trees suggest that some interventions may work best at particular

developmental phases.

Acknowledgements

D.F. and J.B. are supported by the UK Medical Research Council (MRC). I.L.S.-K. is supported

by the Cambridge Trust. R.A.K. is supported by the Sir Henry Wellcome Trust (107392/Z/15/Z)

and MRC Programme Grant (MC-A060-5PR60). The Centre for Attention Learning and

Memory (CALM) research clinic is based at and supported by funding from the MRC

Cognition and Brain Sciences Unit, University of Cambridge. The Principal Investigators are

Joni Holmes (Head of CALM), Susan Gathercole (Chair of CALM Management Committee),

Duncan Astle, Tom Manly and Rogier Kievit. Data collection is assisted by a team of

researchers and PhD students at the CBU that includes Sarah Bishop, Annie Bryant, Sally

Butterfield, Fanchea Daily, Laura Forde, Erin Hawkins, Sinead O'Brien, Cliodhna O'Leary,

Joseph Rennie, and Mengya Zhang. The authors wish to thank the many professionals

working in children's services in the South-East and East of England for their support, and to

the children and their families for giving up their time to visit the clinic. We would also like to

thank all NKI-RS participants and researchers. We are grateful to Amber Ruigrok for helpful

suggestions regarding pubertal development.

References

Ackerman PL, Beier ME, Boyle MO. 2005. Working memory and intelligence: The same or
different constructs? Psychol Bull. 131:30–60.

Alloway TP. 2007. Automated Working Memory Assessment. London, UK: The Psychological
Corporation.

Bathelt J, Johnson A, Zhang M, Astle DE. 2018. Data-driven brain-types and their cognitive
consequences. bioRxiv.

Blum D, Holling H. 2017. Spearman's law of diminishing returns. A meta-analysis. Intelligence.
65:60–66.

Boker S, Neale M, Maes H, Wilde M, Spiegel M, Brick T, Spies J, Estabrook R, Kenny S, Bates T,
Mehta P, Fox J. 2011. OpenMx: An open source extended Structural Equation
Modeling framework. Psychometrika. 76:306–317.

Brandmaier AM, von Oertzen T, McArdle JJ, Lindenberger U. 2013. Structural Equation Model Trees. Psychol Methods. 18:71–86.

Byrne ML, Whittle S, Vijayakumar N, Dennison M, Simmons JG, Allen NB. 2017. A systematic review of adrenarche as a sensitive period in neurobiological development and mental health. Sensitive Periods Dev. 25:12–28.

Cannon TD, Keller MC. 2006. Endophenotypes in the genetic analyses of mental disorders. Annu Rev Clin Psychol. 2:267–290.

Chevalier N, Kurth S, Doucette MR, Wiseheart M, Deoni SCL, Dean DC III, O'Muircheartaigh J, Blackwell KA, Munakata Y, LeBourgeois MK. 2015. Myelination is associated with processing speed in early childhood: Preliminary insights. PLOS ONE. 10:e0139897.

Chuderski A. 2013. When are fluid intelligence and working memory isomorphic and when are they not? Intelligence. 41:244–262.

Cox SR, Ritchie SJ, Tucker-Drob EM, Liewald DC, Hagenaars SP, Davies G, Wardlaw JM, Gale CR, Bastin ME, Deary IJ. 2016. Ageing and brain white matter structure in 3,513 UK Biobank participants. Nat Commun. 7:13629.

Coyle TR, Pillow DR, Snyder AC, Kochunov P. 2011. Processing speed mediates the development of general intelligence (g) in adolescence. Psychol Sci. 22:1265–1269.

de Mooij SMM, Henson RNA, Waldorp LJ, Kievit RA. 2018. Age differentiation within grey matter, white matter and between memory and white matter in an adult lifespan cohort. J Neurosci.

Delis DC, Kramer JH, Kaplan E, Holdnack J. 2004. Reliability and validity of the Delis-Kaplan Executive Function System: An update. J Int Neuropsychol Soc. 10:301–303.

Dorn LD. 2006. Measuring puberty. J Adolesc Health. 39:625–626.

Dunning DL, Holmes J, Gathercole SE. 2013. Does working memory training lead to generalized improvements in children with low working memory? A randomized controlled trial. Dev Sci. 16:915–925.

Fedorenko E, Duncan J, Kanwisher N. 2013. Broad domain generality in focal regions of frontal and parietal cortex. Proc Natl Acad Sci. 201315235.

Ferrer E, Whitaker K, Steele J, Green C, Wendelken C, Bunge S. 2013. White matter maturation supports the development of reasoning ability through its influence on processing speed. Dev Sci. 16:941–951.

Finkel D, Reynolds CA, McArdle JJ, Pedersen NL. 2005. The longitudinal relationship between processing speed and cognitive ability: Genetic and environmental influences. Behav Genet. 35:535–549.

Fry AF, Hale S. 2000. Relationships among processing speed, working memory, and fluid intelligence in children. Biol Psychol. 54:1–34.

Fukuda K, Vogel E, Mayr U, Awh E. 2010. Quantity not quality: The relationship between fluid intelligence and working memory capacity. Psychon Bull Rev. 17:673–679.

Gale CR, Batty GD, Tynelius P, Deary IJ, Rasmussen F. 2010. Intelligence in early adulthood and subsequent hospitalisation and admission rates for the whole range of mental disorders: Longitudinal study of 1,049,663 men. Epidemiol Camb Mass. 21:70–77.

Gallagher A, Frederickson N. 1995. The Phonological Assessment Battery (PhAB): An initial
assessment of its theoretical and practical utility. Educ Child Psychol.

Garrett HE. 1946. A developmental theory of intelligence. Am Psychol. 1:372–378.

Gignac GE, Szodorai ET. 2016. Effect size guidelines for individual differences researchers.
Personal Individ Differ. 102:74–78.

Gur RC, Ragland JD, Moberg PJ, Turner TH, Bilker WB, Kohler C, Siegel SJ, Gur RE. 2001.
Computerized Neurocognitive Scanning: I. methodology and validation in healthy
people. Neuropsychopharmacology. 25:766.

Holmes J, Bryant A, Gathercole SE. 2018. A transdiagnostic study of children with problems of
attention, learning and memory (CALM). bioRxiv.

Hua K, Zhang J, Wakana S, Jiang H, Li X, Reich DS, Calabresi PA, Pekar JJ, van Zijl PCM, Mori S.
2008. Tract probability maps in stereotaxic spaces: analyses of white matter anatomy
and tract-specific quantification. NeuroImage. 39:336–347.

Johnson MH. 2000. Functional brain development in infants: Elements of an interactive
specialization framework. Child Dev. 71:75–81.

Johnson MH. 2011. Interactive Specialization: A domain-general framework for human
functional brain development? Dev Cogn Neurosci. 1:7–21.

Jones DK, Knösche TR, Turner R. 2013. White matter integrity, fiber count, and other fallacies:
The do's and don'ts of diffusion MRI. NeuroImage. 73:239–254.

Jöreskog KG. 1999. How large can a standardized coefficient be?

Jung RE, Haier RJ. 2007. The Parieto-Frontal Integration Theory (P-FIT) of intelligence:
Converging neuroimaging evidence. Behav Brain Sci. 30:135–154.

Kail RV, Lervåg A, Hulme C. 2015. Longitudinal evidence linking processing speed to the
development of reasoning. Dev Sci. 19:1067–1074.

Kail RV, Salthouse TA. 1994. Processing speed as a mental capacity. Acta Psychol (Amst).
86:199–225.

Kamali A, Flanders AE, Brody J, Hunter JV, Hasan KM. 2014. Tracing superior longitudinal
fasciculus connectivity in the human brain using high resolution diffusion tensor
tractography. Brain Struct Funct. 219:10.1007/s00429-012-0498-y.

Kaufman AS. 1975. Factor analysis of the WISC-R at 11 age levels between 61/2 and 161/2
years. J Consult Clin Psychol. 43:135–147.

Kievit R, Fuhrmann D, Borgeest G, Simpson-Kent I, Henson R. 2018. The neural determinants
of age-related changes in fluid intelligence: a pre-registered, longitudinal analysis in
UK Biobank [version 2; referees: 3 approved]. Wellcome Open Res. 3.

Kievit RA, Davis SW, Griffiths J, Correia MM, CamCAN, Henson RN. 2016. A watershed model
of individual differences in fluid intelligence. Neuropsychologia. 91:186–198.

Kievit RA, Davis SW, Griffiths J, Correia MM, Cam-CAN, Henson RN. 2016. A watershed model
of individual differences in fluid intelligence. Neuropsychologia. 91:186–198.

Kievit RA, Romeijn JW, Waldorp LJ, Wicherts JM, Scholte HS, Borsboom D. 2011. Mind the
gap: A psychometric approach to the reduction problem. Psychol Inq. 22:67–87.

Kline RB. 2015. Principles and practice of structural equation modeling. Guilford publications.

Knoll LJ, Fuhrmann D, Sakhardande AL, Stamp F, Speekenbrink M, Blakemore SJ. 2016. A
window of opportunity for cognitive training in adolescence. Psychol Sci. 27:1620–
1631.

Kyllonen PC, Christal RE. 1990. Reasoning ability is (little more than) working-memory
capacity?! Intelligence. 14:389–433.

Lövdén M, Laukka EJ, Rieckmann A, Kalpouzos G, Li T, Jonsson T, Wahlund L, Fratiglioni L,
Bäckman L. 2013. The dimensionality of between-person differences in white matter
microstructure in old age. Hum Brain Mapp. 34:1386–1398.

Lu PH, Lee GJ, Raven EP, Tingus K, Khoo T, Thompson PM, Bartzokis G. 2011. Age-related
slowing in cognitive processing speed is associated with myelin integrity in a very
healthy elderly sample. J Clin Exp Neuropsychol. 33:1059–1068.

Mackey AP, Hill SS, Stone SI, Bunge SA. 2011. Differential effects of reasoning and speed
training in children. Dev Sci. 14:582–590.

MacPherson SE, Cox SR, Dickie DA, Karama S, Starr JM, Evans AC, Bastin ME, Wardlaw JM,
Deary IJ. 2017. Processing speed and the relationship between Trail Making Test-B
performance, cortical thinning and white matter microstructure in older adults.
Cortex. 95:92–103.

Manly T, Anderson V, Nimmo-Smith I, Turner A, Watson P, Robertson IH. 2001. The
differential assessment of children's attention: The Test of Everyday Attention for
Children (TEA-Ch), normative sample and ADHD performance. J Child Psychol
Psychiatry. 42:1065–1081.

Mills KL, Goddings A-L, Herting MM, Meuwese R, Blakemore S-J, Crone EA, Dahl RE, Güroğlu
B, Raznahan A, Sowell ER, Tamnes CK. 2016. Structural brain development between
childhood and adulthood: Convergence across four longitudinal samples.
NeuroImage. 141:273–281.

Mills KL, Tamnes CK. 2014. Methods and considerations for longitudinal structural brain
imaging analysis across development. Dev Cogn Neurosci. 9:172–190.

Noack H, Lövdén M, Schmiedek F. 2014. On the validity and generality of transfer effects in
cognitive training research. Psychol Res. 78:773–789.

Nooner K, Colcombe S, Tobe R, Mennes M, Benedict M, Moreno A, Panek L, Brown S, Zavitz
S, Li Q, Sikka S, Gutman D, Bangaru S, Schlachter RT, Kamiel S, Anwar A, Hinz C, Kaplan
M, Rachlin A, Adelsberg S, Cheung B, Khanuja R, Yan C, Craddock C, Calhoun V,
Courtney W, King M, Wood D, Cox C, Kelly C, DiMartino A, Petkova E, Reiss P, Duan N,
Thompsen D, Biswal B, Coffey B, Hoptman M, Javitt D, Pomara N, Sidtis J, Koplewicz H,
Castellanos F, Leventhal B, Milham M. 2012. The NKI-Rockland Sample: A model for
accelerating the pace of discovery science in psychiatry. Front Neurosci. 6:152.

Oberauer K, Schulze R, Wilhelm O, Süß H-M. 2005. Working memory and intelligence - Their
correlation and their relation: Comment on Ackerman, Beier, and Boyle (2005).
Psychol Bull. 131:61–65.

Orben A, Przybylski AK. 2019. The association between adolescent well-being and digital
technology use. Nat Hum Behav.

Parlatini V, Radua J, Dell'Acqua F, Leslie A, Simmons A, Murphy DG, Catani M, Thiebaut de
Schotten M. 2017. Functional segregation and integration within fronto-parietal
networks. NeuroImage. 146:367–375.
Penke L, Maniega SM, Murray C, Gow AJ, Valdés Hernández MC, Clayden JD, Starr JM,
Wardlaw JM, Bastin ME, Deary IJ. 2010. A general factor of brain white matter
integrity predicts information processing speed in healthy older people. J Neurosci.
30:7569.
Pfefferbaum A, Sullivan EV, Hedehus M, Lim KO, Adalsteinsson E, Moseley M. 2000. Age-
related decline in brain white matter anisotropy measured with spatially corrected
echo-planar diffusion tensor imaging. Magn Reson Med. 44:259–268.
Primi R, Ferrão ME, Almeida LS. 2010. Fluid intelligence as a predictor of learning: A
longitudinal multilevel approach applied to math. Learn Individ Differ. 20:446–451.
R core team the. 2015. R: A language and environment for statistical computing.
Ritchie SJ, Tucker-Drob EM. 2018. How much does education improve intelligence? A meta-
analysis. Psychol Sci. 0956797618774253.
Rosseel Y. 2012. lavaan: An R package for Structural Equation Modeling. J Stat Softw. 48:1–
36.
Roth B, Becker N, Romeyke S, Schäfer S, Domnick F, Spinath FM. 2015. Intelligence and
school grades: A meta-analysis. Intelligence. 53:118–137.
Salthouse TA. 1996. The processing-speed theory of adult age differences in cognition.
Psychol Rev. 103:403–428.
Schermelleh-Engel K, Moosbrugger H, Müller H. 2003. Evaluating the fit of Structural
Equation Models: Tests of significance and descriptive goodness-of-fit measures.
Methods Psychol Res. 8:23–74.
Schubert A-L, Hagemann D, Frischkorn GT. 2017. Is general intelligence little more than the
speed of higher-order processing? J Exp Psychol Gen. 146:1498–1512.
Schwaighofer M, Fischer F, Bühner M. 2015. Does working memory training transfer? A
meta-analysis including training conditions as moderators. Educ Psychol. 50:138–166.
Scott KM, Kline M. 2019. Enabling Confirmatory Secondary Data Analysis by Logging Data
Checkout. Adv Methods Pract Psychol Sci. 2515245918815849.
Sheppard LD, Vernon PA. 2008. Intelligence and speed of information-processing: A review of
50 years of research. Personal Individ Differ. 44:535–551.
Simpson-Kent IL, Fuhrmann D, Bathelt J, the CALM team, Kievit RA. 2019. Cross-sectional
evidence for age differentiation-dedifferentiation in crystallized and fluid intelligence
and their white matter substrates during childhood and adolescence. Prep.
Sisk CL, Zehr JL. 2005. Pubertal hormones organize the adolescent brain and behavior. Front
Neuroendocrinol. 26:163–174.
Tamnes CK, Roalf DR, Goddings A-L, Lebel C. 2017. Diffusion MRI of white matter
microstructure development in childhood and adolescence: Methods, challenges and
progress. Dev Cogn Neurosci.

Tucker-Drob EM. 2009. Differentiation of cognitive abilities across the lifespan. Dev Psychol.
45:1097–1118.

Vestergaard M, Madsen KS, Baaré WF, Skimminge A, Ejersbo LR, Ramsøy TZ, Gerlach C,
Åkeson P, Paulson OB, Jernigan TL. 2011. White matter microstructure in superior
longitudinal fasciculus associated with spatial working memory performance in
children. J Cogn Neurosci. 23:2135–2146.

Volkow ND, Koob GF, Croyle RT, Bianchi DW, Gordon JA, Koroshetz WJ, Pérez-Stable EJ, Riley
WT, Bloch MH, Conway K, Deeds BG, Dowling GJ, Grant S, Howlett KD, Matochik JA,
Morgan GD, Murray MM, Noronha A, Spong CY, Wargo EM, Warren KR, Weiss SRB.
2018. The conception of the ABCD study: From substance use to a broad NIH
collaboration. Adolesc Brain Cogn Dev ABCD Consort Ration Aims Assess Strategy.
32:4–7.

Vollmar C, O'Muircheartaigh J, Barker GJ, Symms MR, Thompson P, Kumari V, Duncan JS,
Richardson MP, Koepp MJ. 2010. Identical, but not the same: intra-site and inter-site
reproducibility of fractional anisotropy measures on two 3.0T scanners. NeuroImage.
51:1384–1394.

Wagenmakers E-J, Farrell S. 2004. AIC model selection using Akaike weights. Psychon Bull
Rev. 11:192–196.

Wagenmakers E-J, Wetzels R, Borsboom D, van der Maas HLJ, Kievit RA. 2012. An agenda for
purely confirmatory research. Perspect Psychol Sci. 7:632–638.

Wandell BA. 2016. Clarifying human white matter. Annu Rev Neurosci. 39:103–128.

Wechsler D. 2011. Wechsler Abbreviated Scale of Intelligence -Second Edition. San Antonio,
TX: Pearson.

Weston SJ, Ritchie SJ, Rohrer JM, Przybylski AK. 2018. Recommendations for increasing the
transparency of analysis of pre-existing datasets. PsyArXiv.