Integrating Vectors for Genetic Studies in the Rare Actinomycete *Amycolatopsis marina*

Hong Gao[1,*], Buvani Murugesan[1], Janina Hossbach[1], Stephanie K. Evans[1], W. Marshall Stark[2] and Margaret C.M. Smith[1]

1 Department of Biology, University of York, York, North Yorkshire, UK, YO10 5DD

2 Institute of Molecular, Cell and Systems Biology, University of Glasgow, Glasgow, UK, G12 8QQ

* Hong Gao: hong.gao@york.ac.uk

Abstract:

Few natural product pathways from rare Actinomycetes have been studied due to the difficulty in applying molecular approaches in these genetically intractable organisms. In this study, we sought to identify integrating vectors, derived using phage *int/attP* loci, that would efficiently integrate site-specifically in the rare Actinomycete, *Amycolatopsis marina* DSM45569. Genomic analysis of the genome of *A. marina* DSM45569 indicated the presence of *attB*-like sites for TG1 and R4 integrases. The TG1 and R4 *attB*s were active in *in vitro* recombination assays with their cognate purified integrases and *attP* loci. Integrating vectors containing either the TG1 or R4 *int/attP* loci yielded a high frequency of exconjugants in conjugation assays from *E. coli* to *A. marina* DSM45569. Site-specific recombination of the plasmids into the host TG1 or R4 *attB* sites was confirmed by sequencing. The presence of homologous TG1 and R4 *attB* sites in other species of this genus indicated that vectors based on TG1 and R4 integrases could be widely applicable.

Keywords: Rare Actinomycetes, Amycolatopsis, integrating vectors, TG1 integrase, R4 integrase

Introduction

Streptomyces bacteria are widely exploited for their abundant bioactive natural products[1]. After 70 years of exploitation, however, the rate of discovery of new *Streptomyces* derived bioactive products has declined, and interest in other potential non-Streptomycete sources, such as rare Actinomycetes has grown[2, 3].

Amongst rare Actinomycetes, the *Amycolatopsis* genus generates particular interest for its production of important antibiotics such as vancomycin[4] and rifamycin[5], as well as a diverse range of active natural products[6-8]. The publicly available NCBI database contains nearly 70 genomes of *Amycolatopsis* strains, covering more than 40 species from this genus. Similar to *Streptomyces*, the genome of each *Amycolatopsis* contains, on average, over 20 secondary metabolic gene clusters[9]. The mining of these metabolic clusters, much like *Streptomyces*, offers great potential for novel antibiotic discovery as the gene products of these clusters are largely uncharacterised. The lack of widely available and efficient genetic tools for these rare species, however, impedes their potential in research and development.

Phage-encoded serine and tyrosine integrases catalyse site-specific integration of a circularized phage genome into the host chromosome as part of the process to establish a lysogen. DNA integration mediated by serine integrases occurs between short (approximately 50 bp) DNA substrates that are located on the phage genome, the phage attachment site (*attP*) and the host genome, the bacterial attachment site (*attB*). The product of *attP* x *attB* recombination is an integrated phage genome flanked by two new sites, *attL* and *attR*, each of which contain half sites from *attP* and *attB*. During phage induction integrase in the presence of a recombination directionality factor (RDF) again mediates site-specific recombination but this time between *attL* and *attR* to excise the phage genome, which can then be replicated during a lytic cycle. The mechanism of recombination and the factors that control integration versus excision have been elucidated in recent years[10-12].

Integrating vectors based on the *Streptomyces* phage φC31 integrase and *attP* locus

are the best known and most widely used in Actinomycete genome engineering[13-16].

In addition to the phage recombination machinery (*int/attP*), integrating vectors

contain a replicon for maintenance in *E. coli*, an *oriT* for conjugal transfer and a

marker or markers for selection in *E. coli* and in the recipient. They are powerful

genome engineering tools that act in an efficient, highly controllable and predictable

way[17].

The use of integrating vectors such as those derived from serine integrases requires

no additional phage or host functions for integration, thus makes it particularly

salient when the system is applied to other organisms that cannot be infected by the

phages. This hallmark makes serine integrases an promising tool used in

heterologous systems[11, 18]. However, these integration systems have not been fully

explored in rare Actinomycetes, e.g. *Amycolatopsis*. There is only one example of a

conjugation system based on φC31 integrase in *Amycolatopsis japonicum*

MG417-CF17[19], and it has been reported that other *Amycolatopsis* species lack φC31

attB sites in their chromosomes[20]. The φBT1 *attB* sites have been more commonly

identified in *Amycolatopsis*, and there have been successful conjugative transfer of

vectors based on φBT1 *int/attP* in *A. orientalis*[20] and *A. mediterranei*[21]. Despite these

successes, electroporation still remains the most applicable method used to transfer

integrative plasmids into this genus[20, 21].

In this paper, we chose to study *Amycolatopsis marina* DSM45569, a particular

Amycolatopsis species isolated from an ocean-sediment sample collected in the

South China Sea[22]. We explored the application of bacterial genetic engineering

using serine integrases, and developed conjugative and integrating vectors for use in

this species. We present evidence that suggests this vector could be applied to

various other species in this genus, opening up the prospect for more versatile

genetic manipulation of *Amycolatopsis*.

Materials and Methods

Bacterial strains and culture conditions

Plasmid propagation and subcloning was conducted using *E. coli* Top10 (F- *mcrA* Δ(*mrr-hsdRMS-mcrBC*) φ80*lacZ*ΔM15 *ΔlacX74 nupG recA1 araD139* Δ(*ara-leu)7697 galE15 galK16 rpsL*(Str^R) *endA1* λ⁻). Plasmid conjugations from *E. coli* to *Streptomyces* were carried out using *E. coli* ET12567(pUZ8002) containing the plasmid to be transferred as the donor [23]. *E. coli* strains were grown in Luria-Bertani broth (LB) or on LB agar at 37°C.

Amycolatopsis marina DSM45569 was purchased from the German Collection of Microorganisms and Cell Cultures (DSMZ, Germany), and maintained on Soya Mannitol (SM) agar at 30°C. Harvested spores were maintained long term in 20% glycerol at -80°C. Conjugations were plated on SM agar containing 10 mM MgCl₂ and ISP2 medium[24] was used for the preparation of genomic DNA[25].

DNA manipulation

E. coli transformation and gel electrophoresis were carried out as described previously[26]. Genomic DNA preparation from *Streptomyces* was performed following the salting out procedure in the *Streptomyces* manual[25]. Plasmids from *E. coli* were prepared using QIAprep® Spin Miniprep Kit (Qiagen, Germany) following the manufacturer's instructions. Polymerase Chain Reaction (PCR) was carried out using Phusion® High-Fidelity DNA Polymerase (NEB, USA) according to the manufacturer's instructions. The primers used in this study are listed in Table 1. DNA samples were purified by the QIAquick Gel Extraction Kit (Qiagen, Germany).

Plasmid construction

The integrating plasmid pHG4 contains the TG1 *int/attP* locus and the apramycin-resistance gene (*aac3(IV)*) for selection (Figure 1). The fragment containing *oriT*, *aac3(IV)* and TG1 *int/attP* was amplified from plasmid pBF20[27] using the primer pair pHG4-for/pHG4-rev. The fragment was joined via In-Fusion cloning to the 3344 bp HindIII-PacI fragment from pBF22[27] (containing the *E. coli* plasmid replication origin, the *bla* gene encoding resistance to ampicillin and the

actII-orf4/act1p expression cassette) to form the plasmid pHG4.

To construct the integrating plasmid pJH1R4 (Figure 1), pSET152[28] was cut with AatII

and PvuI to remove the φC31 *attP* site and integrase gene. R4 phage lysate was used

as template in a PCR with the primers pJH1R4-for and pJH1R4-rev to amplify the R4

attP site and integrase coding region. The PCR product was joined to the AatII-PvuI

fragment from pSET152 via In-Fusion cloning.

The plasmid pHG1 was used as template to amplify attB-containing sequences for in

vitro recombination assays. This plasmid was originally constructed for the

expression of *EryF*. The eryF gene was amplified from *Saccharopolyspora erythraea*

BIOT-0666 genomic DNA using the primer pair pHG1A-for/pHG1A-rev, and inserted

by In-Fusion cloning into pBF20[27] cut with NheI and PacI to form the plasmid pHG1A.

The 3785 bp fragment containing the φC31 int/attP and hygromycin resistance gene

was amplified from plasmid pBF27C[27], using the primer pair pHG1-for and pHG1-rev.

Plasmid pHG1A was digested with XbaI and NheI, the 5668 bp fragment was ligated

with the 3785 bp PCR fragment from pBF27C by In-Fusion to give the plasmid pHG1

(Figure 1).

In Vitro Recombination Assays

In vitro recombination assays were performed using PCR amplified DNA fragments

containing the *attB* and *attP* attachment sites located at the ends. Recombination

between the *attP* and *attB* sites yielded a fusion of the two fragments to give a

product, whose length was almost the sum of the substrates (Figure 3A). To generate

the *attB* containing substrates, the forward primer, TG1-attB-Am-for, contained the

closest match in the *A. marina* genome to the characterised TG1 *attB* site from *S.*

avermitilis, TG1 *attB*[Sa] [29](Figure 2). TG1-attB-Am-for also had a sequence identical to

the 3' end of the act1p element from plasmid pHG1, which was used as a template

for PCR (Figure 1). Similarly, the forward primer R4-attB-Am-for contained the closest

match in the *A. marina* genome to the characterized R4 *attB* site from *S. parvulus*, R4

attB[Sp] [30] (Figure 2). R4-attB-Am-for also had a sequence identical to the 3' end of

ActII-orf4 element from the template plasmid pHG1 (Figure 1). Forward primers

TG1-attB-Sa-for and R4-attB-Sp-for were used to create positive control

recombination substrates containing the TG1 and R4 *attB* sites originally found in *S.*

avermitilis[29] and *S. parvulus*[30] respectively. The reverse primer used to generate all

the *attB* containing substrates (attB-rev) was located within the *hyg* gene of pHG1

and the amplified products were 1627 bp (TG1 attB^{Am}), 1035 bp (TG1 attB^{Sa}), 1854 bp

(R4 attB^{Am}) and 1855 bp (R4 attB^{Sp}). The DNA fragments containing the *attP* sites

were prepared as follows; the TG1-attP fragment (2471 bp) was amplified using the

primer pair TG1-attP-for/TG1-attP-rev with pHG4 as the template, and the R4-attP

fragment (990 bp) was amplified using the primer pair R4-attP-for/R4-attP-rev with

pJH1R4 as the template (Figure 1). Note that other than the *attB* and *attP* sites none

of the substrates contained any DNA that should interact with the integrases other

than non-specifically. Moreover each fragment was designed to be easily identifiable

by molecular weight.

The integrases were purified as described previously[31, 32]. All recombination reactions

were in 20 µl final volume. Recombination reaction of TG1 substrates was carried out

in TG1 RxE buffer (20 mM Tris [pH 7.5], 25 mM NaCl, 1 mM dithiothreitol [DTT], 10

mM spermidine, 10 mM EDTA, 0.1 mg/ml bovine serum albumin [BSA])[33], and

recombination reaction of R4 substrates was carried out in buffer containing 20 mM

Tris-HCl (pH 7.5), 50 mM NaCl, 10 mM spermidine, 5 mM CaCl$_2$ and 50 mM DTT[32].

Integrase was added at the concentrations indicated. Recombination substrates were

used at 50 ng each per reaction. Reactions were incubated at 30°C overnight, and

then heated (10 min, 75°C) to denature integrase. The reaction mixtures were loaded

on a 0.8% agarose gel in Tris/Borate/EDTA (TBE) buffer (90 mM Tris base, 90 mM

boric acid and 2 mM EDTA) containing ethidium bromide for electrophoretic

separation.

Results

Identification of possible *attB* sequences from the genome of *Amycolatopsis*

***marina* DSM45569**

The sequences of *attB* sites recognised by a variety of integrases (φ31[34], φJoe[35],

Bxb1[36], R4[32], SPBc[37], SV1[38], TG1[39] and TP901[40]) were used in BLAST searches of the

genome sequence of *Amycolatopsis marina* DSM45569 (NCBI Genome Database

NZ_FOKG00000000)(Table 2). The most significant hits for R4 and TG1 *attB* sites had

the highest identities and lowest *E*-value. The R4 *attB* site is located within a gene

predicted to encode a fatty-acyl-CoA synthase (SFB62308.1) and the TG1 *attB* site is

located within a gene predicted to encode a putative succinyldiaminopimelate

transaminase (WP_091671332.1). The BLAST analysis was extended to other species

of *Amycolatopsis* to assess the conservation of *attB* sites in the genus (Figure 2). Both

R4 and TG1 *attB* sites were highly conserved to the *attB* sites (84% for R4 and 62%

for TG1) originally identified from *S. parvulus*[30] and *S. avermitilis*[29].

***In Vitro* Recombination**

In each recombination reaction, substrates containing *attP* and the putative *attB* site

were mixed in cognate pairs with different concentrations of purified R4 or TG1 in

the corresponding buffer and incubated overnight at 30°C, as described in Materials

and Methods. The expected recombination events and the nature of the products

are shown in Figure 3A-3C. TG1 catalysed the recombination between substrates

more efficiently than R4 (Figure 3D). As expected because neither phage is an

Amycolatopsis phage, the recombination efficiencies for each integrase were

observably better when the *Streptomyces attB* sites were used (Figure 3E) compared

to the *A. marina attB* sites (Figure 3D), particularly for TG1 integrase. Nevertheless,

the presence of recombination activity indicated that both *A. marina attB* sites were

functional in a recombination reaction and were likely to be active integration sites

for integrative conjugation vectors.

***In Vivo* Integration**

A. marina DSM45569 was unable to grow in the presence of apramycin so integrating

plasmids, pHG4 and pJH1R4, containing the apramycin resistance determinant,

aac3(IV) were constructed. Following the standard *Streptomyces* conjugation

protocol (see Materials and Methods), a frequency of approximately 160

exconjugants/10^8 spores was obtained for transfer of pHG4 (encoding TG1 integrase),

while the conjugation efficiency of pJH1R4 (R4 integrase) was only 20

exconjugants/10^8 spores (Table 3). For each integration, six exconjugants were picked

at random and streaked on MS agar containing apramycin first, then genomic DNA

was prepared and used as template in PCR reactions, in which the primer pairs of

TG1-attL-Am-for/rev and R4-attL-Am-for/rev were used to test the occurrence of

recombination at the expected TG1 and R4 *attB* sites (Figure 4). All PCR reactions

using exconjugants as templates gave the expected band sizes. Sequencing (GATC,

Germany) of the PCR products with the primers TG1-attL-Am-for and R4-attL-Am-for

confirmed that the plasmids had integrated into the predicted *attB* sites for TG1 or

R4 integrase within *A. marina* DSM45569 (Figure 5).

Discussion

The lack of effective genetic engineering tools is considered one of the greatest

hindrances in the search for novel natural products from rare Actinomycetes[41-43].

Previous studies have largely focused on the use of the well-characterised ϕC31

based integration vectors, and have mostly overlooked other phage integrases in rare

Actinomycete species[44-46]. Additionally, the conjugation methods used widely in

Streptomyces gene transfer have shown little success in rare Actinomycetes,

including the genus *Amycolatopsis* species, thus electroporation has been the

long-preferred method of gene transfer in this species[5, 9, 47, 48]. However, the growing

interest in the use of serine integrases for synthetic biology applications[11] has led to

the further research into expanding the pool of available enzymes and their

potentials as genetic tools[49-51]. Therefore, within this study, we explored whether

integrating vectors derived from eight different serine integrases could be employed

for genetic engineering of *A. marina* DSM45569. Sequence analysis of the *A. marina*

DSM45569 genome identified close matches to the *attB* sites used by TG1 and R4

integrases. Although conjugation frequencies were low, integrating plasmids based

on the TG1 and R4 recombination systems successfully integrated into the expected

attB sites in *A. marina* DSM45569. Conservation between the *attB* sites for TG1 and

R4 in a number of *Amycolatopsis* species is high suggesting that the plasmids derived

from these phages should be widely useful in this genus.

As is common with serine integrase mediated recombination, the attB sites are

located within open reading frames and potentially disrupt the gene. TG1 $attB^{Am}$ site

is located within a gene predicted to encode an aminotransferase, and the R4 $attB^{Am}$

site is located within a gene predicted to encode a fatty-acyl-CoA synthase.

Compared to the wild type (unintegrated) strain, the strains with integrated pHG4 or

pJH1R4 did not show any difference in growth. However, further study is required to

investigate the effects of TG1 or R4 plasmid recombination on both of primary and

secondary metabolism as, for example, the integration of φC31 derived plasmids has

been shown to have pleiotropic effects on bacterial physiology[52].

Currently, the two most commonly used methods of bacterial gene transfer are

conjugation and electroporation, and both come with several advantages and

disadvantages. Electroporation is the introduction of pores within bacterial

membranes via an electric current to allow mass, unrestricted transfer of genetic

material into species[53, 54]. The efficiency of transformation, however, is species

dependent[53]. Unlike electroporation, conjugation is not limited by the size of vectors

that can be transformed, and has been used to successfully transfer entire genomes

in *E. coli*[55]. Additionally, conjugation uses the transfer of DNA as a single strand that is

relatively insensitive to the majority of the restriction systems of the cells[56]. This

demonstrates the advantageous use of conjugation over electroporation. In this

study, we successfully integrated a plasmid via the *attB* sites for TG1 and R4

integrases by conjugation, thus supplementing the potential gene transfer methods

that could be used in the genus *Amycolatopsis*.

In conclusion, we have identified the highly conserved sequences of the *attB* sites for

TG1 and R4 integrases within the genus *Amycolatopsis* and demonstrated their use in

conjugative DNA transfer. The *A. marina* DSM45569 *attB* sites showed slightly lower

recombination efficiencies *in vitro* than the previously identified *attB* sites from

Streptomyces spp. although this is not likely enough to explain the order of

magnitude reductions in conjugation frequencies observed with *A. marina* compared

to *Streptomyces* spp. (Table 3). By optimising conjugation conditions the conjugation

frequencies could be increased. Alternatively, efficiently used *attB* sites for the

widely used vectors, such as those based on phiC31 *int/attP* could be incorporated in

to the *Amylcolatopsis* genome using TG1 or R4 integrating plasmids described here.

In short this work shows that integrative vectors are viable and promising tools for

the genetic engineering of rare actinomycetes.

Acknowledgements

272 This work was supported by the Biotechnology and Biological Sciences Research

273 Council project grant BB/K003356/1, and Buvani Murugesan acknowledges the

274 receipt of a summer studentship from the Department of Biology, University of York.

References

1.   Jones, S.E. et al. Streptomyces exploration is triggered by fungal interactions and volatile
signals. *Elife* **6**, e21738 (2017).
2.   Zarins-Tutt, J.S. et al. Prospecting for new bacterial metabolites: a glossary of approaches for
inducing, activating and upregulating the biosynthesis of bacterial cryptic or silent natural
products. *Natural product reports* **33**, 54-72 (2016).
3.   Jose, P.A. & Jebakumar, S.R.D. Non-streptomycete actinomycetes nourish the current
microbial antibiotic drug discovery. *Frontiers in microbiology* **4** (2013).
4.   Jung, H.-M., Kim, S.-Y., Moon, H.-J., Oh, D.-K. & Lee, J.-K. Optimization of culture conditions
and scale-up to pilot and plant scales for vancomycin production by Amycolatopsis orientalis.
Applied microbiology and biotechnology **77**, 789-795 (2007).
5.   Li, C. et al. RifZ (AMED_0655) is a pathway-specific regulator for rifamycin biosynthesis in
Amycolatopsis mediterranei. *Applied and Environmental Microbiology* **83**, e03201-03216
(2017).
6.   Hashizume, H. et al. Valgamicin C, a novel cyclic depsipeptide containing the unusual amino
acid cleonine, and related valgamicins A, T and V produced by Amycolatopsis sp. ML1-hF4.
The Journal of Antibiotics, ja2017135 (2017).
7.   Li, X., Wu, X., Zhu, J. & Shen, Y. Amexanthomycins A–J, pentangular polyphenols produced by
Amycolatopsis mediterranei S699Δ rifA. *Applied Microbiology and Biotechnology*, 1-14
(2017).
8.   Schwarz, P.N. et al. The Immunosuppressant Brasilicardin: Determination of the Biosynthetic
Gene Cluster in the Heterologous Host Amycolatopsis japonicum. *Biotechnology Journal*
(2017).
9.   Kumari, R., Singh, P. & Lal, R. Genetics and genomics of the genus Amycolatopsis. *Indian
journal of microbiology* **56**, 233-246 (2016).
10.   Colloms, S.D. et al. Rapid metabolic pathway assembly and modification using serine

integrase site-specific recombination. *Nucleic acids research* **42**, e23-e23 (2013).

11. Fogg, P.C., Colloms, S., Rosser, S., Stark, M. & Smith, M.C. New applications for phage
integrases. *Journal of molecular biology* **426**, 2703-2716 (2014).

12. Smith, M.C. in Mobile DNA III 253-272 (American Society of Microbiology, 2015).

13. Sosio, M. et al. Artificial chromosomes for antibiotic-producing actinomycetes. *Nature*
biotechnology **18**, 343-345 (2000).

14. Hong, Y. & Hondalus, M.K. Site-specific integration of Streptomyces ΦC31 integrase-based
vectors in the chromosome of Rhodococcus equi. *FEMS microbiology letters* **287**, 63-68
(2008).

15. Saha, S. et al. Activation and characterization of a cryptic gene cluster reveals a cyclization
cascade for polycyclic tetramate macrolactams. *Chemical Science* **8**, 1607-1612 (2017).

16. Thorpe, H.M. & Smith, M.C. In vitro site-specific integration of bacteriophage DNA catalyzed
by a recombinase of the resolvase/invertase family. *Proceedings of the National Academy of*
Sciences **95**, 5505-5510 (1998).

17. Rutherford, K. & Van Duyne, G.D. The ins and outs of serine integrase site-specific
recombination. *Current opinion in structural biology* **24**, 125-131 (2014).

18. Baltz, R.H. Streptomyces temperate bacteriophage integration systems for stable genetic
engineering of actinomycetes (and other organisms). *Journal of industrial microbiology &*
biotechnology **39**, 661-672 (2012).

19. Stegmann, E., Pelzer, S., Wilken, K. & Wohlleben, W. Development of three different gene
cloning systems for genetic investigation of the new species Amycolatopsis japonicum
MG417-CF17, the ethylenediaminedisuccinic acid producer. *Journal of biotechnology* **92**,
195-204 (2001).

20. Lei, X. et al. Complete genome sequence of Amycolatopsis orientalis CPCC200066, the
producer of norvancomycin. *Journal of biotechnology* **247**, 6-10 (2017).

21. Li, C., Zhou, L., Wang, Y., Zhao, G. & Ding, X. Conjugation of ΦBT1-derived integrative plasmid
pDZL802 in Amycolatopsis mediterranei U32. *Bioengineered* **8**, 549-554 (2017).

22. Bian, J. et al. Amycolatopsis marina sp. nov., an actinomycete isolated from an ocean
sediment. *International journal of systematic and evolutionary microbiology* **59**, 477-481
(2009).

23. MacNeil, D.J. Characterization of a unique methyl-specific restriction system in Streptomyces
avermitilis. *Journal of bacteriology* **170**, 5607-5612 (1988).

24. Shirling, E.t. & Gottlieb, D. Methods for characterization of Streptomyces species1.
International Journal of Systematic and Evolutionary Microbiology **16**, 313-340 (1966).

25. Kieser, T., Bibb, M.J., Buttner, M.J., Chater, K.F. & Hopwood, D.A. Practical Streptomyces
Genetics. (The John Innes Foundation, Norwich; 2000).

26. Sambrook, J. & Russell, D.W. Molecular cloning: a laboratory manual. (Cold Spring Harbor
Laboratory Press, New York; 2001).

27. Fayed, B. et al. Multiplexed integrating plasmids for engineering the erythromycin gene
cluster for expression in Streptomyces and combinatorial biosynthesis. *Applied and*
environmental microbiology, AEM. 02403-02415 (2015).

28. Wilkinson, C.J. et al. Increasing the efficiency of heterologous promoters in actinomycetes.
Journal of molecular microbiology and biotechnology **4**, 417-426 (2002).

29. Morita, K. et al. The site-specific recombination system of actinophage TG1. *FEMS*

345   *microbiology letters* **297**, 234-240 (2009).

346   30.   Shirai, M., Nara, H., Sato, A., Aida, T. & Takahashi, H. Site-specific integration of the
347   actinophage R4 genome into the chromosome of Streptomyces parvulus upon lysogenization.
348   *Journal of bacteriology* **173**, 4237-4239 (1991).

349   31.   Rowley, P.A., Smith, M.C., Younger, E. & Smith, M.C. A motif in the C-terminal domain of ΦC31
350   integrase controls the directionality of recombination. *Nucleic acids research* **36**, 3879-3891
351   (2008).

352   32.   Miura, T. et al. In vivo and in vitro characterization of site-specific recombination of
353   actinophage R4 integrase. *The Journal of general and applied microbiology* **57**, 45-57 (2011).

354   33.   Morita, K. et al. Site-specific genome integration in alphaproteobacteria mediated by TG1
355   integrase. *Applied microbiology and biotechnology* **93**, 295-304 (2012).

356   34.   Combes, P., Till, R., Bee, S. & Smith, M.C. The Streptomyces genome contains multiple
357   pseudo-attB sites for the φC31-encoded site-specific recombination system. *Journal of*
358   *bacteriology* **184**, 5746-5752 (2002).

359   35.   Fogg, P.C., Haley, J.A., Stark, W.M. & Smith, M.C. Genome integration and excision by a new
360   Streptomyces bacteriophage, ΦJoe. *Applied and environmental microbiology*, AEM.
361   02767-02716 (2016).

362   36.   Ghosh, P., Pannunzio, N.R. & Hatfull, G.F. Synapsis in phage Bxb1 integration: selection
363   mechanism for the correct pair of recombination sites. *Journal of molecular biology* **349**,
364   331-348 (2005).

365   37.   Lazarevic, V. et al. Nucleotide sequence of the Bacillus subtilis temperate bacteriophage
366   SPβc2. *Microbiology* **145**, 1055-1067 (1999).

367   38.   Fayed, B., Younger, E., Taylor, G. & Smith, M.C. A novel Streptomyces spp. integration vector
368   derived from the S. venezuelae phage, SV1. *BMC biotechnology* **14**, 51 (2014).

369   39.   Morita, K. et al. In vitro characterization of the site-specific recombination system based on
370   actinophage TG1 integrase. *Molecular genetics and genomics* **282**, 607 (2009).

371   40.   Christiansen, B., Johnsen, M., Stenby, E., Vogensen, F. & Hammer, K. Characterization of the
372   lactococcal temperate phage TP901-1 and its site-specific integration. *Journal of bacteriology*
373   **176**, 1069-1076 (1994).

374   41.   Li, X., Zhou, X. & Deng, Z. Vector systems allowing efficient autonomous or integrative gene
375   cloning in Micromonospora sp. strain 40027. *Applied and environmental microbiology* **69**,
376   3144-3151 (2003).

377   42.   Marcone, G.L., Carrano, L., Marinelli, F. & Beltrametti, F. Protoplast preparation and reversion
378   to the normal filamentous growth in antibiotic-producing uncommon actinomycetes. *The*
379   *Journal of antibiotics* **63**, 83-88 (2010).

380   43.   Wolf, T. et al. Targeted genome editing in the rare actinomycete Actinoplanes sp. SE50/110 by
381   using the CRISPR/Cas9 System. *Journal of biotechnology* **231**, 122-128 (2016).

382   44.   Wagner, N., Oßwald, C., Biener, R. & Schwartz, D. Comparative analysis of transcriptional
383   activities of heterologous promoters in the rare actinomycete Actinoplanes friuliensis. *Journal*
384   *of biotechnology* **142**, 200-204 (2009).

385   45.   Anzai, Y. et al. Production of rosamicin derivatives in Micromonospora rosaria by introduction
386   of d-mycinose biosynthetic gene with ΦC31-derived integration vector pSET152. *Journal of*
387   *industrial microbiology & biotechnology* **36**, 1013 (2009).

388   46.   Kim, D.-Y., Hwang, Y.-I. & Choi, S.-U. Cloning of metK from Actinoplanes teichomyceticus

389   ATCC31121 and effect of its high expression on antibiotic production. *Journal of microbiology*
390   *and biotechnology* **21**, 1294-1298 (2011).

391   47.   Shen, Y., Huang, H., Zhu, L., Luo, M. & Chen, D. Type II thioesterase gene (ECO-orf27) from
392   Amycolatopsis orientalis influences production of the polyketide antibiotic, ECO-0501 (LW01).
393   *Biotechnology letters* **34**, 2087-2091 (2012).

394   48.   Lee, K. et al. Increased vancomycin production by overexpression of MbtH-like protein in

395   Amycolatopsis orientalis KFCC10990P. *Letters in applied microbiology* **63**, 222-228 (2016).

396   49.   Mandali, S., Dhar, G., Avliyakulov, N.K., Haykinson, M.J. & Johnson, R.C. The site-specific
397   integration reaction of Listeria phage A118 integrase, a serine recombinase. *Mobile DNA* **4**,
398   2-2 (2013).

399   50.   Yoon, B., Kim, I., Nam, J.-A., Chang, H.-I. & Ha, C.H. In vivo and in vitro characterization of
400   site-specific recombination of a novel serine integrase from the temperate phage EFC-1.
401   *Biochemical and biophysical research communications* **473**, 336-341 (2016).

402   51.   Yang, L. et al. Permanent genetic memory with> 1-byte capacity. *Nature methods* **11**,
403   1261-1266 (2014).

404   52.   Talà, A. et al. Pirin: A novel redox-sensitive modulator of primary and secondary metabolism
405   in Streptomyces. *Metabolic engineering* **48**, 254-268 (2018).

406   53.   Bekker, V., Dodd, A., Brady, D. & Rumbold, K. Tools for metabolic engineering in Streptomyces.
407   *Bioengineered* **5**, 293-299 (2014).

408   54.   Cadoret, F., Soscia, C. & Voulhoux, R. Gene transfer: transformation/electroporation.
409   *Pseudomonas Methods and Protocols*, 11-15 (2014).

410   55.   Isaacs, F.J. et al. Precise manipulation of chromosomes in vivo enables genome-wide codon
411   replacement. *Science* **333**, 348-353 (2011).

412   56.   Dominguez, W. & O'Sullivan, D.J. Developing an efficient and reproducible conjugation-based
413   gene transfer system for bifidobacteria. *Microbiology* **159**, 328-338 (2013).

Figures

Figures

Figure 1. Plasmids used in this study. The primer binding sites are indicated.

Figure 2. Alignment of R4 and TG1 *attB* sites in *A. marina* DSM45569 and other

Amycolatopsis species. GenBank accession nos. of DNA sequences: *A. alba*

(NMQU01000019.1), *A. antarctica* (NKYE01000021.1), *A. azurea* (MUXN01000005.1),

A. kentuckyensis (MUMI01000226.1), *A. lurida* (FNTA01000004.1), *A. nigrescens*

(ARVW01000001.1), *A. niigatensis* (PJMY01000003.1), *A. orientalis*

(ASXH01000007.1), *A. pretoriensis* (MUMK01000092.1), *A. regifaucium*

(LQCI01000034.1), *A. rubida* (FOWC01000001.1), *A. sacchari* (FORP01000010.1), *A.*

thailandensis (NMQT01000219.1), *A. thermoflava* (AXBH01000004.1), *A.*

tolypomycina (FNSO01000004.1), *A. xylanica* (FNON01000002.1), *Streptomyces*

avermitilis (NC_003155.5) and *S. parvulus* (CP015866.1).

Figure 3. (A) Design and synthesis of DNA substrate *attB*. The *attB* sites were fused at

the end of forward primers, to amplify a sequence flanked with attB from a

unrelevant DNA template, pHG1 in this case. (B) Design and synthesis of DNA

substrate attP. The attP sites were amplified directly from integrating vectors carrying

TG1/attP (pHG4) or R4/attP (pJH1R4). (C) Recombination substrates and their

expected products. (D) *In vitro* recombination between DNA fragments containing

TG1 *attB^{Am}* (1627 bp) and TG1 *attP* (2471 bp; left) and R4 *attB^{Am}* (1854 bp) and R4

attP (990 bp; right). The expected product of the TG1 integrase mediated reaction

was a 4.1 kb DNA fragment containing the *attR^{Am}* site and a 53 bp fragment

containing *attL^{Am}* (not observed). For the R4 integrase recombination the expected

product was a 2.8 kb containing *attR^{Am}* fragment and an 51 bp *attL^{Am}* fragment (not

observed). (E) *In vitro* recombination between DNA fragments containing TG1 *attB^{Sa}*

(1035 bp) and TG1 *attP* (2471 bp; left) and R4 *attB^{Sp}* (1855) and R4 *attP* (990 bp;

right). The expected product fragments were 3.5 kb containing *attR^{Sa}* for the TG1

reaction and 2.8 kb containing *attR^{Sp}* for the R4 reaction. M: Fast DNA Ladder (NEB,

USA).

Figure 4. PCR confirmation of site-specific integration in the exconjugants. (A) Design

of primers. (B) PCR (using primers TG1-attL-Am-for/rev) of the expected TG1 *attL*

containing fragment from *A. marina* DSM45569:pHG4. (C) PCR (using primers

R4-attL-Am-for/rev) of the expected R4 *attL* containing fragment from *A. marina*

DSM45569:pJH1R4. M: Fast DNA Ladder. Colonies 1 to 6 are independent

exconjugants.

Figure 5. The insertion sites of TG1 and R4 integration plasmids in *A. marina*

DSM45569. Sequencing (using primers TG1-attL-Am-for or R4-attL-Am-for) of PCR

products containing *attL* from exconjugants validated the site-specific recombination

of the TG1 and R4 *attB* sites in *A. marina* DSM45569 after introduction of pHG4 or

pJH1R4, respectively.

Figure 1

Figure 2.

A. R4 *attB* sites

A marina   GGTTGCCCATCACCATGCCGAAGCAGTGATAGAAGGGAACCGGGATGCAG
A kentuckyensis   GGTTGCCCATCACCATGCCGAAGCAGTGGTAGAAGGGCACCGGGATGCAG
A nigrescens   GGTTGCCCATCACCATGCCGAAGCAGTGGTAGAAGGGCACCGGGATGCAG
A niigatensis   GGTTGCCCATCACCATGCCGAAGCAGTGGTAGAAGGGCACCGGGATGCAG
A orientalis   GGTTGCCCATCACCATGCCGAAGCAGTGGTAGAAGGGCACCGGGATGCAG
A pretoriensis   GGTTGCCCATCACCATGCCGAAGCAGTGGTAGAAGGGCACCGGGATGCAG
A rubida   GGTTGCCCATCACCATGCCGAAGCAGTGGTAGAAGGGCACCGGGATGCAG
A tolypomycina   GGTTGCCCATCACCATGCCGAAGCAGTGGTAGAAGGGCACCGGGATGCAG
A xylanica   GGTTGCCCATCACCATGCCGAAGCAGTGATAGAAGGGCACCGGGATGCAG
S parvulus   AGTTGCCCATGACCATGCCGAAGCAGTGGTAGAAGGGCACCGGCAGACAC

B. TG1 *attB* sites

A marina   TCGATCTCCAGTGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A alba   TCGATCTCCAGCGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A antarctica   AGCATCTCCAGTGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A azurea   TCGATCTCCAGCGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A lurida   TCGATCTCCAGCGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A regifaucium   TCGATCTCCAGCGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A sacchari   TCGATCTCCAGCGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A thaitandensis   TCGATCTCCAGCGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
A thermoflava   TCGATCTCCAGCGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGG
S avermitilis   TCGATCAGCTCCGCGGGCAAGACCTTCTCCTTCACGGGGTGGAAGGTCGG

Figure 3.

463 **Figure 4.**

Figure 5.

A

```
      attB - A. marina GGTTGCCCATCACCATGCCGAAGCAGTG------------------------
attL - exconjugant GGTTGCCCATCACCATGCCGAAGCAGTGGTACTGCTTGTGGGTACACTCT
      attP - pJH1R4 -------------------GAAGCAGTGGTACTGCTTGTGGGTACACTCT
```

B

```
      attB - A. marina GATCTCCAGTGCGGGCAAGACGTT--------------------------
attL - exconjugant GATCTCCAGTGCGGGCAAGACGTTGCTCTTACCCAGTTGGGCGGGATAGC
      attP - pHG4 ----------------------TTGCTCTTACCCAGTTGGGCGGGATAGC
```

 TABLE 1 Oligonucleotides used in this study

Oligonucleotide	Sequence (5'-3')
pHG1A-for	CGAACGCATCGATTAATTAAGGAGGATCGTATGACGACCGTTCCCG
pHG1A-rev	CGTGGTGGGCGCTAGCCTCCTCTAGTCATCCGTCG
pHG1-for	ACTAGAGGAGGCTAGCTTCAATGGAGGAGATGATCGAGG
pHG1-rev	GCAGGTCGACTCTAGATCTCGCTACGCCGCTACG
pHG4-for	CGAACGCATCGATTAATTAAGCGGCCGCCATATGGAATTCGGTACCGCATGCAGATCTAGGAACTTCGAAGTTCCCGC
pHG4-rev	TGATTACGCCAAGCTTTCGACTCTAGAGTAAGCGTCACGG
pJH1R4-for	CTAGCGATTGCCATGACGTCGGAGCTGCTTACCAATGTC
pJH1R4-rev	AAGAGGCCCGCACCGATTCCAAGAGGCCGGCAACTAC
TG1-attB-Am-for	<u>TCGATCTCCAGTGCGGGCAAGACGTTCAACTGCACCGGCTGGAAGATCGGGACCA</u>CCGGACGAACGCA
TG1-attB-Sa-for	TCGATCAGCTCCGCGGGCAAGACCTTCTCCTTCACGGGGTGGAAGGTCGGCGGTGGAGCTCGGAGA
R4-attB-Am-for	<u>GGTTGCCCATCACCATGCCGAAGCAGTGATAGAAGGGAACCGGGATGCAGGTGA</u>GAAGGTGCTCGTGT
R4-attB-Sp-for	AGTTGCCCATGACCATGCCGAAGCAGTGGTAGAAGGGCACCGGCAGACACGG
attB-rev	CTGCATCTCAACGCCTTCCGG
TG1-attP-for	AACCTTCACGCTCATGCC
TG1-attP-rev	GTCGAGATTCTCCGTCTCCTG
R4-attP-for	GATCGGTCTTGCCTTGCTC
R4-attP-rev	ACCCGCAGAGTGTACCCA
TG1-attL-Am-for	ACAACCCCACCGGCACCGTCTTCA
TG1-attL-Am-rev	AGTATAGGAACTTCGAAGCAGCTC
R4-attL-Am-for	CGGCCGGTGATGTTGACGT
R4-attL-Am-rev	TCGGCCGTCACGATGGTCA

469 The possible *attB* sequences identified from the genome sequence of *Amycolatopsis*

471 *marina* DSM45569 are shown underlined.

473   **TABLE 2**. The original *attB* sites for integrases and results of BLAST search

Integrase	*attB* sites	Reference	Best Hit from BLAST	
			Homology (%)	*E*-value
φ31	CGGTGCGGGTGCCAGGGCGTGCCCTTGG GCTCCCCGGGCGCGTACTCCACC	33	41	0.015
φJoe	ATCTGGATGTGGGTGTCCATCTGCGGGCA GACGCCGCAGTCGAAGCACGG	34	30	0.60
Bxb1	TCGGCCGGCTTGTCGACGACGGCGGTCTC CGTCGTCAGGATCATCCGGGC	35	48	0.014
R4	AGTTGCCCATGACCATGCCGAAGCAGTGG TAGAAGGGCACCGGCAGACAC	31	84	3e-11
SPBc	AGTGCAGCATGTCATTAATATCAGTACAGAT AAAGCTGTATCTCCTGTGAACACAATGGGT GCCA	36	No hits	
SV1	ATCAGGGCGGTCAGGCCGTAGATGTGGAA GAACGGCAGCACGGCGAGGAC	37	32	0.17
TG1	TCGATCAGCTCCGCGGGCAAGACCTTCTC CTTCACGGGGTGGAAGGTCGG	38	62	0.001
TP901	ATGCCAACACAATTAACATCTCAATCAAGG TAAATGCTTTTTGCTTTTTTTGC	39	25	8.1

476   **TABLE 3**. Conjugation efficiency of pHG4 and pJH1R4 in different species

Exconjugants/10^8 spores	pHG4	pJH1R4
A. marina	160	20
S. coelicolor	1.47×10^3	3.28×10^4
S. lividans	1.56×10^3	3.33×10^4