A role for VPS35 in *Brucella* intracellular replication revealed by genome-

wide siRNA screening

Alain Casanova[a,c], Shyan Huey Low[a,d], Maxime Québatte[a], Jaroslaw Sedzicki[b,e], Therese Tschon[a],

Kevin Smith[a,f], Mario Emmenlauer[a,g], Houchaima Ben-Tekaya[a,h], and Christoph Dehio[a,#]

[a]Focal Area Infection Biology, Biozentrum, University of Basel, Switzerland

[b]C-CINA, Biozentrum, University of Basel, Switzerland

Running Head: VPS35 is required for *Brucella* trafficking

#Address correspondence to Christoph Dehio, christoph.dehio@unibas.ch

[c]Present address: Novartis AG, Basel, Switzerland

[d]Present address: Lee Kong Chian School of Medicine, Nanyang Technological University

[e]Present address: Focal Area Infection Biology, Biozentrum, University of Basel, Switzerland

[f]Present address: Science for Life Laboratory, KTH Royal Institute of Technology, School of Computer

Science and Communication, Solna, Sweden

[g]Present address: BioDataAnalysis GmbH, Munich, Germany

[h]Present address: University Hospital Basel, Basel, Switzerland

A.C, S.L., and M.Q. contributed equally to this work

Word count for the abstract: 235

Word count for the text: 6940

Key words: *Brucella*, siRNA, retromer, VPS35, VPS26, RAB7A, replicative niche

Abstract

Brucella, the causing agent of brucellosis, is a major zoonotic pathogen with worldwide distribution.

Brucella resides and replicates inside infected host cells in membrane-bound compartments called

BCVs (*Brucella*-containing vacuoles). Following uptake, *Brucella* resides in eBCVs (endosomal

BCVs) that gradually mature from early to late endosomal features. Through a poorly understood

process that is key to the intracellular lifestyle of *Brucella,* the eBCV escapes fusion with lysosomes by

transitioning to the rBCV (replicative BCV), a replicative niche directly connected to the endoplasmic

reticulum (ER). Despite the notion that this complex intracellular lifestyle must depend on a multitude

of host factors, a holistic view on which of these components control *Brucella* cell entry, trafficking

and replication is still missing. Here we used a systematic cell-based siRNA knockdown screen in HeLa

cells infected with *Brucella abortus* and identified 426 components of the human infectome for *Brucella*

infection. These include multiple components of pathways involved in central processes such as cell

cycle, actin cytoskeleton dynamics or vesicular trafficking. Using assays for pathogen entry,

knockdown complementation and co-localization at single-cell resolution, we identified the

requirement of VPS35 for *Brucella* to escape the lysosomal degradative pathway and to establish its

intracellular replicative niche. We thus validated a component of the retromer as novel host factor

critical for *Brucella* intracellular trafficking. Further, our genome-wide data shed light on the interplay

between central host processes and the biogenesis of the *Brucella* replicative niche.

Importance

With about 500'000 new cases of human brucellosis annually, *Brucella* is regarded as the most important zoonotic bacterial pathogen worldwide. The causing agent of brucellosis resides inside host cells within vacuoles termed *Brucella* containing vacuoles (BCVs). Although few host components required to escape the degradative lysosomal pathway and to establish the ER-derived replicative BCV (rBCV) have already been identified, the global understanding of this highly coordinated process is still partial and many factors remain unknown. To gain a deeper insight into these fundamental questions we performed a genome-wide RNA interference (RNAi) screen aiming at discovering novel host factors involved in *Brucella* intracellular cycle. We identified 426 host proteins that contribute to *Brucella* cellular entry, intracellular trafficking, and replication. Together, this study sheds light on previously unknown host pathways required for *Brucella* infection cycle and highlights the retromer component VPS35 as a critical factor for the establishment of *Brucella* replicative niche.

Introduction

Cellular invasion is a common strategy shared by many bacterial pathogens of human and animals (i.e. *Salmonella*, *Shigella*, *Legionella*, or *Brucella*) in order to escape host defenses and establish a protected replicative niche (1-3). Knowledge of the host cellular pathways that are subverted by these pathogenic bacteria in order to reach and/or establish their intracellular replicative niches can be highly instructive for the development of new treatment strategies. *Brucella* is a facultative intracellular zoonotic pathogen causing animal and human brucellosis. With about 500'000 new cases of human brucellosis every year, *Brucella* is regarded as the most important zoonotic bacterial pathogen worldwide (4, 5). There is currently no effective vaccination for humans and even prolonged combinatory antibiotic treatments do not fully protect against relapses (6). Therefore, *Brucella* remains a significant threat to public health and to the economy in endemic areas, and thus new treatment strategies to circumvent *Brucella* infections are highly needed.

At the cellular level, *Brucella* invades both phagocytic and non-phagocytic cells where bacteria persist and replicate inside membrane-bound compartments – the *Brucella* containing vacuoles (BCVs). BCVs sequentially interact with components of the host early and late endocytic pathway (eBCVs) then transit to establish the replicative niche (rBCVs) in vesicles that are directly connected to the endoplasmic reticulum (ER) (7) and harbor ER-specific markers, such as Sec61 and calnexin (7-9). Several critical steps for the intracellular journey of *Brucella* and their associated host factors have been resolved. For instance, adherence to the host cell surface is mediated via interaction with sialic acid residues or binding to fibronectin and vitronectin (10, 11). Internalization requires actin remodeling via the activity of the small GTPases Rac, Rho and direct activation of Cdc42 (12). Upon internalization, bacteria are contained within eBCVs that successively associates with a subset of endosomal markers, starting with RAB5, the early endosomal antigen (EEA1), the transferrin receptor (TfR), as well as the lipid rafts component flotillin-1 (8, 13-15). Next, the eBCV associates with the late endosomal markers RAB7, RILP (RAB7's effector RAB interacting lysosomal protein), LAMP-1 (Lysosomal-associated membrane protein 1), and transiently with autophagosomal markers (8, 9). As it evolves late endosomal characteristics, acidification of the eBCV serves as a trigger for the expression of the VirB type IV secretion system (T4SS, (16, 17)). This major pathogenicity factor is required to prevent the complete

fusion of eBCVs with lysosomes, consequently allowing a fraction of the internalized *Brucella* to avoid

host-mediated degradation, and promotes the maturation of the eBCV towards rBCV (15, 17).

Noteworthy, the T4SS effectors responsible for this escape remain largely elusive, despite a growing

repertoire of identified candidates (recently reviewed in (18)). Most recently, it was discovered that

subversion of both anterograde and retrograde transport and recruitment of Conserved Oligomeric Golgi

(COG) tethering complex-dependent vesicles to the BCV promotes the establishment of the *Brucella*

replicative niche (19). Importantly, despite all these findings, the precise mechanism(s) resulting in

diversion of eBCVs from the endolysosomal pathway towards the ER-associated replicative

compartment (rBCV) is still largely unresolved. The same holds true for host factors required for

maintenance of the replicative niche.

In this study, we took a systems-level approach to gain a deeper insight into these fundamental

questions. Using a genome-wide RNA interference (RNAi) screening approach, we identified 426 host

proteins whose knockdown either increases (202) or decreases (224) *Brucella* intracellular replication.

Beside the rediscovery of several previously identified host targets, that validates our approach, data

reveals numerous novel candidate components that can modulate *Brucella* cellular entry, trafficking,

and/or replication. Among these targets, we identified VPS35 and VPS26A, two components of the

trimeric retromer, which are required for the diversion of BCVs from the endolysosomal pathway and

the establishment of the intracellular replicative niche.

Results

A genome-wide siRNA screen reveals novel host pathways involved in *Brucella* infection

To identify novel host factors important for *Brucella* intracellular infection, we performed a genome-

wide siRNA perturbation screen on the human epithelial cell line HeLa (ATCC© CCL-2) combined

with bacterial infection at biosafety-level 3. Infections of the siRNA-treated cells were performed with

a GFP expressing strain of *B. abortus* and the outcome was analyzed after 48 h infection using

automated fluorescence microscopy (20, 21). Infection rates were determined with a tailored high-

content analysis workflow (Figure 1 and Materials and Methods). In brief, a model of *Brucella*

replication was fitted to the pathogen intensity distribution to gain an infection classification

independent of absolute fluorescence intensity. Further, we implemented an image intensity

normalization coupled with a novel pathogen-to-cell association approach, which enabled quantitative

measurement of the pathogen intensity distribution (Materials and Methods). To account for the well-

known confounding off-target effects associated with siRNA technology (recently reviewed in (22)),

we applied a multiple orthologous RNAi reagents approach (MORR (23)) with $n \geq 5$ perturbations per

host genes. Further, we applied the Redundant siRNA Analysis (RSA) algorithm (24) on the entire

screening data to reduce the number of false positives caused by off-target effects of single siRNAs and

to favors genes with a reproducible phenotype confirmed by independent siRNAs. As a result, we

identified 426 hits significantly affecting *Brucella* infection (224 down-hits, Table S1, and 202 up-hits,

Table S2, RSA p-value ≤ 0.01). A panel of representative images from the screen is presented in Figure

2A. The hit genes were further stratified by gene-annotation enrichment analysis and functional

annotation clustering using DAVID (25), protein-protein interaction network using the STRING

database (26), as well as manual datamining. The functional categories enriched in our hit lists are

presented in Figure 2B-D together with the high confidence protein-protein interaction network for

targets that reduced (Fig. 2E) or increased (Fig. 2F) *Brucella* infection upon knockdown. Gene ontology

and functional clustering analysis indicated a rather small overlap in enriched pathways when

considering up or down hits (Fig. 2B-D). The most prominent clusters that positively affected infection

upon knockdown comprised components involved in the control or the modulation of central cellular

processes such as protein synthesis, transcription and mRNA processing, and cell cycle progression, as

well as clathrin-mediated endocytosis (Fig. 2 and Table S2). The most prominent clusters that

negatively affected infection upon knockdown comprised signaling pathways involved in actin-

remodeling and phagocytosis. These included most core components of the Actin-related protein-2/3

complex (Arp2/3: ArpC2, ArpC3, ACTR2 and ACTR3), and one of its main modulator, the WASP

regulatory complex (WRC: CYFIP1, WASF3, NCKAP1, and ABI3). Down-hits also comprised

multiple components involved in TGF-β and Eph signaling as well as further vesicular/endocytic

pathways (Fig. 2 and Table S1). Among all these factors we can highlight the Ras related protein

RAB7A, which is needed for *Brucella* trafficking to the replicative niche (9), the small GTPases Rac1

and Cdc42, which are involved in *Brucella* internalization into non-phagocytic cells (12) as well as the

transmembrane glycoprotein SLC3A2 (CD98hc), involved in both bacterial uptake and intracellular

multiplication (27). Since the role of these individual components has already been described in the

context of *Brucella* infection, they can be considered as benchmark to our results, and globally validate

our systems-level perspective of the human infectome for *Brucella* infection.

Pathogen entry assay identifies a role for VPS35, VPS26A, Sec61γ and ARMCX3 in *Brucella* post-

entry trafficking

To further dissect the role of the identified genes in the progression of *Brucella* infection, we took

advantage of a pathogen entry assay previously developed in our laboratory (28). Briefly, after 4 h of

infection, extracellular *Brucella* were killed by gentamicin concomitantly to the induction of a plasmid-

encoded reporter by a cell permeable inducer for another 4 h. This approach allowed us to robustly

identify individual intracellular bacteria and quantify the bacterial load before intracellular replication

is initiated (Fig. 3A and Material and Methods). For this assay, we selected a number of genes from the

different pathways identified in the genome-wide screen as well as additional genes supplementing

them (Table S3). The results of this entry assay were plotted against a matching endpoint assay

(intracellular replication at 48 hpi). Strikingly, most of the tested genes displayed a direct correlation

between the results of the entry and the endpoint assay (r^2 =0.763). This was the case for components

involved in the actin-remodeling pathway (Cdc42, Rac1, ACTR3, CYFIP1) or those involved in the

TGFβ signaling (SMAD4, TGFBR1, TGFBR2), which strongly reduced both entry and subsequent

intracellular replication (Fig. 3B and Table S3). Similarly, the components of the clathrin pathway GAK

and AP2S1 both increased bacterial entry and replication (Fig. 3B and Tables S3-4). These results thus

indicate the involvement of these pathways in *Brucella* entry into non-phagocytic cells, without

excluding an additional role at any further stage of the infection.

In order to identify genes predominantly involved in post-entry processes, we focused on those that

mainly affected intracellular replication (>50% reduction in the endpoint assay) while having little

effect on bacterial entry (< 50% in the entry assay). From our dataset, only 4 genes matched this criterion

(Fig 3B, orange box). Those are the vacuolar protein sorting associated proteins VPS35 and VPS26A -

two essential components of the retromer complex (recently reviewed in (29, 30)), Sec61γ - a central

element of the ER-protein translocation machinery (see for instance (31)) and ARMCX3, encoding

Alex3, a protein involved in mitochondria dynamics in neural cells (32). These genes and associated

pathway(s) thus likely represent novel components controlling the post-entry trafficking of *Brucella*

towards its replicative niche and/or are themselves required for the establishment or maintenance of

the replicative niche. For the present study, we further focused on the role of VPS35 in *Brucella*

trafficking as it was the most prominent hit in our entry assay.

The VPS retromer is required for *Brucella* intracellular replication

The retromer complex orchestrates the recycling of numerous transmembrane proteins from early and

maturing endosomes either to the trans-Golgi network (TGN) or back to the plasma membrane. Formed

by a heterotrimeric complex consisting of VPS26, VPS29, and VPS35, the retromer is conserved from

yeast to human. However, the individual retromer sub-complexes have functionally diverged to

organize multiple distinct sorting pathways, depending on the association with different accessory

factors (29, 30). To further decipher the role of the retromer in *Brucella* trafficking and intracellular

replication we specifically browsed our genome-wide siRNA data for retromer-associated proteins (Fig.

4 A and B). Further to VPS35 and VPS26A, already identified both in the genome wide screen and in

the entry screen (Fig. 2E and 3B), knockdown of VPS26B (the paralogue of VPS26A) resulted in a

significant reduction in *Brucella* infection (Fig. 4B). Depletion of VPS29, the third core component of

the retromer, resulted only in a mild reduction of *Brucella* infection and did not reach significance due

to the wide spread of data obtained for the cohort of 9 individual siRNAs tested (suggestive of strong

off-target effects). Next to the retromer component, knockdown of the small GTPase RAB7A showed

the strongest reduction in intracellular *Brucella*(Fig 4B). SNX3, that together with the trimeric retromer

forms the SNX3-retromer or SNX27, another retromer-associated component involved in endosome-

to-plasma membrane trafficking (33, 34), did not show a significant effect (Fig. 4B). Depletion of SNX1

and SNX5, two of the four sorting nexins of the SNX-BAR retromer (35), seem to enhance *Brucella*

infection (although not significantly) while the two others, SNX2 and SNX6, showed no effect (Fig.

4B). Noteworthy, the functional association of the SNX-BAR sorting nexins with the VPS retromer has

been challenged by two recent publications which rather support a VPS-independent action of SNX-

BAR (36, 37).

To validate the requirement of VPS35 on *Brucella* infection and to rule out any off-target effect, we

performed a complementation experiment using a VPS35 cDNA insensitive to a co-expressed shRNA

(38). While shRNA knockdown of endogenous VPS35 inhibited *Brucella* infection, as detected in our

genome-wide approach, ectopic expression of the shRNA-insensitive cDNA of VPS35 rescued the

phenotype (Fig. 4C and D), confirming that depletion of VPS35 indeed negatively affects *Brucella*

infection. Together, these results corroborate the importance of the retromer for *Brucella* to reach and

maintain its replicative niche. Beyond VPS26A and RAB7A, the identity of the VPS-associated proteins

required to support this function, if any, remains to be elucidated.

VPS35 knockdown prevents *Brucella* escape from the lysosomal pathway

Transient association with the lysosomal marker LAMP-1 is a well-established signature of BCV

trafficking during the first hours of infection. This association is eventually lost for those bacteria that

manage to escape the host degradative pathway. Thus, to investigate the role of VPS35 in *Brucella*

trafficking and to assess at which stage it could be required for the establishment of the intracellular

replicative niche, we quantified *Brucella* co-localization with LAMP-1 in siRNA-treated and control

cells. To this end, we analyzed *Brucella*-infected HeLa cells at 6 and 18 hpi and determined the

percentage of LAMP-1 co-localization for each detected bacteria, combining immunostaining and

confocal microscopy (Fig. 5 and Materials and Methods). At 6 hpi, most *Brucella* were found within

LAMP-1-positive vesicles in both control and siRNA-treated cells (Fig. 5A-D, F), indicating that

VPS35 function is not required for the early trafficking of the BCVs. However, loss of LAMP-1

association at 18 hpi was mainly detected in control cells whereas most *Brucella* remained in a LAMP-

1 positive compartment upon VPS35 knockdown (Fig 5B, C). Accordingly, VPS35 depletion strongly

reduced intracellular replication of *Brucella* 18 hpi (Fig. 5G) compared to control cells (Fig. 5E).

Altogether, our single cell co-localization analysis demonstrates the requirement of VPS35, and thus of

a functional retromer, for the diversion of BCVs from the lysosomal pathway and for the subsequent

establishment of a successful replicative niche.

Discussion

The different membrane-bound organelles that compose the secretory pathway and the endo-lysosomal system of eukaryotic cells constitute targets of choice for many intracellular pathogens, which have evolved highly diverse strategies to hijack and/or subvert these trafficking pathways to their benefit (see for instance (2, 3)). It that context, the importance of retrograde trafficking for the infection cycle of a number of human pathogens (viruses and bacteria) has been recognized in the past years (e.g. (39, 40)). Particularly, some pathogens have been shown to specifically target the trimeric vacuolar protein sorting (VPS) complex (termed retromer hereafter), using or subverting its function to their advantage. For instance, the Hepatitis C virus interacts with VPS35 through its protein NS5A. This viral protein is recognized as retromer cargo and its interaction with VPS35 supports viral replication (41). Among bacterial pathogens, the best-studied example to date is the subversion of retromer function by the T4SS effector RidL of *Legionella pneumophila.* RidL was shown to interact with VPS29, inhibiting retromer activity by competing out the binding of the retromer regulator TBC1d5 and thereby promoting *Legionella* intracellular replication (42-44). In this study, we report the involvement of the retromer in *Brucella* intracellular cycle. More specifically, we show that the retromer integrity is required for *Brucella* to escape the host degradative pathway, as supported by the inability of the eBCV to mature upon VPS35 knockdown towards a LAMP-1 negative rBCV. We further show that this VPS35-dependent process takes place after internalization and early trafficking, between 6 h and 18 h post-infection, matching the estimated timing of the transition from eBCV to rBCV. It is thus tempting to speculate that VPS35, together with VPS26A and possibly RAB7A, plays a role in this essential (yet elusive) branching point of *Brucella* intracellular trafficking from the lysosomal pathway towards the ER-associated replicative niche. An alternative, yet non-exclusive hypothesis is that the retromer is involved in the establishment or maintenance of the rBCV, possibly by providing host factors and/or membranes that follow retrograde trafficking. With the accumulation of functional data and the increasing number of described interactors, the retromer is nowadays largely appreciated as recruiting hubs that orchestrates the retrograde endosomal trafficking of numerous cargos to the TGN or the plasma membrane (29, 30, 45). This versatility however obscures the identification of the underlying

mechanism(s) by which VPS35 and the retromer may contribute to *Brucella* intracellular fate. Further

browsing our dataset for the effect of known retromer interactors failed resolving the retromer-

dependent pathway(s) – if any – that is required for *Brucella* intracellular replication. The only retromer

interactor that we unambiguously identified is the small GTPase RAB7A, which is essential for the

recruitment of the retromer to endosomal membranes (46, 47). Importantly, association of this upstream

interactor to the eBCV is a well-established hallmark of early *Brucella* intracellular trafficking (8, 9,

15). Moreover, over-expression of a RAB7 dominant negative allele (RAB7^{T22N}) impairs the

establishment of the *Brucella* replicative niche (9). This finding that was recapitulated by our siRNA

knockdown approach, strengthening the role of RAB7 in controlling BCVs' fate, albeit by an unknown

mechanism. Considering that the recruitment of the retromer to endosomal membranes is strictly

dependent on the presence of RAB7, it is conceivable to assume that depletion of RAB7 prevents the

recruitment of VPS35 to the BCV, consequently explaining the RAB7-dependency observed for

Brucella replication. Alternatively, very recent findings have established that retromer depletion in Hela

cells actually results in the hyper-activation of RAB7, which causes an overall depletion of the RAB7

pool on endo-membranes (48). That drastic consequence could also imply an indirect effect of the

observed retromer requirement for *Brucella* trafficking, by acting at the level of RAB7 activity and its

availability for the BCV maturation. However, our results indicate that the effect of RAB7A or VPS35

siRNA knockdown are not entirely congruent. Whereas depletion of either factor impairs *Brucella*

intracellular replication, only RAB7A knockdown showed a marked effect on pathogen entry whereas

VPS35 appears to be only required at a later stage of the infection. The implications of the newly

described feedback signaling on RAB7 triggered by the retromer depletion, as well as the relative

contribution of the VPS-retromer and RAB7 for the transition of the eBCVs to rBCVs should be

addressed in future studies.

Besides the retromer complex, our study pinpointed the involvement of several host pathways in

Brucella infection, which had not yet been associated with this process. The most prominent cluster

negatively affecting infection upon knockdown comprises factors involved in actin remodeling and

actin dynamics as well as associated signaling pathways. Apart from an early association of Arp2/3

with BCVs (49), surprisingly little is known about the exact role played by the Arp2/3 complex or the

WASP regulatory complex and their regulators during and possibly after *Brucella* internalization. Here

as well, further studies will be needed to decipher the precise nature of their involvement. Finally, we

also found that members of the TGF-β and FGF signaling pathways promote *Brucella* infection as their

depletion resulted in decreased *Brucella* infection. Interestingly, it has previously been reported that

patients with brucellosis show higher TGF-β1 serum levels, a finding that is correlated with depressed

T cell function (50). Further, B cells were also shown to produce TGF-β at early stages of infection with

Brucella in mice (51). A possible immunosuppressive role for this pathway during *Brucella* infection

should be further investigated.

Summing-up, we believe that the genes and pathways identified in this study constitute a rich resource

towards the understanding of *Brucella* intracellular trafficking, which ultimately should allow

development of new approaches to controlling *Brucella* infections in human.

Material & Methods

Cell lines and plasmid constructs

All experiments were performed in the human cervical carcinoma epithelial cell line (Hela) ATCC©

CCL-2. Infections were performed using *Brucella abortus* 2308 carrying the constitutive GFP

expression plasmid pJC43 (*aphT::GFP* (52)), pAC042.08 for entry assay (*apht::dsRed,tetO::tetR-GFP*

(28)) or pAC037 (*apht::cerulean*, this study) for rescue experiments. Cells and bacteria were grown as

described in (21, 28). pAC037 was constructed by replacing dsRed from pJC44 (9) with Cerulean from

pCERC-1 (53). Cerulean was amplified using prAC082 (TGGATCCGAAAGGAGGTTTATTAAATGG-

TGAGCAAGGGCGAGGAGC) and prAC083 (TCTAGAGCTAGCTTACTTGTACAGCTCGTC) and cloned into

pJC44 by restriction/ligation using BamHI and XbaI. The ribosomal binding site which was lost on

pJC44 using the above restriction was re-introduced on prAC082.

siRNA reverse transfection

Reverse siRNA transfection was performed as described in (21, 28) with minor adjustments. In brief,

Genome-wide screens were performed with Dharmacon ON-TARGETplus SMART pool (pool of 4

siRNA per gene) and Qiagen Human Whole Genome siRNA Set HP GenomeWide (QU, 4 individual

siRNAs for each target). Further validation screens included Ambion Silencer and Ambion Silencer

Select custom libraries (with up to 6 additional siRNAs for about 1000 genes) and Sigma MISSION esiRNA libraries for 1900 genes. All screening experiments were conducted in a 384-well plate format. Each plate contained general siRNA controls for transfection efficiency and toxicity (e.g. Kif11) as well as positive controls (e.g. Cdc42, Rac1) that are known to have an effect on *Brucella* infection (12). In addition, negative controls such as mock (transfection reagent only) and scrambled (non-targeting siRNA) were added to each plate. The following specifications apply to all siRNA screens except the QU siRNA library where specifications are given in brackets. RNAiMAX in DMEM without fetal calf serum (FCS) was added to each well containing 1.6 pmol siRNA (QU: 1 pmol) or 15 ng esiRNA. Screening plates were then incubated at room temperature (RT) for 1 h. Following incubation, 500 HeLa cells were seeded per well in DMEM (FCS 10% final). Plates were incubated at 37°C and 5% CO_2 for 72 h prior to infection. For assays in 96- and 24-well formats reverse transfections were performed in 6-well plates and subsequently reseeded in the respective plate format. On-target or control siRNAs were added to reach a final siRNA concentration of 20 nM together with RNAiMAx transfection reagent in DMEM without FCS. After 30 min of complex formation at room temperature, 110'000 HeLa cells in DMEM/10% FCS were added to each well. After 48 h transfection, cells were harvested by trypsinization and reseeded in DMEM/10%FCS (96-well plates: 2'800 cells per well; 24-well plates: 50'000 cells per well). The next day cells were infected as described hereafter.

The following siRNAs used for the colocalization experiments were purchased from Qiagen (Hilden/Germany): hVPS35-2 (ID: SI00760690); all star negative (ID: 0001027281); all star death kif11 (ID: 0001027299).

Infection

For the genome-wide and confirmation screens, infections were performed in 384-well plates as described in (21, 28). In short, *B. abortus* 2308 pJC43 (*aphT::GFP* (52)) was grown in TSB medium containing 50 µg/ml kanamycin at 37°C to an OD of 0.8- 1.1. Bacteria were then diluted in DMEM/10% FCS and added at a final MOI of 10'000. Plates were centrifuged at 400 x *g* for 20 min at 4°C to synchronize bacterial entry. After 4 h incubation at 37°C and 5% CO_2, extracellular bacteria were killed by exchanging the infection medium by DMEM/10% FCS supplemented with 100 µg/ml gentamicin. After a total infection time of 44 h, cells were fixed with 3.7% PFA for 20 min at RT. For the entry

assay, infections were performed as described in (28). In brief, transfected cells were infected with *B. abortus* 2308 pAC042.08 for 4 h after what GFP expression was induced for 4 h by the addition of Anhydrotetracycline (100 ng/ml) during the gentamicin killing of extracellular bacteria. Follow-up experiments and colocalization assays were performed according to the above-described protocol in 96-well and 24-well plates, respectively. For the colocalization assay cells were infected at a MOI 2'000. 2 hpi cells were washed three times with DMEM/10% FCS containing gentamicin (100 μg/ml). After the indicated incubation time cells were washed three times with PBS and finally fixed for 20 min in 3.7% PFA (in PBS).

Imaging with high-throughput microscopy

Microscopy was performed with Molecular Devices ImageXpress microscopes. MetaXpress plate acquisition wizard with no gain, 12 bit dynamic range, 9 sites per well in a 3x3 grid was used with no spacing and no overlap and laser-based focusing. DAPI channel was used for imaging nucleus, GFP for bacteria, and RFP for F-actin or dsRed of bacteria in the entry assay. Robotic plate handling was used to load and unload plates (Thermo Scientific). The objective was a 10X S Fluor with 0.45NA. The Site Autofocus was set to "All Sites" and the initial well for finding the sample was set to "First well acquired". Z-Offset for Focus was selected manually and manual correction of the exposure time was applied to ensure a wide dynamic range with low overexposure. Images from the different siRNA screens are available upon request.

Image analysis

Images were analyzed with the screeningBee analysis framework from BioDataAnalysis GmbH. To correct for uneven illumination inherent in wide-field microscopic imaging, an illumination correction model was computed for every plate using Cidre (54). To ensure that the Cidre-corrected image intensities fall within the range [0.0, 1.0] a linear transformation for pixel intensities was computed that maps the 0.001-quantile to 0.01 and the 0.999-quantile to 0.99 post-illumination correction. Illumination correction and intensity scaling were performed as pre-processing steps for every image prior to analysis.

To reduce the signal of *Brucella* DNA in the DAPI channel, a linear transform of the GFP channel was subtracted from the DAPI channel, with the linear transformation parameters f, o estimated in the

following way: a mapping of GFP pixels to DAPI pixels was constructed so that for all intensities in

the GFP images, the list of corresponding intensities in the DAPI images were recorded. For every list

of DAPI intensities, only the mean intensities were retained. This creates a mapping of GFP intensities

to their corresponding mean DAPI intensities. A linear regression was performed to obtain the linear

parameters f, o that map the GFP channel image to the DAPI channel image. Cleaned DAPI images

with a reduced *Brucella* signal were obtained by subtracting the linear transform of the GFP channel

from the DAPI channel $I'_{DAPI} = I_{DAPI} - (f\ I_{GFP} + o)$ as pre-processing steps for every image prior to

analysis. On a random subset of 128 images, CellProfiler (55) was executed to identify Nucleus objects

using "OTSU Global" segmentation in the DAPI channel, and the median, lower quartile and upper

quartile segmentation thresholds of the images were retained as T_{DAPI-m}, $T_{DAPI-lq}$ and $T_{DAPI-uq}$. On the

same images, the GFP background intensity B_{GFP} was obtained as the position of the peak in the GFP

intensity histogram, the dynamic range of the histogram D_{GFP} was obtained as the difference between

the 99% quantile and the 1% quantile of intensity values, and the Bacteria segmentation threshold was

computed as $T_{GFP} = B_{GFP} + \frac{2}{100} D_{GFP}$. On all images, screeningBee CellProfiler was executed to perform

object segmentation and measurements with the following steps: (a) Nuclei were detected as primary

objects using manual threshold setting. For each plate it was manually chosen to use T_{DAPI-m}, $T_{DAPI-lq}$ or

$T_{DAPI-uq}$, depending on visual inspection of the segmentation results. Using the same threshold on all

images improved site-to-site comparability. (b) Cells were detected as secondary objects around the

Nuclei, with "OTSU Global" segmentation in the RFP channel. (c) Bacteria were detected as primary

objects using manual threshold setting with threshold T_{GFP}. Using the fixed background intensity as a

reference for T_{GFP} allowed for segmenting even rather dim objects while avoiding site-to-site variability.

In order to accurately measure infection scoring, a reliable method to associate pathogen colonies to

individual cells is necessary. A straightforward approach is to assume that pathogen colonies must be

contained within the body of the host cell. However, high cell confluence can make actin channel-based

cell body segmentation inaccurate. Single microcolonies are often split into pieces that are incorrectly

assigned to neighboring cells using this approach (Fig. 1B). To address this issue, we developed a novel

algorithm to intelligently assign pathogen colonies to robust nucleus objects (Fig. 1C). First,

inexpensive 'bridge' and 'majority' morphological operations were applied to the pathogen objects to

connect broken clumps. Next, a weighted distance metric was used to measure an attraction score $a_{N,P}$

between a pathogen P and individual nuclei N within a close proximity d_{prox}. The attraction score is

computed as the surface integral of the nucleus area in a continuous field emanating from the pathogen

defined by an exponential function that is strongest within the microcolony itself, and decays

exponentially as distance from the microcolony increases: $a_{N,P} = \sum_{n \in N} e^{-\lambda\, d_{n,p}}$, where n is an

element (pixel) belonging to nucleus object N, $d_{n,p}$ is the distance transform from the edge of

microcolony P to n, and λ is a parameter controlling the strength of the decay. Attraction scores for all

nuclei proximate to microcolony P are normalized such that the strongest nucleus attraction score is 1,

$\underline{a}_{N,P} = \frac{a_{N,P}}{max(a_{N,P})} \,\forall\, N \quad s.t.\, d_{n,p} < d_{prox}$. Nuclei objects with normalized attraction scores above a

threshold a_{min} are associated with the pathogen microcolony. In the case that multiple nuclei are

associated with the same microcolony, the microcolony is split so that each element is associated to the

nearest nuclei. Large microcolonies are encouraged to split with greater ease than small microcolonies

by weakening the minimum attraction score linearly according to area of the microcolony $a'_{min} =$

$0.5\, a_{min}$ if $A_P < A_{large}$, or $a'_{min} = 1 - 0.5\, \frac{A_P - A_{large}}{A_{large}} a_{min}$ if $A_{large} \leq A_P \leq 2A_{large}$, and $a'_{min} =$

$0.5\, a_{min}$ otherwise (where A_P is the area of the pathogen microcolony). Parameters settings $d_{prox} =$

45, $\lambda = 0.2$, $a_{min} = 0.5$, and $A_{large} = 8{,}000$ were optimized by grid search on a dataset of 7,566 hand-

labeled segmentations resulting in a 95.58% correct association rate.

Nucleus to pathogen microcolonies associations were aggregated. The area and integrated intensity of

the pathogen objects associated to each cell and the mean intensity of the Nuclei in the GFP channel

was computed as readout.

Infection scoring for endpoint assays

Wells that contain only 32 Cells or less were excluded from infection scoring. In the remaining wells,

Bacteria were filtered in a decision tree (DT) classification to exclude objects of only one-pixel area.

Based on the relation of Bacteria to Nuclei, for the remaining Bacteria objects, the integrated GFP

intensity was integrated over all Bacteria relating to a Cell. To reduce the impact of background

intensity, an estimate for GFP background was computed using the 1% lower quantile of mean GFP

intensity in the Nuclei. For every Cell, the estimated GFP background intensity was multiplied with the

area of Bacteria relating to this cell, and the result was subtracted from the integrated Bacteria GFP

intensity of the Cell, to arrive at a background-free estimate of "bacterial load" in each Cell. The value

range for this intensity was zero for Cells with no segmented Bacteria objects, and higher than zero for

all other Cells. This integrated GFP intensity was then log2-scaled, to reflect the exponential growth of

replicating *Brucella*. Before log2-scaling, a small epsilon value of 2^{-20} was added to every Cell, so that

the log2 value of Cells with no segmented Bacteria will not be negative infinity. The arbitrary value 2^{-20}

is by a large margin smaller than the smallest actual intensity of our assays, but large enough to be

used in histogram binning. For every plate, the histogram of the log2-scaled integrated cellular GFP

intensity was computed (Fig. 1D) with a bin size of 0.025. The histograms were normalized to an

arbitrary "virtual plate cell count" of 10^{10}. To extrapolate a continuous distribution from the possibly

sparse histogram, kernel density estimation (KDE) was used with a manually optimized Gaussian kernel

of standard deviation 16. The histogram shows a bimodal distribution. By correlating the plate

histogram distributions to selected images from the plate, we could identify that the first mode of the

distribution is composed of cells with a low number of infection events ranging from single *Brucella* to

small clusters (denoted as), whereas the second mode is composed of large colonies (denoted as). The

two peak positions of the bimodal distribution were identified. Normal distributions G_S and G_L were

fitted to the peak positions for small and large colonies, respectively. For the fitting of G_S and G_L, the

mean was given by the position of the peak, the height was given by the height of the peak, and the

variance was optimized such that the distance between the KDE and the sum of G_S and G_L became

minimal. To arrive at a binary infection scoring threshold, we identified a suitable value three standard

deviations below the mean of G_L. This threshold includes 99.8% of the events in G_L. Cells with an

integrated GFP intensity exceeding this threshold were considered true positive infections and were

labelled infected. The infection index was computed as the ratio of infected Cells to the total number of

Cells in the well.

Redundant siRNA Analysis (RSA)

Redundant siRNA Analysis RSA, (24) ranks genes by iteratively assigning hypergeometric p-values to

each of the multiple siRNAs targeting the same gene and picking the minimum value within a given

group to represent this gene. The ranking score indicates whether the distribution of ranks corresponding

to a gene is shifted towards the top, thereby aggregating the information provided by independent

siRNA sequences with the same target in a robust manner. Individual siRNAs from the Qiagen library

and the averages of independent replicates of the Dharmacon, Ambion, and Sigma libraries (repeated

experiments with identical siRNA) were used as input. Data was filtered prior to RSA by removing

wells that do not pass quality control, control wells and wells where cell count was below the initially

seeded cell number. As both up- and down hits are of interest to this analysis, RSA was run twice, once

with Z-scored infection scores ranked from low to high values and once ranked oppositely.

Additionally, instead of relying solemnly on manufacturer information and in order to ensure

compaRABility between libraries, siRNA targets were re-identified by searching against ENSEMBL

cDNA and the REFSEQ mRNA nucleotide data. Cases where matching failed were also excluded from

this analysis.

Infection scoring for entry assays

Wells that contain only 32 Cells or less were excluded from infection scoring. In the remaining wells,

Bacteria were filtered in a decision tree (DT) classification to exclude objects of only one pixel area.

The remaining Bacteria were filtered in a DT classification to exclude objects of less than a manually

set threshold on the upper quartile of the object intensity. The remaining Bacteria were considered true

positive infections. Based on the relation of Bacteria to Nuclei, Cells were labeled infected if and only

if a true positive Bacteria is related to the Cells Nuclei. The infection index was computed as the ratio

of infected Cells to the total number of Cells in the well. For quantification of bacterial load in infected

cells, the median of integrated GFP intensity of all true positive Bacteria was computed. The final

infection readout was the product of the infection rate and bacterial load, which gives a robust

approximation of the amount of intracellular bacteria (28).

Rescue experiment

The shRNA suppression/rescue constructs for VPS35 were kind gifts from Daniel Billadeau (38). HeLa

cells were seeded in a 6-well plate and transfected 4 h later with 0.9 µg of plasmid DNA using Fugene

HD according to the manufacturer's protocol. 72 h post-transfection cells were reseeded into a 96-well

plate (2'800 cells / well) and infected on the following day. Cells were infected with *Brucella abortus*

carrying pAC037 for 48 h. After PFA fixation and staining, cells were analyzed by image analysis.

Infection scoring was performed on YFP positive cells - indicative of successful transfection.

Immunofluorescence for LAMP-1 co-localization

Following fixation with 3.7% PFA in PBS for 20 min, HeLa cells were incubated in PBS containing

250 mM glycine for 20 min to quench remaining aldehyde residues. Cells were then permeabilized with

saponin buffer (PBS containing 0.2% saponin and 3% bovine serum albumin) for 1 h. Immunostaining

was performed by incubating coverslips with saponin buffer containing antibodies against LAMP-1

(Abcam ab25630) and *Brucella abortus* LPS polyclonal RABbit serum (kind gift from Xavier De Bolle

(56)) overnight in a humidified chamber at 4°C. The coverslips were then washed three times with PBS

and incubated with saponin buffer containing respective fluorophore-conjugated secondary antibodies:

goat anti-mouse Alexa Fluor 488 (Thermo Fisher A11029) and anti-RABbit Alexa Fluor 647 (Cell

Signaling #4414) for 3 h in a humidified chamber at room temperature. The coverslips were then washed

three times with PBS and mounted onto glass slides using Vectashield H-100 Antifade Mounting

Medium (Vector Laboratories) and sealed with nail polish.

Confocal microscopy for single cell data

The images were captured with the LSM-800 Confocal Microscope (Carl Zeiss) using a 63x oil

objective. For each condition, 40 images were obtained at random locations across the coverslip,

representing more than 50 individual cells per conditions. The images were deconvolved using Huygens

software (Scientific Volume Imaging). The presence of LAMP-1 signal around the bacteria was

quantified from the images by assessing the overlap between the anti-LPS and the anti-LAMP-1 staining

for each individual bacterium (more than 400 per condition).

Acknowledgments

We want to thank Julia Feldmann and Dr. Sonia Borrell from the SwissTPH, Basel, Switzerland for

great assistance and support with the BSL-3 environment. We are thankful to the Imaging Core Facility

of the Biozentrum, Basel for excellent assistance and advice with confocal microscopy. We also thank

Dr. Daniel Billadeau for sharing the VPS35 shRNA suppression/rescue constructs. Finally, we thank

Dr Simone Eicher for help with the RSA analysis and critical discussions. The genome sequence used

in this research was derived from a HeLa cell line. Henrietta Lacks, and the HeLa cell line that was

established from her tumor cells without her knowledge or consent in 1951, have made significant

contributions to scientific progress and advances in human health. We are grateful to Henrietta Lacks,

now deceased, and to her surviving family members for their contributions to biomedical research. This

study was reviewed by the NIH HeLa Genome Data Access Working Group

(http://acd.od.nih.gov/hlgda.htm). The genomic datasets used for analysis described in this manuscript

were obtained from the database of Genotypes and Phenotypes (dbGaP) through dbGaP accession

number phs000640.v1.p1. This work was supported by grants 31003A_173119 to C.D. from the Swiss

National Science Foundation (SNSF, www.snf.ch), advanced grant 340330 to C.D. (FicModFun) from

the European Research Council (ERC), grant 51RTP0_151029 to C.D. for the Research and Technology

Development (RTD) project TargetInfectX in the frame of SystemsX.ch (www.systemX.ch), the Swiss

Initiative for System Biology. A.C. and S.L. AC and SH were fellows of the 'Fellowship of Excellence'

International PhD program of the Biozentrum, University of Basel, Switzerland. H.B. benefited from

the SNSF Flexibility Grant.

A.C. and S.L. performed all genome-wide siRNA screening, A.C. performed the entry assays, M.E.,

K.S., and A.C. designed and performed the image analysis for the screening and entry assays. M.Q.,

A.C., and H.B. performed the analysis of the genome-wide datasets. M.Q. and A.C. analyzed the entry

assay datasets. T.T., S.L., and H.B. designed and performed the VPS35 rescue experiment. T.T. and

J.S. designed and performed the single cell experiment. M.Q. and J.S. analyzed the single cell data.

M.Q., A.C. and C.D. wrote the manuscript. C.D. provided strategic leadership for the project, designed

experiments, and discussed data. The authors declare no competing interests.

References

1.   Creasey EA, Isberg RR. 2014. Maintenance of vacuole integrity by bacterial pathogens. Curr
Opin Microbiol 17:46-52.
2.   Di Russo Case E, Samuel JE. 2016. Contrasting lifestyles within the host cell. Microbiol Spectr
4.

3.   Weber MM, Faris R. 2018. Subversion of the endocytic and secretory pathways by bacterial
effector proteins. Front Cell Dev Biol 6:1.

4.   Moreno E. 2014. Retrospective and prospective perspectives on zoonotic brucellosis. Front
Microbiol 5:213.

5.   Pappas G, Papadimitriou P, Akritidis N, Christou L, Tsianos EV. 2006. The new global map of
human brucellosis. Lancet Infect Dis 6:91-9.

6.   Ariza J, Bosilkovski M, Cascio A, Colmenero JD, Corbel MJ, Falagas ME, Memish ZA,
Roushan MR, Rubinstein E, Sipsas NV, Solera J, Young EJ, Pappas G, International Society
of C, Institute of Continuing Medical Education of I. 2007. Perspectives for the treatment of
brucellosis in the 21st century: the Ioannina recommendations. PLoS Med 4:e317.

7.   Sedzicki J, Tschon T, Low SH, Willemart K, Goldie KN, Letesson JJ, Stahlberg H, Dehio C.
2018. 3D correlative electron microscopy reveals continuity of *Brucella*-containing vacuoles
with the endoplasmic reticulum. J Cell Sci 131.

8.   Pizarro-Cerda J, Meresse S, Parton RG, van der Goot G, Sola-Landa A, Lopez-Goni I, Moreno
E, Gorvel JP. 1998. *Brucella abortus* transits through the autophagic pathway and replicates in
the endoplasmic reticulum of nonprofessional phagocytes. Infect Immun 66:5711-24.

9.   Starr T, Ng TW, Wehrly TD, Knodler LA, Celli J. 2008. *Brucella* intracellular replication
requires trafficking through the late endosomal/lysosomal compartment. Traffic 9:678-94.

10.   Castaneda-Roldan EI, Avelino-Flores F, Dall'Agnol M, Freer E, Cedillo L, Dornand J, Giron
JA. 2004. Adherence of *Brucella* to human epithelial cells and macrophages is mediated by
sialic acid residues. Cell Microbiol 6:435-45.

11.   Castaneda-Roldan EI, Ouahrani-Bettache S, Saldana Z, Avelino F, Rendon MA, Dornand J,
Giron JA. 2006. Characterization of SP41, a surface protein of *Brucella* associated with
adherence and invasion of host epithelial cells. Cell Microbiol 8:1877-87.

12.   Guzman-Verri C, Chaves-Olarte E, von Eichel-Streiber C, Lopez-Goni I, Thelestam M,
Arvidson S, Gorvel JP, Moreno E. 2001. GTPases of the Rho subfamily are required for
Brucella abortus internalization in nonprofessional phagocytes: direct activation of Cdc42. J
Biol Chem 276:44435-43.

13.   Arellano-Reynoso B, Lapaque N, Salcedo S, Briones G, Ciocchini AE, Ugalde R, Moreno E,
Moriyon I, Gorvel JP. 2005. Cyclic beta-1,2-glucan is a *Brucella* virulence factor required for
intracellular survival. Nat Immunol 6:618-25.

14.   Bellaire BH, Roop RM, 2nd, Cardelli JA. 2005. Opsonized virulent *Brucella abortus* replicates
within nonacidic, endoplasmic reticulum-negative, LAMP-1-positive phagosomes in human
monocytes. Infect Immun 73:3702-13.

15.   Celli J, de Chastellier C, Franchini DM, Pizarro-Cerda J, Moreno E, Gorvel JP. 2003. *Brucella*
evades macrophage killing via VirB-dependent sustained interactions with the endoplasmic
reticulum. J Exp Med 198:545-56.

16.   Boschiroli ML, Ouahrani-Bettache S, Foulongne V, Michaux-Charachon S, Bourg G, Allardet-
Servent A, Cazevieille C, Liautard JP, Ramuz M, O'Callaghan D. 2002. The *Brucella suis virB*
operon is induced intracellularly in macrophages. Proc Natl Acad Sci U S A 99:1544-9.

17.   Comerci DJ, Martinez-Lorenzo MJ, Sieira R, Gorvel JP, Ugalde RA. 2001. Essential role of
the VirB machinery in the maturation of the *Brucella abortus*-containing vacuole. Cell
Microbiol 3:159-68.

18.   Ke Y, Wang Y, Li W, Chen Z. 2015. Type IV secretion system of *Brucella* spp. and its effectors.
Front Cell Infect Microbiol 5:72.

19.   Miller CN, Smith EP, Cundiff JA, Knodler LA, Bailey Blackburn J, Lupashin V, Celli J. 2017.
A *Brucella* type IV effector targets the COG tethering complex to remodel host secretory traffic
and promote intracellular replication. Cell Host Microbe 22:317-329 e7.

20.   Franceschini A, Meier R, Casanova A, Kreibich S, Daga N, Andritschke D, Dilling S, Ramo P,
Emmenlauer M, Kaufmann A, Conde-Alvarez R, Low SH, Pelkmans L, Helenius A, Hardt
WD, Dehio C, von Mering C. 2014. Specific inhibition of diverse pathogens in human cells by
synthetic microRNA-like oligonucleotides inferred from RNAi screens. Proc Natl Acad Sci U
S A 111:4548-53.

21.   Ramo P, Drewek A, Arrieumerlou C, Beerenwinkel N, Ben-Tekaya H, Cardel B, Casanova A,
Conde-Alvarez R, Cossart P, Csucs G, Eicher S, Emmenlauer M, Greber U, Hardt WD,

Helenius A, Kasper C, Kaufmann A, Kreibich S, Kuhbacher A, Kunszt P, Low SH, Mercer J,
Mudrak D, Muntwiler S, Pelkmans L, Pizarro-Cerda J, Podvinec M, Pujadas E, Rinn B, Rouilly
V, Schmich F, Siebourg-Polster J, Snijder B, Stebler M, Studer G, Szczurek E, Truttmann M,
von Mering C, Vonderheit A, Yakimovich A, Buhlmann P, Dehio C. 2014. Simultaneous
analysis of large-scale RNAi screens for pathogen entry. BMC Genomics 15:1162.
22. Quebatte M, Dehio C. 2017. Systems-level interference strategies to decipher host factors
involved in bacterial pathogen interaction: from RNAi to CRISPRi. Curr Opin Microbiol 39:34-
41.
23. Zhu J, Davoli T, Perriera JM, Chin CR, Gaiha GD, John SP, Sigiollot FD, Gao G, Xu Q, Qu H,
Pertel T, Sims JS, Smith JA, Baker RE, Maranda L, Ng A, Elledge SJ, Brass AL. 2014.
Comprehensive identification of host modulators of HIV-1 replication using multiple
orthologous RNAi reagents. Cell Rep 9:752-66.
24. Konig R, Chiang CY, Tu BP, Yan SF, DeJesus PD, Romero A, Bergauer T, Orth A, Krueger
U, Zhou Y, Chanda SK. 2007. A probability-based approach for the analysis of large-scale
RNAi screens. Nat Methods 4:847-9.
25. Huang da W, Sherman BT, Lempicki RA. 2009. Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. Nat Protoc 4:44-57.
26. Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, Santos A, Doncheva NT,
Roth A, Bork P, Jensen LJ, von Mering C. 2017. The STRING database in 2017: quality-
controlled protein-protein association networks, made broadly accessible. Nucleic Acids Res
45:D362-D368.
27. Keriel A, Botella E, Estrach S, Bragagnolo G, Vergunst AC, Feral CC, O'Callaghan D. 2015.
Brucella intracellular life relies on the transmembrane protein CD98 heavy chain. J Infect Dis
211:1769-78.
28. Casanova A, Low SH, Emmenlauer M, Conde-Alvarez R, Salcedo SP, Gorvel JP, Dehio C.
2016. Microscopy-based assays for high-throughput screening of host factors involved in
Brucella infection of Hela cells. J Vis Exp doi:10.3791/54263.
29. Gallon M, Cullen PJ. 2015. Retromer and sorting nexins in endosomal sorting. Biochem Soc
Trans 43:33-47.
30. Wang J, Fedoseienko A, Chen B, Burstein E, Jia D, Billadeau DD. 2018. Endosomal receptor
trafficking: Retromer and beyond. Traffic 19:578-590.
31. Linxweiler M, Schick B, Zimmermann R. 2017. Let's talk about Secs: Sec61, Sec62 and Sec63
in signal transduction, oncology and personalized medicine. Signal Transduct Target Ther
2:17002.
32. Lopez-Domenech G, Serrat R, Mirra S, D'Aniello S, Somorjai I, Abad A, Vitureira N, Garcia-
Arumi E, Alonso MT, Rodriguez-Prados M, Burgaya F, Andreu AL, Garcia-Sancho J, Trullas
R, Garcia-Fernandez J, Soriano E. 2012. The Eutherian Armcx genes regulate mitochondrial
trafficking in neurons and interact with Miro and Trak2. Nat Commun 3:814.
33. Steinberg F, Gallon M, Winfield M, Thomas EC, Bell AJ, Heesom KJ, Tavare JM, Cullen PJ.
2013. A global analysis of SNX27-retromer assembly and cargo specificity reveals a function
in glucose and metal ion transport. Nat Cell Biol 15:461-71.
34. Temkin P, Lauffer B, Jager S, Cimermancic P, Krogan NJ, von Zastrow M. 2011. SNX27
mediates retromer tubule entry and endosome-to-plasma membrane trafficking of signalling
receptors. Nat Cell Biol 13:715-21.
35. Seaman MN, McCaffery JM, Emr SD. 1998. A membrane coat complex essential for
endosome-to-Golgi retrograde transport in yeast. J Cell Biol 142:665-81.
36. Kvainickas A, Jimenez-Orgaz A, Nagele H, Hu Z, Dengjel J, Steinberg F. 2017. Cargo-
selective SNX-BAR proteins mediate retromer trimer independent retrograde transport. J Cell
Biol 216:3677-3693.
37. Simonetti B, Danson CM, Heesom KJ, Cullen PJ. 2017. Sequence-dependent cargo recognition
by SNX-BARs mediates retromer-independent transport of CI-MPR. J Cell Biol 216:3695-
3712.
38. Liu TT, Gomez TS, Sackey BK, Billadeau DD, Burd CG. 2012. Rab GTPase regulation of
retromer-mediated cargo export during endosome maturation. Mol Biol Cell 23:2505-15.

39. Elwell C, Engel J. 2018. Emerging role of retromer in modulating pathogen growth. Trends
Microbiol 26:769-780.
40. Personnic N, Barlocher K, Finsel I, Hilbi H. 2016. Subversion of retrograde trafficking by
translocated pathogen effectors. Trends Microbiol 24:450-462.
41. Yin P, Hong Z, Yang X, Chung RT, Zhang L. 2016. A role for retromer in hepatitis C virus
replication. Cell Mol Life Sci 73:869-81.
42. Yao J, Yang F, Sun X, Wang S, Gan N, Liu Q, Liu D, Zhang X, Niu D, Wei Y, Ma C, Luo ZQ,
Sun Q, Jia D. 2018. Mechanism of inhibition of retromer transport by the bacterial effector
RidL. Proc Natl Acad Sci U S A 115:E1446-E1454.
43. Romano-Moreno M, Rojas AL, Williamson CD, Gershlick DC, Lucas M, Isupov MN,
Bonifacino JS, Machner MP, Hierro A. 2017. Molecular mechanism for the subversion of the
retromer coat by the *Legionella* effector RidL. Proc Natl Acad Sci U S A 114:E11151-E11160.
44. Barlocher K, Hutter CAJ, Swart AL, Steiner B, Welin A, Hohl M, Letourneur F, Seeger MA,
Hilbi H. 2017. Structural insights into *Legionella* RidL-Vps29 retromer subunit interaction
reveal displacement of the regulator TBC1D5. Nat Commun 8:1543.
45. Harbour ME, Breusegem SY, Antrobus R, Freeman C, Reid E, Seaman MN. 2010. The cargo-
selective retromer complex is a recruiting hub for protein complexes that regulate endosomal
tubule dynamics. J Cell Sci 123:3703-17.
46. Harrison MS, Hung CS, Liu TT, Christiano R, Walther TC, Burd CG. 2014. A mechanism for
retromer endosomal coat complex assembly with cargo. Proc Natl Acad Sci U S A 111:267-72.
47. Seaman MN, Harbour ME, Tattersall D, Read E, Bright N. 2009. Membrane recruitment of the
cargo-selective retromer subcomplex is catalysed by the small GTPase Rab7 and inhibited by
the Rab-GAP TBC1D5. J Cell Sci 122:2371-82.
48. Jimenez-Orgaz A, Kvainickas A, Nagele H, Denner J, Eimer S, Dengjel J, Steinberg F. 2018.
Control of RAB7 activity and localization through the retromer-TBC1D5 complex enables
RAB7-dependent mitophagy. EMBO J 37:235-254.
49. Lee JJ, Kim DG, Kim DH, Simborio HL, Min W, Lee HJ, Her M, Jung SC, Watarai M, Kim
S. 2013. Interplay between clathrin and Rab5 controls the early phagocytic trafficking and
intracellular survival of *Brucella abortus* within HeLa cells. J Biol Chem 288:28049-57.
50. Elfaki MG, Al-Hokail AA. 2009. Transforming growth factor beta production correlates with
depressed lymphocytes function in humans with chronic brucellosis. Microbes Infect 11:1089-
96.
51. Goenka R, Parent MA, Elzer PH, Baldwin CL. 2011. B cell-deficient mice display markedly
enhanced resistance to the intracellular bacterium *Brucella abortus*. J Infect Dis 203:1136-46.
52. Celli J, Salcedo SP, Gorvel JP. 2005. *Brucella* coopts the small GTPase Sar1 for intracellular
replication. Proc Natl Acad Sci U S A 102:1673-8.
53. Thanbichler M, Iniesta AA, Shapiro L. 2007. A comprehensive set of plasmids for vanillate-
and xylose-inducible gene expression in *Caulobacter crescentus*. Nucleic Acids Res 35:e137.
54. Smith K, Li Y, Piccinini F, Csucs G, Balazs C, Bevilacqua A, Horvath P. 2015. CIDRE: an
illumination-correction method for optical microscopy. Nat Methods 12:404-6.
55. Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang IH, Friman O, Guertin DA, Chang
JH, Lindquist RA, Moffat J, Golland P, Sabatini DM. 2006. CellProfiler: image analysis
software for identifying and quantifying cell phenotypes. Genome Biol 7:R100.
56. Deghelt M, Mullier C, Sternon JF, Francis N, Laloux G, Dotreppe D, Van der Henst C, Jacobs-
Wagner C, Letesson JJ, De Bolle X. 2014. G1-arrested newborn cells are the predominant
infectious form of the pathogen *Brucella abortus*. Nat Commun 5:4366.
687

Figure legend (figure included for review)

Figure 1

Overview of the high-content screening and analysis. (**A**) Summary of RNAi screening workflow.

Reverse transfection of HeLa cells was performed in 384-well format for 72 h followed by 48 h infection

with GFP expressing *B. abortus*, PFA fixation, and staining of HeLa cells with DAPI and DY-547-

Phalloidin before automated imaging. (**B**) Image analysis was performed with CellProfiler to segment

nuclei and bacteria and to extract measurements. (**C**) Accurate association of segmented bacteria to

nuclei enables quantitative single cell measurements. The naive association (middle image) of

segmented pathogen can be affected by over-splitting in dense cell populations (left image). Our

proposed solution (right image) based on a nucleus attraction score. (**D**) The plate histogram shows the

bimodal distribution of integrated GFP intensity corresponding to *Brucella* replication. Intensity on the

X-axes is log2-scaled to account for exponential growth. The normal distribution fitted (red curve) to

the Kernel Density Estimation of the histogram allows to compute a robust binary infection threshold

(dashed line). Associated are samples of single-cell images corresponding to the intervals of the

intensity distribution (for more details see Materials and Methods).

Figure 2

The *human* infectome for *Brucella* infection determined by genome-wide siRNA screening. (A)

Representative merged images from the genome wide screen showing nuclei (DAPI) and intracellular

replication of GFP expressing *Brucella abortus* for either control conditions (scramble) or a panel of

identified hits (RAB7A, FGFR1, ActR3, TGFBR2, VPS35, ANLN and AP2S1). Scale bar = 100 μm.

(B, C, D) Results of gene ontology enrichment analysis (DAVID) for the entire hit list (B), the down-

(C) or up-hits (D). The -$\log_{10}$ of P-value associated to the different categories are indicated (cut-off:

Benjamini corrected P-value ≤ 0.01) as well as the number of individual components associated to the

displayed categories are indicated. In (B), the first number refers to down-hits and the second to up-

hits. **(E, F)** High confidence protein-protein interaction networks ($x \geq 0.8$) for the 224 RSA down- (A)

or 202 up-hits (B) determined using the STRING database. Clusters with common predicted cellular

function are colored and their prominent function are indicated. For clarity, disconnected components

are not displayed.

 Figure 3

Entry assay identifies new components required for post-entry processes during *Brucella* infection. (A) Representative images from the entry assay showing nuclei (DAPI) of HeLa cells and intracellular *Brucella abortus* (GFP) for control condition (mock) and cells treated with siRNAs against Cdc42 or VPS35. Scale bar = 100 μm. HeLa cells were infected with *B. abortus* expressing GFP under a tetracycline inducible system for 8 h (see Material and Methods). **(B)** Scatter plot in double logarithmic scale showing infection rates measured for the entry assay (8 h) versus endpoint assay (48 h), normalized to the respective mock dataset (Table S3). For the entry assay, cells containing single bacteria were considered as infected and the final readout is proportional to the median number of bacteria per infected cells. For the endpoint assay, only cells containing replicating bacteria were considered as infected (Fig.1 and Materials and Methods). Each data point corresponds to the average of all siRNAs or esiRNAs targeted against the gene of interest (n=3). Dotted lines indicate (+/- 50%) infection threshold for both assays. The straight fit (oblique line, $r^2 = 0.763$) indicates a globally high

734 correlation between both assays. Highlighted (orange box) are the genes showing an average endpoint

735 infection score < 50% and an entry infection score <50% For ease of visualization, only the averages

736 over all RNAi products targeting a given gene are displayed.

 Figure 4

The VPS retromer is a key component of *Brucella* intracellular trafficking. (**A**) Schematic

representation of the retromer components and their prominent interactors. (**B**) Dot box representation

of the z-scored infection index for components of the retromer and interactors, including the positive

control ARPC3. Asterisks indicate statistically significant difference to scrambled siRNA-treated

bacteria (Ctrl) as determined by one-way ANOVA and Dunnett's multiple comparisons test (** Pval ≤

0.001, ns not significant). (**C**) Schematic representation of the shRNA constructs used in (D). The gray

box on the shVPS35 / Rescue construct indicates the silent mutations that prevent recognition by to the

co-expressed shRNA (38). (**D**) Infection index of transfected cells infected with *Brucella abortus* for

48 h. Displayed are the averaged infection index and associated standard deviation, normalized to the

YFP only condition (n=3). Asterisks indicate statistically significant difference to scrambled YPF only

condition as determined by paired t-test (* Pval ≤ 0.01, ns not significant).

Figure 5

VPS35 is required for *Brucella* to escape the lysosomal pathway. (**A**) Immunofluorescence approach

used to quantify localization of *Brucella* within LAMP-1 positive vesicles, illustrated with a

representative example of control-treated cells 6 hpi. Individual channels and merged picture are

presented. Arrows indicate examples of co-localization of bacteria with LAMP-1 positive

compartments. Staining of *Brucella* LPS (αLPS) was used to confirm the presence of LAMP-1 in direct

proximity of the bacterial surface. Asterisks indicate examples of LAMP-1 negative *Brucella*. (**B**)

Representative images from *Brucella* infected cells either mock transfected (ctrl) or after VPS35

knockdown (siVPS35). Samples were fixed 6 and 18 hpi. For clarity only the LAMP-1 (cyan) and

dsRed (magenta) channels are presented. Scale bar: 5 μm. (**C**) Global quantification of LAMP-1

negative *Brucella*. Displayed are the average and associated standard deviation for more than 500

bacteria and more than 50 Hela cells per time point and condition (n=3). (**D**) Single-cell data

representation of the data presented in (C). Displayed is the distribution of LAMP-1 positive *Brucella*

per cell as a function of the total number of bacteria counted in that given cell.

Supporting information legends

Supporting Table S1

List of the 224 down-hits from the genome-wide siRNA screen selected based on the RSA analysis with

associated p-value and infection score (cut-off: RSA p-value <0.01, S1_Table.xlsx).

Supporting Table S2

List of the 202 up-hits from the genome-wide siRNA screen selected based on the RSA analysis with

associated p-value and infection score (cut-off: RSA p-value <0.01, S2_Table.xlsx).

Supporting Table S3

Aggregated image analysis data from the entry assay used for Figure 3, with associated infection scores

for both entry and endpoint assays. Single data points and averaged data per siRNA and per genes (as

displayed in Fig. 3B) are indicated (S3_Table.xlsx).