The dynamics of spawning acts by a semelparous fish and its associated energetic expenses

Cédric TENTELIER[1,a], Colin BOUCHARD[1,2,b], Anaïs BERNARDIN[1], Amandine TAUZIN[1],

Jean-Christophe AYMES[1,c], Frédéric LANGE[1,d], Charlotte RECAPET[1,e], Jacques RIVES[1,f]

[1] Universite de Pau et des Pays de l'Adour, E2S UPPA, INRAE, ECOBIOP, Saint-Pée-sur-Nivelle, France

[2] Tour du Valat, Research Institute for the Conservation of Mediterranean Wetlands, Arles, France

[a] ORCID: 0000-0003-2178-4900
[b] ORCID: 0000-0003-0845-9492
[c] ORCID: 0000-0002-1181-3010
[d] ORCID: 0000-0001-8561-1879
[e] ORCID: 0000-0001-5414-8412
[f] ORCID: 0000-0001-6074-3488

Corresponding author: Cédric TENTELIER,

Univ. Pau & Pays Adour

Collège STEE – Campus de la côte basque

1 allée du parc Montaury

64 600 Anglet, France.

cedric.tentelier@univ-pau.fr

Abstract

1.   During the reproductive season, animals have to manage both their energetic and

gametic budgets. In particular, for semelparous capital breeders with determinate fecundity and

no parental care other than gametic investment, the depletion of energetic stock must match the

depletion of gametic stock, so that individuals get exhausted just after their last egg is laid.

Although these budgets are managed continuously, monitoring the dynamics of mating acts and

energy expenditure at a fine temporal scale in the wild is challenging.

2.   This study aimed to quantify the individual dynamics of spawning acts and the

concomitant energy expenditure of female Allis shad (*Alosa alosa*) throughout the mating

season.

3.   Using eight individual-borne accelerometers for one month, we collected tri-axial

acceleration, temperature, and pressure data that we analysed to i) detect the timing of spawning

acts, ii) estimate energy expenditure from tail beat frequency and water temperature, and iii)

monitor changes in body roundness from the position of the dorsally-mounted tag relative to

the vertical.

4.   Female shad had a higher probability to spawn during warmer nights, and their

spawning acts were synchronized within each active night. They underwent warmer

temperature during the day, when they stayed deeper, and swam faster at night, when they spent

more energy. Over one month of spawning, they performed on average 15.75 spawning acts,

spent on average 6 277 kJ and died with a significant portion of residual oocytes. The

acceleration-based indicator of body roundness was correlated to condition coefficient

measured at capture, and globally decreased through the spawning season, although the

indicator was noisy and was not correlated to changes in estimated energy expenditure.

5.   Despite significant individual variability, our results indicate that female shad exhausted

their energetic stock faster than their egg stock. Water warming might accentuate the mismatch

between energetic and gametic stocks. Although perfectible, the three complimentary analyses

of acceleration data are promising for *in situ* monitoring of energy expenditure related to

specific behaviour.

56 Keywords

57 Accelerometer, biologging, clupeid, egg retention, energy budget, reproductive effort,

58 semelparity, temperature, thinning.

Introduction

The energy acquired by living organisms is allocated to survival and reproduction, and natural selection is expected to favour optimal allocation, resulting in life-histories that maximize Darwinian fitness in the environment where evolution occurs (Pianka 1976; Stearns 1992; Roff 1993). When adult survival is low compared to juvenile survival, extreme reproductive effort (in gametogenesis, mating behaviour and parental care) can be selected and semelparity may arise, in which individuals die after their first and only breeding season. Semelparity is often accompanied with capital breeding, so that reproduction relies on energy reserves constituted before the breeding season (Bonnet et al. 1998). Although semelparity is sometimes quoted as "big bang reproduction", its broad definition encompasses cases where individuals breed in several bouts within a breeding season (Kirkendall and Stenseth 1985; Hughes 2017). Furthermore, in species with no other parental care than gametic investment, the optimal allocation should result in individuals dying just after their last progeny is produced, and deviation from optimality consists in individuals either surviving after their last egg is laid, or dying of exhaustion while still bearing unlaid eggs (Heimpel and Rosenheim 1998). Hence, the schedule of breeding events and the dynamics of energy expenditure during the breeding season, which may be steep in semelparous capital breeders, is a crucial aspect of reproductive strategy. Beside energy expenditure, the temporal distribution of breeding events within a season may be linked to social factors that synchronize breeding and affect the mating system of the population (Emlen and Oring 1977).

Tracking the schedule of breeding events and the dynamics of energy expenditure of breeding individuals in the wild is technically challenging. The timing of breeding events along the breeding season requires thorough observation of repeatedly detectable individuals, for example through video supervision of highly sedentary individuals (e.g. Borgia 1985). Energy expenditure has been monitored at the population level, by quantifying energy reserves on

different individuals sampled at different stages of the breeding season (e.g. Hendry and Berg

1999). In some cases, the same individuals were captured at the beginning and at the end of the

breeding season, and the energy expenditure quantified with difference in weight (Anderson

and Fedak 1985; Rands et al. 2006), body composition (Hendry and Beall 2004; Casas et al.

2005), or concentration of plasma metabolites (Gauthey et al. 2015). Likewise, in terrestrial

animals, the turnover of ^{18}O and ^{2}H from doubly labelled water quantifies the average field

metabolic rate at the individual level between two sampling occasions (Nagy et al. 1999). All

these methods give a good idea of the total energy expenditure over the whole breeding season

but often lack the temporal resolution to document the longitudinal dynamics of energy

expenditure all along the breeding season.

With the advent of bio-logging, the temporal resolution of individual data collected in

the field has tremendously increased. Among bio-loggers, accelerometers, often coupled with

other sensors such as thermometers, have been increasingly used, and advances in their

energetic efficiency now allows to record high-frequency data for weeks or months, making

them valuable tools for longitudinal studies of breeding behaviour in many organisms. In this

context, acceleration data can quantify both the overall movement of tagged individuals, which

can be translated in energy through laboratory calibration (Wilson et al. 2006; Groscolas et al.

2010; Collins et al. 2016; Hicks et al. 2017 Jan 1), and the number and temporal distribution of

key behaviours such as mating events, if these produce a typical acceleration pattern (Brown et

al. 2013).

The presented study aimed at describing the activity and change in body condition of

female semelparous fish throughout their spawning season. Using accelerometers, we

investigated three main aspects:

1) The schedule of spawning events was identified using a characteristic acceleration. We

tested whether the timing of spawning events was influenced by temperature (as usual

in fishes) and whether it was synchronized within and among females.

2) The global activity was quantified as tail beat frequency, and converted to energy

expenditure through an energetic model including water temperature. This was used to

identify variability in energy expenditure between individuals and between periods of

the spawning season.

3) We tested the possibility to use the angle between the dorsally-mounted accelerometer

and the vertical as an indicator of body roundness of the fish. This would allow

monitoring the thinning process through the spawning season, and relate it to periods of

high energy expenditure.

Egg load at the death of individuals was also measured to test the relationship between

spawning dynamics, energy expenditure and quantity of unspent eggs.

Methods

Characteristics of the species studied, and predictions

Allis shad (*Alosa alosa* L.) is an anadromous clupeid fish distributed along the Atlantic

coast of Europe, from Portugal to the British Isles, with the main populations dwelling in the

French rivers Garonne, Dordogne and Loire (Baglinière and Elie 2000). Across its distribution,

Allis shad is considered as a semelparous species, although spawning marks on scales suggest

that a very small proportion of individuals may spawn on two consecutive years (Mennesson-

Boisneau and Boisneau 1990; Taverny 1991). After having spent a few months in freshwater

as juveniles and four to six years at sea, shad undertake freshwater upstream migration during

which they fast, thus qualifying as capital breeders. Gonad maturation occurring during

migration leads females to bear between 13 000 and 576 000 eggs, reaching an average gonad

mass of 221 g and a gonad / somatic weight ratio of 15% on spawning grounds (Cassou-Leins

and Cassou-Leins 1981; Taverny 1991). Spawning typically occurs at night in a 0.3 to 3 meter-

deep glide. The spawning act consists of at least one male and one female swimming side by

side during five to ten seconds while describing three to five circles of one meter in diameter

and beating the water surface vigorously with their tail (Baglinière and Elie 2000). The typical

splashing noise (35 dB at one meter) can be heard and recorded from the river bank, and the

number of splashes recorded during the spawning season is often used as an indicator of the

number of spawners in a population (Chanseau et al. 2004).

Although very few data are available, the number of splashes recorded upstream from

dams where migrating shad were counted suggest that females perform on average five to

twelve spawning acts during the season (Fatin and Dartiguelongue 1996; Acolas et al. 2006

Apr). Moreover, based on the dynamics of ovary index and oocyte diameter measured on

individual caught and dissected across the season, Cassou-Leins & Cassou-Leins (1981)

postulated that shad mature their eggs in five to seven batches. We therefore expected the

number of spawning acts recorded by the individual-borne accelerometers to be between five

and twelve, separated with a regular delay corresponding to the maturation time of each egg

batch.

The spawning season spans from early-May to late July, but dead individuals can be

collected downstream from spawning ground as soon as late May, so we expected individual

reproductive lifespan to be around one month. The number of splashes heard on spawning

ground increases with increasing water temperature (Baglinière and Elie 2000; Paumier et al.

2019). This could be due to either more individuals being sexually active or individuals to be

more active at higher temperature. Hence, we expect a positive effect of water temperature

during a night on either the individual probability of performing at least one splash during that

night or the number of splashes performed by each individual active on that night. Within a

night, the temporal distribution of splashes typically follows a Gaussian distribution centred on

2 AM and spanning from 10 PM to 6 AM (Cassou-Leins and Cassou-Leins 1981). However,

this was observed mainly for large populations with thousands of spawners and hundreds of

splashes per night. In the case of a small population, the temporal distribution may depart from

this particular Gaussian distribution. In particular, the aggregative behaviour of shad, which

shoal even during the spawning season (Baglinière and Elie 2000), suggests that females may

synchronize their splashes within a night.

Shad being capital breeders, samples of different individuals caught at different times

during the migration have shown dramatic changes in somatic mass and tissue composition

(Cassou-Leins and Cassou-Leins 1981; Bengen 1992), but no longitudinal data exist at the

individual level. We expected that shad would get thinner as the season progresses, and all the

more so during periods of high energy expenditure due to warm water (recorded by individual-

borne thermometers) or intense activity (recorded by individual-borne accelerometers). If the

angle of the accelerometer with the vertical is a good indicator of the individual's roundness, it

should be correlated to individual coefficient of condition when fish are measured and weighed.

It should also decrease throughout the spawning season as the fish thins, and its shift during a

period of the season should be correlated to the energy expenditure during this period.

Because shad are nocturnal and capital breeders, they were expected to rest during the

day in order to save energy and survive until they have laid all their eggs. We therefore predicted

that shad should swim slowly and stay in deeper, cooler water during the day, and be more

active and in shallow water during the night (Baglinière and Elie 2000). Accordingly, their

energy expenditure should be lower during the day than during the night, and their body

condition was expected to decrease faster during the night than during the day. Finally, facing

the risk of death before having laid all eggs, shad which have well managed their energy were

expected to die with less residual eggs. We therefore predicted that the mass of residual eggs at

death should be linked negatively with the number of spawning acts and energy expenditure

during the night, and positively with energy expenditure during the day.

Fieldwork

We conducted this study in spring 2017 and 2018 in the Nivelle, a 39 km long coastal

river situated in the Northern Basque Country, France, and draining a 238 km² basin. The

downstream limit of the study zone was the impassable Uxondoa weir, situated 12 km upstream

from the river mouth (43°21'40.64"N, 1°35'13.99"W), and equipped with a vertical slot fishway

and a trap where a yearly average of 230 (min = 26; max = 688) migrating shad have been

counted since 1996, with less than 30 individuals per year since 2015. Five kilometres upstream

from Uxondoa stands another impassable weir equipped with a fishway and a trap, where shad

have almost never been captured.

The Uxondoa fish trap was controlled daily throughout spring (ECP 2018). Nine female

shad were captured and tagged in 2017, and 15 in 2018. The tagging procedure was quite similar

to Breine (2017) for twaite shad (*Alosa fallax*). We anesthetized each individual in a bath of 15

mg/L benzocaine diluted in river water before to weigh it, measure its fork length, and finally

tag it with a radio transmitter emitting at a unique frequency (F2020, ATS, Isanti, MN, USA)

and a three-dimensional accelerometer (WACU, Atesys-Montoux, Huguenau, France). For

tagging, each fish was kept in a 20 L tank filled with anaesthetic solution. A water pump placed

in the fish's mouth and a stone bubbler in the bath ensured a good circulation of aerated water

during the tagging procedure. Adjustable plastic plates covered with foam were placed

vertically against both flanks to maintain the fish in an upright position, with only its back being

above water surface. The radio transmitter and the accelerometer were cleaned with a povidone

iodine solution (Betadine®) and dried with surgical cotton. The two Teflon-coated metallic

wires of the radio transmitter were inserted approximately 1 cm under the dorsal fin through

sterile hollow needles. The hollow needles were then removed, and the metallic wires were

passed through holes drilled in the accelerometer's lug, secured with plastic eyelets and

aluminium sleeves, and the extra length was cut. The radio transmitter and the accelerometer

weighed 8.6 g and 9 g, respectively, so the weight was balanced on both sides of the fish. After

tagging, each fish was placed in a 50 L box filled with river water, and released upstream from

the weir upon waking.

From its release at Uxondoa weir until its death at the end of the spawning season, each

tagged shad was localized twice a day using a mobile receiver (R2100, ATS, Isanti, MN, USA)

and a loop antenna, in order to check whether it was still alive and to obtain its position. The

radio transmitters were set to double their pulsing rate after 8 h of total immobility. Within two

days after double pulse was detected, the dead fish and the tags were recovered by snorkelling,

and the whole fish and its ovaries were immediately weighed.

Analysis of acceleration data

The loggers used in this study recorded acceleration between -8 g and +8 g (1 g = 9.81

m.s^{-2}) in each of the three dimensions at an average frequency of 50 logs per second. Every

second, they also recorded the temperature, pressure, date, time and exact number of

acceleration logs within the past second. Both dynamic and static (gravitational) acceleration

were used to estimate three types of variables: body roundness through the angle of the

accelerometer with the vertical, tail beat frequency and occurrence of spawning acts.

The static, gravitational, component on each axis i (x, y and z) was extracted from the

raw signal at each time t by replacing each data point $\overrightarrow{A_{i,t}}$ by the average of the points within a

window of width w centred on t, so that:

$$\overrightarrow{SA_{i,t}} = \frac{1}{w} \sum_{j=t-\frac{w}{2}}^{j=t+\frac{w}{2}} \overrightarrow{A_{i,j}}$$

Unlike what is usually done, these data were not used to assess the posture of the fish,

as no known behaviour in shad involves change of posture. Instead, we hypothesized that it

may reflect the shape of the fish. Indeed, since the accelerometer was pinned on the dorsal part

of the fish's flank, the angle θ between the x-axis of the tag and the vertical may be linked to

the roundness of the fish, a proxy of its energetic reserves. This angle could be computed at any

data point of static acceleration, as $\theta_t = cos^{-1}(\overrightarrow{SA_{x,t}})$, where θ_t is the angle at time t and $\overrightarrow{SA_{x,t}}$

is the static acceleration computed on the x-axis at time t (Fig. 1.A). To track change in body

roundness through the season, θ was estimated for each fish from $\overrightarrow{SA_{x,t}}$ computed over

w=180 000 points (*i.e.* 1 h at a 50Hz sampling frequency) every eight hours (6AM, 2PM,

10PM), from its release until its death.

        A   B

243
244

Figure 1. Allis shad were tagged with a radio transmitter on the left flank and an accelerometer

on the right flank. The accelerometer continuously recorded acceleration on three axes. A. The

static component of acceleration recorded on the x-axis at time t, $\overrightarrow{SA_{x,t}}$, corresponds to the

orthogonal projection of gravity, which is vertical by definition, on this axis. Hence, the angle

θ between the x-axis and the vertical can be computed for any time t as $\theta_t = cos^{-1}(\overrightarrow{SA_{x,t}})$. B.

The dynamic component of acceleration (smoothed with a 250-point wide median filter) can be

used to detect spawning acts, characterized by increased acceleration on both x (red), y (green)

and y (blue) axes, resulting in a peak in the norm of the 3-dimensional acceleration vector

(black), and accompanied by a decrease in hydrostatic pressure (purple).

The dynamic component of acceleration was computed at every time point on each

dimension i ($\overrightarrow{DA_{i,t}}$) as the raw signal minus the static acceleration computed over w=250 points

(=5 s at 50 Hz). This was used to quantify the activity of the fish through Tail Beat Frequency

(TBF), computed as the number of zero-crossings of $\overrightarrow{RDA_{z,t}}$ per second. TBF can be computed

at every instant, but for further analysis we used average computed for every minute.

We estimated the energy consumption for every minute of each individual's

reproductive season, using its tail beat frequency (computed as described above), the

temperature recorded by the individual logger, and equations derived from the data of Leonard

et al. (1999) and Castro-Santos & Letcher (2010) on American shad, *Alosa sapidissima*.

Leonard et al. (1999) measured oxygen consumption and tail beat frequency of 18 American

shad in a swimming respirometer with water temperature varying between 13°C and 24°C. On

their data, we fitted a linear mixed model (lmer function in lme4 package; Bates et al. 2014)

with individual random intercept and fixed effects of temperature and TBF, on log-transformed

oxygen demand (MO_2, in mmol O_2.kg^{-1}.h^{-1}). Then, assuming that 0.4352 kJ of somatic energy

are burnt per mmol O_2 (Brett and Groves 1979), we used the three parameters of the mixed

model (average intercept = 1.0064, slope of temperature = 0.0531, and slope of TBF = 0.2380)

to compute the amount of energy $E_{i,t}$ (in kJ) consumed by each fish i for each minute t of its

spawning period, from its body mass $M_{i,t}$ (in kg), the average temperature recorded by its logger

during that minute $T_{i,t}$ (in °C), and its average tail beat frequency during that minute $TBF_{i,t}$ (in

beat per second):

$E_{i,t} = \frac{1}{60} \times 0.4352 \times M_{i,t} \times e^{1.0064+0.0531 \times T_{i,t}+0.2380 \times TBF_{i,t}}$ (Equation 1)

Since energy consumption continually reduces fish mass, we iteratively modelled fish mass and

energy expenditure at each minute of the spawning season. For this, we assumed that shad get

69% of their energy from lipids and 31% from proteins to fuel their metabolism (Leonard &

McCormick, 1999), and that one gram of fat yields 39.54 kJ, against 23.64 kJ per gram of protein

(Craig et al. 1978), so fish mass at the t^{th} minute was:

$M_{i,t} = M_{i,t-1} - \frac{E_{i,t}}{0.69 \times 39.54 + 0.31 \times 23.64} . 10^{-3}$ (Equation 2)

The dynamic acceleration was also used to count the number of spawning events

performed by each tagged shad. To do this, the acceleration pattern typical of spawning act was

first described in a controlled experiment and then searched in the field-collected data. To

characterize the typical pattern associated with spawning act, a male and a female shad were

captured at Uxondoa trap, tagged as described above, and observed for one month (June 2016)

in a basin (400 m², 0.6 m water depth) at the INRAE experimental facilities in St Pée sur Nivelle

(ECP 2018). Eight spawning acts were recorded on video or audio, which could be paralleled

with acceleration data. The typical pattern of acceleration associated to the spawning act was a

rise in the norm of the 3D acceleration vector, which stayed above 3 g for at least 3 seconds,

mainly driven by acceleration on the z-axis corresponding to TBF reaching up to 15 beats per

second. A decrease of hydrostatic pressure was typically associated to this pattern, due to the

fish reaching the water surface during the spawning act (Fig. 1.B). These criteria for

identification of spawning acts were robust, as they were always detected when spawning was

visually observed (no false negative) and never observed when spawning was not visually

observed (no false positive). These criteria were implemented in an algorithm to scan the

accelerograms collected in the field in 2017. This automated identification was validated by

visual comparison of accelerogram sequences automatically identified in 2017 with sequences

of eight acknowledged spawning acts recorded in 2016.

Finally, the exact timing of each individual's death was detected as a clear change in the

acceleration signal, where the gravitational component indicated that the fish switched from

upright posture to lying on its side or upside down, and the only dynamic component left was

the tenuous jiggling due to water flowing over the tag.

Statistical analysis

All analyses were performed on R (R Development Core Team 2008). A zero-inflated

mixed regression (package glmmTMB; Brooks et al. 2017) was used to test the effect of

temperature on the number of spawning acts performed by each individual on each night. The

Binomial component of the model informed on the effect of temperature on the probability to

perform any spawning act, while the Poisson component tested the effect of temperature on the

number of spawning acts. Individual was used as a random effect on both components. At a

finer timescale, we tested whether females spawning in the same night synchronized their

spawning acts. To do this, we performed 10 000 permutations of the hour of spawning acts

recorded along all nights, and compared the median of the delay to the nearest spawning act for

actual data and for simulated data. Because our aim was not to test the synchrony between acts

performed by the same female, only the first act performed by each female on a given night

was considered in this analysis.

As mentioned above, TBF was computed for every minute of each individual's

spawning season, to track its energy expenditure. As an indicator of fish activity, average TBF

was also aggregated over eight-hour time windows corresponding to three periods of the day:

morning (6AM-2PM), afternoon (2PM-10PM) and night (10PM-6AM). The limits of the night

period correspond to the earliest and latest hour of spawning activity classically recorded on

spawning grounds (Cassou-Leins and Cassou-Leins 1981). To test whether shad adopted a

different behaviour at different periods of the day, linear mixed models with individual random

intercept and period of the day as a fixed effect were fitted to TBF, temperature and depth. For

TBF, we also added a binary covariate indicating whether the fish had been tagged for more or

less than three days, to test whether recent tagging impaired activity (Føre et al. 2020 Apr 21).

Depth was estimated by computing hydrostatic pressure ($1hPa.cm^{-1}$) as the difference of

pressure recorded every second by each tag and the atmospheric pressure recorded every 20

minutes by a meteorological station situated at the INRAE facilities, less than one kilometre

from the spawning ground. All mixed models were fitted using the package lme4 for R (Bates

et al. 2014), significance of fixed effects was tested using the likelihood ratio test (LRT χ^2)

between the model including the fixed effect and the nested model excluding it. Marginal and

conditional R^2 (R^2_m and R^2_c), indicating the proportion of variance explained by the fixed effects

and by the whole model, respectively, where computed using R package MuMIn (Barton 2009).

To our knowledge, static acceleration has never been used to assess changes in animal

body roundness. We tested three predictions resulting from the hypothesis that the angle θ

between the x-axis of an individual's accelerometer and the vertical was an indicator of body

roundness. First, we fitted a Pearson correlation between Fulton condition coefficient

($100*$weight (in grams)/length (in centimetres)^3) and θ, both measured at the initial capture and

at death. Second, a linear mixed model, with individual random intercept and slope was used to

test whether θ decreased with time since individual release. Third, to test if change in θ over

the breeding season was proportional to change in condition, a correlation was performed

between the differences in θ and in Fulton condition coefficient from release to death. Finally,

to test whether the decrease in θ could be linked to temperature, activity, energy expenditure or

the moment of the days, we fitted linear mixed models with the difference in θ between the first

and the last hour of each 8-hour period and an individual random intercept. Four models were

tested, 1) the null model with only random intercept, 2) a model with average temperature and

TBF during the 8-hour period as fixed effects, 3) a model with cumulated energy expenditure

during the 8-hour period as a fixed effect, and 4) a model with the moment of the period

(morning, afternoon, night) as a fixed effect. To compare mixed models corresponding to

alternative hypotheses, we used Akaike Information Criterion corrected for small sample bias

(AICc) on R package MuMIn (Barton 2009).

Results

The nine female shad tagged in 2017 survived between 20 and 37 days (mean=26 days),

and all tags were retrieved within one or two days after fish died, although one tag stopped

recording data after ten days. The 2018 campaign was much less successful: two fish died one

week after tagging, before any spawning acts were recorded; two fish lost their tags three and

four days after tagging. The eleven remaining fish were radio tracked throughout the spawning

season, until two exceptional floods (on June 7[th] and 16[th]) flushed them down to the estuary and

the ocean where high water conductivity prevented further radio tracking, hence tag retrieval.

Eventually, among the 25 tagged shad, only eight fully exploitable and one partially exploitable

accelerograms could be collected. This sample of eight females represented half of the females

that passed the Uxondoa weir in 2017, but the power of Spearman correlations to detect even a

strong correlation of 0.5 between variables observed at the individual level was only 0.22.

According to their accelerograms, the eight female shad performed 7, 9, 12, 14, 14, 17,

24 and 26 spawning acts (mean=15.75). The total number of spawning acts performed by each

female was correlated to none of the individual variables tested (Fig. 2).

Figure 2. Spearman coefficient of correlation (expressed as percentage) between biometric and reproductive variables of female Allis shad. In order, day of capture, day of death, number of days between capture and death, body length, initial mass, initial Fulton coefficient of condition, final mass, final Fulton coefficient of condition, final gonad mass, number of spawning acts, number of nights with at least one spawning act.

For each female, spawning acts were distributed in three to six nights (mean=4.87) each separated by zero to eight nights (mean=3.55) without spawning acts, resulting in individual spawning seasons ranging from 14 to 24 nights (mean=18.12) from the first spawning act to the last (Fig. 3.A). The tagged females performed their first spawning act zero to eight days after tagging (mean=3). Only on eight occasions did a female perform a single spawning act during a night, so the 137 remaining acts were performed in volleys. At the individual level, an active night comprised from two to eight acts (mean=3.23) performed in two to 84 minutes (mean=29.7). The mixed zero-inflated Poisson regression indicated that water temperature during the night had a positive effect on the probability that a female performed at least one

spawning act (negative effect on the zero inflation; z=-1.95, p=0.05) but no effect on the number

of spawning acts in the volley (the Poisson component; z=-1.44, p=0.15). Cumulated over all

the season, the temporal distribution of spawning acts within the night followed a Normal

distribution, centred on 3AM, with 95% of spawning acts occurring between 0:30AM and

5:30AM. However, the permutation test on the hour of spawning acts indicated synchrony

between acts performed by different females on the same night: the median time lag to the

nearest act was one hour for observed data, which corresponds to the first percentile of

simulated data (Fig. 3.B).

Figure 3. The temporal distribution of spawning acts by eight Allis shad females in the river

Nivelle in spring 2017. A. Cumulated number of spawning acts for each night of the season.

Each colour corresponds to an individual. The line above the bar plot represents the average

water temperature measured each night between 10PM and 7AM. B. Synchrony of spawning

acts performed by different females within a night (only the first act of the night, for each

female). The red vertical line represents the median time lag between nearest spawning acts for

observed data. The histogram represents the same thing for 10 000 permutation of the hour of

the acts.

Average tail beat frequency (TBF), temperature and depth were computed for 312 840

minutes across all individuals' spawning seasons (eight complete and one partial). Tail beat

frequency ranged from 1.3 to 9.5 beats per second (mean=3.2), temperature ranged from 13.5

to 23.8°C (mean=18.3), and depth ranged from 0 to 400 cm (mean=186).

From equations (1) and (2), the estimated instantaneous rate of energy expenditure

ranged from 0.09 to 0.89 kJ.min^{-1} (mean=0.17), and energy expenditure cumulated by each of

the eight individuals from initial capture to death ranged from 4 395 to 8 361 kJ (mean=6 277;

Fig. 4.A). The corresponding weight loss was estimated to range from 127 to 241 g (mean=181),

making from 9% to 17% of initial weight (mean=12%). The shad died 44 to 182 hours

(mean=98.62) after their last spawning act. They had lost between 33% and 53% of their weight

(mean=42%), and their ovaries weighed 25.9 to 141.5 g (mean=79.7). No correlation was found

between weight loss and ovary weight (Spearman S=89.3; rho=-0.06; p=0.888). The predicted

and observed weight lost during the season were positively correlated, (Spearman S=8.55;

rho=0.9; p=0.002; Fig. 4.B), but the predicted weight loss was on average 1.5 times less than

the actual loss (479 g difference on average).

Figure 4. A. Energy expenditure over the spawning season for nine female Allis shad, estimated from temperature and TBF using equations (1) and (2). B. Observed weight loss and weight loss modelled with equations (1) and (2) over the spawning season. The straight line in B is the one on which points should lie if the modelled weight loss would fit the observations. Colours are as in Fig. 3, with black line in A for the individual whose tag stopped recording before the end of the experiment.

Aggregating TBF, temperature and pressure data over 8-hour periods (morning: 6AM-2PM, afternoon: 14PM-10PM, night: 10PM-6AM) produced 649 8-hour periods (Fig. 5). Shad stayed closer to the surface at night than during the morning and afternoon (LRT χ^2=238.9; p<0.0001; R^2_m=0.14; R^2_c=0.69), and underwent warmer temperature in the afternoon than in the night than in the morning (LRT χ^2=15.8; p<0.0001; R^2_m=0.02; R^2_c=0.05). TBF was higher at night than in the afternoon than in the morning (LRT χ^2=26.8; p<0.0001) and was also higher during the first nine periods (three days) just after tagging than afterwards (LRT χ^2=37.6;

p<0.0001; R^2_m=0.06; R^2_c=0.36 for the model including both effects). The estimated energy

expenditure was the highest at night, followed by afternoon and morning (LRT χ^2=24.8;

p<0.0001; R^2_m=0.03; R^2_c=0.34). The mass of eggs remaining at death was not correlated to the

energy expenditure cumulated across mornings (Spearman S=66; rho=0.21, p=0.619),

afternoons (Spearman S=64; rho=24; p=0.582), nights (Spearman S=76; rho=0.09; p=0.84), or

the whole season (Spearman S=64; rho=0.24; p=0.582).

Figure 5. Depth (A), temperature (B), tail beat frequency (C) and estimated energy spent (D)

by nine female allis shad during the morning (6AM:2PM), afternoon (2PM:10PM) and night

(10PM:6AM) of their spawning season. Box plots show the distribution of the variables

averaged (A, B, C) or cumulated (D) across 8-hour periods for each individual (colours as in

Fig. 3, black for the individual whose tag stopped recording before the end of the experiment).

In C, energy spent was computed from temperature and TBF using equations (1) and (2).

452   As expected, the θ angle between the x-axis of the accelerometer and the vertical was

453 positively correlated to the individual's Fulton coefficient of condition (Pearson's r=0.63;

454 p=0.006), although the effect seemed to be due to within-individual difference in condition and

455 θ at the beginning and at the end of the season rather than inter-individual variability in

456 condition and θ (Fig. 6.A). Moreover, the θ angle globally decreased over time for all

457 individuals (LRT χ^2=516.9; p=0.001; R^2_m=0.29; R^2_c=0.78; Fig. 6.B), with a slope of -0.21

458 degree per eight hours, although it increased during some 8-hours periods. However, contrary

459 to our expectation, the difference of the θ angle between initial capture and death was not related

460 to change in body condition between initial capture and final recapture (Spearman S=76,

461 rho=0.09, p=0.84). The best mixed model to explain the shift in θ during a 8-hour period was

462 the one with the moment (morning, afternoon or night) of the period as the independent factor

463 (LRT χ^2=56.7; p<0.0001; R^2_m=0.08; R^2_c=0.08), followed by the null model including

464 individual random effect only (ΔAICc=51), the model including estimated energy expenditure

465 as the independent variable (ΔAICc=56), and the model including both temperature and TBF

466 as independent variables (ΔAICc=58). According to the best model, the shift in θ was negative

467 during mornings (mean $\pm$ standard deviation -1.25° $\pm$ 3.08), slightly negative during afternoons

468 (-0.23° $\pm$ 2.36), and positive during the nights (0.89° $\pm$ 3.15).

Figure 6. The θ_t angle between the x-axis of the accelerometer and the vertical as an indicator of fish body roundness A. Temporal evolution of θ computed each day along the spawning season for nine Allis shad. B. Relationship between Fulton coefficient of condition and θ_t angle at initial release (triangles) and at recapture after death (points). Each individual is represented with the same colour as in Fig. 3, and black is for the individual whose tag stopped recording before the end of the experiment.

Discussion

In this study, we used acceleration data to quantify the dynamics of spawning acts, energy expenditure and thinning of Allis shad females throughout their spawning season. The results indicate that the spawning schedule of shad females, although constrained by the serial maturation of oocyte batches, was also influenced by temperature and social factors: individual females had a higher probability to spawn during warmer nights, spawned repeatedly during most of their active nights and females that spawned on the same nights synchronized their

spawning acts. Tail beat frequency and water temperature recorded by the loggers showed that

energy expenditure was slightly higher during night time than during daytime, and may have

been too high to allow shad to spawn all their eggs. Finally, the original use of gravitational

acceleration to monitor fish thinning, although perfectible, seems a promising method in animal

ecology.

Dynamics of spawning acts and energy expenditure

The number of spawning acts performed by female Allis shad ranged from 7 to 26, with

an average of 15.75. So far, the average individual number of spawning acts was indirectly

estimated to be either five (Acolas et al., 2006) or 12 (Fatin and Dartiguelongue 1996) by

counting the number of acts heard upstream a dam where all passing individuals were censused.

Such method provides no estimate of interindividual variability. In a first attempt to work at the

individual level, Acolas et al. (2004) marked three females and three males with acoustic tags,

and based on the number of detections of the tags near the surface (assessed by the reception

power by hydrophones immersed at different depth), estimated that zero, one and two acts were

performed by the females and three, 38 and 60 acts by the males. Here, despite a small sample,

we estimated a more representative distribution of the number of acts per individual, as their

detection on accelerograms is certainly more reliable than the method used by Acolas et al.

(2004). This distribution, as well as the timing of spawning acts, is crucial to build a model

estimating the number of shad in a river from the acoustic survey of spawning acts, as it is

routinely done in European rivers (e.g. Chanseau et al. 2004). Such a model, simulating shad

spawning behaviour in an Approximate Bayesian Computation framework (ABC; Csilléry et

al. 2010) is available in French here: https://ctentelier.shinyapps.io/alose_abc/. The number of

acts per female could be correlated to none of the few biometric variables collected, but the

very small sample implied a weak statistical power.

Beside oocyte maturation and temperature, the temporal distribution of spawning acts

was aggregated within and among females. While each female was active for most of the

population's spawning season, the individual spawning season was punctuated by three to six

nights of activity generally corresponding to volleys of two to eight acts performed in a few

tens of minutes, and separated by on average 3.5 nights of inactivity. The schedule of spawning

acts must be constrained by the fragmented maturation of oocytes (Cassou-Leins and Cassou-

Leins 1981; Olney et al. 2001), but also depended on temperature. Interestingly, while data

collected at the population level indicate that spawning activity increases with temperature

(Paumier et al. 2019), our data collected at the individual level showed that temperature

increased the probability that a female performed some acts during the night, but not the number

of acts it performed. In fact, two results suggest that the dynamics of spawning acts within a

night might be influenced by social factors. First, the temporal proximity of acts performed by

a given female in a given night in this study, and the high proportion of acts performed without

oocyte expulsion reported by Langkau et al. (2016) remind of female trout's 'false orgasm'

(Petersson 2001) or female lamprey's 'sham mating' (Yamazaki and Koizumi 2017). For both

of these species, it has been suggested that repeated spawning simulations enable the female to

exert mate choice, by both exhausting the sperm stock of an unwanted courtier and signalling

its mating activity to peripheral males. Second, spawning acts of different females in a given

night were more synchronous than expected from the hour of spawning acts across all nights of

the season. This suggests that a female which is ready to spawn (mature batch of oocytes) in a

propitious night (warm temperature) may trigger its spawning acts when another female does

so. Such fine scale synchrony may again affect sexual selection, reducing the environmental

potential for polygyny by making it difficult for the same male to monopolize several females

at the same time (Emlen and Oring 1977).

The final point in the spawning schedule of semelparous organisms is death, which

struck female shad two to seven days after their last spawning act, and before they laid all their

eggs, which suggests that energetic reserves were exhausted before egg stock. Combining

acceleration and temperature data collected in the field to equations parametrized in laboratory

experiments, we estimated that the energetic expenditure of spawning shad during three weeks

of spawning ranged from 4 395 to 8 361 kJ (mean=6 277). This is surprisingly of the same order

as female American shad, which entered the Connecticut river with a stock of 12 000 kJ, of

which approximately 5 000 kJ were consumed along their 228-km and seven-week long

upstream migration in waters warming from 10 to 22°C (Leonard & McCormick, 1999).

Extreme pre-spawning energy expenditure caused by long migration, obstacle crossing or warm

water can reduce the reproductive lifespan of semelparous females and hinder their ability to

spawn their full egg stock (Hruska et al. 2011). While the spawning ground used by shad in our

study is only 13 km from the river mouth with only one obstacle to pass (Uxondoa, where we

captured them), the median daily temperature recorded in the Nivelle for May and June 2017

was 18.4°C, the second warmest spring since the beginning of record in 1984, the maximum

being 18.5°C in 1989, the minimum 14.6°C in 2013, and the median 16.9°C. According to our

analysis of data from Leonard et al. (1999; our equation 1), an increase in temperature from

16.9 to 18.4°C would raise the energy expenditure of 8.3%, giving both a possible explanation

for the inability of shad to spawn all their eggs before they die and a hint of the impact of global

warming on the breeding performance of such species.

As expected from visual observations on spawning grounds, shads were more active at

night than during the day, and stayed deeper during daytime when water was warmer. Although

nocturnal spawning is usually interpreted as a strategy to decrease predation risk on spawners

or eggs (Robertson 1991), resting in deeper, hence fresher, water during the warm hours of the

day may also save energy for spawning at night. However, contrary to our predictions, the

individuals who spent less energy during mornings or afternoons and more during the night did

not have fewer residual eggs at death.

Methodological considerations

According to calculated energy expenditure, the energetic model predicted that shad

should have lost on average 12% of their initial weight, which was much less than the observed

42% loss. Of course, the energetic model did not account for egg expulsion, which must

represent a large proportion of weight loss, given that ovaries can represent up to 15% of

somatic weight at the onset of the spawning season (Cassou-Leins and Cassou-Leins 1981;

Taverny 1991). Moreover, the equations used to convert TBF and temperature to energy

expenditure and weight loss were not parametrized with data obtained on *A. alosa* but on *A.*

sapidissima (Castro-Santos & Letcher, 2010; Leonard et al., 1999; Leonard & McCormick,

1999) which is the species most closely related to *A. alosa* for which such reliable information

exist. Although American shad and Allis shad have the same morphology and ecology, some

elements suggest that results on American shad could not exactly apply to our study on Allis

shad. First, although the range of temperatures used by Leonard et al. (1999) were similar to

the temperatures encountered by shad in our study, TBF of Allis shad in the field exceeded the

fastest swim tested by Leonard et al. (1999). In particular, the brief bouts of very high TBF

which correspond to spawning acts may represent anaerobic efforts, which are more costly than

the aerobic effort observed by Leonard et al. (1999). The volleys of spawning acts performed

by females in a few tens of minutes could even lead to sexual selection of traits affecting the

recovery after sprint (Kieffer 2000). Second, the relationship linking swim speed to oxygen

demand (MO_2) varies between species, even closely related, and American shad seems to have

a particularly high metabolism compared to other clupeids (Leonard et al. 1999). This would

have led to an overestimation of energy expense for *A. alosa*. Third, the relative contribution of

lipids and proteins as metabolic fuel may differ between the iteroparous American shad and the

semelparous Allis shad. Indeed, as migratory iteroparous fishes have to spare proteins,

especially in the red muscle, for their downstream migration, semelparous species can exhaust

their protein stock to complete spawning (Schultz 1999). Also, Leonard & McCormick (1999)

measured the relative expenditure of lipids and proteins in *migrating* American shad, while we

monitored *spawning* Allis shad. It has been shown that most anadromous fishes catabolise

proteins only when lipid reserves have been exhausted, in a way that lipids mainly fuel

migration while proteins fuel spawning (Idler and Bitners 1958; Beamish et al. 1979; Jonsson

et al. 1997; Hendry and Berg 1999). Given that proteins have a lower energy density than lipids,

the surprising high energy consumption and the low weight loss predicted by the energetic

model may be explained by spawning Allis shad relying more on proteins than on lipids.

While the presence of the radio tag and the accelerometer may have imposed an

energetic cost on shad due additional weight (less than 2% of initial fish weight) and drag,

several indicators suggest that our tagging method did not impact shad behaviour as heavily

than the commonly gastric implants which result in high mortality rate or long post-tagging

downstream movements (Steinbach et al. 1986; Verdeyroux et al. 2015; Tétard et al. 2016).

Such downstream movements were only observed in 2018, after exceptional floods which have

probably also wiped away untagged shad, especially after the spawning season. Although the

dead individuals we retrieved in 2017 showed clear depigmentation, neither severe abrasion nor

fungal proliferation was observed, even after one month in water at 18° C. Moreover, the eight

females for which complete accelerograms were retrieved all spawned, three of them spawned

the night directly following tagging, and the median of 14 spawning acts was above the five to

twelve acts per individuals estimated by Acolas *et al.* (2006 Apr) and Fatin & Dartiguelongue

(1996). Tagging did not seem to impair swimming activity either, as tail beat frequency (TBF)

was not lower in the three days following tagging than in the remaining of the season. TBF was

actually higher during the first three days, probably because the fish were finishing their

upstream migration, before settling near spawning grounds (unpublished radio tracking data).

This absence of negative impact suggests that external tagging under the dorsal fin, provided

fish are continually kept immerged and rapidly handled, is a suitable tagging technique for Allis

shad (and Jepsen et al. 2015 for other fishes; see also Breine et al. 2017 for Twaite shad).

613   On top of the estimation of energetic costs from dynamic acceleration, the continuous

estimation of body roundness from static acceleration is a promising yet perfectible application.

Gravitational acceleration is commonly used to infer the posture of the animal, assuming that

the position of the accelerometer on the animal is constant (Brown et al. 2013). Here, assuming

that shad stayed upright for most of the time, gravitational acceleration was used as an indicator

of change in body shape. In particular, we hypothesized that the angle θ between the x axis of

the accelerometer and the vertical may indicate body roundness. For all individuals, θ was

higher at the beginning of the season than at the end, when the fish had thinned, and globally

decreased along the spawning season. The accelerometer was attached under the dorsal fin,

where the cross-section is the broadest, which probably maximized the sensitivity of

acceleration data to fish thinning. Furthermore, shad mainly store energy for migration and

spawning as subdermal fat and interstitial fat in the white muscle (Leonard and McCormick

1999), whereas the limited consumption of visceral and liver fat may not have affected θ.

However, interindividual variability in θ was not correlated to interindividual variability in

body condition, and the shift in θ was not correlated to individual weight loss during the

spawning season. This could be due to interindividual differences in the exact position of the

accelerometers on the fish, or in the relationship between body condition and roundness. Indeed,

weight loss due to gamete expulsion was probably undetectable by the accelerometer attached

to the back of the fish. Given that ovaries can represent up to 15% of a shad weight at arrival

on spawning grounds (Cassou-Leins and Cassou-Leins 1981; Taverny 1991), a significant loss

of weight was probably not reflected in the rotation of the accelerometer across the spawning

season. In fact, shift of θ across eight hour periods increased with neither TBF, temperature,

nor estimated energy expenditure, contrary to what was expected from the relationship linking

TBF and temperature to energy consumption in American shad (Leonard et al. 1999).

Additionally, θ increased during some periods (especially nights), and fluctuated a lot

especially at the end of the season, so it was clearly affected by other processes than fish

thinning. In particular, while the attachment wires were tightened during the tagging procedure,

fish thinning loosened them so the accelerometer must have jiggled more and more as fish

thinned, thereby increasing the noise in acceleration data. Hence, using the angle of the

accelerometer to quantitatively track change in weight or body roundness will certainly require

further tuning, such as laboratory experiment with repeated weighing and image analysis of the

fish's cross-section, and drastic standardization of attachment procedure, but we consider it a

promising method to monitor individual condition in the field. Once refined, this approach

could be used to detect the many ecologically or behaviourally relevant changes in animal shape

beyond thinning due to energy consumption, such as parturition, massive food intake in large

predators (Cuyper et al. 2019), inflammatory swelling (Duncan et al. 2016), or inflation as a

courtship or defence behaviour (Wainwright and Turingan 1997). Combined with dynamic

acceleration, such data could be used to test whether these changes in shape are associated to

global activity or to the expression of specific behaviours.

Acknowledgements

We thank B. Liquet, F. Luthon, B. Larroque and E. Bouix for their help in the initial exploration

of data, and J. Leonard and T. Castro-Santos for sharing their data on American shad. All

experimental procedures comply with French and European legislation, and were approved by

the legal representative (prefectural decree #64-2017-04-25-004) and the ethical committee for

birds and fishes in the French region Nouvelle Aquitaine (authorization #2016020116037869).

This study was financed by the French Office for Biodiversity, the Adour-Garonne Water

Agency, and the French region Nouvelle Aquitaine.

References

Acolas M, Veron V, Jourdan H, Begout M, Sabatie M, Bagliniere JL. 2006. Upstream

migration and reproductive patterns of a population of allis shad in a small river (L'Aulne,

Brittany, France). doi:DOI:10.1016/j.icesjms.2005.05.022.

Acolas ML, Anras MLB, Véron V, Jourdan H, Sabatié MR, Baglinière JL. 2004. An assessment

of the upstream migration and reproductive behaviour of allis shad (*Alosa alosa* L.) using

acoustic tracking. ICES J Mar Sci. 61(8):1291–1304. doi:10.1016/j.icesjms.2004.07.023.

Anderson SS, Fedak MA. 1985. Grey seal males: energetic and behavioural links between size

and sexual success. Animal Behaviour. 33(3):829–838. doi:10.1016/S0003-3472(85)80017-8.

Baglinière J-L, Elie P. 2000. Les Aloses (*Alosa alosa* et *Alosa fallax* spp.): écobiologie et

variabilité des populations. Editions Quae.

Barton K. 2009. MuMIn: Multimodel inference. https://cran.r-

project.org/web/packages/MuMIn/index.html.

Bates D, Maechler M, Bolker B, Walker S. 2014. lme4: Linear mixed-effects models using

Eigen and S4.

Beamish FWH, Potter IC, Thomas E. 1979. Proximate composition of the adult anadromous

sea lamprey, *Petromyzon marinus,* in relation to feeding, migration and reproduction. Journal

of Animal Ecology. 48(1):1–19. doi:10.2307/4096.

Bengen GSH. 1992. Suivi de la maturation gonadique des aloses, *Alosa alosa* L. lors de leur

migration en Garonne. PhD Thesis, Toulouse, France: Institut national polytechnique.

Bonnet X, Bradshaw D, Shine R. 1998. Capital versus income breeding: an ectothermic

perspective. Oikos. 83(2):333–342.

Borgia G. 1985. Bower quality, number of decorations and mating success of male satin

bowerbirds (*Ptilonorhynchus violaceus*): an experimental analysis. Animal Behaviour.

33(1):266–271. doi:10.1016/S0003-3472(85)80140-8.

Breine J, Pauwels IS, Verhelst P, Vandamme L, Baeyens R, Reubens J, Coeck J. 2017.

Successful external acoustic tagging of twaite shad *Alosa fallax* (Lacépède 1803). Fisheries

Research. 191(Supplement C):36–40. doi:10.1016/j.fishres.2017.03.003.

Brett JR, Groves TDD. 1979. Physiological energetics. In: Hoar WS, Randall DJ, Brett JR,

editors. Bioenergetics and growth. Vol. 8. Academic Press. New York. (Fish physiology). p.

279–352.

Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, Skaug HJ,

Mächler M, Bolker BM. 2017. glmmTMB balances speed and flexibility among packages for

zero-inflated generalized linear mixed modeling. The R Journal. 9(2):378–400.

Brown DD, Kays R, Wikelski M, Wilson R, Klimley AP. 2013. Observing the unwatchable

through acceleration logging of animal behavior. Animal Biotelemetry. 1(1):1–16.

doi:10.1186/2050-3385-1-20.

Casas J, Pincebourde S, Mandon N, Vannier F, Poujol R, Giron D. 2005. Lifetime nutrient

dynamics reveal simultaneous capital and income breeding in a parasitiod. Ecology. 86(3):545–

554. doi:10.1890/04-0812.

Cassou-Leins F, Cassou-Leins J-J. 1981. Recherches sur la biologie et l'halieutique des
migrateurs de la Garonne et principalement l'alose : *Alosa alosa*. PhD thesis. Institut National
Polytechnique de Toulouse.

Castro-Santos T, Letcher BH. 2010. Modeling migratory energetics of Connecticut River
American shad (*Alosa sapidissima*): implications for the conservation of an iteroparous
anadromous fish. Can J Fish Aquat Sci. 67(5):806–830. doi:10.1139/F10-026.

Chanseau M, Castelnaud G, Carry L, Martin-Vandembulcke D, Belaud A. 2004. Essai
d'évaluation du stock de géniteurs d'alose *Alosa alosa* du bassin versant Gironde-Garonne-
Dordogne sur la période 1987-2001 et comparaison de différents indicateurs d'abondance.
Bulletin Français de la Pêche et de la Pisciculture.(374):1–19. doi:10.1051/kmae:2004023.

Collins PM, Halsey LG, Arnould JPY, Shaw PJA, Dodd S, Green JA. 2016. Energetic
consequences of time-activity budgets for a breeding seabird. J Zool. 300(3):153–162.
doi:10.1111/jzo.12370.

Craig JF, Kenley MJ, Talling JF. 1978. Comparative estimations of the energy content of fish
tissue from bomb calorimetry, wet oxidation and proximate analysis. Freshwater Biology.
8(6):585–590. doi:10.1111/j.1365-2427.1978.tb01480.x.

Csilléry K, Blum MGB, Gaggiotti OE, François O. 2010. Approximate Bayesian Computation
(ABC) in practice. Trends in Ecology & Evolution. 25(7):410–418.
doi:10.1016/j.tree.2010.04.001.

Cuyper AD, Clauss M, Carbone C, Codron D, Cools A, Hesta M, Janssens GPJ. 2019. Predator
size and prey size–gut capacity ratios determine kill frequency and carcass production in
terrestrial carnivorous mammals. Oikos. 128(1):13–22. doi:10.1111/oik.05488.

Duncan AE, Torgerson-White LL, Allard SM, Schneider T. 2016. An evaluation of infrared

thermography for detection of bumblefoot (pododermatitis) in penguins. J Zoo Wildl Med.

47(2):474–485. doi:10.1638/2015-0199.1.

ECP.   2018.   Ecology   and   fish   population   biology   facility.

doi:10.15454/1.5572402068944548E12. http://cnue-pierroton.pierroton.inra.fr:5000/dataset/i-

e-ecologie-comportementale-des-poissons.

Emlen ST, Oring LW. 1977. Ecology, sexual selection, and the evolution of mating systems.

Science. 197(4300):215–223. doi:10.1126/science.327542.

Fatin D, Dartiguelongue J. 1996. Etude préliminaire de la reproduction des aloses en 1995 entre

Tuilières et Mauzac sur la Dordogne.

Føre M, Svendsen E, Økland F, Gräns A, Alfredsen JA, Finstad B, Hedger R, Uglem I. 2020

Apr 21. Heart rate and swimming activity as indicators of post-surgical recovery time of

Atlantic salmon (*Salmo salar*). doi:10.21203/rs.3.rs-23506/v1.

Gauthey Z, Freychet M, Manicki A, Herman A, Lepais O, Panserat S, Elosegi A, Tentelier C,

Labonne J. 2015. The concentration of plasma metabolites varies throughout reproduction and

affects offspring number in wild brown trout (*Salmo trutta*). Comparative Biochemistry and

Physiology   Part   A: Molecular   &   Integrative   Physiology.   184:90–96.

doi:10.1016/j.cbpa.2015.01.025.

Groscolas R, Viera V, Guerin N, Handrich Y, Côté SD. 2010. Heart rate as a predictor of energy

expenditure in undisturbed fasting and incubating penguins. Journal of Experimental Biology.

213(1):153–160. doi:10.1242/jeb.033720.

Heimpel GE, Rosenheim JA. 1998. Egg limitation in parasitoids: A review of the evidence and

a case study. Biological Control. 11(2):160–168. doi:10.1006/bcon.1997.0587.

Hendry AP, Beall E. 2004. Energy use in spawning Atlantic salmon. Ecology of Freshwater

Fish. 13(3):185–196. doi:10.1111/j.1600-0633.2004.00045.x.

Hendry AP, Berg OK. 1999. Secondary sexual characters, energy use, senescence, and the cost

of reproduction in sockeye salmon. Can J Zool. 77(11):1663–1675. doi:10.1139/z99-158.

Hicks O, Burthe S, Daunt F, Butler A, Bishop C, Green JA. 2017. Validating accelerometry

estimates of energy expenditure across behaviours using heart rate data in a free-living seabird.

Journal of Experimental Biology.:jeb.152710. doi:10.1242/jeb.152710.

Hruska KA, Hinch SG, Patterson DA, Healey MC. 2011. Egg retention in relation to arrival

timing and reproductive longevity in female sockeye salmon (*Oncorhynchus nerka*). Can J Fish

Aquat Sci. 68(2):250–259. doi:10.1139/F10-153.

Hughes PW. 2017. Between semelparity and iteroparity: Empirical evidence for a continuum

of modes of parity. Ecol Evol. 7(20):8232–8261. doi:10.1002/ece3.3341.

Idler DR, Bitners I. 1958. Biochemical studies on Sockeye salmon during spawning migration:

Ii. Cholesterol, fat, protein, and water in the flesh of standard fish. Can J Biochem Physiol.

36(8):793–798. doi:10.1139/o58-084.

Jepsen N, Thorstad EB, Havn T, Lucas MC. 2015. The use of external electronic tags on fish:

an evaluation of tag retention and tagging effects. Animal Biotelemetry. 3(1):49.

doi:10.1186/s40317-015-0086-z.

Jonsson N, Jonsson B, Hansen LP. 1997. Changes in proximate composition and estimates of

energetic costs during upstream migration and spawning in Atlantic salmon *Salmo salar*.

Journal of Animal Ecology. 66(3):425–436. doi:10.2307/5987.

Kieffer JD. 2000. Limits to exhaustive exercise in fish. Comp Biochem Physiol, Part A Mol

Integr Physiol. 126(2):161–179.

Kirkendall LR, Stenseth NChr. 1985. On defining 'breeding once'. The American Naturalist.

125(2):189–204.

Langkau MC, Clavé D, Schmidt MB, Borcherding J. 2016. Spawning behaviour of Allis shad

Alosa alosa: new insights based on imaging sonar data. J Fish Biol. 88(6):2263–2274.

doi:10.1111/jfb.12978.

Leonard JBK, McCormick SD. 1999. Effects of migration distance on whole-body and tissue-

specific energy use in American shad (*Alosa sapidissima*). Can J Fish Aquat Sci. 56(7):1159–

1171. doi:10.1139/f99-041.

Leonard JBK, Norieka JF, Kynard B, McCormick SD. 1999. Metabolic rates in an anadromous

clupeid, the American shad (*Alosa sapidissima*). J Comp Physiol B. 169(4–5):287–295.

doi:10.1007/s003600050223.

Mennesson-Boisneau C, Boisneau P. 1990. Migration, répartition, reproduction,

caractéristiques biologiques et taxonomie des aloses (*Alosa sp*) dans le bassin de la Loire. PhD

thesis. Université de Rennes et Paris XII.

Nagy KA, Girard IA, Brown TK. 1999. Energetics of free-ranging mammals, reptiles, and birds.

Annu Rev Nutr. 19:247–277. doi:10.1146/annurev.nutr.19.1.247.

Olney JE, Denny SC, Hoenig JM. 2001. Criteria for determining maturity stage in female

American shad, Alosa sapidissima, and a proposed reproductive cycle. Bull Fr Pêche

Piscic.(362–363):881–901. doi:10.1051/kmae:2001025.

Paumier A, Drouineau H, Carry L, Nachón DJ, Lambert P. 2019. A field-based definition of

the thermal preference during spawning for allis shad populations (*Alosa alosa*). Environmental

Biology of Fishes. 102(6):845–855. doi:10.1007/s10641-019-00874-7.

Petersson E. 2001. 'False orgasm' in female brown trout: trick or treat? Animal Behaviour.

61(2):497–501. doi:10.1006/anbe.2000.1585.

Pianka ER. 1976. Natural Selection of Optimal Reproductive Tactics. American Zoologist.

16(4):775–784.

R Development Core Team. 2008. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org.

Rands SA, Houston AI, Cuthill IC. 2006. Measurement of mass change in breeding birds: A

bibliography and discussion of measurement techniques. Ringing & Migration. 23(1):1–5.

doi:10.1080/03078698.2006.9674337.

Robertson DR. 1991. The Role of Adult Biology in the Timing of Spawning of Tropical Reef

Fishes. In: Sale PF, editor. The Ecology of Fishes on Coral Reefs. San Diego: Academic Press.

p. 356–386.

Roff D. 1993. Evolution of life histories: Theory and analysis. Springer Science & Business

Media.

Schultz DL. 1999. Comparison of lipid levels during spawning in annual and perennial darters

of the subgenus *Boleosoma*, *Etheostoma perlongum*, and *Etheostoma olmstedi*. Copeia.

1999(4):906–916. doi:10.2307/1447966.

Stearns SC. 1992. The evolution of life histories. OUP Oxford.

Steinbach P, Gueneau P, Autuoro A, Broussard D. 1986. Radio-pistage de grandes aloses

adultes en Loire. Bulletin Français de la Pêche et de la Pisciculture.(302):106–117.

doi:10.1051/kmae:1986007.

Taverny C. 1991. Contribution à la connaissance de la dynamique des populations d'aloses :

Alosa alosa et *Alosa fallax* dans le système fluvio-estuarien de la Gironde : pêche, biologie et

écologie : étude particulière de la dévalaison et de l'impact des activités humaines. PhD thesis.

Bordeaux 1.

Tétard S, Feunteun E, Bultel E, Gadais R, Bégout M-L, Trancart T, Lasne E. 2016. Poor oxic

conditions in a large estuary reduce connectivity from marine to freshwater habitats of a

diadromous   fish.   Estuarine,   Coastal   and   Shelf   Science.   169:216–226.

doi:10.1016/j.ecss.2015.12.010.

Verdeyroux P, Guerri O, Chanseau M, Cazeaux J, Fauvel F, Bogun F, Desmoulin A, Tarrene

C, Nicole T, Dubois A, et al. 2015. Radio-telemetry study of the Allis shad (*Alosa alosa*)

migration at Bergerac and Tuilières along the Dordigne river and at Golfech along the Garonne

river from 2011 to 2014. Epidor.

Wainwright PC, Turingan RG. 1997. Evolution of Pufferfish Inflation Behavior. Evolution.

51(2):506–518. doi:10.1111/j.1558-5646.1997.tb02438.x.

Wilson RP, White CR, Quintana F, Halsey LG, Liebsch N, Martin GR, Butler PJ. 2006. Moving

towards acceleration for estimates of activity-specific metabolic rate in free-living animals: the

case of the cormorant. J Anim Ecol. 75(5):1081–1090. doi:10.1111/j.1365-2656.2006.01127.x.

Yamazaki C, Koizumi I. 2017. High frequency of mating without egg release in highly

promiscuous nonparasitic lamprey *Lethenteron kessleri*. J Ethol. 35(2):237–243.

doi:10.1007/s10164-017-0505-0.