Length regulation of multiple flagella that self-assemble from a shared pool of components

Thomas G. Fai[1] **, Lishibanya Mohapatra**[2] **, Jane Kondev**[2] **, Ariel Amir**[3*]

*For correspondence:
arielamir@seas.harvard.edu (AA)

[1] Brandeis University, Department of Mathematics, Waltham, MA 02454, USA; [2] Brandeis University, Department of Physics, Waltham, MA 02454, USA; [3] Harvard University, Paulson School of Engineering and Applied Sciences, Cambridge MA 02138, USA

Abstract   Control of organelle size is a problem that has intrigued cell biologists for at least a century. The single-celled green algae *Chlamydomonas reinhardtii* with its two flagella has proved to be a very useful model organism for studies of size control. Numerous experiments have identified motor-driven transport of tubulin to the growing ends of microtubules at the tip of the flagella as the key component of the machinery responsible for controlling their length. Here we consider a model of flagellar length control whose key assumption is that proteins responsible for the intraflagellar transport (IFT) of tubulin are present in limiting amounts. We show that this limiting-pool assumption and simple reasoning based on the law of mass action leads to an inverse relationship between the rate at which a flagellum grows and its length, which has been observed experimentally, and has been shown theoretically to provide a mechanism for length control. Experiments in which one of the two flagella are severed have revealed the coupled nature of the growth dynamics of the two flagella, and we extend our length-control model to two flagella by considering different mechanisms of their coupling. We describe which coupling mechanisms are capable of reproducing the observed dynamics in severing experiments, and why some that have been proposed previously are not. Within our theoretical framework we conclude that if tubulin and IFT proteins are freely exchanged between flagella simultaneous length control is not possible if the disassembly rate is constant. However, if disassembly depends on the concentration of IFT proteins at the tip of the flagellum, simultaneous length control can be achieved. Finally, we make quantitative predictions for experiments that could test this model.

Introduction

The size regulation of cellular organelles is a fundamental problem in biology *Marshall* (*2016*); *Milo and Phillips* (*2015*). For example, nuclear size is tightly coupled with cell size across a wide range of species *Hara and Merten* (*2015*) and the loss of this coupling mechanism is implicated in various types of cancer *Zink et al.* (*2004*).

A striking example of organelle size control in eukaryotes is the single-celled algae *Chlamydomonas reinhardtii* (*Figure 1*(a)), which uses two flagella to move through its aqueous environment. The lengths of these flagella are tightly controlled; mutants with longer flagella have decreased swimming velocities and beat frequencies *Khona et al.* (*2013*) compared to wild type cells and mutants with unequal flagellar lengths are observed to spin around in circles *Tam et al.* (*2003*).

It is known from experiments on *Chlamydomonas Rosenbaum et al.* (*1969*) that (i) its two flagella reach steady-state lengths of about 10 μm, and that (ii) the flagellar lengths are correlated since, if

Figure 1. (a) Electron microscopy images of the biflagellate green algae *Chlamydomonas* and its flagella captured by Elisa Vannuccini and Pietro Lupetti (University of Siena, Italy) and reproduced from *Morga and Bastin* (*2013*) (Creative Commons Attribution License CC BY). The inset shows the whole organism (scale bar 5 μm) and the close-up shows the flagellar basal body (BB), transition zone (TZ), and cell wall (CW) (scale bar 1 μm). (b) Severing experiments: after one flagellum is severed, the two flagella equalize at a shorter length and then grow together to the original steady-state length. This is demonstrated in experimental data from *Ludington et al.* (*2012*) courtesy of William Ludington.

one flagellum is severed, the remaining flagellum shortens until it reaches the length of the growing,
previously severed flagellum (*Figure 1*(b)). This latter protocol is typically referred to as a "severing"
or "long-zero" experiment.

A key process contributing to the assembly of flagella is the continual transport of proteins
from the flagellar base to tip and back. The original evidence for this intraflagellar transport (IFT)
was provided around 25 years ago by experimental observations in *Chlamydomonas* of particles
moving processively along the flagellum at constant speed *Kozminksi et al.* (*1993*). In the time since,
a significant body of work has emerged with the goal of understanding the many proteins and
biochemical pathways that coordinate this complex process in *Chlamydomonas* and other organisms
such as *C. elegans*, as described in several review articles *Prevo et al.* (*2017*); *Scholey* (*2008*, 2003);
Cole (*2003*); *Rosenbaum et al.* (*1999*); *Rosenbaum and Witman* (*2002*); *Rosenbaum et al.* (*1999*).

Already at the time of its discovery, IFT was hypothesized to play a role in flagellar length
control by transporting proteins required for assembly at the flagellar tip *Kozminksi et al.* (*1993*).
Subsequent work has showed that the proteins in the flagellum are continually exchanged with
those in the basal body (*Figure 1*(a)) *Song and Dentler* (*2001*); *Marshall and Rosenbaum* (*2001*);
Buisson et al. (*2013*). IFT particles containing tubulin are transported along the flagellum by two
different motor proteins: kinesin-2 transports IFT particles from the flagellar base to tip (the
anterograde direction) whereas dynein carries IFT particles from the tip to the base (the retrograde
direction).

These observations motivated the development of mathematical models of flagellar length
dynamics such as the balance point model *Marshall and Rosenbaum* (*2001*); *Marshall et al.* (*2005*).
In the balance point model, the steady-state flagellar length is achieved when the assembly and
disassembly processes, which are in continual competition, come into a balance set by the finite pool
of tubulin shared between the flagellum and basal body. The balance point model yields a length-
dependent assembly rate by assuming a constant number of transport complexes moving along the
flagellum. This is consistent with experimental evidence that the flagellar assembly rate decreases
with length *Rosenbaum and Child* (*1967*); *Marshall and Rosenbaum* (*2001*). Notably, however,
the balance-point does not explain how the cell could maintain a constant number of transport

complexes on the flagellum. Identifying the underlying mechanism of length control remains an
open problem, and a subsequent theoretical studies proposed several potential mechanisms that
give rise to length-dependent assembly rates *Ludington et al.* (*2015*); *Hendel et al.* (*2018*).

In this work, we propose a simple theoretical framework for the dynamics of flagellar assembly
based on the assumption that one or more IFT proteins are present in a limiting amount. Our
model captures the essential features of IFT, namely the transport of cargo by molecular motors,
and it includes the balance point model as a limiting case. For concreteness, we will assume in our
presentation that the rate-limiting protein is a molecular motor; however, the resulting formulas
are valid for any rate-limiting IFT protein, as made precise in the Discussion.

We perform a detailed theoretical analysis of the resulting equations in the case of two flagella,
considering various possible scenarios for how the finite pools of molecular motors and tubulin
may be shared. Whereas a single flagellum reaches a well-defined steady state given finite pools of
motors and tubulin, we find that if all pools of biomolecules are shared between multiple flagella
and the disassembly rate is length-independent, length control is not achieved regardless of the
specific transport mechanisms or model parameters.

To recover simultaneous length control of multiple flagella in the context of a constant disas-
sembly rate, we first explore models in which either the motors or tubulin are shared between
flagella, but not both. Supported by experimental observations of microtubule-depolymerizing
proteins *Piao et al.* (*2009*); *Luo et al.* (*2011*); *Hilton et al.* (*2013*), we subsequently consider a model
in which the disassembly rate is dependent on the local concentration of a depolymerizing IFT
protein. Overall, our exploration results in five viable models of flagellar length control (*Figure 6*)
that each generalize to the case of more than 2 flagella while maintaining their salient features.

Four of the five viable models require that the flagella have separate pools for at least one
biomolecule. Such a barrier to exchange seems incompatible with the biological evidence of diffusive
coupling between basal bodies. It is possible, however, that some proteins are exchanged while
others (e.g. tubulin or motors) are not. The fifth viable model, which includes length-dependent
depolymerization as an essential ingredient (caused by diffusion and transport within the flagellum
of a depolymerizing factor such as kinesin-13 *Piao et al.* (*2009*)), allows for all biomolecules to be
exchanged between flagella. Depending on the diffusion constant, the assembly rate in this fifth
viable model may scale with inverse length as reported in the experimental literature *Marshall and*
Rosenbaum (*2001*). In the Discussion, we comment further on the merits of the viable models given
the currently available experimental data.

For the five viable models analyzed in this study, we consider potential experimental methods of
model validation and/or discrimination. Many cellular processes, including intraflagellar transport,
are inherently noisy, and the fluctuations in flagellar length contain valuable information about the
underlying biomolecular processes *Amir and Balaban* (*2018*). The essential model parameters may
be inferred from measurements of the steady-state length and the fluctuations about steady-state.
We use Monte Carlo simulations to estimate the resolution and quantity of time series data that
is required for statistically robust data fitting. These estimates may provide useful guidelines for
future experiments.

Results

Single flagellum

The key biochemical variables represented in our model are the tubulin dimers that form the
flagellar microtubules and the molecular motors kinesin and dynein that transport IFT particles
from the base to the tip and back. IFT particles combine in the basal body with kinesin, tubulin and
dynein to form a complex that is injected into the flagellum *Cole et al.* (*1998*). See the schematic of
Figure 2(a).

We assume that the injection of IFT particles into the flagellum is rate-limited by the supply
of molecular motors. The total number of molecular motors is assumed to be conserved, and

Figure 2. Single flagellar growth dynamics in *Chlamydomonas* in the case of diffusive return of kinesin. (a) IFT particles assemble with kinesin, dynein, and tubulin in the basal body and are injected into the flagellum. In steady-state, the fluxes balance and the injection flux J_i, ballistic IFT flux J_b, and diffusive IFT flux J_d are all equal. (b) Numbers of molecular motors undergoing ballistic and diffusive motion on the flagellum at a given length, obtained using the parameter values in *Table 1*. There is a crossover at $L \approx 1.5$ µm (inset) after which diffusive motion is dominant. (c) IFT particle flux vs. length for several choices of the on-rate of molecular motors k_{on} (see (3)). (d) Model simulations show that an initially severed flagellum grows until it reaches steady-state, consistent with the behavior observed experimentally in *Ludington et al.* (*2012*). The shaded red area illustrates the mean and standard deviation computed from 50 realizations of the Langevin equation (see (29) in Materials and methods).

we consider the case that the rate-limiting motor undergoes ballistic motion only (as for dynein
Toropova et al. (*2017*)) and the case that the rate-limiting motor undergoes ballistic transport in
the anterograde direction and diffusive motion in the retrograde direction (as for kinesin *Chien*
et al. (*2017*)). The total amount of tubulin is also conserved in our model. The flagellar assembly
rate, i.e. the rate of addition of tubulin dimers to the microtubule ends at the tip of the flagellum,
is assumed proportional to the flux of IFT particles from the basal body into the flagellum times
the amount of free tubulin in the basal body. This assumption is a consequence of mass-action
kinetics of IFT particle assembly. (Note that we consider the cell volume to be fixed, in which case
the amount and concentration of a given biomolecule may be used interchangeably.)

The ballistic and diffusive IFT dynamics are fast compared to changes in length. The timescale
of flagellar length dynamics e.g. the recovery time after severing is of the order of 10 minutes,
whereas the molecular motors involved in IFT take at most tens of seconds to traverse the length
of the flagellum by either ballistic or diffusive motion. Based on this separation of timescales, we
follow *Hendel et al.* (*2018*) by neglecting the time delay between injection and tubulin delivery and
by treating IFT as a quasi-steady state process.

We use a stochastic differential equation to model the flagellar length dynamics $L(t)$. We are motivated to include stochasticity because of experimental observations of flagellar length fluctuations and since measuring fluctuations about steady-state can be used to discriminate between candidate models. Our stochastic model follows *Wemmer and Marshall* (*2018*), a combined experimental and theoretical study that illustrates how noise in the flagellar length control system affects biological function. Whereas the theoretical component of this previous study focused on the balance point model, here we include fluctuations within a more general theoretical framework. We model the growth of a *single* flagellum by

$$\frac{\mathrm{d}L}{\mathrm{d}t} = \gamma J T_f - d + \eta, \tag{1}$$

where J is the flux of IFT particles (to be determined), T_f is the amount of free tubulin in the basal
body, d is the disassembly rate (assumed constant for now), and γ is a constant. To model the
molecular noise due to the inherent stochasticity of binding and unbinding events, we add a noise
term η. Mathematically, it represents white noise forcing of magnitude Υ, i.e. η is a random variable
with autocorrelation $\langle \eta(t)\eta(t') \rangle = \Upsilon^2 \delta(t - t')$. See the explanation preceding (29) in Materials and
methods for details.

In the context of our model, the quasi steady-state assumption implies an injection rate J that exactly balances the ballistic and diffusive fluxes. As shown in (33) and (42) of Materials and methods, the fluxes satisfy

$$\begin{cases} J = \dfrac{k_{\mathrm{on}}M}{1 + 2k_{\mathrm{on}}L/v}, & \text{Ballistic return} \tag{2} \\[2ex] J = \dfrac{k_{\mathrm{on}}M}{1 + k_{\mathrm{on}}L/v + k_{\mathrm{on}}L^2/2D}, & \text{Diffusive return} \tag{3} \end{cases}$$

where k_{on} is the rate constant of motor injection, M is the total number of motors, v is the speed of
ballistically-moving motors in IFT, and D is the diffusion constant of motors in the flagellum (with
parameter values given in *Table 1*). Note that in (3) and throughout the manuscript, terms written
in the form x/yz are shorthand for $x/(yz)$, e.g. $k_{\mathrm{on}}L^2/2D$ is to be interpreted as $k_{\mathrm{on}}L^2/(2D)$.

Near the steady-state length in the case of diffusive return, we find that most motors on the
flagellum undergo diffusive motion, although at very small lengths ballistic motion is dominant
(*Figure 2*(b)). The flux decreases with length (*Figure 2*(c)). On a qualitative level, this decreasing
trend is supported by experimental measurements *Dentler* (*2005*), although more data is needed to
make a quantitative comparison. In the appropriate limits, this expression recovers existing theories
in which the length dependence goes purely as $1/L$ or $1/L^2$. In the case of ballistic return, or for
diffusive return in the limit $D \gg Lv$, the $1/L$ scaling of *Marshall and Rosenbaum* (*2001*) is recovered.

For diffusive return in the limit $D \ll Lv$, the $1/L^2$ scaling of **Hendel et al.** (**2018**) is recovered. A
distinction between our formula for the flux and those in the aforementioned paper is the presence
of the constant term in the denominator, which follows from injection according to the law of
mass action. Consequently, the flux does not become singular at $L = 0$ in our formulation. It is
interesting to note that the flux has a similar functional form to the familiar substrate production
rate in Michaelis-Menten enzyme kinetics **Fall** (**2002**); this is because of the separation of timescales
assumption invoked in both derivations.

Substituting the above expressions for the flux into the growth rate equation (1) results in

$$
\begin{cases}
\dfrac{\mathrm{d}L}{\mathrm{d}t} = \dfrac{\gamma k_{\mathrm{on}} M}{1 + 2k_{\mathrm{on}} L/v}(T - L) - d + \eta, & \text{Ballistic return,} & (4) \\[2ex]
\dfrac{\mathrm{d}L}{\mathrm{d}t} = \dfrac{\gamma k_{\mathrm{on}} M}{1 + k_{\mathrm{on}} L/v + k_{\mathrm{on}} L^2/2D}(T - L) - d + \eta, & \text{Diffusive return,} & (5)
\end{cases}
$$

157 where $T_f = T - L$ by conservation of the total tubulin pool T. Each of the above models is consistent
158 with the results of flagellum severing experiments, i.e. when the severing experiment is simulated by
159 setting the initial length to zero, the length increases until it reaches its steady-state L_{ss} (**Figure 2**(d)).
160 In the case of ballistic return, the steady-state length is given by

$$
L_{ss} = \left(T - \frac{d}{\gamma k_{\mathrm{on}} M} \right) \Big/ \left(1 + \frac{2d}{\gamma M v} \right). \tag{6}
$$

161 In the case of diffusive return, solving (5) for the steady-state results in a quadratic equation for L_{ss}.
162 One root is always negative, leaving the solution

$$
L_{ss} = \left(\frac{D}{v} + \frac{\gamma DM}{d} \right) \left(-1 + \sqrt{1 + \left(\frac{2dT}{\gamma MD} \right) \frac{1 - d/\gamma k_{\mathrm{on}} MT}{(1 + d/\gamma M v)^2}} \right), \tag{7}
$$

163 which is positive provided that $T > d/\gamma kM$. If $T \leq d/\gamma kM$, the flagellar length shrinks to zero.
164 (We reiterate the convention that terms such as $d/\gamma k_{\mathrm{on}} MT$ and $d/\gamma M v$ are to be interpreted as
165 $d/(\gamma k_{\mathrm{on}} MT)$ and $d/(\gamma M v)$, respectively.)

Expanding to first order in $\Delta L := L - L_{ss}$, we find that the steady-state is stable, i.e. $\mathrm{d}(\Delta L)/\mathrm{d}t = -\lambda(\Delta L)$ with $\lambda > 0$ given in (36) of Materials and methods. Based on the parameters estimated in
Appendix 1 for diffusive return, the associated timescale $\tau := 1/\lambda$ is approximately 15 min, which is
consistent with experiment. This timescale is long compared to the few tens of seconds needed
for molecular motors to traverse the flagellum in IFT, which justifies *a posteriori* our approximation
of IFT as a quasi-steady state process. We next consider the parameter space associated with
the length dynamics. Introducing the nondimensional length $\widetilde{L} = L/L_{ss}$, time $\widetilde{t} = t\gamma k_{\mathrm{on}} MT/L_{ss}$,
and random forcing $\widetilde{\eta} = \eta/\gamma k_{\mathrm{on}} MT$ we may rewrite (4) and (5) in terms of the nondimensional
parameters $\pi_1 = d/\gamma k_{\mathrm{on}} MT$, $\pi_2 = L_{ss}/T$, $\pi_3 = k_{\mathrm{on}} L_{ss}/v$, and $\pi_4 = k_{\mathrm{on}} L_{ss}^2/2D$ as

$$
\begin{cases}
\dfrac{\mathrm{d}\widetilde{L}}{\mathrm{d}\widetilde{t}} = \dfrac{1 - \pi_2 \widetilde{L}}{1 + 2\pi_3 \widetilde{L}} - \pi_1 + \widetilde{\eta}, & \text{Ballistic return} & (8) \\[2ex]
\dfrac{\mathrm{d}\widetilde{L}}{\mathrm{d}\widetilde{t}} = \dfrac{1 - \pi_2 \widetilde{L}}{1 + \pi_3 \widetilde{L} + \pi_4 \widetilde{L}^2} - \pi_1 + \widetilde{\eta}, & \text{Diffusive return} & (9)
\end{cases}
$$

We may interpret these numbers as follows: π_1 is the ratio of disassembly and assembly rates, π_2 is
the fraction of the tubulin pool taken up by the flagellum at steady-state, $\pi_3 = \tau_b/\tau_i$ is the ratio of the
ballistic timescale $\tau_b := L_{ss}/v$ of IFT transport to the timescale $\tau_i := k_{\mathrm{on}}^{-1}$ of injection, and $\pi_4 = \tau_d/\tau_i$ is
an analogous ratio involving the diffusive timescale $\tau_d = L_{ss}^2/2D$. (We could equivalently think of π_3
and π_4 as ratios of lengthscales related to the same physical processes.)
We next show that, in a special limit, the length dynamics with diffusive return are nearly equiv-
alent to the model of **Hendel et al.** (**2018**). In terms of the experimentally measured parameters
and those estimated in **Appendix 1**, we find $\pi_1 \approx 0.4$, $\pi_2 \approx 0.2$, $\pi_3 \approx 0.1$, and $\pi_4 \approx 0.8$. The relatively

Table 1. Parameter values and definitions

Symbol	Definition	Value	Units	References
Parameters				
L_{ss}	Steady-state length	10–12	μm	*Marshall and Rosenbaum* (*2001*); *Rosenbaum et al.* (*1969*)
T/N	Tubulin pool per flagellum	38–47	μm	*Marshall et al.* (*2005*)
d	Disassembly rate	0.5	μm/min	*Marshall and Rosenbaum* (*2001*); *Ludington et al.* (*2012*)
v	IFT speed	2.5–3	μm/s	*Kozminksi et al.* (*1993*); *Buisson et al.* (*2013*)
D	Diffusion coefficient	1.7	μm^2/s	*Chien et al.* (*2017*)
$\gamma k_{on} M/N$	Assembly rate per tubulin	2.3×10^{-2}–3.6×10^{-2}	min^{-1}	Fit
k_{on}	Injection rate constant	0.8–4.5	min^{-1}	Fit
γ	Prefactor in (1)	2.5×10^{-4}		Estimate (*Appendix 1*)
Variables				
N	Number of flagella			
T_f	Free tubulin		μm	
M	Total motors			
M_f	Free motors			
M_b	Ballistic motors			
M_d	Diffusing motors			
J	Flux		min^{-1}	
$c_d(x)$	Motor concentration		μm^{-1}	
$\overline{c_d}$	Average concentration		μm^{-1}	

174 small values of π_2 and π_3 suggest that it may be useful to consider the limit $\pi_2 \to 0$ (i.e. no tubulin
175 depletion) and $\pi_3 \to 0$ (i.e. instantaneous ballistic motion). In this limit, we have

$$\frac{\mathrm{d}L}{\mathrm{d}t} = \frac{\gamma k_{on} MT}{1 + k_{on} L^2/2D} - d + \eta, \tag{10}$$

176 recovering the model of *Hendel et al.* (*2018*) with the only distinction that, as mentioned above, in
177 our model there is an additional constant term in the denominator. In this limit, the steady-state
178 length satisfies

$$L_{ss} = \sqrt{\frac{2\gamma DMT}{d}} \sqrt{1 - \frac{d}{\gamma kMT}}, \tag{11}$$

179 which carries an extra factor of $\sqrt{1 - d/\gamma kMT}$ in comparison with the formula of *Hendel et al.*
180 (*2018*).

181 **Two flagella**

In the case of two flagella with lengths $L_1(t)$ and $L_2(t)$ the length dynamics are given by

$$\frac{\mathrm{d}L_1}{\mathrm{d}t} = \gamma J_1 T_{f,1} - d + \eta_1 \tag{12}$$

$$\frac{\mathrm{d}L_2}{\mathrm{d}t} = \gamma J_2 T_{f,2} - d + \eta_2, \tag{13}$$

where J_i and $T_{f,i}$ for $i = 1, 2$ denote the fluxes and free amounts of tubulin for the two flagella, and
η_1 and η_2 are independent white noise forcing. Severing experiments illustrate that the flagella are
coupled. We consider various modes of coupling, illustrated in *Figure 3*(a), that give rise within our
model to different forms of J_i and $T_{f,i}$ and investigate their consequences for length control by
focusing on the stability of solutions to the steady-state deterministic equations

$$0 = \gamma J_1 T_{f,1} - d \tag{14}$$

$$0 = \gamma J_2 T_{f,2} - d. \tag{15}$$

Figure 3. Length dynamics of two flagella assembling from shared pool of building blocks. The case of diffusive motor return is shown; similar results are obtained with ballistic return. (a) Flagellar assembly in *Chlamydomonas reinhardtii* and modes of coupling between basal bodies. (b) Simulations of severing experiment by models **Ø**, **M**, **T**, and **TM**. (i) In the case of separate pools of both motors and tubulin, the unsevered flagellum does not decrease in length. (ii) When only motors are shared, the flagella reach equal steady-state lengths (inset shows longer times). (iii) When only tubulin is shared, the two flagellar lengths reach a shared steady-state on a timescale consistent with experiment. (iv) When both tubulin and motors are shared, the lengths do not equalize. The difference between the two flagellar lengths undergoes a random walk, as shown by the mean square displacement (inset, average over 50 runs. The deviation from linearity is caused by the non-negativity of flagellar lengths, which limits the maximum difference.) In all the above simulations, to capture the loss of the tubulin from severing and the gradual recovery by intracellular regulation mechanisms, we initialize the tubulin pool with 14 μm less of tubulin and linearly increase it over the first 20 minutes to reach the steady-state value of T.

Tubulin separate, motors separate (∅)

The case of individual tubulin pools and individual motor pools leads to two uncoupled instances of (5), and therefore following the logic of the single flagellum case yields steady state lengths given by (7). While these uncoupled equations trivially yield simultaneous length control, they are fundamentally inconsistent with the coupling observed between flagella in severing experiments, and in particular the significant decrease in the length of the unsevered flagellum (*Figure 3*(b)(i)). Therefore we exclude this model as a viable candidate.

Tubulin separate, motors shared (**M**)

When motors are shared through a common pool, IFT particles are injected into either flagellum with equal probability. Therefore $J_1 = J_2 =: J$ and it can be shown by a straightforward generalization of (2) and (3) (see Model (**M**) in Materials and methods) that the flux satisfies

$$
\begin{cases}
J = \dfrac{k_{on}M/2}{1 + k_{on}(L_1 + L_2)/v}, & \text{Ballistic return} & (16) \\[3ex]
J = \dfrac{k_{on}M/2}{1 + k_{on}(L_1 + L_2)/2v + k_{on}(L_1^2 + L_2^2)/4D}, & \text{Diffusive return.} & (17)
\end{cases}
$$

The model yields simultaneous length control (*Figure 3*(b)(ii)). Because the tubulin pools are separate the steady-state equations are given by

$$0 = \gamma J(T - L_{1,ss}) - d \tag{18}$$

$$0 = \gamma J(T - L_{2,ss}) - d, \tag{19}$$

and it follows that $L_{1,ss} = L_{2,ss}$ after subtracting the above equations. With M rescaled to $M/2$, the steady-state equations are identical to the corresponding steady-state equations (4) and (5) for a single flagellum. Therefore the steady-state lengths for ballistic and diffusive return satisfy (6) and (7), respectively, upon rescaling $M \to M/2$. See Model (**M**) in Materials and methods for detailed derivations. The stability of the steady-state solution is proven in *Appendix 2* in the context of diffusive return. (The proof for ballistic return is essentially identical.)

Given the large amount of turnover and interaction between different cellular compartments, the assumption of separate tubulin pools in this model is questionable. There is experimental evidence that tubulin is exchanged between each flagellum and its basal body *Song and Dentler* (*2001*); *Marshall and Rosenbaum* (*2001*); *Buisson et al.* (*2013*). However, we cannot rule out the possibility of separate tubulin pools, as we are not aware of experimental evidence of tubulin exchange between flagella. Therefore we retain this as a viable model.

Tubulin shared, motors separate (**T**)

We next consider the case in which tubulin is shared but the motor pools are separate. The separate motor pools yield decoupled fluxes so that the flux is nearly identical to that of model ∅. However, in this case the equations are coupled through the shared tubulin pool term $T - L_1 - L_2$. The model equations (see (75)–(76) in Materials and methods) have a similar form to existing models *Marshall et al.* (*2005*); *Hendel et al.* (*2018*), in which the assembly rates involve a factor of $T - L_1 - L_2$ and either a $1/L_i$ or $1/L_i^2$-dependence in the denominator, for $i = 1, 2$. In fact, the present model recovers these two dependences on L_i in the appropriate limits, as discussed in Results in the context of a single growing flagellum.

As illustrated in *Figure 3*(b)(iii), this model yields simultaneous length control and the flagella reach their steady-state lengths on a timescale consistent with experiment. Based on this analysis of the **M** and **T** models, sharing tubulin *or* motors is in principle sufficient to attain simultaneous length control. This is independent of ballistic versus diffusive return of motors to the basal body. As mentioned above, in light of the constant turnover and biochemical interactions between different cellular compartments, the assumption of separate motor pools may be incorrect. Indeed, there

is preliminary evidence of IFT protein exchange between flagella based on experiments using photoconvertible fluorophores *Marshall* (*2018*). It is generally believed that both tubulin and motor pools are shared between flagella, and we next consider this case.

Tubulin shared, motors shared (**TM**)

The flux resulting from the shared motor assumption is the same as in (17), and we are left with the steady-state equations

$$0 = \gamma J(T - L_{1,ss} - L_{2,ss}) - d \tag{20}$$

$$0 = \gamma J(T - L_{1,ss} - L_{2,ss}) - d. \tag{21}$$

Because the two steady-state equations are identical, there is only a single equation for the two unknowns $L_{1,ss}$ and $L_{2,ss}$, which makes the lengths indeterminate. The model in which all biomolecules are shared between flagella does not yield simultaneous length control! There are infinitely many solutions to the equations, none of which is asymptotically stable (since performing a linear stability analysis corresponding yields a zero eigenvalue, as shown in *Appendix 2*). The same issue also appears in the dynamics; since $dL_1/dt = dL_2/dt$ this model cannot possibly explain the simultaneous positive and negative growth rates for the two flagella observed in the severing experiment.

Note that this conclusion is *independent* of the parameters. That is, regardless of whether the transport is diffusive, ballistic, or some combination, the model **TM** with shared tubulin and shared motors with constant disassembly does not yield simultaneous length control of two flagella (*Figure 3*(b)(iv)), and therefore is not a viable model.

The above analysis shows that to obtain length control in our framework under the assumption of constant disassembly, either tubulin or motors may be shared, but not both. However, the two viable models **M** and **T** identified above are incompatible with the widely-held belief that biomolecules are continually exchanged between flagella. It follows that either the conventional wisdom of shared pools is wrong or something is missing from the space of models considered. This motivates us to extend our study beyond the shared pool models considered so far.

Tubulin shared, motors shared and concentration-dependent disassembly (**TM***)

Figure 4. Simultaneous length control is achieved in a model with shared tubulin, shared motors, and a disassembly rate that depends on the concentration of molecular motors. Here we set $k_{on} = 4.5 \, min^{-1}$, $d_0 = 0.1$ µm/min, and d_1 is computed by (25) to obtain the observed steady-state length.

We next consider a model that allows for full biomolecule exchange and replaces the constant disassembly assumption with a concentration-dependent disassembly rate. The assumption of a constant rate of disassembly was based on experiments on mutants in which IFT was disabled *Marshall et al.* (*2005*). However, these experiments do not preclude motor-dependent disassembly since they were performed in the absence of motor dynamics. Indeed, subsequent experiments in organisms with intact IFT led to 50-fold greater disassembly rates than those measured in the absence of IFT *Ludington et al.* (*2012*).

Experimental observations that some kinesin species (e.g. kinesin-13) participate in microtubule disassembly *Piao et al.* (*2009*) provide a potential biochemical basis for IFT-dependent disassembly, and in what follows we take the disassembly rate to depend on motor concentration. We replace the constant disassembly by a disassembly rate of the form $d_0 + d_1 c_d(L)$,

where $c_d(L)$ is the concentration of diffusing motors at the tip of the flagellum. In general the disassembly rate may be a complicated function of concentration, and in that case the model is only appropriate near steady-state, where the first-order term in the Taylor expansion dominates.

Under the assumption of ballistic return this model does not yield length control; diffusive return is required to generate a non-constant concentration along the length of the flagellum. Assuming diffusive return the flux and concentration at the flagellar tip are related by $c_d(L) = JL/D$ (see (37) in Materials and methods), we may rewrite the governing equations as

$$\frac{dL_1}{dt} = \gamma J(T - L_1 - L_2) - d_0 - d_1 \frac{JL_1}{D} + \eta_1 \tag{22}$$

$$\frac{dL_2}{dt} = \gamma J(T - L_1 - L_2) - d_0 - d_1 \frac{JL_2}{D} + \eta_2, \tag{23}$$

where as before in the case of shared motors

$$J = \frac{k_{on}M/2}{1 + k_{on}(L_1 + L_2)/2v + k_{on}(L_1^2 + L_2^2)/4D}. \tag{24}$$

The concentration-dependent disassembly model with shared tubulin and shared motors (**TM***) exhibits simultaneous length control (*Figure 4*). Subtracting (23) from (22), it follows immediately that $L_{1,ss} = L_{2,ss} =: L_{ss}$, and solving for the steady-state results in

$$L_{ss} = \left(\frac{D}{v} + \frac{\gamma DM}{d_0} + \frac{Md_1}{2d_0} \right) \left(-1 + \sqrt{1 + \left(\frac{d_0 T}{\gamma MD} \right) \frac{1 - 2d_0/\gamma k_{on}MT}{\left(1 + d_0/\gamma Mv + d_1/2\gamma D \right)^2}} \right). \tag{25}$$

In the Appendix it is shown that this solution is stable. Therefore, concentration-dependent disassembly allows for simultaneous length control when all biomolecules are shared between flagella, and results in a third viable model. Diffusive return is a critical ingredient in model **TM***; with ballistic return it would not yield simultaneous length control. This is because, as remarked above, the non-constant diffusive motor concentration profile is necessary to make the growth rates of L_1 and L_2 distinct, which provides two independent equations for the steady-state lengths.

Generalization to $N > 2$ flagella

Taking inspiration from the eukaryotic parasite Giardia, which has four pairs of flagella *Dawson and House* (*2010*); *McInally and Dawson* (*2016*), we demonstrate the simultaneous length control attained for $N = 8$ flagella (*Figure 5*).

We may generalize the concentration-dependent disassembly model **TM*** to arbitrary flagellar number N. (The other viable models **T** and **M** may be generalized in a similar manner.) Because motors are shared, the injection fluxes are equal so that $J_i = J$ for all $i = 1 \ldots N$, with J satisfying

$$J = \frac{k_{on}M/N}{1 + k_{on}(\sum_{i=1}^{N} L_i)/Nv + k_{on}(\sum_{i=1}^{N} L_i^2)/2ND} \tag{26}$$

and length dynamics given by

$$\frac{dL_i}{dt} = \gamma J \left(T - \sum_{j=1}^{N} L_j \right) - d_0 - d_1 \frac{JL_i}{D} + \eta_i, \quad i = 1 \ldots N, \tag{27}$$

Figure 5. The viable models generalize to arbitrary flagellar number N. We demonstrate using the concentration-dependent disassembly model **TM*** with $N = 8$ flagella and the larger shared pool $T = 336$, $d_0 = 0.1$, and $d_1 = 120$; otherwise all parameters are as in the biflagellate case.

280 Taking any pairwise difference between the i^{th} and j^{th} equations at steady-state yields immediately

281 $L_{i,ss} = L_{j,ss}$, so that the steady-state lengths are equal to

$$L_{i,ss} = \left(\frac{D}{v} + \frac{\gamma DM}{d_0} + \frac{Md_1}{Nd_0} \right) \left(-1 + \sqrt{1 + \left(\frac{2d_0 T}{N\gamma MD} \right) \frac{1 - Nd_0/\gamma k_{\text{on}} MT}{\left(1 + d_0/\gamma Mv + d_1/N\gamma D \right)^2}} \right) \quad (28)$$

282 for all $i = 1, \ldots, N$. Stability follows from analyzing the linearized equations, as shown in *Appendix 2*.

Discussion

In this work, we explore models of flagellar length dynamics in the context of organelle size
regulation. We capture the essential aspects of IFT relevant to length control, i.e. the motor-driven
transport of tubulin across the flagellum. As in *Hendel et al.* (*2018*), since the timescale of transport
by IFT is a few seconds, whereas changes in flagellar length take place on a timescale of a few
minutes, we treat IFT as a quasi-steady-state process in which the anterograde and retrograde
fluxes of motors are balanced.

It has been well-established that the biomolecules represented by our model, i.e. tubulin and
molecular motors, are recycled within each flagellum and its basal body. There is also the possibility
of turnover *between* flagella. We have used our theoretical model to investigate the consequences
of sharing pools of tubulin and/or motors between flagella on their length dynamics. We take the
assembly rate to be proportional to the flux of IFT particles times the amount of free tubulin. In our
initial exploration, we take the disassembly rate to be constant. We find that sharing both tubulin
and motors leads to an indeterminate system of equations regardless of the details of the model,
whereas sharing either tubulin or motors, but not both, results in simultaneous length control of
both flagella. Interestingly, the model in which only tubulin is shared (model (**T**)) recovers essential
features of the balance point model *Marshall and Rosenbaum* (*2001*), in particular the decay with
$1/L$ in the growth rate.

This theoretical exploration illustrates the dramatic consequences in behavior that can occur
when biomolecules are shared between compartments, and highlights the importance of knowing
which proteins are exchanged between the basal bodies in the context of flagellar length control.
In particular, having a protein that is *not* exchanged can provide simultaneous length control by a
limited pool mechanism. Future experiments to identify which proteins are exchanged between
flagella, e.g. by tagging with photoconvertible fluorophores as in *Marshall* (*2018*), are needed to
assess the validity of this mechanism.

We investigated additional models that are consistent with the widely-held belief that biomolecules
are exchanged continually between flagella, e.g. in which both the motor and tubulin pools are
shared. In particular, we considered the possibility that disassembly is not constant but depends on
the local concentration of molecular motors. Because the diffusive gradient is linear at steady-state
and the basal body is taken to act as a diffusive sink, the concentration at the flagellar tip depends
on length. This leads to simultaneous length control regardless of whether the disassembly rate is
strongly or weakly dependent on concentration. We conclude that the title of *Hendel et al.* (*2018*) is
apt; the linear concentration gradient generated by diffusion indeed acts as a ruler! However, it
cannot function on its own. Diffusion must be combined with a mechanism such as concentration-
dependent disassembly; regardless of whether the motor dynamics are ballistic, diffusive, or some
combination, sharing all biomolecules does not yield length control within our theoretical frame-
work if the disassembly rate is constant. To put it another way, within the space of the five viable
models (*Figure 6*), diffusion is not essential for length control in the **M** and **T** models, whereas it *is*
essential for model **TM***.

Note that, although the present discussion has assumed that a molecular motor is the rate-
limiting IFT protein, our formulation is actually more general. If a non-motor IFT protein is rate-
limiting for injection, the same considerations and resulting formulas apply with the number of
motors replaced by the rate-limiting protein number and accounting for the relevant mode of

Viable models	Steady-state length and linearization parameters
🔴 Rate limiting protein (**M**) 🔸 Tubulin (**T**)	$$\frac{\mathrm{d}}{\mathrm{d}t}\begin{pmatrix}\Delta L_1\\\Delta L_2\end{pmatrix} = -\begin{pmatrix}a+b & a\\a & a+b\end{pmatrix}\begin{pmatrix}\Delta L_1\\\Delta L_2\end{pmatrix}+\begin{pmatrix}\eta_1\\\eta_2\end{pmatrix}$$

M (ballistic)

Tubulin separate, IFT protein shared

$$L_{ss} = \frac{T - 2d/(\gamma k_{\mathrm{on}}M)}{1 + 4d/(\gamma Mv)}$$

$$a = \frac{\gamma k_{\mathrm{on}}M(T - L_{ss})(k_{\mathrm{on}}/v)}{2(1 + 2k_{\mathrm{on}}L_{ss}/v)^2}$$

$$b = \frac{\gamma k_{\mathrm{on}}M/2}{1 + 2k_{\mathrm{on}}L_{ss}/v}$$

M (diffusive)

Tubulin separate, IFT protein shared

$$L_{ss} = \left(\frac{D}{v} + \frac{\gamma DM}{2d}\right)\left(-1 + \sqrt{1 + \left(\frac{4dT}{\gamma MD}\right)\frac{1 - 2d/\gamma k_{\mathrm{on}}MT}{(1 + 2d/\gamma Mv)^2}}\right)$$

$$a = \frac{\gamma k_{\mathrm{on}}M(T - L_{ss})(k_{\mathrm{on}}/2v + k_{\mathrm{on}}L_{ss}/2D)}{2(1 + k_{\mathrm{on}}L_{ss}/v + k_{\mathrm{on}}L_{ss}^2/2D)^2}$$

$$b = \frac{\gamma k_{\mathrm{on}}M/2}{1 + k_{\mathrm{on}}L_{ss}/v + k_{\mathrm{on}}L_{ss}^2/2D}$$

T (ballistic)

Tubulin shared, IFT protein separate

$$L_{ss} = \frac{T/2 - d/(2\gamma k_{\mathrm{on}}M)}{1 + d/(\gamma Mv)}$$

$$a = \frac{\gamma k_{\mathrm{on}}M}{1 + 2k_{\mathrm{on}}L_{ss}/v}$$

$$b = \frac{\gamma k_{\mathrm{on}}M(T - 2L_{ss})(2k_{\mathrm{on}}/v)}{(1 + 2k_{\mathrm{on}}L_{ss}/v)^2}$$

T (diffusive)

Tubulin shared, IFT protein separate

$$L_{ss} = \left(\frac{D}{v} + \frac{2\gamma DM}{d}\right)\left(-1 + \sqrt{1 + \left(\frac{dT}{2\gamma MD}\right)\frac{1 - d/\gamma k_{\mathrm{on}}MT}{(1 + d/2\gamma Mv)^2}}\right)$$

$$a = \frac{\gamma k_{\mathrm{on}}M}{1 + k_{\mathrm{on}}L_{ss}/v + k_{\mathrm{on}}L_{ss}^2/2D}$$

$$b = \frac{\gamma k_{\mathrm{on}}M(T - 2L_{ss})(k_{\mathrm{on}}/v + k_{\mathrm{on}}L_{ss}/D)}{(1 + k_{\mathrm{on}}L_{ss}/v + k_{\mathrm{on}}L_{ss}^2/2D)^2}$$

TM*(diffusive)

Tubulin shared, IFT protein shared
Diffusive return of depolymerizing IFT protein

$$L_{ss} = \left(\frac{D}{v} + \frac{\gamma DM}{d_0} + \frac{Md_1}{2d_0}\right)\left(-1 + \sqrt{1 + \left(\frac{d_0 T}{\gamma MD}\right)\frac{1 - 2d_0/\gamma k_{\mathrm{on}}MT}{(1 + d_0/\gamma Mv + d_1/2\gamma D)^2}}\right)$$

$$a = \frac{\gamma k_{\mathrm{on}}M/2}{1 + k_{\mathrm{on}}L_{ss}/v + k_{\mathrm{on}}L_{ss}^2/2D}\left(1 + \frac{(k_{\mathrm{on}}/2v + k_{\mathrm{on}}L_{ss}/2D)(T - (2 + d_1/\gamma D)L_{ss})}{1 + k_{\mathrm{on}}L_{ss}/v + k_{\mathrm{on}}L_{ss}^2/2D}\right)$$

$$b = \frac{d_1 k_{\mathrm{on}}M/2D}{1 + k_{\mathrm{on}}L_{ss}/v + k_{\mathrm{on}}L_{ss}^2/2D}$$

Figure 6. Our theoretical exploration results in five viable models

transport (i.e. ballistic or diffusive return). This abstracted view of the rate-limiting protein is represented schematically in *Figure 6*.

Diffusion is an essential ingredient for length control in model **TM***, and depending on the diffusion constant the growth rate decays as either $1/L$ or $1/L^2$. Using the diffusion constant measured for kinesin results in a $1/L^2$ decay; although the experimentally-determined growth rate was originally fit to a curve with $1/L$ decay *Marshall et al.* (*2005*), we find that fitting the same data to a curve with $1/L^2$ decay or to the second-order polynomial decay of model **TM*** lead to fits of comparable quality.

Among the viable models discussed here, only model **TM*** is consistent with both full biomolecule exchange and simultaneous length control. This model requires a rate-limiting IFT protein coupled to microtubule disassemby, shared between flagella, and diffusing back from the flagellar tip. Further experiments are needed to establish whether a protein meeting these criteria exists. One promising candidate is aurora-like kinase CALK, which has been shown to influence disassembly through its state of phosphorylation *Luo et al.* (*2011*); *Cao et al.* (*2013*). Another promising candidate is CNK2, a NIMA-related protein kinase known to localize to flagella *Bradley and Quarmby* (*2005*). It has been shown that a long-flagella Chlamydomonas mutant lacking a functional CNK2 exhibits decreased disassembly rates *Hilton et al.* (*2013*), so that this protein would also be compatible with the concentration-dependent disassembly assumption. Future experiments will determine whether flagellar lengths are controlled by the diffusion of CALK, CNK2, or some other protein, or whether additional physical mechanisms beyond those included in this theoretical framework are required to explain the observed phenomena.

On the side of theory, a promising avenue to discriminate between the potential models is by studying length fluctuations about steady-state. The fluctuation spectra of each model provides a signature that can be used to assess both the general model framework and to test the autocorrelation timescales predicted by each model *Amir and Balaban* (*2018*). In *Appendix 3* we have described the spectra predicted by the viable models. We have mimicked experiments by performing stochastic simulations with various sampling rates and durations to provide constraints on the experimental parameters necessary to obtain robust and statistically significant results through an analysis of fluctuations. We find that while significant amounts of data are needed in order to extract information from fluctuations, the sampling intervals and time durations needed to obtain 10% error are within the realm of experimental plausibility (see *Appendix 3*).

Materials and methods

We use a stochastic ordinary differential equation model for the flagellar length dynamics. The key biochemical variables represented in our model are the tubulin monomers that form the flagellar axoneme and the molecular motor that carries IFT particles along the flagellum.

Single flagellum

To introduce the model in a simpler setting, we consider first the case of a single flagellum with time-dependent length $L(t)$. In our model the flagellar assembly rate is proportional to the flux J of IFT particles times the size of an IFT particle, which is in turn proportional to the amount of free tubulin T_f in the basal body. In other words, once the flux of IFT motors has been established, forming a type of conveyor belt, tubulin dimers have a constant probability per unit time of joining with a motor, i.e. of jumping onto the conveyor belt. (Equivalently, one may think of a sequence of biochemical reactions: first, IFT particles are generated at a rate J, and second, tubulin forms a complex with IFT at a rate proportional to JT_f.) We take the disassembly rate to be equal to a constant d for now. Finally, to account for random fluctuations we include a white noise forcing term η with autocorrelation $\langle \eta(t)\eta(t') \rangle = |\Upsilon|^2\,\delta(t-t')$, where $\delta(t-t')$ represents the Dirac delta function. This yields a Langevin equation for the growth rate:

$$\frac{\mathrm{d}L}{\mathrm{d}t} = \gamma J T_f - d + \eta. \tag{29}$$

We assume that the rate of injection of IFT particles into the flagellum is rate-limited by the supply of
molecular motors. The molecular motors are assumed to be conserved, and we consider two cases
separately: first the case in which the rate-limiting IFT motor undergoes ballistic motion in both
directions on the flagellum, and second the case in which the motor undergoes ballistic motion
from the flagellar base to tip and diffusive motion in the retrograde direction. Note that, although
there is evidence that IFT loading occurs in trains of various sizes *Wren et al.* (*2013*); *Ludington*
et al. (*2013*), we have not included differential loading into our model as the simpler model in which
dimers load independently is sufficient for agreement with the severing experiments.

Ballistic return

Motors are conserved having total number $M = M_f + M_b$, where M_f is the number of motors freely
available in the basal body and M_b is the number of motors moving ballistically on the flagellum in
IFT particles. In our model the flux, or injection rate, is proportional to the number of free molecular
motors M_f so that $J = k_{on}M_f$ according to mass action kinetics, with first-order rate constant k_{on}.
We assume tubulin is conserved with total amount $T = T_f + T_b + L$, where T_f is the amount of
tubulin freely available in the basal body, T_b is the amount moving ballistically while attached to IFT
particles, and L is the amount incorporated in the flagellum. As the flagellum grows it incorporates
more tubulin and the size of the free tubulin pool decreases. We assume that the amount of tubulin
undergoing IFT is negligible, i.e. $T_b \ll T_f + L$, so that $T \approx T_f + L$. This approximation is justified since
it can be shown by the quasi steady-state assumption that $T_b/T_f < 2\gamma k_{on}M L_{ss}/v$, and consequently
$T_b/T_f < 2.6 \times 10^{-3}$ in terms of the parameters contained in *Table 1*.
By mass action and conservation of motors, the flux of motors may be expressed as

$$J = k_{on}(M - M_b). \tag{30}$$

Given velocities v_a and v_r in the anterograde and retrograde directions, respectively, in ballistic motion the flux is related to the corresponding concentrations in a simple manner. The flux satisfies $J = c_a v_a = c_r v_r$, where c_a and c_r are the concentrations of motors moving in the anterograde and retrograde directions. (It follows from the quasi steady-state assumption that the anterograde and retrograde fluxes are balanced, so that there is a single flux J.) Therefore

$$M_b = Lc_a + Lc_r = LJ/v_a + LJ/v_r \tag{31}$$
$$= 2Lc = 2LJ/v, \tag{32}$$

383 where $c = (c_a + c_r)/2$ is the arithmetic mean and $v = 2(1/v_a + 1/v_r)^{-1}$ is the harmonic mean (i.e. the
384 time-averaged velocity of motion along the flagellum *Marshall and Rosenbaum* (*2001*)). Substituting
385 (32) into (30) and solving for the flux results in

$$J = \frac{k_{on}M}{1 + 2k_{on}L/v}, \tag{33}$$

386 which from (29) and the relation $T_f = T - L$ yields the growth rate

$$\frac{dL}{dt} = \frac{\gamma k_{on}M}{1 + 2k_{on}L/v}(T - L) - d + \eta, \tag{34}$$

387 where we neglect time delays and treat IFT as a quasi steady-state process based on the timescale
388 separation discussed earlier between IFT transit and flagellar length dynamics. The steady-state
389 $dL/dt = 0$ is reached at

$$L_{ss} = \left(T - \frac{d}{\gamma k_{on}M}\right) \Big/ \left(1 + \frac{2d}{\gamma M v}\right). \tag{35}$$

390 Expanding to first order in $\Delta L = L - L_{ss}$ yields

$$\frac{d(\Delta L)}{dt} = -\lambda(\Delta L), \qquad \lambda = \frac{\gamma k_{on}M}{1 + 2k_{on}L_{ss}/v}\left(1 + \frac{2k_{on}(T - L_{ss})/v}{1 + 2k_{on}L_{ss}/v}\right), \tag{36}$$

so that the steady-state is stable since $\lambda > 0$. The steady-state length L_{ss} is positive provided that
$TM > d/\gamma k_{on}$. This sets a theoretical lower limit on the product of total motors and tubulin needed
to observe stable flagella, which makes sense given the constant disassembly term that must be
balanced to reach steady-state.

Diffusive return

Direct experimental evidence shows that kinesin motors do not move ballistically through all phases
of IFT but rather, after ballistic anterograde motion, diffuse back upon reaching the tip of the
flagellum *Chien et al.* (*2017*). This mechanism has been considered in a recent theoretical study
Hendel et al. (*2018*). We next examine how the results above obtained under the assumption of
purely ballistic motion change when the rate-limiting IFT protein must return from tip to base by
diffusion. In this case, there is an additional variable M_d that represents the number of motors
diffusing from the tip to the base, and $M = M_f + M_b + M_d$.

We assume that ballistic and diffusive IFT dynamics are fast compared to changes in length
and are well-approximated by a quasi-steady state process. This implies the diffusive flux $D\partial c_d/\partial x$
equals the injection rate J, and moreover the steady-state condition $D\partial^2 c_d/\partial x^2 = 0$ implies the
concentration profile $c_d(x)$ of the diffusing motors is linear, i.e.

$$c_d(x) = \overline{c_d} + \frac{J}{D}\left(x - \frac{L}{2}\right), \tag{37}$$

407 in which $\overline{c_d}$ is the average concentration along the flagellum and D is the diffusion constant. The
408 flagellar base is taken to act as a diffusive sink so that $c_d(0) = 0$. This boundary condition implies
409 that [1]

$$\overline{c_d} = \frac{JL}{2D}. \tag{39}$$

410 Upon including diffusion, we may replace (30) in the above derivation for the purely-ballistic case
411 by

$$J = k_{on}M_f = k_{on}(M - M_b - M_d), \tag{40}$$

412 where $M_b = JL/v$ given diffusive motion from base to tip and moreover

$$M_d = \overline{c_d}L = \left(\frac{JL}{2D}\right) \cdot L = \frac{JL^2}{2D} \tag{41}$$

413 for the diffusively-moving motors. This results in the expression

$$J = \frac{k_{on}M}{1 + k_{on}L/v + k_{on}L^2/2D}. \tag{42}$$

414 The flux is therefore a quadratic function in length. Substituting this expression for the flux into the
415 growth rate equation (29) results in

$$\frac{\mathrm{d}L}{\mathrm{d}t} = \frac{\gamma k_{on}M}{1 + k_{on}L/v + k_{on}L^2/2D}(T - L) - d + \eta. \tag{43}$$

416 The rate-limited flux arising from a finite number of motors therefore provides a mechanistic
417 derivation that can account for either $1/L$ or $1/L^2$-length dependence under different limits of
418 model parameters. Upon introducing the measured parameters and simply computing the terms in
419 (42), we find $k_{on}L_{ss}/v \approx 0.1$ and $k_{on}L_{ss}^2/2D \approx 0.8$, so that diffusion dominates over ballistic motion at
420 steady-state, but none of the terms is negligible over the complete short to long-length dynamics.

---

[1]To further justify the condition $c_d(0) = 0$, consider a microscopic random walk on a one-dimensional lattice with spacing l. If the probability of hopping in either direction is equal for each timestep Δt, the flux of motors hopping off the flagellum is $\frac{1}{2}c_d(l)l/\Delta t = c_d(l)D/l$, where we have used the relation $D = l^2/(2\Delta t)$ between the continuous diffusion coefficient and parameters of the discrete random walk. For the fluxes onto and off of the flagellum to balance,

$$J = c_d(l)D/l, \tag{38}$$

so that in the limit $l \to 0$ we must have $c_d(0) = 0$ in order to obtain finite J.

421   Solving (43) for the steady-state results in a quadratic equation for L_{ss}. One root is always
422 negative, leaving the positive solution

$$L_{ss} = \left(\frac{D}{v} + \frac{\gamma D M}{d} \right) \left(-1 + \sqrt{1 + \left(\frac{2dT}{\gamma M D} \right) \frac{1 - d/\gamma k_{on} M T}{(1 + d/\gamma M v)^2}} \right) \tag{44}$$

423 We can evaluate the stability of this solution by linearizing about L_{ss}. Expanding to first order in
424 $\Delta L := L - L_{ss}$, we find $\mathrm{d}(\Delta L)/\mathrm{d}t = -\lambda(\Delta L)$ with λ a positive constant given by

$$\lambda = \left(\frac{\gamma k_{on} M}{1 + k_{on} L_{ss}/v + k_{on} L_{ss}^2/2D} \right) \left(1 + \frac{k_{on}/v + k_{on} L_{ss}/D}{1 + k_{on} L_{ss}/v + k_{on} L_{ss}^2/2D} (T - L_{ss}) \right). \tag{45}$$

Two flagella
425
In the case of two flagella with lengths $L_1(t)$ and $L_2(t)$ the length dynamics are given by

$$\frac{\mathrm{d}L_1}{\mathrm{d}t} = \gamma J_1 T_{f,1} - d + \eta_1 \tag{46}$$

$$\frac{\mathrm{d}L_2}{\mathrm{d}t} = \gamma J_2 T_{f,2} - d + \eta_2, \tag{47}$$

where J_i and $T_{f,i}$ for $i = 1, 2$ denote the fluxes and free amounts of tubulin for the two flagella,
which may be equal if they are shared. The fluctuation terms η_1 and η_2 are taken to be uncorrelated
white noise forcing having the covariance

$$\langle \eta_1(t)\eta_1(t') \rangle = \langle \eta_2(t)\eta_2(t') \rangle = |\Upsilon|^2 \delta(t - t') \tag{48}$$

$$\langle \eta_1(t)\eta_2(t') \rangle = 0. \tag{49}$$

We will consider various modes of coupling between the flagella giving rise to different forms of J_i
and $T_{f,i}$ and their consequences for length control. To assess whether a particular length control
model is viable we focus on stability of solutions to the steady-state deterministic equations

$$0 = \gamma J_1 T_{f,1} - d \tag{50}$$

$$0 = \gamma J_2 T_{f,2} - d. \tag{51}$$

Tubulin separate, motors separate (∅)
426
The presence of individual tubulin pools and individual motor pools leads to two uncoupled
instances of the single flagellum dynamics, that is

$$\begin{cases} \dfrac{\mathrm{d}L_1}{\mathrm{d}t} = \dfrac{\gamma k_{on} M}{1 + 2k_{on} L_1/v} (T - L_1) - d + \eta_1, & (52) \\[2ex] \dfrac{\mathrm{d}L_2}{\mathrm{d}t} = \dfrac{\gamma k_{on} M}{1 + 2k_{on} L_2/v} (T - L_2) - d + \eta_2, & (53) \end{cases} \qquad \text{Ballistic return}$$

and

$$\begin{cases} \dfrac{\mathrm{d}L_1}{\mathrm{d}t} = \dfrac{\gamma k_{on} M/2}{1 + k_{on} L_1/v + k_{on} L_1^2/2D} (T - L_1) - d + \eta_1, & (54) \\[2ex] \dfrac{\mathrm{d}L_2}{\mathrm{d}t} = \dfrac{\gamma k_{on} M/2}{1 + k_{on} L_2/v + k_{on} L_2^2/2D} (T - L_2) - d + \eta_2. & (55) \end{cases} \qquad \text{Diffusive return}$$

427 This leads to steady state lengths given by (35) and (44).

Tubulin separate, motors shared (M)
428
429 In the case of individual tubulin pools, we have $T_{f,1} = T - L_1$ and $T_{f,2} = T - L_2$. We next compute
430 the motor flux in the cases of ballistic and diffusive return.

Ballistic return

We have

$$J = \frac{1}{2} k_{\text{on}} M_f = \frac{1}{2} k_{\text{on}} (M - M_b), \tag{56}$$

where the factor of one-half comes from assuming equal injection probability into either flagellum. Further, $M_b = 2J(L_1 + L_2)/v$ so that

$$J = \frac{k_{\text{on}} M/2}{1 + k_{\text{on}} (L_1 + L_2)/v}, \tag{57}$$

and consequently

$$\begin{cases} \dfrac{\mathrm{d}L_1}{\mathrm{d}t} = \dfrac{\gamma k_{\text{on}} M/2}{1 + k_{\text{on}} (L_1 + L_2)/v} (T - L_1) - d + \eta_1, & (58) \\[4mm] \dfrac{\mathrm{d}L_2}{\mathrm{d}t} = \dfrac{\gamma k_{\text{on}} M/2}{1 + k_{\text{on}} (L_1 + L_2)/v} (T - L_2) - d + \eta_2. & (59) \end{cases} \quad \text{Ballistic return}$$

The steady-state length is given by The resulting equations for the steady-state length are

$$0 = \gamma J(T - L_{1,ss}) - d \tag{60}$$

$$0 = \gamma J(T - L_{2,ss}) - d, \tag{61}$$

which yield

$$L_{1,ss} = L_{2,ss} = \left(T - \frac{2d}{\gamma k_{\text{on}} M} \right) \bigg/ \left(1 + \frac{4d}{\gamma M v} \right). \tag{62}$$

Diffusive return

The flux may be calculated according to

$$J = \frac{1}{2} k_{\text{on}} M_f = \frac{1}{2} k_{\text{on}} (M - M_b - M_d). \tag{63}$$

Since the ballistic motion is only in the base to tip direction $M_b = J(L_1 + L_2)/v$ and

$$M_d = \overline{c_{d,1}} L_1 + \overline{c_{d,2}} L_2 = J \left(\frac{L_1^2}{2D} + \frac{L_2^2}{2D} \right) \tag{64}$$

so that

$$J = \frac{k_{\text{on}} M/2}{1 + k_{\text{on}} (L_1 + L_2)/2v + k_{\text{on}} (L_1^2 + L_2^2)/4D} \tag{65}$$

which yields the flagellar length dynamics

$$\begin{cases} \dfrac{\mathrm{d}L_1}{\mathrm{d}t} = \dfrac{\gamma k_{\text{on}} M/2}{1 + k_{\text{on}} (L_1 + L_2)/2v + k_{\text{on}} (L_1^2 + L_2^2)/4D} (T - L_1) - d + \eta_1, & (66) \\[4mm] \dfrac{\mathrm{d}L_2}{\mathrm{d}t} = \dfrac{\gamma k_{\text{on}} M/2}{1 + k_{\text{on}} (L_1 + L_2)/2v + k_{\text{on}} (L_1^2 + L_2^2)/4D} (T - L_2) - d + \eta_2. & (67) \end{cases} \quad \text{Diffusive return}$$

In the case of diffusive return the steady-state solution is

$$L_{1,ss} = L_{2,ss} = \left(\frac{D}{v} + \frac{\gamma DM}{2d} \right) \left(-1 + \sqrt{1 + \left(\frac{4dT}{\gamma MD} \right) \frac{1 - 2d/\gamma k_{\text{on}} MT}{(1 + 2d/\gamma M v)^2}} \right). \tag{68}$$

441 Tubulin shared, motors separate (**T**)

We next consider the case in which only tubulin is shared. The separate pools of motors yield decoupled fluxes

$$
\begin{cases}
J_1 = \dfrac{k_{\mathrm{on}} M}{1 + 2 k_{\mathrm{on}} L_1 / v} & (69) \\[2ex]
J_2 = \dfrac{k_{\mathrm{on}} M}{1 + 2 k_{\mathrm{on}} L_2 / v}, & (70)
\end{cases}
\qquad \text{Ballistic return}
$$

and

$$
\begin{cases}
J_1 = \dfrac{k_{\mathrm{on}} M}{1 + k_{\mathrm{on}} L_1 / v + k_{\mathrm{on}} L_1^2 / 2D} & (71) \\[2ex]
J_2 = \dfrac{k_{\mathrm{on}} M}{1 + k_{\mathrm{on}} L_2 / v + k_{\mathrm{on}} L_2^2 / 2D}, & (72)
\end{cases}
\qquad \text{Diffusive return}
$$

which leads to the systems of equations

$$
\begin{cases}
\dfrac{dL_1}{dt} = \dfrac{\gamma k_{\mathrm{on}} M}{1 + 2 k_{\mathrm{on}} L_1 / v}(T - L_1 - L_2) - d + \eta_1 & (73) \\[2ex]
\dfrac{dL_2}{dt} = \dfrac{\gamma k_{\mathrm{on}} M}{1 + 2 k_{\mathrm{on}} L_2 / v}(T - L_1 - L_2) - d + \eta_2 & (74)
\end{cases}
\qquad \text{Ballistic return}
$$

and

$$
\begin{cases}
\dfrac{dL_1}{dt} = \dfrac{\gamma k_{\mathrm{on}} M}{1 + k_{\mathrm{on}} L_1 / v + k_{\mathrm{on}} L_1^2 / 2D}(T - L_1 - L_2) - d + \eta_1 & (75) \\[2ex]
\dfrac{dL_2}{dt} = \dfrac{\gamma k_{\mathrm{on}} M}{1 + k_{\mathrm{on}} L_2 / v + k_{\mathrm{on}} L_2^2 / 2D}(T - L_1 - L_2) - d + \eta_2. & (76)
\end{cases}
\qquad \text{Diffusive return}
$$

These systems of equations are nearly identical to those of model Ø given by (52)–(53) and (54)–
(55), with the notable exception that the equations are coupled through the shared tubulin pool
term $T - L_1 - L_2$. This model yields simultaneous length control. The steady-state lengths satisfy
$L_{1,ss} = L_{2,ss}$. The resulting steady-state equations are identical to those from (4) and (5) for a single
flagellum, with T rescaled to $T/2$ and γ rescaled to 2γ. Therefore the steady-state lengths for ballistic
and diffusive return satisfy (6) and (7), respectively, upon rescaling $T \to T/2$ and $\gamma \to 2\gamma$.

Tubulin shared, motors shared (**TM**)

We finally consider the case in which both tubulin and motors are shared through a common pool. By the shared tubulin pool assumption $T_{f,1} = T_{f,2} = T - L_1 - L_2$ and by the shared motor pool assumption the injection rates satisfy $J_1 = J_2 \equiv J$. Regardless of the particular form of the flux, we are left with the steady-state equations

$$
0 = \gamma J(T - L_{1,ss} - L_{2,ss}) - d \tag{77}
$$

$$
0 = \gamma J(T - L_{1,ss} - L_{2,ss}) - d. \tag{78}
$$

There is only a single equation for the two unknowns $L_{1,ss}$ and $L_{2,ss}$ and the lengths are indetermi-
nate. Upon adding random noise, the sum fluctuates around a specific value whereas the difference
undergoes a random walk (inset to *Figure 3*(b)(iv)).

Acknowledgments

We acknowledge useful discussions with Wallace Marshall, William Ludington, Shashank Shekhar,
and Shane Mcinally. We thank Jie Lin and Ethan Levien for reading a draft of this manuscript and
offering helpful feedback. We acknowledge funding support from National Science Foundation
grant DMS-1502851 (TGF), grant DMR-1610737 (JK), MRSEC-1420382 (JK), the Simons Foundation
(JK, LM), the A. P. Sloan Foundation (AA), and the Kavli Institute (AA).

References

Amir A, Balaban NQ. Learning from Noise: How Observing Stochasticity May Aid Microbiology. Trends in Microbiology. 2018; .

Bradley BA, Quarmby LM. A NIMA-related kinase, Cnk2p, regulates both flagellar length and cell size in Chlamydomonas. Journal of Cell Science. 2005; 118(15):3317–3326.

Buisson J, Chenouard N, Lagache T, Blisnick T, Olivo-Marin JC, Bastin P. Intraflagellar transport proteins cycle between the flagellum and its base. Journal of Cell Science. 2013; 126(1):327–338.

Cao M, Meng D, Wang L, Bei S, Snell WJ, Pan J. Activation loop phosphorylation of a protein kinase is a molecular marker of organelle size that dynamically reports flagellar length. Proceedings of the National Academy of Sciences. 2013; 110(30):12337–12342.

Chien A, Shih SM, Bower R, Tritschler D, Porter ME, Yildiz A. Dynamics of the IFT machinery at the ciliary tip. eLife. 2017; 6.

Cole DG. The intraflagellar transport machinery of Chlamydomonas reinhardtii. Traffic. 2003; 4(7):435–442.

Cole DG, Diener DR, Himelblau AL, Beech PL, Fuster JC, Rosenbaum JL. Chlamydomonas kinesin-II–dependent intraflagellar transport (IFT): IFT particles contain proteins required for ciliary assembly in Caenorhabditis elegans sensory neurons. The Journal of Cell Biology. 1998; 141(4):993–1008.

Dawson SC, House SA. Life with eight flagella: flagellar assembly and division in Giardia. Current Opinion in Microbiology. 2010; 13(4):480–490.

Dentler W. Intraflagellar transport (IFT) during assembly and disassembly of Chlamydomonas flagella. The Journal of Cell Biology. 2005; 170(4):649–659.

Fall CP. Computational Cell Biology {Interdisciplinary Applied Mathematics; V. 20}. Springer-Verlag New York Incorporated; 2002.

Gardiner C. Stochastic Methods, vol. 4. Springer Berlin; 2009.

Hara Y, Merten CA. Dynein-based accumulation of membranes regulates nuclear expansion in Xenopus laevis egg extracts. Developmental Cell. 2015; 33(5):562–575.

Hendel NL, Thomson M, Marshall WF. Diffusion as a ruler: modeling kinesin diffusion as a length sensor for intraflagellar transport. Biophysical Journal. 2018; 114(3):663–674.

Hilton LK, Gunawardane K, Kim JW, Schwarz MC, Quarmby LM. The kinases LF4 and CNK2 control ciliary length by feedback regulation of assembly and disassembly rates. Current Biology. 2013; 23(22):2208–2214.

Khona DK, Rao VG, Motiwalla MJ, Varma PS, Kashyap AR, Das K, Shirolikar SM, Borde L, Dharmadhikari JA, Dharmadhikari AK, et al. Anomalies in the motion dynamics of long-flagella mutants of Chlamydomonas reinhardtii. Journal of Biological Physics. 2013; 39(1):1–14.

Kozminksi KG, Johnson KA, Forscher P, Rosenbaum JL. A motility in the eukaryotic flagellum unrelated to flagellar beating. Proceedings of the National Academy of Sciences. 1993; 90(12):5519–5523.

Ludington WB, Ishikawa H, Serebrenik YV, Ritter A, Hernandez-Lopez RA, Gunzenhauser J, Kannegaard E, Marshall WF. A systematic comparison of mathematical models for inherent measurement of ciliary length: how a cell can measure length and volume. Biophysical Journal. 2015; 108(6):1361–1379.

Ludington WB, Shi LZ, Zhu Q, Berns MW, Marshall WF. Organelle size equalization by a constitutive process. Current Biology. 2012; 22(22):2173–2179.

Ludington WB, Wemmer KA, Lechtreck KF, Witman GB, Marshall WF. Avalanche-like behavior in ciliary import. Proceedings of the National Academy of Sciences. 2013; p. 201217354.

Luo M, Cao M, Kan Y, Li G, Snell W, Pan J. The phosphorylation state of an aurora-like kinase marks the length of growing flagella in Chlamydomonas. Current Biology. 2011; 21(7):586–591.

Marshall W. Personal communication. . 2018; .

Marshall WF. Cell geometry: How cells count and measure size. Annual Review of Biophysics. 2016; 45:49–64.

Marshall WF, Qin H, Brenni MR, Rosenbaum JL. Flagellar length control system: testing a simple model based on intraflagellar transport and turnover. Molecular Biology of the Cell. 2005; 16(1):270–278.

Marshall WF, Rosenbaum JL. Intraflagellar transport balances continuous turnover of outer doublet microtubules implications for flagellar length control. The Journal of Cell Biology. 2001; 155(3):405–414.

McInally SG, Dawson SC. Eight unique basal bodies in the multi-flagellated diplomonad Giardia lamblia. Cilia. 2016; 5(1):21.

Milo R, Phillips R. Cell biology by the numbers. Garland Science; 2015.

Mohapatra L, Lagny TJ, Harbage D, Jelenkovic PR, Kondev J. The limiting-pool mechanism fails to control the size of multiple organelles. Cell Systems. 2017; 4(5):559–567.

Morga B, Bastin P. Getting to the heart of intraflagellar transport using Trypanosoma and Chlamydomonas models: the strength is in their differences. Cilia. 2013; 2(1):16.

Piao T, Luo M, Wang L, Guo Y, Li P, Snell WJ, Pan J, et al. A microtubule depolymerizing kinesin functions during both flagellar disassembly and flagellar assembly in Chlamydomonas. Proceedings of the National Academy of Sciences. 2009; 106(12):4713–4718.

Prevo B, Scholey JM, Peterman EJ. Intraflagellar transport: mechanisms of motor action, cooperation, and cargo delivery. The FEBS journal. 2017; 284(18):2905–2931.

Rosenbaum JL, Child F. Flagellar regeneration in protozoan flagellates. The Journal of Cell Biology. 1967; 34(1):345–364.

Rosenbaum JL, Cole DG, Diener DR. Intraflagellar transport: the eyes have it. The Journal of Cell Biology. 1999; 144(3):385–388.

Rosenbaum JL, Moulder JE, Ringo DL. Flagellar elongation and shortening in Chlamydomonas: the use of cycloheximide and colchicine to study the synthesis and assembly of flagellar proteins. The Journal of Cell Biology. 1969; 41(2):600–619.

Rosenbaum JL, Witman GB. Intraflagellar transport. Nature Reviews Molecular Cell Biology. 2002; 3(11):813.

Scholey JM. Intraflagellar transport. Annual Review of Cell and Developmental Biology. 2003; 19(1):423–443.

Scholey JM. Intraflagellar transport motors in cilia: moving along the cell's antenna. The Journal of Cell Biology. 2008; 180(1):23–29.

Song L, Dentler WL. Flagellar protein dynamics in Chlamydomonas. Journal of Biological Chemistry. 2001; 276(32):29754–29763.

Tam LW, Dentler WL, Lefebvre PA. Defective flagellar assembly and length regulation in LF3 null mutants in Chlamydomonas. The Journal of Cell Biology. 2003; 163(3):597–607.

Toropova K, Mladenov M, Roberts AJ. Intraflagellar transport dynein is autoinhibited by trapping of its mechanical and track-binding elements. Nature Structural and Molecular Biology. 2017; 24(5):461.

Wemmer KA, Marshall WF. Biological noise in an organelle size control system. In preparation. 2018; .

Wren KN, Craft JM, Tritschler D, Schauer A, Patel DK, Smith EF, Porter ME, Kner P, Lechtreck KF. A differential cargo-loading model of ciliary length regulation by IFT. Current Biology. 2013; 23(24):2463–2471.

Zink D, Fischer AH, Nickerson JA. Nuclear structure in cancer cells. Nature Reviews Cancer. 2004; 4(9):677–687.

540 **Appendix 1**

541
542 ## Parameter estimation

543 To estimate the model parameters, we fit to experimentally-measured data. First, we use
544 the extrapolated data point from **Marshall et al.** (**2005**) that the growth rate upon severing a
545 flagellum is approximately 0.4 μm/min. This implies

$$\gamma k_{on} MT - d \approx 0.4, \tag{79}$$

where fluctuations have been neglected. We first estimate the product $\gamma k_{on} M$. To do so, we
use the measured disassembly rate $d = 0.5$ μm/min **Marshall and Rosenbaum** (**2001**) and
an estimated instantaneous tubulin pool of $T = 25$–40 μm. This yields $\gamma k_{on} M = 2.3 \times 10^{-2}$–
3.6×10^{-2} min^{-1}. The estimate T for the tubulin pool available to a single flagellum in the
instant after severing is based on the reported amount 76–94 μm for the total tubulin pool
shared between two flagella **Marshall et al.** (**2005**) after accounting for some tubulin lost
after severing. This estimate for $\gamma k_{on} M$ is not expected to be very precise, particularly given
that the total tubulin pool size from **Marshall et al.** (**2005**) was itself obtained by fitting the
parameters of a related model to data on mutants with extra flagella.

We can subsequently estimate k_{on} by fitting it to the steady-state length observed in
the severing experiment. Returning to (43) and once again neglecting fluctuations, we may
express k_{on} in terms of quantities that have already been measured or estimated:

$$k_{on} \approx \left(\frac{L_{ss}}{v} + \frac{L_{ss}^2}{2D} \right)^{-1} \left(\frac{\gamma k M}{d}(T - L_{ss}) - 1 \right). \tag{80}$$

Plugging in the values for $\gamma k M, d$, and T above as well as the measured values $v = 150$
μm/min **Kozminksi et al.** (**1993**); **Buisson et al.** (**2013**), $D = 102$ μm^2/min **Chien et al.** (**2017**),
and $L_{ss} = 10$ μm measured in **Ludington et al.** (**2012**), we obtain $k_{on} = 0.8$–4.5 min^{-1}. See the
parameter values and definitions in **Table 1**.

Appendix 2

Linearization

Stability is established by linearizing about steady-state. We will perform the calculation for
the **M** model, then comment on the general form of viable models and their corresponding
results. We also use the linearization to demonstrate how simultaneous length control
breaks down in the **TM** model. We work out the explicit formulas for the case of diffusive
return; the results for ballistic return can be computed with minor changes (see *Figure 6* for
the resulting formulas).

M model
Let ΔL_1 and ΔL_2 be the deviations from steady-state such that $L_1 = L_{ss} + \Delta L_1$ and $L_2 = L_{ss} + \Delta L_2$. From the Main Text, the flux J is given by

$$J = \frac{k_{on}M/2}{1 + k_{on}((L_{ss} + \Delta L_1) + (L_{ss} + \Delta L_2))/2v + k_{on}(L_{ss} + \Delta L_1)^2 + (L_{ss} + \Delta L_2)^2)/4D}, \tag{81}$$

Keeping only those terms in the denominator up to first order in ΔL_1 and ΔL_2,

$$J = \frac{k_{on}M/2}{1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D + k_{on}(\Delta L_1 + \Delta L_2)/2v + k_{on}L_{ss}(\Delta L_1 + \Delta L_2)/2D}$$
$$+ \mathcal{O}(\Delta L_1^2) + \mathcal{O}(\Delta L_1 \cdot \Delta L_2) + \mathcal{O}(\Delta L_2^2), \tag{82}$$

and taking the first-order Taylor series expansion yields

$$J = \frac{k_{on}M/2}{1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D} \left(1 - \frac{k_{on}(\Delta L_1 + \Delta L_2)/2v + k_{on}L_{ss}(\Delta L_1 + \Delta L_2)/2D}{1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D}\right)$$
$$+ \mathcal{O}(\Delta L_1^2) + \mathcal{O}(\Delta L_1 \cdot \Delta L_2) + \mathcal{O}(\Delta L_2^2). \tag{83}$$

Substituting this expression for the flux into the dynamical equations (12) and (13) we have, to first order,

$$\frac{d(\Delta L_1)}{dt} = \frac{\gamma k_{on}M/2}{1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D}(-\Delta L_1)$$
$$+ \frac{\gamma k_{on}M(T - L_{ss})\left(k_{on}/2v + k_{on}L_{ss}/2D\right)/2}{\left(1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D\right)^2}(-\Delta L_1 - \Delta L_2) + \eta_1 \tag{84}$$

$$\frac{d(\Delta L_2)}{dt} = \frac{\gamma k_{on}M/2}{1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D}(-\Delta L_2)$$
$$+ \frac{\gamma k_{on}M(T - L_{ss})\left(k_{on}/2v + k_{on}L_{ss}/2D\right)/2}{\left(1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D\right)^2}(-\Delta L_1 - \Delta L_2) + \eta_2. \tag{85}$$

Defining

$$a_{\mathbf{M}} = \frac{\gamma k_{on}M(T - L_{ss})\left(k_{on}/2v + k_{on}L_{ss}/2D\right)}{2\left(1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D\right)^2}$$

$$b_{\mathbf{M}} = \frac{\gamma k_{on}M/2}{1 + k_{on}L_{ss}/v + k_{on}L_{ss}^2/2D},$$

we may write this system in the matrix form

$$\frac{d}{dt}\begin{pmatrix} \Delta L_1 \\ \Delta L_2 \end{pmatrix} = -\begin{pmatrix} a_{\mathbf{M}} + b_{\mathbf{M}} & a_{\mathbf{M}} \\ a_{\mathbf{M}} & a_{\mathbf{M}} + b_{\mathbf{M}} \end{pmatrix}\begin{pmatrix} \Delta L_1 \\ \Delta L_2 \end{pmatrix} + \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}. \tag{86}$$

The matrix above is the sum of a rank-1 matrix and a diagonal perturbation. Roughly speaking, $a_{\mathbf{M}}$ corresponds to the shared quantities whereas $b_{\mathbf{M}}$ corresponds to the separate quantities. This system may be diagonalized in terms of the sum $\Sigma = \Delta L_1 + \Delta L_2$ and difference $\Gamma = \Delta L_1 - \Delta L_2$ to yield

$$\frac{\mathrm{d}}{\mathrm{d}t}\begin{pmatrix}\Sigma \\ \Gamma\end{pmatrix} = -\begin{pmatrix}2a_{\mathbf{M}} + b_{\mathbf{M}} & 0 \\ 0 & b_{\mathbf{M}}\end{pmatrix}\begin{pmatrix}\Sigma \\ \Gamma\end{pmatrix} + \begin{pmatrix}\eta_\Sigma \\ \eta_\Gamma\end{pmatrix}, \tag{87}$$

where $\eta_\Sigma = \eta_1 + \eta_2$ and $\eta_\Gamma = \eta_1 - \eta_2$. Note that the eigenvalues

$$\lambda_\Sigma = -(2a_{\mathbf{M}} + b_{\mathbf{M}}), \tag{88}$$

$$\lambda_\Gamma = -b_{\mathbf{M}}, \tag{89}$$

are both negative, so that the steady-state is stable, since $b_{\mathbf{M}}$ is clearly positive and $T > L_{ss}$ implies the positivity of $a_{\mathbf{M}}$. The fact that λ_Σ and λ_Γ are *distinct* is also important as it provides a possible means to extract two independent parameters from experiment.

T model

In fact, the matrix form (86) above applies to all five viable models, with suitably modified parameters. For the case of shared tubulin pool and separate motor pools, we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\begin{pmatrix}\Delta L_1 \\ \Delta L_2\end{pmatrix} = -\begin{pmatrix}a_{\mathbf{T}} + b_{\mathbf{T}} & a_{\mathbf{T}} \\ a_{\mathbf{T}} & a_{\mathbf{T}} + b_{\mathbf{T}}\end{pmatrix}\begin{pmatrix}\Delta L_1 \\ \Delta L_2\end{pmatrix} + \begin{pmatrix}\eta_1 \\ \eta_2\end{pmatrix}. \tag{90}$$

with

$$a_{\mathbf{T}} = \frac{\gamma k_{\mathrm{on}} M}{1 + k_{\mathrm{on}} L_{ss}/v + k_{\mathrm{on}} L_{ss}^2/2D}$$

$$b_{\mathbf{T}} = \frac{\gamma k_{\mathrm{on}} M(T - 2L_{ss})\left(k_{\mathrm{on}}/v + k_{\mathrm{on}} L_{ss}/D\right)}{\left(1 + k_{\mathrm{on}} L_{ss}/v + k_{\mathrm{on}} L_{ss}^2/2D\right)^2}.$$

The same logic used in the previous section implies that both eigenvalues are negative and the dynamics is thus stable.

TM model

Simultaneous length control breaks down if all biomolecules are shared and the disassembly rate is constant. As shown in the Main Text, in this case the steady-state equations provide only a single equation for the two unknowns $L_{1,ss}$ and $L_{2,ss}$:

$$0 = \gamma J(T - L_{1,ss} - L_{2,ss}) - d. \tag{91}$$

This equation admits infinitely-many solutions $L_{1,ss}$ and $L_{2,ss}$. By symmetry, linearizing about any one of these solutions yields a matrix equation of the form

$$\frac{\mathrm{d}}{\mathrm{d}t}\begin{pmatrix}\Delta L_1 \\ \Delta L_2\end{pmatrix} = -\begin{pmatrix}a_{\mathbf{TM}} & a_{\mathbf{TM}} \\ a_{\mathbf{TM}} & a_{\mathbf{TM}}\end{pmatrix}\begin{pmatrix}\Delta L_1 \\ \Delta L_2\end{pmatrix} + \begin{pmatrix}\eta_1 \\ \eta_2\end{pmatrix}. \tag{92}$$

Unlike the analogous matrices for the viable models, the 2×2 matrix above has an vanishing eigenvalue, as we now show. Diagonalizing in terms of the sum $\Sigma = \Delta L_1 + \Delta L_2$ and difference $\Gamma = \Delta L_1 - \Delta L_2$ gives

$$\frac{\mathrm{d}}{\mathrm{d}t}\begin{pmatrix}\Sigma \\ \Gamma\end{pmatrix} = -\begin{pmatrix}2a_{\mathbf{TM}} & 0 \\ 0 & 0\end{pmatrix}\begin{pmatrix}\Sigma \\ \Gamma\end{pmatrix} + \begin{pmatrix}\eta_\Sigma \\ \eta_\Gamma\end{pmatrix}. \tag{93}$$

There is a vanishing eigenvalue associated to the difference of lengths, i.e. perturbations from steady-state in the difference of lengths do not decay on a finite timescale. This is consistent with previous results from stochastic simulations that the **TM** model does not yield simultaneous length control *Mohapatra et al.* (*2017*).

TM* model

For the case of shared tubulin pool and shared motor pools with concentration-dependent disassembly, we have

$$\frac{d}{dt}\begin{pmatrix} \Delta L_1 \\ \Delta L_2 \end{pmatrix} = -\begin{pmatrix} a_{\text{TM}^*} + b_{\text{TM}^*} & a_{\text{TM}^*} \\ a_{\text{TM}^*} & a_{\text{TM}^*} + b_{\text{TM}^*} \end{pmatrix}\begin{pmatrix} \Delta L_1 \\ \Delta L_2 \end{pmatrix} + \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}. \tag{94}$$

with

$$a_{\text{TM}^*} = \frac{\gamma k_{\text{on}} M/2}{1 + k_{\text{on}} L_{ss}/v + k_{\text{on}} L_{ss}^2/2D}\left(1 + \frac{\left(k_{\text{on}}/2v + k_{\text{on}} L_{ss}/2D\right)\left(T - (2 + d_1/\gamma D)L_{ss}\right)}{1 + k_{\text{on}} L_{ss}/v + k_{\text{on}} L_{ss}^2/2D}\right)$$

$$b_{\text{TM}^*} = \frac{d_1 k_{\text{on}} M/2D}{1 + k_{\text{on}} L_{ss}/v + k_{\text{on}} L_{ss}^2/2D}.$$

Arbitrary flagellar number

This matrix form also applies to the case of arbitrary flagellar number N discussed in Results. Let $\Delta L_i := L_i - L_{ss}$ for $i = 1, \ldots, N$ denote the deviation from steady-state for the i^{th} flagella. The linearized equations satisfy

$$\Delta \mathbf{L} = M \Delta \mathbf{L} + \boldsymbol{\eta}, \tag{95}$$

where $\Delta \mathbf{L} = (\Delta L_1, \ldots, \Delta L_N)$, $\boldsymbol{\eta} = (\eta_1, \ldots, \eta_N)$, and the matrix M has the form

$$M = -aR - bI, \tag{96}$$

with $R = \mathbf{1}\mathbf{1}^T$ the rank-one matrix satisfying $R_{ij} = 1$ for all i, j and I is the $N \times N$ identity matrix, e.g. for N=3

$$M = -a\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} - b\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} -a - b & -a & -a \\ -a & -a - b & -a \\ -a & -a & -a - b \end{pmatrix}.$$

M is straightforward to diagonalize. Applying M to the vector $\mathbf{v}_1 := \mathbf{1} = (1, \ldots, 1)^T$ corresponding to the sum of all lengths yields

$$M\mathbf{1} = -aR\mathbf{1} - bI\mathbf{1}$$
$$= -(aN + b)\mathbf{1} \tag{97}$$

Further, for any vector $\mathbf{x} = (x_1, \ldots, x_N)$ such that $\sum_{i=1}^{N} x_i = 0$, we have $R\mathbf{x} = 0$ and

$$M\mathbf{x} = -b\mathbf{x}, \tag{98}$$

therefore it is an eigenvector with eigenvalue $\lambda = -b$. Note that the pairwise differences

$$\mathbf{v}_k = (0, \ldots, 0, \underbrace{1}_{\substack{(k-1)^{\text{st}} \\ \text{entry}}}, \underbrace{-1}_{\substack{k^{\text{th}} \\ \text{entry}}}, 0, \ldots, 0)$$

for $k = 2, \ldots, N$ form a convenient basis for the space of such $\mathbf{x}$ whose components sum to zero. From (97) and (98) above, in terms of the basis $\{\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_N\}$ consisting of the sum and differences in lengths, the evolution equations diagonalize with eigenvalues

$$\lambda_\Sigma(N) = -(aN + b) \tag{99}$$

$$\lambda_\Gamma(N) = -b \tag{100}$$

for the sum and differences, respectively, so that $\lambda_\Sigma(N)$ is an eigenvalue of multiplicity 1 and $\lambda_\Gamma(N)$ is an eigenvalue of multiplicity $N - 1$. Note that in addition to the explicit dependence of $\lambda_\Sigma(N)$ and $\lambda_\Gamma(N)$ on N there is an implicit number-dependence through the steady-state length (and potentially the size of the pool T). Since $\lambda_\Sigma(N) \neq \lambda_\Gamma(N)$, the autocorrelations described in *Appendix 3* will involve two distinct timescales $\tau_\Sigma(N) = \lambda_\Sigma(N)^{-1}$ and $\tau_\Gamma(N) = \lambda_\Gamma(N)^{-1}$ corresponding to these two eigenvalues.

701 **Appendix 3**

702 ## Fluctuation analysis

We return our attention to the case of $N = 2$ flagella relevant to *Chlamydomonas*. As shown
703 in **Appendix 2**, linearizing about steady-state results in a diagonal system in terms of the
704 sum $\Sigma = \Delta L_1 + \Delta L_2$ and difference $\Gamma = \Delta L_1 - \Delta L_2$:
705

706 $$\frac{d}{dt}\begin{pmatrix}\Sigma \\ \Gamma\end{pmatrix} = \begin{pmatrix}\lambda_\Sigma & 0 \\ 0 & \lambda_\Gamma\end{pmatrix}\begin{pmatrix}\Sigma \\ \Gamma\end{pmatrix} + \begin{pmatrix}\eta_\Sigma \\ \eta_\Gamma\end{pmatrix}, \tag{101}$$
707

708
where $\eta_\Sigma = \eta_1 + \eta_2$ and $\eta_\Gamma = \eta_1 - \eta_2$ have the covariance structure
709

710 $$\langle \eta_\Sigma(t)\eta_\Sigma(t) \rangle = \langle \eta_\Gamma(t)\eta_\Gamma(t) \rangle = 2\,|\Upsilon|^2\,\delta(t - t'), \tag{102}$$

711 $$\langle \eta_\Sigma(t)\eta_\Gamma(t) \rangle = 0. \tag{103}$$

712
713 Applying the Fourier transform and solving for $\hat{\Sigma}(\omega)$ and $\hat{\Gamma}(\omega)$ results in a Lorentzian function
714 **Gardiner (2009)**, as we explicitly show here. We first apply the Fourier transform to (101) to
715 obtain
716

717 $$-i\omega\hat{\Sigma} = \lambda_\Sigma\hat{\Sigma} + \hat{\eta}_\Sigma \tag{104}$$

718 $$-i\omega\hat{\Gamma} = \lambda_\Gamma\hat{\Gamma} + \hat{\eta}_\Gamma, \tag{105}$$

719
with λ_Σ and λ_Γ given in **Appendix 2** for the viable models. Solving for $\hat{\Sigma}(\omega)$ and $\hat{\Gamma}(\omega)$ yields
720

721 $$\hat{\Sigma}(\omega) = -\frac{\hat{\eta}_\Sigma}{i\omega + \lambda_\Sigma}, \qquad \hat{\Gamma}(\omega) = -\frac{\hat{\eta}_\Gamma}{i\omega + \lambda_\Gamma}. \tag{106}$$
722

723 Multiplying by complex conjugates and taking the ensemble average results in
724

725 $$\left|\hat{\Sigma}(\omega)\right|^2 = \frac{2\,|\Upsilon|^2}{\omega^2 + \lambda_\Sigma^2}, \qquad \left|\hat{\Gamma}(\omega)\right|^2 = \frac{2\,|\Upsilon|^2}{\omega^2 + \lambda_\Gamma^2}, \tag{107}$$
726

727 where we have used (102) for the spectra. The Wiener-Khinchin Theorem asserts that for
728 a random variable X the autocorrelation function $C_X(\tau) := \langle X(t)X(t + \tau) \rangle$ is equal to the
729 inverse Fourier transform of the spectral density $\left|\hat{X}(\omega)\right|^2$. Taking inverse transforms in (107)
730 gives
731

732 $$C_\Sigma(\tau) = \frac{|\Upsilon|^2}{|\lambda_\Sigma|}e^{\lambda_\Sigma|\tau|}, \qquad C_\Gamma(\tau) = \frac{|\Upsilon|^2}{|\lambda_\Gamma|}e^{\lambda_\Gamma|\tau|}. \tag{108}$$
733

From the linearization in **Appendix 2**, the predicted timescales based on the parameters in
Table 1 are $\tau_\Sigma = 17$ min and $\tau_\Gamma = 63$ min for the **M** model, $\tau_\Sigma = 12$ min and $\tau_\Gamma = 15$ min for the
T model, and $\tau_\Sigma = 17$ min and $\tau_\Gamma = 26$ min for the **TM*** model.

Extracting parameters via fluctuations

As described above, the autocorrelations $C_\Sigma(\tau) := \langle \Sigma(t)\Sigma(t + \tau) \rangle$ and $C_\Gamma(\tau) := \langle \Gamma(t)\Gamma(t + \tau) \rangle$
of the sum and difference in flagellar lengths can be expressed in terms of the timescales
$\tau_\Sigma := |\lambda_\Sigma|^{-1}$ and $\tau_\Gamma := |\lambda_\Gamma|^{-1}$ and the fluctuation magnitudes $|\Upsilon|^2\tau_\Sigma$ and $|\Upsilon|^2\tau_\Gamma$. These
formulas may be used to fit parameters and make testable predictions.

Our models make the prediction $\tau_\Sigma < \tau_\Gamma$ and $\left|\Upsilon_\Sigma\right|^2 = \left|\Upsilon_\Gamma\right|^2$, which provides a useful
test for the overall modeling framework. On a more quantitative level, since $\tau_\Sigma \neq \tau_\Gamma$, the
autocorrelations provide three distinct predictions, i.e. the formulas for τ_Σ, τ_Γ, and Υ in terms
of model parameters. Experimental measurements can be used to test these predictions (or

used to fit any parameters that cannot be independently estimated). For the concentration-dependent disassembly model **TM***, the parameter d_1 is inversely proportional to τ_Γ, making it possible to fit by comparison to the fluctuation spectrum.

To provide a proof of concept, we generate synthetic data using our model **T** to demonstrate that we can use fluctuations to extract the input parameters. We fit the pooled autocorrelation functions to exponential functions $C_\Sigma(\tau) = \tau_\Sigma \left|\Upsilon_\Sigma\right|^2 e^{-\tau/\tau_\Sigma}$, $C_\Gamma(\tau) = \tau_\Gamma \left|\Upsilon_\Gamma\right|^2 e^{-\tau/\tau_\Gamma}$ and find that we are able to extract the input values (***Appendix 3 Figure 1***(a)-(b)).

To quantify how much data would be needed to effectively constrain the model parameters, we have performed Monte Carlo simulations with different sampling frequencies and total time durations. ***Appendix 3 Figure 1*** (c)-(d) shows, for instance, that if the total experiment time is fixed at 1000 minutes (16 hours), sampling intervals of two minutes or less are needed to reliably obtain 10% error in both τ_Σ and τ_Γ. Although the amount of data required is significant, existing protocols are able to maintain healthy cells for over 18 hours using microfluidics ***Ludington et al.*** (***2012***), so that extracting the model parameters from fluctuations should be experimentally tractable.

Appendix 3 Figure 1. Analysis of synthetic data generated near steady-state using model **T**. The autocorrelations in the sum (a) and difference (b) of flagellar lengths are described by exponential functions with distinct timescales. These timescales can be extracted from data by data fitting. To bound the quantity of data required to accurately extract the timescales, we generated synthetic data for 20 pairs of flagella using different sampling intervals and time durations. We find that using a sampling interval of less than 2 minutes and a time duration of greater than 1000 minutes (approx. 16 hrs) allows the timescales τ_Σ (c) and τ_Γ (d) to be extracted to within 10% error.