Sensory experience controls dendritic structure and behavior by distinct pathways involving degenerins

Sharon Inberg[1], Yael Iosilevskii[1], Alba Calatayud-Sanchez[2], Hagar Setty[3,4], Meital Oren-Suissa[3,4], Michael Krieg[2], and Benjamin Podbilewicz[1,*]

[1]Department of Biology, Technion-Israel Institute of Technology, 3200003 Haifa, Israel

[2]ICFO - Institut de Ciencies Fotoniques, The Barcelona Institute of Science and Technology, Castelldefels (Barcelona) 08860, Spain

[3]Department of Brain Sciences, Weizmann Institute of Science, Rehovot 7610001, Israel

[4]Department of Molecular Neuroscience, Weizmann Institute of Science, Rehovot 7610001, Israel

* Corresponding author
Department of Biology,
Technion-Israel Institute of Technology, 3200003 Haifa, Israel
Tel +972 4 8293454
Email: podbilew@technion.ac.il

Competing Interest Statement: None

Classification: Biological Sciences. Neuroscience

Keywords: Dendritic plasticity, *Caenorhabditis elegans,* Epithelial sodium channels (degenerins)

This file includes:

Main Text
Figure Legends
Supplementary Figure Legends
Video Legends
Supplementary File Legend
Titles for the Source Data

Abstract

Dendrites are crucial for receiving information into neurons. Sensory experience affects the structure of these tree-like neurites, which, it is assumed, modifies neuronal function, yet the evidence is scarce, and the mechanisms are unknown. To study whether sensory experience affects dendritic morphology, we use the *Caenorhabditis elegans'* arborized nociceptor PVD neurons, under natural mechanical stimulation induced by physical contacts between individuals. We found that mechanosensory signals induced by conspecifics and by glass beads affect the dendritic structure of the PVD. Moreover, developmentally isolated animals show a decrease in their ability to respond to harsh touch. The structural and behavioral plasticity following sensory deprivation are functionally independent of each other and are mediated by an array of evolutionarily conserved mechanosensory amiloride-sensitive epithelial sodium channels (degenerins). Calcium imaging of the PVD neurons in a micromechanical device revealed that controlled mechanical stimulation of the body wall produces similar calcium dynamics in both isolated and crowded animals. Our genetic results, supported by optogenetic, behavioral, and pharmacological evidence, suggest an activity-dependent homeostatic mechanism for dendritic structural plasticity, that in parallel controls escape response to noxious mechanosensory stimuli.

Significance Statement

Little is known about how neuronal activity and sensory experience influence the structure and function of dendritic trees. Furthermore, how dendritic structures affect neuronal functions remains to a large extent a mystery despite their fundamental functions in neuronal sensory and synaptic activities in health and diseases. Here we show that complex dendritic trees of the nociceptive and proprioceptive PVD neuron in *C. elegans* display a dynamic nature where they undergo pronounced dendritic modifications during adulthood. These modifications are determined by the received sensory signals generated by conspecific worms on the plate or by inert glass beads. We revealed functions for degenerins/Epithelial sodium channels in this phenomenon, using genetic and optogenetic approaches. We found that the degenerin MEC-10 acts cell autonomously to translate environmental mechanical signals into structural and functional modifications in the PVD.

Main Text

Introduction

The general structure of the nervous system has been known for over a century. Groundbreaking studies on synaptic plasticity and its underlying mechanisms have shown that before birth and in adult animals, neuronal activity is needed for synaptic remodeling (Fox and Wong, 2005, Goodman and Shatz, 1993, Katz and Shatz, 1996, Wiesner et al., 2020, Zuo et al., 2005). In contrast, the molecular mechanisms responsible for structural remodeling of dendritic trees, as a result of different sensory inputs (experience), especially during adulthood, are less understood (Kolb and Whishaw, 1998, Tavosanis, 2012, Wong and Ghosh, 2002).

Mechanistic understanding of experience-dependent structural plasticity is primarily focused on activity sensation by calcium channels and N-methyl-D-aspartate (NMDA) receptors. These are known to induce downstream signaling cascades affecting, among others, the Rho family of small GTPases, calcium metabolism and microtubule stability (Ghiretti et al., 2014, Sin et al., 2002, Vaillant et al., 2002, Zhou et al., 2006). Several neurological conditions including autism, Down syndrome, fragile X syndrome, and schizophrenia are characterized by abnormal dendritic spine structures (Hu et al., 2020, Huebschman et al., 2020, Jan and Jan, 2010, Tendilla-Beltran et al., 2019). Uncovering the molecular basis of dendritic tree instability during development and adulthood, may shed light on neurological disease mechanisms and elucidate their behavioral phenotypes.

The dendrite morphology of the *Caenorhabditis elegans'* PVD bilateral neurons is composed of repetitive, stereotypical and spatially organized structural units that resemble candelabra (Figure 1A), making it a useful platform to study dendritic morphogenesis in hermaphrodites and males (Oren-Suissa et al., 2010; Iosilevskii and Podbilewicz, 2024; for a review see (Heiman and Bülow, 2024)). While some of the genetically-programmed molecular mechanisms responsible for the morphogenesis and regeneration of PVD's dendritic trees are known (Dong et al., 2013, Dong et al., 2015, Inberg et al., 2019, Kravtsov et al., 2017, Oren-Suissa et al., 2017, Oren-Suissa et al., 2010, Salzberg et al., 2013, Salzberg et al., 2014, Smith et al., 2010; Heiman and Bülow, 2024), the influence of nurture (e.g. sensory experience) on its structure and function remain unexplored. The PVD mediates several sensory modalities (reviewed in (Goodman and Sengupta, 2019)), notably response to harsh mechanical stimuli (nociception) (Chatzigeorgiou et al., 2010), response to low temperatures (Chatzigeorgiou et al., 2010) and proprioception (Albeg et al., 2011, Tao et al., 2019, reviewed in Krieg, et. al., 2022). While the PVD response to low temperatures is mediated by transient receptor potential (TRP) channels (Chatzigeorgiou et al., 2010), nociception and proprioception are mediated by Degenerins/Epithelial sodium (Na^+) Channels (DEG/ENaCs) expressed in the PVD (Chatzigeorgiou et al., 2010, Husson et

al., 2012), which form homo- and hetero-trimers and are involved in force sensing. In
mammals, some DEG/ENaCs such as ASIC1a participate in synaptic plasticity and
cognitive functions such as learning and memory (Baldin et al., 2020, Bianchi and
Driscoll, 2002, Chen et al., 2015, Gillespie, 2001, Gobetto et al., 2021, Hill and Ben-
Shahar, 2018, Mango and Nistico, 2020, Welsh et al., 2002). As sensory and social
isolation affect the behavior and fitness of diverse animals (Bailey and Moore, 2018,
Kuhlman et al., 2014, Wilbrecht et al., 2010, Yu and Zuo, 2011), including primates
(Harlow et al., 1965), studying adult nematode somatosensory neurons can reveal
possibly-conserved mechanisms of dendritic plasticity, which are induced by sensory
stimuli.

In gentle touch circuits, which are distinct from nociception (Chalfie and Sulston,
1981, Li et al., 2011, Oren-Suissa et al., 2010), Rose and colleagues (Rose et al., 2005)
found that deprivation of mechanosensory stimulation generated by colliding
conspecifics in the growing plate, resulted in modified glutamatergic signaling and
reduced response to tap stimulation. Here, we adapted this mechanosensory
stimulation paradigm (Rose et al., 2005), where the crowded worms are thought to
represent the natural 'default' enriched mechanosensory state, to look into nociceptive
circuits and identify the mechanism that couples mechanosensory experience to
structural and functional dendritic plasticity (Inberg and Podbilewicz, 2018). We focused
on how mechanosensory experience, perceived through DEG/ENaCs, affects structural
plasticity of the PVD dendritic trees in adult *C. elegans*, and whether this entails
behavioral consequences. We find that mechanosensory experience not only alters the
PVD's dendritic structure in the adult, but also affects associated behavioral outputs.
However, and in contrast to the prevalent hypothesis, these structural and behavioral
properties are not correlated.

Results

Sensory isolation induces behavioral plasticity

Rose and colleagues (Rose et al., 2005) have shown that isolation of wildtype (WT) *C.*
elegans causes a decreased response to gentle touch circuits when comparing to
worms raised in a crowded setting. We utilized a similar paradigm to study nociception,
a behavior associated with PVD neuron activation (Oren-Suissa et al., 2010, Way and
Chalfie, 1988), using a behavioral assay that registers escape following prodding with a
platinum wire (harsh touch assay; Figure 1A) (Chalfie and Sulston, 1981, Li et al., 2011,
Oren-Suissa et al., 2010). To study whether mechanosensory deprivation affects the
nociceptive functions of the PVD, we isolated embryos into single plates where they
grew 72 h to adulthood and compared their behavioral response to harsh touch against
same-aged adults that were grown on crowded plates (Figure 1B). We found that ~40%
of isolated WT animals responded to harsh touch, compared with ~80% of animals
grown in crowded plates (Figure 1C). To test whether gentle touch neurons are involved

in this behavioral difference, we studied *mec-4(e1611)* mutants, in which gentle touch
neurons are degenerated (Caneo et al., 2019, Driscoll and Chalfie, 1991, Hedgecock et
al., 1983, Li et al., 2011, Suzuki et al., 2003), and obtained similar results (Figure 1C).
Given that *mec-4(e1611)* exhibits the same experience-dependent behavioral plasticity
as WT, we conclude that this phenomenon is independent from the *mec-4* gentle-touch
neurons. As a negative control, we used *mec-3(e1338)* mutants, that are harsh-touch-
insensitive (Way and Chalfie, 1988) and found no responses for both groups (Figure
1C). Thus, isolation reduces the response to noxious stimuli, in a process that is
independent of gentle touch circuits.

To determine whether the effect of isolation on nociception is related to the
mechanosensory deficit itself (absence of channel gating to detect collisions (Shi et al.,
2016; Chatzigeorgiou et al., 2010)), in contrast to chemosensory stimuli, we used the
mec-4 strain to compare responses of isolated worms to responses of isolated worms
grown in the presence of glass beads (resembling a method used by (Sawin et al.,
2000)). Worms grown in isolation with beads had a similar response compared to
animals grown in crowded plates (Figure 1–figure supplement 1A). While the beads are
sufficient to increase isolated animals' responses, they do not fully recapitulate the
crowded state (Figure 1–figure supplement 1A). To study whether the effect was
chemosensory-mediated, we tested plates pre-stimulated with pheromones as well as
osm-6 mutants (defective in olfactory sensory cilia (Collet et al., 1998)). We found that
isolation induced a reduction in the response to harsh touch, regardless of plate 'odor'
(Figure 1D) or chemosensory function (Figure 1–figure supplement 1B). We note that
the chemosensory defective *osm-6* mutants also have a defective ciliated
mechanosensory PDE neuron (Collet et al., 1998) (which also mediates posterior harsh
touch responses (Li et al., 2011)). These results suggest that adult worms display a
behavioral plasticity in PVD-related harsh touch circuits, which is dependent on
mechanosensory experience, and is independent of the PDE neuron, olfactory function
and gentle touch neurons.

MEC-10 and other mechanosensory channels mediate isolation-induced
behavioral plasticity before adulthood

To study the genetic mechanisms behind the nociceptive response plasticity, we
performed a candidate gene screen for DEG/ENaCs and TRP channels that are
expressed in the PVD, where degenerins mediate mechanosensation and
proprioception, while TRPs sense low temperatures. In all these signal transduction
pathways some of the outcomes are behavioral responses (Chatzigeorgiou et al., 2010,
Huang and Chalfie, 1994). We found that the WT-like isolation-induced reduction in
harsh touch response was also present in *degt-1* DEG/ENaC and *gtl-1* TRP channel
mutants, suggesting that these channels are not directly involved in behavioral plasticity
following isolation (Figure 1E). In contrast, for *del-1*, *asic-1* and *mec-10* DEG/ENaC
mutants the difference between isolated and crowded conditions was undetectable,

indicating that they are required for such behavioral plasticity. Interestingly, while the
harsh touch response of *asic-1* mutants was consistently high, and similar to crowded
WT worms, the response for *mec-10* mutants was consistently low, similar to isolated
WT animals, regardless of sensory experience (Figure 1E). To test whether the
response to harsh touch is dependent on DEG/ENaC activity, we used the DEG/ENaC
blocker amiloride (Ben-Shahar, 2011, Bianchi and Driscoll, 2002) on WT worms that
were grown in crowded plates. We found that the response to harsh touch was not
affected by continuous growth in the presence of amiloride (Figure 1–figure supplement
1C). This result possibly supports the idea that different amiloride-sensitive epithelial
sodium channels may have positive and negative effects on the response to harsh
touch, which are masked by global inhibition (Figure 1E). This is further evident by the
plasticity consequences of some double and triple DEG/ENaC mutant combinations,
which are difficult to align into a coherent and simple epistatic genetic model (Figure 4–
figure supplement 1).

The DEG/ENaC MEC-10 is expressed in the PVD and responds to
mechanosensory signals (Chatzigeorgiou et al., 2010) including shear forces when
expressed in heterologous cells (Shi et al., 2016, Shi, Mutchler et al., 2018). Since
crowded *mec-10* mutants seem isolated-like in their behavioral response (Figure 1E),
we asked how *mec-10* mediates the behavioral plasticity in crowded conditions. To test
whether the activity of MEC-10 is required cell-autonomously in the PVD, we expressed
MEC-10 under a PVD-specific promoter in a *mec-10* mutant background (Oren-Suissa
et al., 2010, Tsalik et al., 2003). We found that expression of MEC-10 in the PVD
rescues the low response to harsh touch in crowded *mec-10* mutants, indicating that it
acts cell autonomously to modulate behavioral plasticity (Figure 1F).

To determine whether isolation affects the response to harsh touch during
development or in adults, we isolated crowded-raised young adults for 24 h and found
no difference in their response to harsh touch (Figure 1G). Thus, isolation-induced
reduction in mechanosensation is determined before early adulthood, is PVD cell-
autonomous, and *mec-10*-dependent.

Experience affects PVD morphology via *mec-10* in adulthood

Altered sensory experience (such as stimuli deprivation) is known to drive synaptic
plasticity in the nervous system (Alvarez and Sabatini, 2007, Fox and Wong, 2005, Zuo
et al., 2005). However, little is known about the effects of mechanosensory deprivation
on the architecture of sensory neurons. To study whether reduced mechanosensory
experience can alter the dendritic structure of the PVD we followed the isolated growth
paradigm (Figure 1B) and examined two morphological features of the PVD (Figure 2A):
The fraction of ectopic (excessive, disordered) branches out of non-ectopic branches
(those that form the 'ideal' candelabrum), and the percentage of straight 4th order
(quaternary) branches, which form the terminal processes of the candelabrum.

When quantifying the branching pattern of the PVD in adulthood, WT (or WT-like
him-5 background) animals which were isolated as embryos showed an increase in
ectopic branching compared with crowded age-matched worms (Figure 2B) and the
quaternary branches assumed a more rippled shape (fewer straight 4ry branches,
Figure 2C), regardless of the neuron marker utilized (see Figure 2–figure supplement
1).

To determine whether chemical stimulation plays a role in the observed
morphological plasticity, animals were isolated onto pheromone-conditioned plates
(Maures et al., 2014). Similarly to the response to harsh touch (Figure 1D), chemical
stimulation of the plates did not rescue the isolation-induced increase in ectopic
branching or the decrease in straight quaternary branches (Figure 2–figure supplement
2). Additionally, we looked at *mec-4* gentle-touch-insensitive mutants (Rose et al., 2005,
Suzuki et al., 2003) and found that isolation caused a similar change to PVD structure
as in the WT (Figure 2–figure supplement 3, compare with Figure 2B and 2C). In
summary, these results suggest that mechanosensory experience controls
morphological plasticity of PVD dendritic trees independently of *mec-4* and chemical
stimulation.

We next sought to determine whether DEG/ENaCs mediate these experience-
driven morphological alterations, by examining PVD morphology in crowded and
isolated *mec-10* mutants. We found that crowded *mec-10* animals were different from
WT crowded worms, and appear more isolated-like in terms of morphological features
(more ectopic branches and fewer straight quaternary branches, compared to crowded
him-5 control animals; Figure 2B and 2C). These results suggest that PVD dendritic
morphology is affected by sensory experience in a *mec-10*-dependent pathway.
Importantly, these results confirm a role for the mechanosensory channel MEC-10, but
not other degenerins (such as ASIC-1), in the plasticity of both the dendritic structure
and the nociceptive function of the PVD in the crowded background.

To study whether *mec-10* acts cell autonomously to mediate morphological
plasticity, we again utilized PVD-specific MEC-10 expression in a crowded *mec-10*
mutant background and found it reduced the fraction of ectopic branches and increased
the percentage of straight quaternary branches, compared to age-matched non-
transgenic animals (Figure 2D-I). Thus, MEC-10 acts in the PVD to mediate
morphological plasticity.

Having identified an isolation-induced morphological plasticity in adults (observed
at 72 h from embryo), we next sought to establish the temporal dependence of this
effect, by comparing animals after only 48 h, as young adults (see Figure 2–figure
supplement 4A). Isolated worms showed a small but significant difference for the
percentage of straight quaternary branches, but not in ectopic branching (Figure 2–
figure supplement 4B-E) when compared with age-matched crowded worms,
suggesting that isolation induces some morphological alterations during

morphogenesis, while some relate to adult-stage maintenance of the neuron. We then
isolated animals grown for 48 h in crowded conditions, and found that isolation of young
crowded adults for 24 h induced changes in both measured parameters (Figure 2–figure
supplement 4F-I), recapitulating isolation of embryos (Figure 2B-C). Thus, isolation of
adults for 24 h is sufficient to affect PVD dendritic tree architecture, in contrast to the
developmentally-established behavioral response (Figure 1G).

To further determine whether the morphological effect on PVD structure is solely
mediated by mechanical cues, we used glass beads under isolated conditions. We
found that while the presence of glass beads did not reduce the fraction of ectopic
branching, it significantly increased the number of straight quaternary branches
(compared with isolated animals without beads, Figure 2J-K). Thus, the
mechanosensory stimuli of inert beads can partially rescue isolation-induced
morphological changes to the PVD.

Isolation triggers dynamic plasticity of the dendritic tree

To pinpoint the precise time interval required to mediate dendritic arborization changes
in adulthood, we examined the effect of varying isolation times on the PVD morphology
of crowded-raised young adults. We found that isolation of adults for 2 h had no
significant consequence on the structure of the PVD, while isolation of adults for 5 h or
longer significantly increased the proportion of ectopic branches and reduced the
percentage of quaternary straight branches compared to crowded worms (Figure 3A-B).

To further study plasticity at the individual level, we compared the dendritic tree
of individual worms grown under crowded conditions before and after 4 h isolation. This
revealed isolation induced more events of ectopic branch growth and fewer events of
branch retraction (Figure 3C). Furthermore, this period was sufficient to induce an
increase in ectopic branching and a decrease in percentage of straight quaternary
branches (Figures 3D, E, and Figure 3–figure supplement 1). Thus, 4 h of isolation are
sufficient to induce changes in the architecture of the adult PVD.

Since the geometry of quaternary branches shows rapid response to the amount
of mechanosensory stimulation (Figure 3A-E), we next asked how dynamic is this
experience-dependent morphological change. We performed a 3 h time lapse imaging
in crowded and isolated animals anesthetized with 1% tricaine (Videos 1 and 2). We
found that while the number of straight quaternary branches in the isolated worms
remained low and stable (~8%), for the crowded animals, we observed a gradual
reduction in straight quaternary branches, starting after ~2 h of isolation of individual
worms on the slide (Figure 3F,G).

In summary, isolation-induced morphological plasticity is apparent within 2-3 h
from the switch from crowded conditions to growth in the absence of mechanical stimuli.
This dynamic process involves simultaneous increase in ectopic branching and
decrease in straight terminal branches.

Activity affects morphology

It has been shown that activity and sodium influx via the DEG/ENaC UNC-8 promotes synapse elimination in *C. elegans* (Miller-Fleming et al., 2016). To test how global inhibition of DEG/ENaCs affects the morphology of the PVD neuron, we evaluated crowded worms that were grown on plates with amiloride compared to control worms. We found that blocking DEG/ENaCs by amiloride increased the fraction of ectopic branching and decreased the percentage of straight quaternary branches (Figure 3–figure supplement 2A-D). Thus, unlike harsh touch response, which is independent of sensory-level global inhibition (Figure 1–figure supplement 1C), global inhibition at the sensory level alters dendritic morphology. These results suggest that activity via degenerins modulates the structure of the PVD.

To test whether manipulations of other neuromodulators can affect the branching dynamics of the PVD, we utilized two anesthetics: tetramisole, which activates the nematode nicotinic acetylcholine receptors inducing muscle contraction (Aceves et al., 1970), and tricaine, which modulates neuronal activity by blocking sodium channels (Katz et al., 2020), and analyzed their effects during 2-3 h of time lapse movies of crowded and isolated animals. We first tested the commonly used mixture of 0.1% tricaine and 0.01% tetramisole (the racemic mixture of the enantiomer levamisole) (Kravtsov et al., 2017) and found that it induces more growth than retraction of ectopic branches in crowded animals (Figure 3–figure supplement 2E-G). In contrast, 1% tricaine alone has the opposite effect- more retractions of ectopic branches (Figure 3–figure supplement 2H-I; Videos 1 and 2). In addition, 1% tricaine induces more growth for isolated compared to crowded worms, indicating that there are activity-dependent intrinsic factors that are different for the two experience states. These results suggest that modulating neuronal activity triggers structural modifications in the PVD, indicating a possible structural sensitivity to the amount of neuronal activity. In summary, we found that mechanosensory experience, DEG/ENaC protein presence and activity, and pharmacological targeting of activity dynamically affect the structure of the PVD.

PVD structure and function are independent

After establishing that mechanosensory experience induces both a behavioral and a structural plasticity of the PVD, we asked whether there is a causal link between the morphology of the dendritic tree of the PVD and its function as a nociceptor (Hall and Treinin, 2011). We followed the isolation protocol described in Figure 1B for seven combinations of DEG/ENaC mutants and analyzed their response to harsh touch (Figure 4–figure supplement 1) and their PVD structure (Figure 4–figure supplement 2). In order to compare the morphological properties of the different DEG/ENaC genotypes under different sensory experience (crowded, isolated; Figure 4–figure supplement 2), we used discriminant analysis. This provides a supervised classification method to

combine all the analyzed morphological phenotypes, including loss of self-avoidance between adjacent candelabra (Figure 4–figure supplement 3), into a certain value characterizing the PVD morphological state (Figure 4A). We then superimposed the behavioral results of the DEG/ENaC mutants' response to harsh touch (Figure 4–figure supplement 1) on the morphological clustering, as a binary-like property (<45% responding for isolated WT versus >65% for crowded WT, as shown in Figure 4–figure supplement 1*). We found no correlation between the morphology of the PVD and the response to harsh touch when testing the different combinations of genotypes and treatments. For example, isolated *mec-10; degt-1* double mutants show crowded-like morphology with isolated-like behavioral response (Figure 4A and Figure 4–figure supplement 1 and 2). These findings suggest that while degenerins are required for isolation-induced plasticity of both traits, response to harsh touch is independent of the structural alteration of the PVD.

An additional line of evidence supporting the independence of the behavioral and morphological phenotypes was demonstrated by isolation of young adult worms for 24 h (Figure 3). This isolation affects the structure of the PVD (Figure 2–figure supplement 4) but has no effect on the response to harsh touch (Figure 1G). Amiloride also has no effect on harsh touch response (Figure 1–figure supplement 1C) but does alter dendritic morphology (Figure 3–figure supplement 2A-D). Finally, to directly demonstrate that these two features are independent, we assayed harsh touch responses of isolated animals and then analyzed each individual animal for its PVD morphology. We then compared the dendritic morphology of responding and non-responding worms, and found that the morphological parameters were similar (Figure 4B-E). Thus, analysis at the level of individual worms failed to demonstrate a correlation between the morphology and the response to harsh touch. In summary, the morphological and behavioral phenotypes were independently affected by sensory experience via degenerins. We cannot exclude the possibility that other functions of the PVD, like the response to low temperatures (Chatzigeorgiou et al., 2010) and proprioception (Albeg et al., 2011) are more tightly associated to the structure of the PVD, nor that the morphological changes induced by isolation are too minor to constitute a difference in neuronal function.

In addition to the isolation-induced changes in the number of ectopic branches and the percentage of straight quaternary branches, we found that worms raised in isolation are also more likely to lose the self-avoidance between two adjacent menorahs (candelabra). This effect is also *mec-10*-dependent, but appears to act cell non-autonomously. In addition, it is chemosensory and amiloride independent (Figure 4–figure supplement 3).

MEC-10 and DEGT-1 localization is experience-dependent

Differential localization of degenerins can affect both the behavioral response to harsh touch and the structural properties of the neuron. When considering the effect of

isolation, we hypothesized that changes in the localization patterns of DEG/ENaC can
account for plasticity at both the behavioral (Figure 1) and the structural level (Figure 2).
Since MEC-10 and DEGT-1 tend to co-localize within the PVD (Chatzigeorgiou et al.,
2010) we analyzed the interaction between these two proteins, under different
mechanosensory experiences. We found that MEC-10 localization in the plasma
membrane, in intracellular vesicular compartments of the axon and in the quaternary
branches was reduced after isolation (Figure 5A-D, Videos 3 and 4). In contrast to
MEC-10 (Figure 5A-D), DEGT-1 localization is reduced only in the cell body following
isolation (Figure 5E-H). Furthermore, *degt-1* mutants show reduced MEC-10 signal, and
more importantly, abrogated the isolation-induced reduction in MEC-10 localization at
the quaternary branches and the axon (Figure 5A-D). In the reciprocal experiment,
DEGT-1 localization was affected in *mec-10* mutants, as isolated worms exhibit
increased localization to the cell body compared with WT isolated worms (Figure 5E-H).
Thus, mechanosensory experience also induces plasticity in the localization pattern of
MEC-10 and DEGT-1. We propose this differential localization may be part of the
mechanism that independently and locally modulates dendritic and axonal properties, to
affect both the structure and the function of the PVD, respectively.

Optogenetic stimulation suggests behavioral plasticity is post-sensory

We have shown that sensory deprivation (isolation) affects the localization of two
different mechanoreceptors, the degenerins MEC-10 and DEGT-1, in the dendrites,
axon and soma of the PVD. It is conceivable that the differential localization of
degenerins in different domains of the PVD may affect the morphology and function of
the PVD. Thus, we decided to activate the PVD independently of the endogenous
mechanoreceptors and study whether the escape behavior is similar in isolated and
crowded animals. We hypothesized that if we circumvent the normal degenerin-
mediated mechano-stimulation, exciting the PVD downstream to sensory activity, the
animals grown in isolation will respond to the same degree as if they were in crowded
conditions and independently of the dendritic tree morphology. Thus, we used
optogenetic stimulation with Channelrhodopsin (ChR2) expressed in the PVD (Husson
et al., 2012) to activate the neuron while bypassing sensory perception. We found that
isolation significantly reduced the percentage of worms responding to optogenetic
stimulation of the PVD (Figure 5I), indicating that the plasticity in the response is acting
downstream to PVD activation, mechanosensory channels and signal transduction
pathways. This activation is sensitive to isolation probably because it is acting pre- or
post-synaptically.

To determine whether optogenetic stimulation of isolated animals can reverse the
morphological changes induced by the absence of mechanical stimuli in PVD, we used
optogenetic stimulation of the PVD and found no significant difference between isolated
and isolated optogenetically-stimulated animals in any of the measured PVD structural
characteristic (Figure 5–figure supplement 1). Thus, optogenetic stimulation on isolated

animals is not sufficient to convert their dendritic trees to crowded-like. In addition, the reduced response of isolated animals to optogenetic stimulation suggests the escape response is not dependent on the structure of dendritic trees but on unknown downstream pathways.

Calcium dynamics in response to harsh touch are similar in isolated and crowded animals

Having shown that PVD optogenetic activation elicits a weaker escape response in isolated animals, we asked whether this occurs at the level of PVD itself, or originates further downstream.

To determine whether spontaneous Ca^{++} dynamics in PVD is different between crowded and isolated animals we first observed the baseline calcium dynamics in animals grown isolated and crowded. We found no significant differences in the calcium levels between the isolated and crowded animals (Figure 6–figure supplement 1). To directly elicit PVD mechanosensory activation in isolated and crowded animals, we performed calcium recordings under mechanical stimuli using a custom microfluidic device (Nekimken et. al., 2017). The device allows for the controlled application of mechanical stimulation, and was previously applied to study gentle-touch receptor neurons (TRNs) (Sanfeliu et al., 2023) and AVG (Setty et al., 2022). Similar devices were used for PVD (Cho et al., 2017; Tao et al., 2019). To emulate harsh touch using our device, a 3 bar pressure caused a 15 µm cuticle displacement, greater than the indentation used for studying TRNs and similar to the tail stimulation paradigm used by Setty and colleagues (Setty et al., 2022). Previously we have shown that TRNs responded most efficiently to a high frequency stimulation but were largely insensitive to stimuli delivered below 2 Hz (Nekimken et. al., 2017). We applied three different stimulus profiles: a single step indentation ('step'), similar to a brief manual harsh-touch prod, as well as a gradual pressure 'ramp' increase ('sawtooth'; in which the pressure increases slowly to 3 bar within 2s) and a 10 Hz 'buzz' vibration (see Materials and Methods). Even though we observed a strong and robust activation of PVD for all three stimuli tested (Figure 6), the response to the 'sawtooth' was significantly weaker (Figure 6A). Moreover, we noticed that the response correlated with the abrupt offset of the force in the 'saw tooth', rather than the gradual onset (Figure 6Aiii). To corroborate this result, we also tested a 20 s square step, and indeed found a separate response at the onset as well as the offset (Figure 6–figure supplement 2, Video 7). Thus, PVD, like TRNs, responds better to faster strain rates and adapt under continuous deformation (Eastwood et al., 2015). Using this analysis, we found that both isolated and crowded animals responded similarly and efficiently across all three types of stimuli (Figure 6, Videos 5, 6). Together, these results suggest that the difference in the behavioral response to external mechanical stimuli is due to a pre and/or postsynaptic effect

following mechanosensory deprivation and not a direct consequence of the
morphological changes in the structure of the PVD or its intrinsic activity at the level of
the cell body.

In summary, our genetic and pharmacological evidence suggest that dendritic
structural plasticity is an autonomous activity-dependent homeostatic mechanism.
Combined with optogenetic testing and calcium imaging for mechanically stimulated
PVD, our results indicate that dendritic tree plasticity is independent from downstream
processes that affect escape behavior in response to harsh touch.

Discussion
From the evolutionary point of view, dendritic trees, their morphogenesis (Heiman and
Bülow, 2024) and their structural complexity remain mysterious objects, despite many
efforts to understand the contribution of arborization complexity to dendritic physiology
and function (Hausser and Mel, 2003).

Previous research has demonstrated both cell autonomous (Aguirre-Chen et al.,
2011, Oren-Suissa et al., 2010, Salzberg et al., 2014) and cell non-autonomous (Dong
et al., 2013, Salzberg et al., 2013) mechanisms that regulate the PVD's dendritic
morphogenesis during development. Some studies have also focused on regeneration
and aging effects on the tree structure of the PVD, revealing plasticity in the adult stage
(Iosilevskii and Podbilewicz, 2021, Kravtsov et al., 2017, Oren-Suissa et al., 2017),
similar to what has been shown for *Drosophila* sensory neurons (DeVault et al., 2018).
We found, by using mutants for DEG/ENaCs, chemosensory stimulation, pharmacology,
optogenetic stimulation and glass beads, that the dendritic structure of the PVD and the
behavioral response to harsh touch are activity- and mechanosensory-dependent, but
appear to be chemosensory-independent. It is still conceivable that chemosensation
can affect nociception and proprioception, in a cross modal plasticity mechanism, but at
the moment we do not have evidence for this scenario. In contrast, we show that
DEG/ENaCs mechanosensory channels (degenerins) affect the architecture of the PVD.

Our results suggest that 'nurture', manifested as mechanosensory experience,
activates mechanotransduction signaling, via DEG/ENaCs amiloride-sensitive activity,
to stabilize the homeostatic structure of the dendritic tree in adults. The sensory
experience induced when one body side of the worm is in contact with the agar plate is
not sufficient for this purpose (Figure 7–figure supplement 1); however, other worms on
the plate, or the presence of glass beads, elicit significant structural alterations.
Moreover, the cell autonomous activity of MEC-10 in the PVD was found to be both
necessary and sufficient to preserve the crowded-experience phenotype in terms of the
simplified structure of the PVD. While there are other multiple plausible explanations,
our failure in remodeling the isolated-state arbor by repeated optogenetic activation of
the PVD may provide further evidence for the necessity of DEG/ENaCs in this structural
plasticity, since neuron activation by ChR2 bypasses the activation of mechanically-

gated DEG/ENaC channels. Figure 7 depicts our working model, where the amount of
mechanosensory stimulation, in crowded or isolated conditions, affects the expression
and localization of MEC-10 and DEGT-1 in different compartments of the PVD.
Localization of MEC-10, probably by forming higher-order complexes with other
DEG/ENaCs, can affect the structure of the PVD at the level of the dendritic tree. The
morphological and behavioral phenotypes shown here were rescued by a PVD-specific
expression of MEC-10, supporting such a cell autonomous mechanism. The
experience-induced structural plasticity seems homeostatic at the individual branch
dynamics level in the adult. In contrast, we show that the behavioral response to harsh
touch is modulated by mechanosensory experience during development and by the
presence of DEG/ENaCs, and perhaps involves other neurons downstream. Based on
optogenetic stimulation, we presume that the behavioral plasticity is a pre- and/or post-
synaptic property, mediated by DEG/ENaCs (Hill and Ben-Shahar, 2018) and related to
neurotransmission modulation, independently of the structure of the PVD dendritic tree.
These structural and behavioral plasticity are separated in time (adulthood vs.
development) and space (dendrite vs. axon; Figure 7). We found no correlation or
causation between the structure and the nociceptive function of the PVD. It is possible
that such a link exists, but our physiological and behavioral outputs are not sensitive
enough to detect it.

The PVD is a polymodal sensory neuron, and it has been shown that its dendrite
structure is related to body posture, while the response to harsh touch is related to
synaptic connection (Tao et al., 2019). Current assays for proprioception in *C. elegans*
largely rely on video movement tracking, and it is possible the changes in the dendrite
structure caused by sensory deprivation are too minor to cause any quantitative
changes in body posture. Elegant work by Tao and colleagues (2019) supports the role
of MEC-10 in PVD-mediated proprioception, however has some discrepancies with our
results with regard to harsh touch response, which may be related to the different
methodologies used, as well as to the animals' age and mechanosensory experience
(Inberg and Podbilewicz., 2018).

Calcium imaging reveals PVD responds to fast changes in mechanical stress,
which includes pressure onset as well as offset (Figs. 6Aiii, Figure 6–figure supplement
2, Video 7). We found PVD to be insensitive to slow stimuli and adapt under constant
mechanical body deformation. This adaptation may facilitate the response to a wide
dynamic range of stimuli applied to the body wall and is thus an essential feature of
mechanical nociception. While TRNs respond to both the application and release of a
step stimulus (Eastwood et al., 2015), other mechanosensory channels in different
mechanosensitive systems respond preferentially to either on or off instead of a
symmetric response to both (Katta et al., 2015). Interestingly, previous calcium imaging
studies on PVD have not described the response to offset (Chatzigeorgiou et al., 2010;
Husson et al., 2012; Tao et al., 2019; Cho et al., 2017, Cho et al., 2018). One possible

explanation to our new observation may be due to the fact that we are the first to apply
'sawtooth' stimuli to the PVD with a slow increase such that the activity is not induced.
An alternative explanation is that in previous work, the onset and offset were too close
(e.g. for a 0.5 s step as in Tao et al., 2019), such that the onset and offset could not be
differentiated.

Our results indicate that sensory experience does not alter the magnitude of the
calcium activation (Figure 6), as tested using three different stimulation profiles. These
findings support additional effects downstream to the dendritic structural plasticity,
which govern behavioral plasticity and add another layer of support to the separation in
time and space between the structure of the PVD and its function.

Somatosensory activation in vertebrates plays a prominent role in shaping the
structural and functional properties of dendritic spines, mainly studied in the central
nervous system (Gyorffy et al., 2018, Holtmaat and Svoboda, 2009, Xu, Yu et al., 2009,
Yang et al., 2009). In contrast to mammalian cortical neurons, much less is known
about sensory neurons' degree of plasticity and the molecular mechanisms utilized
during adulthood. Here we suggest that degenerins mediate mechanosensation-
induced dendritic growth in sensory dendrites. The dendritic plasticity we described
bears resemblance to the activity-dependent effect of glutamatergic signaling and
NMDA receptors. Activity via NMDA affects dendritic spines as an upstream mechanism
of cell signaling, resulting in structural modifications (Nagerl et al., 2004, Star et al.,
2002, Stein et al., 2021, Zhang et al., 2015). It is possible that degenerins mediate
mechanosensory signaling sensation, by activating cationic gradients (Kellenberger and
Schild, 2002), leading to activation of downstream intracellular signaling pathways
(Ghiretti et al., 2014, Sin et al., 2002, Vaillant et al., 2002, Zhou et al., 2006), which in
turn stabilize local, actin-mediated (Halpain, 2018, Luo, 2002, Zuo et al., 2005)
structural plasticity in the PVD dendritic branches. In *Drosophila,* at the epithelium,
mechanosensitive ion channels, together with E-cadherin-catenin complexes and
calcium sensing mechanisms affect epithelial morphogenesis (Choudhury et al., 2021).
In parallel, DEG/ENaCs may also modulate pre- and post-synaptic homeostatic
signaling in the harsh touch circuit, as has been shown in neuromuscular junctions
(Younger et al., 2013). While this study focuses on degenerins activation in the PVD
with emphasis on dendritic structures, future studies may establish the pre- and -post-
synaptic mechanisms which act downstream, on the transcriptional and translational
levels. These directions for future studies have the potential to increase our
understanding of the mechanisms that couple sensory experience, to structural dendritic
plasticity.

In summary, we propose that the combinatorial actions of DEG/ENaCs have
mechano-signaling functions mediating plasticity in sensory dendritic trees, and provide
mechanistic insights into dendritic structural responses to sensory experience in
adulthood and the behavioral consequences of such adaptations during development.

Materials and Methods

Strains
Nematode strains were maintained according to standard protocols (Brenner, 1974). The list of the strains is presented in Supplementary file 1a. Strains of the DEG/ENaCs family obtained from the CGC (JPS282: *asic-1*(*ok415*) I, ZB2551: *mec-10*(*tm1552*) X, VC2633: *degt-1*(*ok3307*) V) were crossed with BP709: *hmnIs133[ser-2Prom3::kaede]*. The validation of F2 homozygotes for the DEG/ENaCs deletions (including single, double and triple mutants) was performed by PCR amplification of the genomic area containing the deleted region.

Primers for Multiplex PCR
The list of the primers used is presented in Supplementary file 1b.

Spinning disk confocal microscopy
Prior to imaging, the worms were mounted on a 10% agar pad placed on a microscope glass slide, in 1 µl polystyrene bead suspension (100 nm diameter; Polysciences, Inc.) for their mechanical restraint, and sealed with a coverslip for complete physical immobilization (Kim et al., 2013). The PVD neuron was visualized using a Yokagawa CSU-X1 spinning disk, Nikon eclipse Ti inverted microscope and iXon3 camera (ANDOR). Images were captured with MetaMorph, version 7.8.1.0. For each worm, a sequential z-series image stack (step size of 0.35 µm) was obtained with an oil Plan Fluor 40X (NA 1.3) lens around the PVD cell body, encompassing approximately 50-100 µm segments both anteriorly and posteriorly.

Data analysis
The analysis of the PVD structure was performed for the area surrounding the cell body. All images were analysed with ImageJ, version 1.48 (NIH), in TIFF format, by producing a maximal intensity z-series projection and converting it to negative form (invert lookup table) for improved visibility. Ectopic branching was defined as described previously (Hausser and Mel, 2003). Briefly, non-ectopic branches form the 'ideal' WT candelabrum of the late L4 stage, whereas excess branches, which create non-quaternary terminal ends, are considered ectopic, as illustrated with dashed lines for ectopic branching in Figure 2A. The total number of ectopic and non-ectopic branches was quantified for each image, and presented as a fraction (ectopic/non-ectopic branches).
The geometry of each quaternary branch was defined in the following manner: Straight geometry- all the pixels that constitute the branch are positioned on a straight line generated with ImageJ. The width of the line (1 pixel) was constant for the entire sets of experiments. The number of straight quaternary branches were divided by the total quaternary branches in the image for each worm and presented as a percentage. The analysis was done only for worms which did not move through the Z series. Moving

612 worms were excluded from the experiment. Self-avoidance defects- the number of
613 events where two adjacent candelabra overlapped (no gap formation), was divided by
614 the total number of gaps between the candelabra within the frame (Figure 2A, Figure 4
615 – figure supplement 3). The self-avoidance values are presented as percentage.

Behavioral procedures

Harsh touch assay
After 72 h, adult worms (both isolated and those from the crowded plate as described
below) were transferred using an eyelash each to a new agar plate, freshly seeded with
150 μl OP50 (about 16 h after seeding). This step is required to avoid a thick edge of
the bacterial lawn, which might interfere with harsh touch response measurement. After
~45 minutes in the plate, the non-moving worms were prodded with a platinum wire
posterior to the vulva, above the interface between worm's body and the agar plate
(Way and Chalfie, 1988), every 10 seconds, and the number of responses to harsh
touch was counted. Animals with a functional PVD moved, sometimes backing up. More
than one response constitutes a responsive animal. The non-responding worms were
defined if two prodding events were observed sequentially without response. The
percentage of responding worms was calculated for each genotype and treatment. The
experimenter was blind to both the genotype and the treatment- crowded or isolated.

Isolation of embryos
To establish the method for conspecifics based mechanosensory stimulation, we
performed several calibration experiments with different population densities (250
embryos, progeny of 15 adults, progeny of 30 adults) to test conditions for the crowded
plate.
We found a gradual increase in the crowding effect and reduced variation in the
measured parameter. Following that, we decided to work with the progeny of 30
crowded adults as a source of mechanosensory signal.
638
The worm isolation procedure was based on previous work (Rose et al., 2005) with a
few modifications, as indicated. Isolated animals were grown on 6 cm agar plate with
150 μl of OP50 *E. coli*, while crowded plate worms were grown on 600-700 μl OP50 to
prevent starvation. The embryos and adult worms were isolated with platinum wire. The
plates were sealed with one layer of Parafilm M and placed into a plastic box, at 20 °C,
for the entire experiment.
Three experimental groups were used for the 72 h (96 h of experiment was performed
only for *mec-4* worms, since they were L4- very young adults at 72 h) isolation
experiment: **(1)** Single isolated embryos **(2)** Crowded worms - the progeny of 30 young,
non-starved, adults (approximately, 7,000-9,000 worms in different developmental
stages, without approaching starvation) **(3)** Crowded adult worms that were isolated for
a certain amount of time as adults.

After 48/72/96 h (according to the experiment), age-matched worms from each group
were transferred to 10% agar pad slides for imaging, as described above (Spinning disk
confocal microscopy) (Rose et al., 2005).

Isolation of adults for 2, 5, 15 or 24 hours
Crowded plates were prepared as described above, with progeny from 30 adult
hermaphrodites. Animals were separated into individual plates for the desired time
window before 72 h have passed (*i.e.* after ~70 h for 2 h isolation, ~67 h for 5 h isolation
etc.). Worms were isolated using an eyelash into new plates containing 150 µl of OP50.
At 72 h, the PVD of isolated animals was imaged and compared with age-matched
animals which remained in the crowded plate of origin (see Figure 1B).

Optogenetics stimulation for harsh touch
Crowded and isolated worms (ZX819: *lite-1(ce314) X;*
zxIs12[pF49H12.4::ChR2::mCherry pF49H12.4::GFP]) were grown on OP50 with 100
µM All Trans Retinal (ATR, Sigma R2500), in order to obtain functional
Channelrhodopsin (Husson et al., 2012) (as a control we tested the response of worms
that were grown on ethanol alone, no response was detected for these worms). After 72
h the worms were singly mounted with eyelash on a chunk (1 cm X 1 cm) of agar that
was mounted on a microscope glass slide. The agar contained fresh but dry OP50, with
100 µM ATR. About 30 minutes following the transfer the worms were tested for the
response to light. Worms were stimulated at 488 nm wavelength, with laser intensity of
40% and exposure time of 100 ms, with 10X Plan Fluor (NA 0.3) for ~1 second and the
forward acceleration response was measured. The microscope system is as described
above. The experimenter determined the presence or the absence of forward
acceleration in response to light activation.

Optogenetics stimulation of isolated worms
Isolated eggs were grown as described at the 'Isolation of embryos' section.
Isolated worms (ZX819: *lite-1(ce314) X; zxIs12[pF49H12.4::ChR2::mCherry*
pF49H12.4::GFP]) were grown on 150 µM OP50 with 100 µM ATR (Husson et al.,
2012) or 0.3% ethanol as control. After 72 h both the crowded and isolated worms were
singly mounted with eyelash onto a cube (with surface area of about approx.1 cm^2,
depth of 0.5 cm) of agar layered with fresh, fully dried, OP50 with 100 µM ATR or 0.3%
ethanol as a control group. The worms were stimulated with 488 nm wavelength, with
laser intensity of 40% and exposure time of 100 ms, with 10X Plan Fluor (NA 0.3) for
~60 seconds, every 5 min during 4 h. Stimulation and recording was performed with the
microscope system described above.
As a control, before each experiment, crowded ZX819 worms were tested for forward
acceleration as a positive control to the functionality of the ChR2. After the end of 4 h
stimulation, the worms were mounted for imaging of the PVD at 40X as described under
"Spinning disk confocal microscopy" section.

Isolation with chemical stimulation

The *glp-4(bn2)* mutants which are sterile at 25°C were used (~40 worms for each plate), in order to prepare conditioned/chemically stimulated plates (Maures et al., 2014) prior to growing isolated animals. The *glp-4* mutants were transferred at early larval stage (L1, L2) to a new agar plate for 96 h at 25°C. After the removal of the *glp-4* mutants, the embryo isolation procedure was used, as described before at 20°C.

Isolation with glass beads

Single embryos were isolated to agar plates with 150 µl OP50 and 2.5 g of glass beads (1 mm diameter) were placed on the OP50 lawn in the middle of the plate. The worms were isolated for 72 h and tested for response to harsh touch as described above.

Pharmacology

Amiloride hydrochloride hydrate (Sigma, #A7410) 1 M stock solution in DMSO was stored at -20°C. A final concentration of 3 mM amiloride in 0.03% DMSO was prepared in OP50 bacteria (LB medium) and seeded on NGM plates. 650 µl of the OP50 mixture was seeded on each plate. As a control, 0.03% DMSO was added to OP50 bacteria. For each plate (control 0.03% DMSO or 3 mM amiloride) 30 non-starved adult worms were added. After 72 h at 20°C the progeny of the 30 adults were tested as young adults for their PVD morphology and their response to posterior harsh touch, as described in the previous sections.

Anesthetics and long-term imaging

In addition to immobilization of worms with 100 nm polystyrene beads as described in the "Spinning disk confocal microscopy" section above, two additional methods for long term imaging and pharmacological effects assays were used: 1% tricaine (Sigma, A5040) in M9 buffer or a mixture of 0.01% tetramisole (Sigma, T-1512) and 0.1% tricaine in M9 buffer were utilized.
Worms were exposed to the anesthetics (in a glass well) for ~20 min until paralyzed, then transferred to 3% agar pads with 1 µl of the anesthetics. Sequential z-axis image series (0.6 µm step-size) were taken, as described above, every 5 min for 2-3 h.

Imaging individual worms- crowded to isolated

Worms were grown at crowded conditions, as described above. At ~24 h of adulthood, individual worms were placed in a 1% tricaine solution in M9 buffer for 10 min until paralyzed and transferred to 3% agar slides with 1 µl 1% tricaine. Following a short PVD imaging session, the worms were recovered from the slide with M9 and isolated to a new plate for 4 h. Each 4 h-isolated animal was anesthetized once again with 1% tricaine for 10 min prior to PVD imaging.

Analysis of DEG/ENaCs localization in the PVD

Two DEG/ENaCs protein constructs, *pF49H12.4*::MEC-10::mCherry and *pF49H12.4::DEGT-1*::mCherry (plasmids kindly provided by W. Schafer's lab, (Chatzigeorgiou et al., 2010)) were analysed for their localization in the PVD, by

comparing worms expressing the co-injection marker *punc-122*::GFP raised in crowded
or isolated conditions, in a similar behavioral assay as described in Figure 1B. A z-
series (step size of 0.35 µm, immobilization performed with polystyrene beads, as
described in the microscopy section) of the area around the cell body of the PVD was
obtained with a 60X Apochromat (NA 1.4) lens. The images (after maximal intensity
projection of the Z-series) were encoded so the analysis was performed in a blind
manner. The presence or absence of fluorescent signal was examined in three
compartments: The cell body, the quaternary branches and the axon of the PVD.

Rescue strains
Worms from BP1022 (*mec-10(tm1552) X*; *hmnIs133[ser-2Prom3::Kaede]*; *him-5(e1490)*
V) were injected into the gonad with a rescuing plasmid for *mec-10*, with a PVD specific
promoter (pWRS825: *ser-2Prom3::mec-10* genomic) kindly provided by W. Schafer's
lab.(Chatzigeorgiou et al., 2010). The injection mix contained *myo-2p::GFP* (20 ng/µl) as
a co-transformation marker and pWRS825 (80 ng/µl). For both behavioral and structural
characterization, the *him-5; mec-10* strain, with and without rescuing plasmid, shared
the same crowded plate and were differentiated by the presence or absence of the co-
injection marker.

Effect of plate on PVD in contact with agar and the opposing side
Embryos were singled into plates with 150 µl OP50 bacteria at 20°C, and
analyzed as young adults 72 h later. Every animal was transferred using an
eyelash into freshly seeded plates (<24 h), and noted for its side orientation.
Once on these experiment plates, animals were observed for their side
orientation at 30-minute intervals for a minimum of 3 h prior to microscopy.
For microscopy of both PVD neurites, while few animals were removed from the slide
and repositioned using an eyelash, most animals were imaged using a coverslip
'sandwich' method (adapted from (Sulston et al., 1983)), whereby the animal is pressed
on an agar pad between two large coverslips, which are subsequently flipped as a
single unit and observed from the other side. Briefly, a thin 3% agar pad was pressed
onto a 24x50 mm coverslip (170±5 µm thick), placed on a standard microscope slide for
ease of handling. Animals were singly placed in a drop of 0.05% tetramisole (Sigma, T-
1512) in M9 on the agar pad using an eyelash, and a second, similar coverslip was
immediately placed above. The two coverslips were sealed shut onto the supporting
slide using two thin strips of marking tape, and once the animal was immobilized the
PVD on the side closest to the objective was imaged as described above (see Spinning
Disk confocal microscopy). For viewing the opposite neurite, the tape was carefully
peeled and the two coverslips flipped as one, taped again, and similarly imaged. Note
that this method inevitably makes the second side imaged to be fainter due to the laser
penetrating through the agar pad layer itself; laser intensities were typically increased
by up to 5% in order to compensate for the decrease in fluorescence.

Micromechanical experiments and calcium image recordings

Micromechanical experiments were performed in microfluidic devices which have
been prepared as described using a 15:1 PDMS base-polymer/curing agent ratio
(Fehlauer et al., 2018; Setty et al., 2022), by replicating a design presented in
(Nekimken et. al., 2017). Prior to the experiments, single animals raised in
isolated conditions or multiple animals raised in the crowded conditions were
loaded into the device as described (Fehlauer et al., 2018). After gently pushing
individual animals into the trapping channel, a sequence of three different
stimulus profiles (2 s each, 3 bar max pressure, immediate 'step', sawtooth (or
gradual 'ramp') or 10 Hz vibration `buzz') were applied (Nekimken, et. al., 2017)
each separated by 20 s. The pressure-deformation relation was previously
characterized (Setty et al., 2022). Accordingly, we found a deformation of 15 μm
with a 3 bar pressure on the channel. In all cases, the animals ended up with the
lateral sides facing the side walls of the chip, providing a ventral or dorsal view of
the body. Stimuli location was within 200 μm of the PVD cell body (either anterior
or posterior), and ipsilateral or contralateral.
Imaging was performed using a 40x (NA1.1) water immersion lens on an inverted
Leica DMi8 equipped with a Hamamatsu Orca Flash 4 camera, and a
fluorescence cube with a multiband dichroic mirror (Semrock Quadband
FF409/493/573/652). Calcium sensitive and insensitive channel were divided into
its spectral components using a Hamamatsu Gemini W-View beamsplitter with a
538 nm edge dichroic (Semrock, FF528-FDi1-25-36) and projected on either half
of the camera chip. A Lumencor Spectra X LED light source (using the cyan-488
and green-555 LED) was used to excite the Ca-sensitive and insensitive dye,
respectively, and slaved to the camera acquisition with an exposure time of 100
ms in HCImage Software (version 4.4.2.7). Likewise, the pressure application
(ElveFlow OB1 with an 8 bar channel) was synchronized to the camera
acquisition using the SMA trigger out, with a delay equal to the pre-stimulus
period.
The whole procedure including loading, focusing, and stimulation did not take
longer than 5-10 min per animals to avoid pre-stimulus adaptation. All animals
remained viable after the procedure, as visualized by vivid and lively thrashing in
the buffer.
801
Analysis of calcium traces after stimulation
All quantifications were performed on the PVD cell body after ipsilateral application of
the mechanical stimulus. No dependence of the calcium signals on the
anterior/posterior position was noticed. Contralaterally applied stimuli lead to a strongly
reduced response, without detectable difference between the two conditions (isolated,
crowded), but were not further analyzed. To extract calcium signals from the cell body, a
procedure described in (Porta-de-la-Riva et al, 2023) was followed. Calcium traces were

background subtracted, baseline corrected, and normalized by the first 10 s preceding
the first mechanical stimulus. Results are displayed as fold change, compared to
baseline calcium levels.
812
PVD basal activity
To image spontaneous calcium traces in PVD, we used worms that were raised
in isolated or crowded conditions, as described above. Using a drop of S-basal
buffer, adult worms were inserted in a dual olfactory chip (Gat et al, 2023; Karimi
et al, 2024), with a flow rate of 0.005 ml/min. Imaging was done with a Zeiss LSM
880 confocal microscope using a 40x magnification water objective. When the
worm was properly located inside the chip with minimal movement, PVD was
imaged for 1:30 minutes, with imaging rate of 6.667 Hz.
For analysis, the GCaMP6s fluorescence intensity was measured using FIJI.
ROIs of PVD soma were picked manually to extract mean intensity values.
Data analysis was performed using Python 3.10. For each worm, the baseline
fluorescent level (F0) was calculated by averaging the mean values of 100 frames in the
beginning of each recording. Then, for each frame, ΔF was calculated by subtracting F0
from the value of that time point, and the result was divided by F0, to normalize the
differences in the fluorescence baseline levels between individuals ($\Delta F/F0$). For Figure
6–figure supplement 1C, the derivative of $\Delta F/F0$ was calculated and averaged across
worms in each time point. Then, a histogram representing the distribution of the average
derivative was plotted.
831
Statistics and data plotting
At least two independent experiments constitute the data set described in each figure.
For the morphological characterization of the PVD the results are expressed as means
(blue circle) ± standard error of the mean (s.e.m.). In the boxplot (first, third quartiles)
the upper whisker extends from the hinge to the highest value that is within 1.5 * IQR
(inter-quartile range), the distance between the first and third quartiles. The lower
whisker extends from the hinge to the lowest value within 1.5 * IQR of the hinge.
The statistical analyses were performed with SPSS software (IBM, version 20) and "R
package". Two-tailed tests were performed for the entire data sets.
Since for many experiments the distribution of the data was not normal, nonparametric
tests were used: Mann-Whitney test for comparison between two independent groups.
Kruskal-Wallis test was used for multiple comparisons for more than two groups.
For proportions (percentage worms) ± standard error of proportion was calculated.
Fisher's exact test was used for analysis of differences in proportions. To estimate the

variability in proportion we calculated the Standard Error of Proportion: $\sqrt{\frac{(1-p)\cdot p}{n}}$

The dot plot figures were prepared with "R package", the bar charts with Microsoft Excel
software. Final figures were prepared with Adobe Illustrator CS version 11.0.0.

Discriminant analysis

Eight different strains (WT and seven DEG/ENaCs), with two treatments (crowded, isolated worms) for each strain were analyzed for Linear discriminant analysis for morphological characteristics, to evaluate similarity between different strains and treatments. Each worm in the data set was characterized by the three morphological characteristics (the fraction of ectopic branching, the percentage straight quaternary branches and the percentage of self-avoidance defects). The centroid for morphological characterization was calculated for each condition and represented by a square. Data from independent harsh touch experiments are shown for each group. The analysis was performed using SPSS 20.

Acknowledgments

We thank current and former lab members for their intellectual and technical support. Veronika Kravtsov, Sagi Levy, Anna Meledin, Tom Shemesh, Shay Stern, Yehuda Salzberg and Alon Zaslaver for critically reading and commenting on the manuscript. Ehud Ahissar, Dan Cassel, Michel Labouesse and Kang Shen for fruitful discussions. William Schafer, Max Heiman, Hannes Bülow, Yehuda Salzberg and Alexander Gottschalk for plasmids and strains. Some strains were provided by the Caenorhabditis Genetics Center (CGC), which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). This work was supported by grants from the Israel Science Foundation (442/12 and 257/17, BP), Adelis Fund (2023479, BP), and the Ministry of Science and Technology (3-13022, BP). MK acknowledges PID2021-123812OB-I00 and the CNS2022-135906 project funded by MCIN /AEI /10.13039/501100011033 / FEDER, UE, and the HFSPO through the RGP021/2023.

References

Aceves J, Erlij D, Martinez-Maranon R (1970) The mechanism of the paralysing action of tetramisole on Ascaris somatic muscle. British journal of pharmacology 38: 602-607

Aguirre-Chen C, Bülow HE, Kaprielian Z (2011) C. elegans bicd-1, homolog of the Drosophila dynein accessory factor Bicaudal D, regulates the branching of PVD sensory neuron dendrites. Development 138: 507-18

Albeg A, Smith C, Chatzigeorgiou M, Feitelson DG, Hall DH, Schafer WR, Miller DM, 3rd, Treinin M (2011) C. elegans multi-dendritic sensory neurons: Morphology and function. Mol Cell Neurosci 46: 308-317

Alvarez VA, Sabatini BL (2007) Anatomical and physiological plasticity of dendritic spines. Annual review of neuroscience 30: 79-97

Bailey NW, Moore AJ (2018) Evolutionary Consequences of Social Isolation. Trends in ecology & evolution 33: 595-607

Baldin JP, Barth D, Fronius M (2020) Epithelial Na+ Channel (ENaC) Formed by One or Two Subunits Forms Functional Channels That Respond to Shear Force. Frontiers in Physiology 11

Ben-Shahar Y (2011) Sensory Functions for Degenerin/Epithelial Sodium
        Channels (DEG/ENaC). Adv Genet 76: 1-26
Bianchi L, Driscoll M (2002) Protons at the gate: DEG/ENaC ion channels help
        us feel and remember. Neuron 34: 337-340
Brenner S (1974) The genetics of *Caenorhabditis elegans*. Genetics 77: 71-94
Caneo M, Julian V, Byrne AB, Alkema MJ, Calixto A (2019) Diapause induces
        functional axonal regeneration after necrotic insult in C. elegans. PLoS
        Genet 15: e1007863
Chalfie M, Sulston J (1981) Developmental genetics of the mechanosensory
        neurons of Caenorhabditis elegans. Dev Biol 82: 358-370
Chatzigeorgiou M, Yoo S, Watson JD, Lee WH, Spencer WC, Kindt KS, Hwang
        SW, Miller DM, 3rd, Treinin M, Driscoll M, Schafer WR (2010) Specific roles
        for DEG/ENaC and TRP channels in touch and thermosensation in C.
        elegans nociceptors. Nature neuroscience 13: 861-868
Chen Y, Bharill S, Isacoff EY, Chalfie M (2015) Subunit composition of a
        DEG/ENaC mechanosensory channel of Caenorhabditis elegans. Proc Natl
        Acad Sci U S A 112: 11690-11695
Cho Y, Porto DA., Hwang H, Grundy LJ, Schafer WR, Lu H (2017) Automated
        and controlled mechanical stimulation and functional imaging: In vivo in C.
        elegans. Lab on a chip, 17, 15, 2609 – 2618
Cho Y, N. Oakland DN, Lee SA, Schafer WR and Lu H (2018) On-chip functional
        neuroimaging with mechanical stimulation in Caenorhabditis elegans larvae
        for studying development and neural circuits. Lab on a chip 18, 601–609
Collet J, Spike CA, Lundquist EA, Shaw JE, Herman RK (1998) Analysis of osm-
        6, a gene that affects sensory cilium structure and sensory neuron function
        in Caenorhabditis elegans. Genetics 148: 187-200
DeVault L, Li T, Izabel S, Thompson-Peer KL, Jan LY, Jan YN (2018) Dendrite
        regeneration of adult Drosophila sensory neurons diminishes with aging and
        is inhibited by epidermal-derived matrix metalloproteinase. Genes Dev 32:
        402-414
Dong X, Liu OW, Howell AS, Shen K (2013) An extracellular adhesion molecule
        complex patterns dendritic branching and morphogenesis. Cell 155: 296-
        307
Dong X, Shen K, Bülow HE (2015) Intrinsic and extrinsic mechanisms of dendritic
        morphogenesis. Annu Rev Physiol 77: 271-300
Driscoll M, Chalfie M (1991) The *mec-4* gene is a member of a family of
        Caenorhabditis elegans genes that can mutate to induce neuronal
        degeneration. Nature 349: 588-593
Eastwood A L., Sanzeni A, Petzold B C, Park S J, Vergassola M, Pruitt B L, and
        Goodman M B. (2015) Tissue mechanics govern the rapidly adapting and
        symmetrical response to touch. PNAS, E6955–E6963
Fehlauer H, Nekimken AL, Kim Anna A, Pruitt BL, Goodman MB, Krieg M (2018)
        Using a Microfluidics Device for Mechanical Stimulation and High
        Resolution Imaging of *C. elegans.* J Vis Exp. (132):56530

Fox K, Wong RO (2005) A comparison of experience-dependent plasticity in the
visual and somatosensory systems. Neuron 48: 465-477
Gat A, Pechuk V, Peedikayil-Kurien S, Karimi S, Goldman G, Sela S, Lubliner J,
Krieg M, and Oren-Suissa M (2023) Integration of spatially opposing cues
by a single interneuron guides decision-making in C. elegans. Cell reports
42, 113075
Ghiretti AE, Moore AR, Brenner RG, Chen LF, West AE, Lau NC, Van Hooser
SD, Paradis S (2014) Rem2 Is an Activity-Dependent Negative Regulator of
Dendritic Complexity In Vivo. J Neurosci 34: 392-407
Gillespie PGW, R. G. (2001) Molecular basis of mechanosensory transduction.
Nature 413: 194-202
Gobetto MN, Gonzalez-Inchauspe C, Uchitel OD (2021) Histamine and
Corticosterone Modulate Acid Sensing Ion Channels (ASICs) Dependent
Long-term Potentiation at the Mouse Anterior Cingulate Cortex. Neurosci
Behav Physiol 460: 145-160
Goodman CS, Shatz CJ (1993) Developmental mechanisms that generate
precise patterns of neuronal connectivity. Cell 72 Suppl: 77-98
Goodman MB, Sengupta P (2019) How Caenorhabditis elegans Senses
Mechanical Stress, Temperature, and Other Physical Stimuli. Genetics 212:
25-51
Gyorffy BA, Kun J, Torok G, Bulyaki E, Borhegyi Z, Gulyassy P, Kis V, Szocsics
P, Micsonai A, Matko J, Drahos L, Juhasz G, Kekesi KA, Kardos J (2018)
Local apoptotic-like mechanisms underlie complement-mediated synaptic
pruning. Proc Natl Acad Sci U S A 115: 6303-6308
Hall DH, Treinin M (2011) How does morphology relate to function in sensory
arbors? Trends Neurosci 34: 443-51
Halpain S (2018) Avoiding Sibling Conflict: Lessons from Dendrite Self-
Avoidance in C. elegans. Neuron 98: 864
Harlow HF, Dodsworth RO, Harlow MK (1965) Total social isolation in monkeys.
Proc Natl Acad Sci U S A 54: 90-97
Hausser M, Mel B (2003) Dendrites: bug or feature? Curr Opin Neurobiol 13:
272-283
Hedgecock EM, Sulston JE, Thomson JN (1983) Mutations affecting
programmed cell deaths in the nematode *Caenorhabditis elegans*. Science
220: 1277-1279
Heiman MG, Bülow HE (2024) Dendrite morphogenesis in Caenorhabditis
elegans. Genetics 227(2), iyae056
Hill AS, Ben-Shahar Y (2018) The synaptic action of Degenerin/Epithelial sodium
channels. Channels 12: 262-275
Holtmaat A, Svoboda K (2009) Experience-dependent structural synaptic
plasticity in the mammalian brain. Nat Rev Neurosci 10: 647-658
Hu HT, Huang TN, Hsueh YP (2020) KLHL17/Actinfilin, a brain-specific gene
associated with infantile spasms and autism, regulates dendritic spine
enlargement. J Biomed Sci 27: 103

Huang M, Chalfie M (1994) Gene interactions affecting mechanosensory transduction in Caenorhabditis elegans. Nature 367: 467-470

Huebschman JL, Corona KS, Guo Y, Smith LN (2020) The Fragile X Mental Retardation Protein Regulates Striatal Medium Spiny Neuron Synapse Density and Dendritic Spine Morphology. Front Mol Neurosci 13: 161

Husson SJ, Costa WS, Wabnig S, Stirman JN, Watson JD, Spencer WC, Akerboom J, Looger LL, Treinin M, Miller DM, 3rd, Lu H, Gottschalk A (2012) Optogenetic analysis of a nociceptor neuron and network reveals ion channels acting downstream of primary sensors. Curr Biol 22: 743-752

Inberg S, Meledin A, Kravtsov V, Iosilevskii Y, Oren-Suissa M, Podbilewicz B (2019) Lessons from worm dendritic patterning. Annual review of neuroscience 42: 365-383

Inberg S, Podbilewicz B (2018) Sensory experience controls dendritic structure and behavior by distinct pathways involving degenerins. bioRxiv [Preprint]. doi: 10.1101/436758)

Inberg S, Iosilevskii Y, Podbilewicz B (2021) Sensory experience controls dendritic structure and behavior by distinct pathways involving degenerins. bioRxiv [Preprint]. doi: 10.1101/436758)

Iosilevskii Y, Podbilewicz B (2021) Programmed cell fusion in development and homeostasis. Curr Top Dev Biol 144: 215-244

Iosilevskii Y, Podbilewicz B (2024) The PVD neuron has male-specific structure and mating function in *C. elegans.* bioRxiv 2024.05.19.594847

Jan YN, Jan LY (2010) Branching out: mechanisms of dendritic arborization. Nat Rev Neurosci 11: 316-328

Karimi S, Gat A, Agazzi C, Oren-Suissa M, and Krieg M (2024) Automated dual olfactory device for studying head/tail chemosensation in Caenorhabditis elegans. APL Bioeng. 8(2), 026104. https://doi.org/10.1063/5.0187441

Katta S, Krieg M, Goodman MB. (2015) Feeling force: physical and physiological principles enabling sensory mechanotransduction. Annu Rev Cell Dev Biol. 31:347-371.

Katz EM, Chu DK, Casey KM, Jampachaisri K, Felt SA, Pacharinsak C (2020) The Stability and Efficacy of Tricaine Methanesulfonate (MS222) Solution After Long-Term Storage. J Am Assoc Lab Anim Sci. 12: 393-400

Katz LC, Shatz CJ (1996) Synaptic activity and the construction of cortical circuits. Science 274: 1133-1138

Kellenberger S, Schild L (2002) Epithelial Sodium Channel/Degenerin Family of Ion Channels: A Variety of Functions for a Shared Structure. Physiol Rev 82: 735-767

Kim E, Sun L, Gabel CV, Fang-Yen C (2013) Long-Term Imaging of Caenorhabditis elegans Using Nanoparticle-Mediated Immobilization. PLoS One 8: e53419

Kolb B, Whishaw IQ (1998) Brain plasticity and behavior. Annu Rev Psychol 49: 43-64

Kravtsov V, Oren-Suissa M, Podbilewicz B (2017) The fusogen AFF-1 can
rejuvenate the regenerative potential of adult dendritic trees via self-fusion.
Development 144: 2364-2374
Krieg M, Pidde A, Das R (2022) Mechanosensitive body–brain interactions
in *Caenorhabditis elegans.* Current Opinion in Neurobiology 75:102574
Kuhlman SJ, O'Connor DH, Fox K, Svoboda K (2014) Structural Plasticity within
the Barrel Cortex during Initial Phases of Whisker-Dependent Learning. J
Neurosci 34: 6078-6083
Li W, Kang L, Piggott BJ, Feng Z, Xu XZ (2011) The neural circuits and sensory
channels mediating harsh touch sensation in Caenorhabditis elegans. Nat
Commun 2: 315
Luo L (2002) Actin Cytoskeleton Regulation in Neuronal Morphogenesis and
Structural Plasticity. Annu Rev Cell Dev Biol 18: 601-635
Mango D, Nistico R (2020) Role of ASIC1a in Normal and Pathological Synaptic
Plasticity. Reviews of physiology, biochemistry and pharmacology 177: 83-
100
Maures TJ, Booth LN, Benayoun BA, Izrayelit Y, Schroeder FC, Brunet A (2014)
Males shorten the life span of C. elegans hermaphrodites via secreted
compounds. Science 343: 541-4
Miller-Fleming TW, Petersen SC, Manning L, Matthewman C, Gornet M, Beers A,
Hori S, Mitani S, Bianchi L, Richmond J, Miller DM (2016) The DEG/ENaC
cation channel protein UNC-8 drives activity-dependent synapse removal in
remodeling GABAergic neurons. eLife 5
Nagerl UV, Eberhorn N, Cambridge SB, Bonhoeffer T (2004) Bidirectional
Activity-Dependent Morphological Plasticity in Hippocampal Neurons.
Neuron 44: 759-767
Nekimken A, Holger F, Kim A, Manosalvas-KS, Ladpl P, Memon F, Gopisetty D,
Sanchez V, Goodman MB, Pruitt BL, and Krieg M (2017) Pneumatic
stimulation of C. elegans mechanoreceptor   neurons in a microfluidic trap,
Lab Chip 17(6): 1116–1127
Oren-Suissa M, Gattegno T, Kravtsov V, Podbilewicz B (2017) Extrinsic Repair of
Injured Dendrites as a Paradigm for Regeneration by Fusion in
Caenorhabditis elegans. Genetics 206: 215-230
Oren-Suissa M, Hall D, Treinin M, Shemer G, Podbilewicz B (2010) The Fusogen
EFF-1 Controls Sculpting of Mechanosensory Dendrites. Science 328:
1285-1288
Porta-de-la-Riva M, Gonzalez AC, Sanfeliu-Cerdán N, Karimi S, Malaiwong,
Pidde A, Curiel LFM, Fernandez P, González-Bolívar S, Hurth C, Krieg M
(2023) Neural engineering with photons as synaptic transmitters. Nature
methods 20, 761–76
Rose JK, Sangha S, Rai S, Norman KR, Rankin CH (2005) Decreased sensory
stimulation reduces behavioral responding, retards development, and alters
neuronal connectivity in Caenorhabditis elegans. J Neurosci 25: 7159-68
Roy Choudhury A, Groflhans J, Kong D (2021) Ion Channels in Epithelial
Dynamics and Morphogenesis. Cells 10: 2280

Salzberg Y, Diaz-Balzac CA, Ramirez-Suarez NJ, Attreed M, Tecle E, Desbois M, Kaprielian Z, Bülow HE (2013) Skin-Derived Cues Control Arborization of Sensory Dendrites in Caenorhabditis elegans. Cell 155: 308-20

Salzberg Y, Ramirez-Suarez NJ, Bülow HE (2014) The proprotein convertase KPC-1/furin controls branching and self-avoidance of sensory dendrites in Caenorhabditis elegans. PLoS genetics 10: e1004657

Sanfeliu-Cerdán N, Català-Castro F, Mateos B, Garcia-Cabau C, Ribera M, Ruider I, Porta-de-la-Riva Montserrat Canals-Calderón A, Wieser S, Salvatella X, Krieg M (2023). A MEC-2/stomatin condensate liquid-to-solid phase transition controls neuronal mechanotransduction during touch sensing. Nature cell biology, 25(11), 1590–1599

Sawin ER, Ranganathan R, Horvitz HR (2000) C. elegans locomotory rate is modulated by the environment through a dopaminergic pathway and by experience through a serotonergic pathway. Neuron 26: 619-631

Setty H, Salzberg Y, Karimi S, Berent-Barzel E, Krieg M, Oren-Suissa M (2022) Sexually dimorphic architecture and function of a mechanosensory circuit in C. elegans. Nature communications 13:6825

Shi S, Luke CJ, Miedel MT, Silverman GA, Kleyman TR (2016) Activation of the Caenorhabditis elegans Degenerin Channel by Shear Stress Requires the MEC-10 Subunit. J Biol Chem 291: 14012-14022

Shi S, Mutchler SM, Blobner BM, Kashlan OB, Kleyman TR (2018) Pore-lining residues of MEC-4 and MEC-10 channel subunits tune the Caenorhabditis elegans degenerin channel's response to shear stress. J Biol Chem 293: 10757-10766

Sin WC, Haas K, Ruthazer ES, Cline HT (2002) Dendrite growth increased by visual activity requires NMDA receptor and Rho GTPases. Nature 419: 475-480

Smith CJ, Watson JD, Spencer WC, O'Brien T, Cha B, Albeg A, Treinin M, Miller DM, 3rd (2010) Time-lapse imaging and cell-specific expression profiling reveal dynamic branching and molecular determinants of a multi-dendritic nociceptor in C. elegans. Dev Biol 345: 18-33

Star EN, Kwiatkowski DJ, Murthy VN (2002) Rapid turnover of actin in dendritic spines and its regulation by activity. Nat Neurosci 5: 239-246

Stein IS, Park DK, Claiborne N, Zito K (2021) Non-ionotropic NMDA receptor signaling gates bidirectional structural plasticity of dendritic spines. Cell reports 34: 108664

Sulston JE, Schierenberg E, White JG, Thomson JN (1983) The Embryonic Cell Lineage of the Nematode *Caenorhabditis elegans*. Dev Biol 100: 64-119

Suzuki H, Kerr R, Bianchi L, Frokjaer-Jensen C, Slone D, Xue J, Gerstbrein B, Driscoll M, Schafer WR (2003) In vivo imaging of C. elegans mechanosensory neurons demonstrates a specific role for the MEC-4 channel in the process of gentle touch sensation. Neuron 39: 1015-1017

Tao L, Porto D, Li Z, Fechner S, Lee SA, Goodman MB, Xu XZS, Lu H, Shen K (2019) Parallel Processing of Two Mechanosensory Modalities by a Single Neuron in C. elegans. Dev Cell 51: 617-631 e3

Tavosanis G (2012) Dendritic structural plasticity. Dev Neurobiol 72: 73-86
Tendilla-Beltran H, Antonio Vazquez-Roque R, Judith Vazquez-Hernandez A,
Garces-Ramirez L, Flores G (2019) Exploring the Dendritic Spine Pathology
in a Schizophrenia-related Neurodevelopmental Animal Model.
Neuroscience 396: 36-45
Tsalik EL, Niacaris T, Wenick AS, Pau K, Avery L, Hobert O (2003) LIM
homeobox gene-dependent expression of biogenic amine receptors in
restricted regions of the C. elegans nervous system. Dev Biol 263: 81-102
Vaillant AR, Zanassi P, Walsh GS, Aumont A, Alonso A, Miller FD (2002)
Signaling mechanisms underlying reversible, activity-dependent dendrite
formation. Neuron 34: 985-998
Way JC, Chalfie M (1988) mec-3, a homeobox-containing gene that specifies
differentiation of the touch receptor neurons in C. elegans. Cell 54: 5-16
Welsh MJ, Price MP, Xie J (2002) Biochemical Basis of Touch Perception:
Mechanosensory Function of Degenerin/Epithelial Na+ Channels. J Biol
Chem 277: 2369-2372
Wiesner T, Bilodeau A, Bernatchez R, Deschenes A, Raulier B, De Koninck P,
Lavoie-Cardinal F (2020) Activity-Dependent Remodeling of Synaptic
Protein Organization Revealed by High Throughput Analysis of STED
Nanoscopy Images. Front Neural Circuits 14: 57
Wilbrecht L, Holtmaat A, Wright N, Fox K, Svoboda K (2010) Structural Plasticity
Underlies Experience-Dependent Functional Plasticity of Cortical Circuits. J
Neurosci 30: 4927-4932
Wong RO, Ghosh A (2002) Activity-dependent regulation of dendritic growth and
patterning. Nat Rev Neurosci 3: 803-812
Xu T, Yu X, Perlik AJ, Tobin WF, Zweig JA, Tennant K, Jones T, Zuo Y (2009)
Rapid formation and selective stabilization of synapses for enduring motor
memories. Nature 462: 915-919
Yang G, Pan F, Gan WB (2009) Stably maintained dendritic spines are
associated with lifelong memories. Nature 462: 920-924
Younger MA, Muller M, Tong A, Pym EC, Davis GW (2013) A Presynaptic ENaC
Channel Drives Homeostatic Plasticity. Neuron 79: 1183-1196
Yu X, Zuo Y (2011) Spine plasticity in the motor cortex. Curr Opin Neurobiol 21:
169-174
Zhang Y, Cudmore RH, Lin DT, Linden DJ, Huganir RL (2015) Visualization of
NMDA receptor–dependent AMPA receptor synaptic plasticity in vivo. Nat
Neurosci 18: 402-407
Zhou Z, Hong EJ, Cohen S, Zhao WN, Ho HY, Schmidt L, Chen WG, Lin Y,
Savner E, Griffith EC, Hu L, Steen JA, Weitz CJ, Greenberg ME (2006)
Brain-Specific Phosphorylation of MeCP2 Regulates Activity-Dependent
Bdnf Transcription, Dendritic Growth, and Spine Maturation. Neuron 52:
255-269
Zuo Y, Yang G, Kwon E, Gan WB (2005) Long-term sensory deprivation
prevents dendritic spine loss in primary somatosensory cortex. Nature 436:
261-265

Figure legends

Figure 1. Mechanosensory deprivation during development reduces the behavioral response of the PVD neuron. (A) The PVD neuron dendritic tree, cell body (CB) and axon. The red arrow corresponds to the contact point with a platinum wire during posterior harsh touch, while the black arrow represents the behavioral escape response of the worm. The red dashed square represents the analyzed region around the CB. One representative candelabrum is colored by branch orders: blue primary (1°), purple secondary (2°), cyan tertiary (3°) and green quaternaries (4°). A- anterior; P- posterior; V- ventral; D-dorsal. **(B)** Schematic of the isolation protocol followed at 72 h by posterior harsh touch assay or PVD imaging of adult worms. **(C)** Isolation of embryos reduced the percentage of worms responding to harsh touch at adulthood. Crowded- black bars, Isolated- red bars. Wildtype (WT) N2 worms (Crowded, n=32; Isolated, n=32), *mec-4* (Crowded, n=52; Isolated, n=46), *mec-3* (Crowded, n=12; Isolated, n=12). *mec-4* animals were assayed as adults after 96 h to account for their slower rate of growth. **(D)** Growth in plates with chemical cues from adult hermaphrodites did not alter the reduced response rate of isolated *mec-4* animals to harsh touch (Crowded, n=33; Isolated, n=28). **(E)** Harsh touch response in crowded and isolated conditions for mutants of different DEG-ENaCs and the TRP channel *gtl-1*. WT worms (same set of worms as in Figure 1C. Crowded, n=32; Isolated, n=32), *asic-1* (Crowded, n=46; Isolated, n=30), *mec-10* (Crowded, n=38; Isolated, n=31), *degt-1* (Crowded, n=37; Isolated, n=31), *unc-8* (Crowded, n=18; Isolated, n=15), *del-1* (Crowded, n=31; Isolated, n=27), *gtl-1* (Crowded, n=27; Isolated, n=25). **(F)** MEC-10 expression in the PVD rescues *mec-10* mutants' crowded-specific reduction in response to harsh touch. All strains contain the *ser-2Prom3::Kaede* PVD marker construct, and were tested in the crowded conditions. (WT, n=42; *mec-4,* n=17; *him-5,* n=34 (*him-5* was used as WT background for several strains after cross); *him-5; mec-10,* n=35; *him-5; mec-10*; *ser2Prom3::mec-10*, n=33). **(G)** Isolation for 24 h in adulthood did not affect the response to harsh touch. Worms grown under crowded conditions were isolated for 24 h as young adults and compared against their crowded age-matched counterparts in their response to harsh touch (Crowded, n=25; Isolated for 24 h, n=26). The proportion of responding worms (percentage) ± standard error of proportion is shown. Fisher's exact test, *$p<0.05$, **$p<0.01$, ***$p<0.001$, n.s. not significant.

Figure 2. Mechanosensory deprivation and cell autonomous activity of *mec-10* affects the architecture of the PVD. (A) Schematic representation of the PVD dendritic structure, marking the morphological characteristics of interest: ectopic branches and quaternary branching geometry. Dashed lines represent ectopic branching at each order. Colors correspond to Figure 1A. (**B-G**) Both isolation and *mec-10* affect the structure of the PVD: **(B)** Isolation increases the fraction of ectopic branching, *mec-10* mutation increases ectopic branching in the crowded state. **(C)** Isolation decreases the percentage of straight quaternary branches, *mec-10* mutation decreases the percentage of straight quaternary branches in the crowded state. Crowded- black dots, isolated- red dots; WT (Crowded, n=28; Isolated n=26), *him-5* (Crowded, n=27; Isolated, n=25; *him-5* was used as WT background for several strains after cross), *him-5; mec-10* (Crowded, n=30; Isolated, n=24). **(D)** Expression of *mec-10* in the PVD on the background of *him-5; mec-10* in the crowded state reduces ectopic branching and **(E)** increases the percentage of straight quaternary branches (Crowded *him-5*, n=28; Crowded *him-5:mec-10*, n=36; Crowded *him-5:mec-10*;PVD::MEC-10, n=27). **(F-I)** Representative PVD images of WT,

mec-10 and *mec-10*; PVD::*mec-10* in different growth conditions. (Scale bar, 50 µm). **(J)**
Embryo isolation with glass beads did not affect ectopic branching. **(K)** Embryo isolation with
glass beads increased the percentage of straight quaternary branches (Crowded, n=28; Isolated
n=26; Isolated with beads, n=22. Crowded and isolated WT are the same set of worms as in
Figure 2B,C). Each dot represents a single worm. The mean ± s.e.m. are shown in blue. Box
plot with median and hinges for the first and third quartile. The whiskers represent an estimated
calculation of the 95% confidence interval. Kruskal-Wallis test, # # $p < 0.01$, # # # $p < 0.001$, Mann
Whitney test with Bonferroni Correction α=0.0167. *$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, n.s. not
significant.
1211
Figure 3. Adult isolation for less than 4 h affects the structure of the PVD.
(A) Isolation of crowded worms induces a time-dependent increase in the proportion of ectopic
branches, and **(B)** a decrease in the number of straight quaternary branches (2 h: Crowded,
n=24, Isolated, n=22; 5 h: Crowded, n=24, Isolated, n=27; 15 h: Crowded, n=3, Isolated, n=3;
24 h: Crowded, n=20, Isolated, n=21; animals imaged using polystyrene beads for
immobilization, see Materials and Methods). **(C)** Isolation for 4 h induces more growth of new
branches than retraction of existing ones. Ectopic branch dynamics were compared for crowded
animals against their state after 4 h isolation. Imaging used 1% tricaine, see Materials and
Methods. **(D)** Isolation of crowded worms for 4 h increases the ratio between ectopic/non-
ectopic branches in individual animals and **(E)** reduces the percentage of straight quaternary
branches (n=6). **(F)** Crowded, but not isolated worms, show dynamic reduction in the
percentage of straight quaternary branches during a 3 h time lapse movie in 1% tricaine; see
Videos 1 and 2 (Crowded, n=5, Isolated, n=6). **(G)** Representative time lapse frames of a
crowded worm in 1% tricaine. Thick and thin arrows represent straight and non-straight
quaternary branches, respectively. For **(D, E)** the lines connecting two points refer to the same
crowded specimen at time 0 and after 4 h isolation. Box plot representation as in Figure 2. For
(A-B) Mann-Whitney test, * $p < 0.05$, *** $p < 0.001$. For **(C-E)** Wilcoxon test (for two related
samples), * $p < 0.05$. The mean ± s.e.m. are shown for panels A-C, and F.
1230
Figure 4. Response to harsh touch and PVD morphology are independent.
(A) Discriminant analysis shows independence of harsh touch response from the PVD's
morphological classification. Squares indicate the centroids for morphological characteristics
analyzed in Figure 4–figure supplement 2A-C. The response to harsh touch (Figure 4–figure
supplement 1) is illustrated by its magnitude (low, <45% in yellow star; high, >65% in light blue
circles). The different genotypes are numbered in the list on the right. Crowded- black, Isolated-
red. **(B-E)** Isolated animals show similar PVD morphology regardless of their touch response:
(B) isolated responding and non-responding animals are not different with regard to the fraction
of ectopic branching, and **(C)** the geometry of quaternary branches (Isolated non-responding,
n=18; Isolated responding, n=20). Box plot representation as in Figure 2. Mann-Whitney test.
n.s. not significant. **(D, E)** Representative PVD images of responding and non-responding
worms. (Scale bar, 50 µm).
Figure 5. Mechanosensory-dependent localization of degenerins and
optogenetics. (A) PVD::MEC-10::mCherry localization (shown and labeled in purple,
also in Figure 7) is reduced in the quaternary branches and the axon following isolation,
in a *degt-1*-dependent manner (WT: Crowded, n=31; Isolated, n=27; *degt-1* mutants:

Crowded, n=23; Isolated, n=22). **(B-D)** Representative images and reconstructions for
PVD::*mec-10*::mCherry localization for crowded and isolated WT worms, and crowded
degt-1 worms. (Scale bar, 50 µm). **(E)** PVD::*degt-1::*mCherry localization level is
reduced at the cell body, but not in the quaternary branches or the axon, following
isolation, in a *mec*-10-dependent-manner (WT: Crowded, n=33; Isolated, n=21; *mec-10*
mutants: Crowded, n=28; Isolated, n=22). The percentage of expressing worms ±
standard error of proportion is shown. **(F-H)** Representative images and reconstructions
for PVD::*degt-1*::mCherry localization for crowded and isolated WT worms and crowded
mec-10 worms. **(I)** Isolation leads to a reduced escape response following optogenetic
photoactivation of Channelrhodopsin 2 in the PVD (Crowded, n=30; Isolated, n=24,
grown on All Trans Retinal. No response was observed for worms grown without All
Trans Retinal). Fisher exact test, *$p<0.05$, **$p<0.01$, ***$p<0.01$ n.s. not significant.
1259
Figure 6. The calcium response to mechanical stimulation shows no significant
difference between crowded and isolated worms.
Mechanically induced calcium recordings of PVD of animals grown as **(A)** isolated
individuals or **(B)** in crowded conditions. **i)** Stimulus profile displaying a single step,
sawtooth indentation or 10 Hz `buzz'. Bottom panels show examples of negative images
from experiments at pre- and post-stimulation; note PVDs' cell bodies about 200 µm to
the right of the location of the mechanical stimuli. Scale bar, 100 µm. **(ii)** Kymograph of
individual response for N=17 animals and N=27 animals for the crowded conditions.
Green bars, time of application of a mechanical stimulus. **(iii)** Average GCaMP6s signal
derived from the cell body of PVD for the duration of the recordings, black line. N =
number of recordings. Mean +/- standard deviation, grey region. Inset in panel iii shows
the running p-value, comparing the crowded and isolated conditions (Dashed line
indicates alpha level of significance a=0.05).
1273
Figure 7. Model of experience-induced structural and behavioral plasticity. During
adulthood, mechanosensory signals maintain the structure of the PVD in crowded animals, with
straight quaternary branches and fewer ectopic branches. Sensory deprivation (in isolated
animals) results in ectopic dendrites, wavy branches and defects in candelabra self-avoidance.
During development, mechanosensory experience alters the response to harsh touch and the
crawling gait of the worm (Inberg et al., 2021), possibly through changes in MEC-10 localization
in the axon and mediated by other DEG/ENaCs. Mechanosensory stimuli are a driving force for
changes in the compartment specific localization of MEC-10 and DEGT-1 in the PVD, which
may affect the structure of the PVD. MEC-10 is represented in blue, DEGT-1 is represented in
green.
1284
1285

Supplementary figure legends

Figure 1–figure supplement 1. The reduction in response to harsh touch following isolation is PVD-dependent and chemosensory-independent.
(A) Isolation of gentle-touch-insensitive *mec-4* worms with glass beads partially rescues the isolation-induced reduction in harsh touch response. Experiment performed after 96 h to account for slower growth rate. Crowded, n=39; Isolated, n=42; Isolated with beads, n=31). **(B)** Isolation of *osm-6* mutants, defective in ciliated chemosensory neurons and the PDE nociceptor, resulted in reduced response to harsh touch, when compared to worms from a crowded plate. (Crowded, n=34; Isolated, n=27). **(C)** Continuous exposure to amiloride does not affect the WT response to harsh touch in the crowded conditions (Crowded with DMSO, n=54; Crowded with 3 mM amiloride, n=52). The proportion of responding worms (percentage) ± the standard error of proportion is shown. Fisher exact test, * $p<0.05$, ** $p<0.01$, n.s. not significant.

Figure 2–figure supplement 1. **The effect of isolation on the structure of the PVD is independent from the identity of the promoter driving expression of the reporter in the PVD.**
Worms expressing the PVD fluorescent marker, *dzIs53* [*pF49H12.4*::mCherry] were isolated as in Figure 1B. **(A)** Embryo isolation for 72 h increased total ectopic branching, and **(B)** reduced the percentage of straight quaternary branches (Crowded, n=20; Isolated, n=20). **(C, D)** Representative PVD images of *dzIs53* [*pF49H12.4*::mCherry] crowded and isolated adults. (Scale bar, 50 μm). Box plot description as in Figure 2. Mann Whitney test, *** $p<0.001$. One representative candelabrum is colored as in Figure 1A.

Figure 2–figure supplement 2. The PVD undergoes isolation-dependent structural plasticity in the presence of pheromonal signals in the plate.
Both crowded and isolated worms were pre-exposed to pheromonal signals, released by *glp-4* worms (~40 worms for 96 h. The *glp-4* worms were taken out of the plate and the isolation protocol described in Figure 1B was used). **(A)** Chemical stimulation of plates with *glp-4*, followed by egg isolation, increased the fraction of ectopic branching, and **(B)** decreased straight quaternary branches (Crowded, n=24; Isolated, n=20). **(C, D)** Representative PVD images for crowded and isolated worms with chemical stimulation of the plates. White arrow, ectopic branching. (Scale bar, 50 μm). Box plot representation as in Fig 2. Mann Whitney test, *** $p<0.001$. One representative candelabrum is colored as in Figure 1A.

Figure 2–figure supplement 3. The effect of isolation on the structure of PVD is not mediated by the gentle touch mechano-sensory neurons.
(A) Description of the protocol for the isolation of *mec-4* eggs for 96 h (the experiment was performed for 96 h since the worms were L4s or very young adults at 72 h), followed by PVD imaging. **(B)** Isolation of the gentle touch impaired strain, *ser-2Prom3*::Kaede; *mec-4*, increased total ectopic branching, and **(C)** reduced the percentage of straight quaternary branching (Crowded, n=20; Isolated, n=21). **(D, E)** Representative PVD images derived from crowded worms and isolated eggs. White arrow, ectopic branching (Scale bar, 50 μm). Box plot representation as in Figure 2. Mann Whitney test with Bonferroni correction ** $p<0.01$. One representative candelabrum is colored as in Figure 1A.

1335
Figure 2–figure supplement 4. Isolation of eggs for 48 h and adults for 24 h is
sufficient to induce changes in the structure of the PVD.
(A) Schematic protocol for the isolation of WT eggs for 48 h and adults for 24 h, followed
by PVD imaging. **(B)** Isolation of eggs for 48 h (resulted in young adult worms) did not
affect the number of ectopic branches, and **(C)** decreased the percentage of straight
quaternary branching (Crowded, n=30; Isolated, n=24). **(D, E)** Representative PVD
images for crowded worms and isolated eggs. (Scale bar, 50 µm). **(F)** Isolation of young
adult worms for 24 h increased the fraction of ectopic branching, and **(G)** reduced the
percentage of straight quaternary branches (Crowded, n=20; Isolated adults for 24 h,
n=21). **(H, I)** Representative pictures from the PVD images for crowded and matched
animal isolated as young adult for 24 h. A small arrow marks a self-avoidance defect.
Box plot representation as in Figure 2. Mann Whitney test, ** $p<0.01$, *** $p<0.001$, n.s.
not significant. One representative candelabrum is colored as in Figure 1A.

Figure 3–figure supplement 1. Four hours are sufficient to induce an increase in
ectopic branching and a decrease in percentage of straight branches.
Left image is from a young adult grown in crowded conditions. Following confocal
imaging this worm was isolated for 4 h and imaged again (right image). Straight terminal
branches (black arrows, left image) became non-straight and there are new ectopic
branches (white arrows, right image; see quantitative analyses in Figure 3D,E).

Figure 3–figure supplement 2. Activity-modulating pharmacological agents affect
the structure and the dynamics of PVD branch growth and retraction.
(A-D) Blocking DEG/ENaC with 3 mM amiloride in crowded plate conditions affects the
structure of the PVD. **(A)** Increased the fraction of ectopic branches, and **(B)** decreased
the percentage of straight quaternary branches, (Crowded with DMSO, n=23; Crowded
with amiloride n=27). **(C, D)** Representative PVD images in DMSO- and amiloride-
treated worms. One representative candelabrum is colored as in Figure 1A (Scale bar,
25 µm). **(E-G)** A mixture containing 0.01% tetramisole and 0.1% tricaine induced more
dendritic growth than retraction in crowded conditions **(E;** Crowded: n=5; Isolated, n=5).
Representative images from time lapse movies demonstrating dynamic outgrowth of
branching under tricaine-tetramisole anesthetic mixture for crowded **(F)** and isolated
worms **(G). (H-J)** 1% tricaine induced more dendritic retraction than growth. In addition,
more dendritic growth was measured for isolated worms, compared to crowded (**H**;
Crowded: n=6; Isolated, n=6). Representative time lapse movies demonstrating dynamic
retraction of branching for crowded **(I)** and isolated **(J)** worms**;** see Videos 1 and 2. Blue
arrowheads, branch growth; red arrowheads, branch retraction; bar for panel 'I', 25 µm.
Box plots are as represented in Figure 2. Mann Whitney test * $p<0.05$, ** $p<0.01$. For
(E,H) t-Test. * $p<0.05$, *** $p<0.001$, n.s. not significant.

Figure 4–figure supplement 1. The DEG/ENaC *asic-1* and *mec-10* mediate
experience-dependent behavioral plasticity following isolation.
Posterior harsh touch assay was performed for crowded and isolated WT and DEG-
ENaC mutants, similar to the procedure described in Figure 1B. Seven combinations of
DEG-ENaC mutants were tested for the isolation effect on response to harsh touch.
Isolated worms were compared to crowded worms from the same strain. N2 worms (The
same set of worms as in Figure 1C. Crowded, n=32; Isolated, n=32), *asic-1* (Crowded,
n=46; Isolated, n=30), *mec-10* (Crowded, n=38; Isolated, n=31), *degt-1* (Crowded, n=37;
Isolated, n=31), *asic-1; mec-10* (Crowded, n=20; Isolated, n=22), *mec-10; degt-1*

1385 (Crowded, n=40; Isolated, n=36), *asic-1*; *degt-1* (Crowded, n=35; Isolated, n=34), *asic-1*;
1386 *mec-10; degt-1* (Crowded, n=52; Isolated, n=50). The proportion (percentage) of
1387 responding worms ± the standard error of proportion is shown. Fisher exact test,
1388 **p*<0.05, *** *p*<0.001, n.s. not significant.
1389
Figure 4–figure supplement 2. The DEG-ENaCs, *mec-10* and *degt-1*, mediate
mechanosensory dependent structural changes in the PVD.
Mutants of DEG-ENaC were isolated and were compared to crowded animals from the
same strain, similar to the procedure described in Figure 1B. **(A)** Three DEG-ENaC
mutants: *degt-1, mec-10; degt-1*, and *asic-1; mec-10; degt-1* show no difference
between crowded and isolated worms in terms of ectopic branching. In addition,
crowded *mec-10* and *degt-1* animals increased the number of ectopic branches
compared to crowded WT worms. **(B)** *mec-10; degt-1* mutants retain a high percentage
of straight quaternary branches following isolation, while *mec-10* and *degt-1* single
mutants have fewer straight branches compared with crowded WT animals, which
decrease further upon isolation. **(C)** The increase in self-avoidance defects following
isolation is DEG-ENaC dependent. WT (The same set of worms as shown in Figure 2A,
B. Crowded, n=28; Isolated, n=26), *asic-1* (Crowded, n=22; Isolated, n=25), *mec-10*
(Crowded, n=27; Isolated, n=21. The same set of worms as shown in Figure A, B), *degt-*
1 (Crowded, n=28; Isolated, n=24), *asic-1; mec-10* (Crowded, n=30; Isolated, n=31),
mec-10; degt-1 (Crowded, n=26; Isolated, n=28), *asic-1; degt-1* (Crowded, n=23;
Isolated, n=22), *asic-1; mec-10; degt-1* (Crowded, n=23; Isolated, n=21). **(D)**
Representative PVD images for crowded and isolated worms of WT and seven different
DEG-ENaC mutants. (Scale bar, 50 µm). The mean ± s.e.m. are shown. Mann Whitney
test, **p*<0.05, ** *p*<0.01, *** *p*<0.001, n.s. not significant. One representative
candelabrum is colored as in Figure 1A.
1411
Figure 4–figure supplement 3. Isolation induces increase in loss of self-avoidance
defects, in a *mec-10* dependent manner. (A) Schematic representation of the PVD
dendritic structure, marking a self-avoidance defect, in which two candelabra overlap.
Colors correspond to Figure 2A. **(B)** Isolation increases the percentage of self-avoidance
defects between two adjacent candelabra. WT (Crowded, n=28; Isolated n=26), *him-5*
(Crowded, n=27; Isolated, n=25; *him-5* was used as WT background for several strains
after cross), *him-5; mec-10* (Crowded, n=30; Isolated, n=24). **(C)** Overexpression of
MEC-10 in the PVD does not affect the percentage of self-avoidance defects seen in
mec-10 crowded animals. Crowded *him-5* (n=28), crowded *him-5; mec-10* (n=36),
crowded *him-5; mec-10;* PVD::MEC-10 (n=27). **(D)** Isolation with glass beads reduced
the percentage of self-avoidance defects, compared to isolated worms without beads
(Crowded, n=28; Isolated n=26; Isolated with beads, n=22. Crowded and isolated are
the same worms as used in Figure 2B-C). **(E)** Isolation during adulthood increases the
percentage of self-avoidance defects in a time-dependent manner (2 h: Crowded, n=24,
Isolated adults for 2 h, n=22; 5 h: Crowded, n=24, Isolated adults for 5 h, n=27; 15 h:
Crowded, n=3, Isolated adults for 15 h, n=3; 24 h: Crowded, n=20, Isolated adults for 24
h, n=21). **(F)** The percentage of self-avoidance defects is not correlated with the
response to harsh touch (Isolated non-responding, n=18; Isolated responding, n=20).
(G) Worms from a different strain and fluorescent marker, *dzls53* [*pF49H12.4*::mCherry],
were isolated as eggs for 72 h and compared to crowded worms from the same strain**.**

Isolated worms increased the percentage of self-avoidance defects between two
adjacent candelabra (Crowded, n=20; Isolated, n=20). **(H)** Blocking DEG/ENaC does not
affect the percentage of self-avoidance defects (Crowded DMSO, n=23; Crowded 3 mM
amiloride, n=27). **(I)** Both crowded and isolated worms were pre-exposed to pheromonal
signals, released by *glp-4* worms (~40 worms for 96 h). The *glp-4* worms were taken out
of the plate and the isolation protocol described in Figure 1B was used. Isolation
increased the percentage of self-avoidance defects (Crowded, n=24; Isolated, n=20). **(J)**
Isolation of the gentle touch impaired strain, *ser-2Prom3*::Kaede; *mec-4* increased the
percentage of self-avoidance defects after isolation of eggs. (Crowded, n=20; Isolated,
n=21). **(K)** Isolation of eggs for 48 h did not affect the percentage of self-avoidance
defects (Crowded, n=30; Isolated, n=24). **(L)** Isolation of young adult worms for 24 h
increased the percentage of self-avoidance defects (Crowded, n=20; Isolated adults for
24 h, n=21). Each dot represents a single worm. Box plots are as represented in Figure
2. Kruskal-Wallis test, # $p<0.05$, # # # $p<0.001$, Mann Whitney test with Bonferroni
Correction α=0.0167. *$p<0.05$, ** $p<0.01$, *** $p<0.001$, n.s. not significant.

Figure 5–figure supplement 1. Optogenetic stimulation of isolated worms does not
affect the morphology of the PVD. Worms were grown isolated with (n=8) and without
(n=10) all trans retinal and subjected to 4 h light stimulation at 488 nm. Light stimulation
with all trans retinal does not affect the amount of ectopic branching **(A)**, the geometry of
the quaternary branches **(B)** or the the proportion of self-avoidance defects **(C)**. n.s. not
significant. The mean ± s.e.m. are shown for panels A-C.

Figure 6–figure supplement 1. Basal Calcium dynamics are indistinguishable for
crowded and isolated worms. The 90-second calcium dynamics of the PVD were
measured in crowded (n=16) and isolated (n=17) worms. **(A)** Heatmaps representing the
calcium activity for each worm. Note that the third isolated worm had a spontaneous
Ca^{2+} activation. **(B)** The population averages for calcium dynamics. **(C)** The distribution
of neuronal activity change across the populations of crowded and isolated worms.

Figure 6–figure supplement 2. Calcium response in the PVD is sensitive to both
onset and offset of the mechanical stimulus signals. Mechanically induced calcium
recordings of PVD of animals in a micromechanical device. **(A)** The GCaMP6s signal
derived from the PVD before, during and after application of a mechanosensory force
within a micromechanical device. The cell body of the PVD is surrounded by a dashed
circle. Scale bars, 100 μm. Note, to pressure channels fingers were actuated
simultaneously in this particular experiment. **(B)** Quantification of GCaMP6s signal with
the corresponding time frames indicated by arrows. Green region shows time of single
step stimulus. See Video 7.

1468

Video legends

Video 1. **Time lapse of cell body and arborizations of a wildtype young adult grown under crowded conditions and anesthetized with 1% tricaine.** Red arrows point to retracting branches. See Figure 3 - figure supplement 2I for more details.

Video 2. **Time lapse of cell body and arborizations of a wildtype young adult grown in isolated conditions and anesthetized with 1% tricaine.** Red arrows point to retracting branches. See Figure 3 - figure supplement 2J for more details.

Video 3. **Localization pattern of MEC-10::mCherry in the PVD crowded adult.** MEC-10 is localized in moving vesicles, indicated by red arrow. Six z-stack series (~60 optical slices for each) were taken around the cell body every 3 minutes.

Video 4. **Localization pattern of MEC-10::mCherry in the PVD for crowded adult.** MEC-10 is localized in moving vesicles, indicated by red arrow. At time 00:00 quaternary branches can be seen. Six z-stack series (~60 optical slices for each) were taken around the cell body every 2 minutes.

Video 5. Calcium dynamics in the PVD of an isolated worm after mechanosensory stimulation. PVD cell body received ipsilateral application of a mechanical stimulus, the calcium responses from the PVD were measured. See Figure 6 for more details. Scale bar, 40 µm.

Video 6. Calcium dynamics in the PVD of a representative crowded worm after mechanosensory stimulation. PVD cell body received ipsilateral application of a mechanical stimulus, the calcium responses from the PVD were measured. See Figure 6 for more details. Scale bar, 40 µm.

Video 7. Calcium dynamics in the PVD is sensitive to both onset and offset signals. PVD cell body received ipsilateral application of a mechanical stimulus and responded by increase in calcium signal for both the onset and offset stimulations. See Figure S14 for more details. Scale bar, 40 µm.

Supplementary file legend

Supplementary file 1a. List of strains and transgenes used in this work

Supplementary file 1b. List of primers used in this work

Supplementary References

Titles for the Source Data

Figure 1-source data 1.
Original data file for Figure 1 graphs on mechanosensory deprivation during development reduces the behavioral response of the PVD neuron.

Figure 1-figure supplement 1-source data 1.
Original data file for Figure 1-figure supplement 1 on the reduction in response to harsh touch following isolation is PVD-dependent and chemosensory-independent.

Figure 2-source data 1.
Original data file for Figure 2 graphs on mechanosensory deprivation and cell autonomous activity of mec-10 affects the architecture of the PVD.

Figure 2-figure supplement 1-source data 1.
Data for Figure 2-figure supplement 1 on the effect of isolation on the structure of the PVD is independent from the promoter driving expression of the reporter in the PVD.

Figure 2-figure supplement 2-source data 1.
Original data for Figure 2-figure supplement 2 on the PVD undergoes isolation-dependent structural plasticity in the presence of pheromonal signals in the plate.

Figure 2-figure supplement 3-source data 1.
Original data for Figure 2-figure supplement 3 graphs on the effect of isolation on the structure of PVD is not mediated by the gentle touch mechano-sensory neurons.

Figure 2-figure supplement 4-source data 1.
Original data for Figure 2-figure supplement 4 graphs on isolation of eggs for 48 h and adults for 24 h is sufficient to induce changes in the structure of the PVD.

Figure 3-source data 1.
Original data file for Figure 3 graphs on adult isolation for less than 4 h affects the structure of the PVD.

Figure 3-figure supplement 2-source data 1.
Original data for Figure 3 supplement 2 graphs on activity-modulating pharmacological agents affect the structure and the dynamics of PVD branch growth and retraction.

Figure 4-source data 1.
Original data file for Figure 4 graphs on response to harsh touch and PVD morphology are independent.

Figure 4-figure supplement 1-source data 1.
Original data for Figure 4-figure supplement 1 on the DEG/ENaC *asic-1* and *mec-10* mediate experience-dependent behavioral plasticity following isolation.

Figure 4-figure supplement 2-source data 1.

1558 **Original data for Figure 4-figure supplement 2 on the DEG-ENaCs, *mec-10* and**
1559 ***degt-1*, mediate mechanosensory dependent structural changes in the PVD.**
1560
1561 **Figure 4-figure supplement 3-source data 1.**
1562 **Original data for Figure 4-figure supplement 3 on isolation induces increase in loss of**
1563 **self-avoidance defects, in a *mec-10* dependent manner.**
1564
1565 **Figure 5-source data 1.**
1566 **Original data file for Figure 5 graphs on mechanosensory-dependent localization of**
1567 **degenerins and optogenetics.**
1568
1569 **Figure 5-figure supplement 1-source data 1.**
1570 **Original data file for Figure 5-figure supplement 1 on optogenetic stimulation of**
1571 **isolated worms does not affect the morphology of the PVD.**
1572
1573 **Figure 6-source data 1.**
1574 **Original data file for Figure 6 graphs on calcium response to mechanical stimulation.**
1575
1576 **Figure 6-figure supplement 1-source data 1.**
1577 **Original data file for Figure 6-figure supplement 1 on basal Calcium dynamics are**
1578 **indistinguishable for crowded and isolated worms.**
1579
1580 **Figure 6-figure supplement 2-source data 1.**
1581 **Original data file for Figure 6-figure supplement 2 on Calcium response in the PVD is**
1582 **sensitive to both onset and offset of the mechanical stimulus signals.**
1583
1584 **Figure 7-figure supplement 1-source data 1.**
1585 **Original data file for Figure 7-figure supplement 1 on side of plate contact alone over**
1586 **several hours is insufficient to elicit a difference in ectopic branching.**
1587

A Morphological clustering

Isolated (red) Crowded (black)

1. **WT** 2. **WT**
3. *asic-1* 4. *asic-1*
5. *mec-10* 6. *mec-10*
7. *degt-1* 8. *degt-1*
9. *asic-1;mec-10* 10. *asic-1;mec-10*
11. *asic-1;degt-1* 12. *asic-1;degt-1*
13. *mec-10;degt-1* 14. *mec-10;degt-1*
15. asic-1; *mec-10;degt-1* 16. asic-1;*mec-10;degt-1*

Response to harsh touch
<45% response   >65% response

B n.s.

C n.s.

D Isolated non responding — Self avoidance defects — Ectopic branching — Non straight

E Isolated responding — Non straight — Self avoidance defects

A

ectopic/non-ectopic branches

***

● Crowded
● Isolated

1.25
1.00
0.75
0.50
0.25
0

Crowded   Isolated

B

% straight 4ry branches

***

Crowded   Isolated

C

Crowded
pF49H12.4::mCherry

Straight

D

Isolated pF49H12.4::mCherry

Ectopic
branching

A

ectopic/non-ectopic branches

- ● Crowded
- ● Isolated

***

Crowded   Isolated

B

% straight 4ry branches

***

Crowded   Isolated

C

Crowded with chemical stimulaton

Straight

D

Isolated with chemical stimulation

Self avoidance defects

A *mec-4*

0 h crowded /isolated eggs | 72 h young adults | 96 h adults | PVD imaging

B ** Crowded ● Isolated ●

ectopic/non-ectopic branches

Crowded   Isolated

C **

% straight 4ry branches

Crowded   Isolated

D Straight   Crowded

mec-4

E Isolated

Self avoidance defects

mec-4

A 0 h crowded /isolated eggs — 48 h young adults — 72 h adults — PVD imaging

B ectopic/non-ectopic branches — n.s. — Crowded — Isolated — ● Crowded ● Isolated

C % straight 4ry branches — ** — Crowded — Isolated

D Straight

E Non straight — Isolated 48 h from egg

F ectopic/non-ectopic branches — *** — Crowded — Isolated 24 hr

G % straight 4ry branches — *** — Crowded — Isolated 24 hr

H Straight — Crowded

I Non straight — Ectopic — Isolated adults for 24 h

A

ectopic/non-ectopic branches

** (bracket over WT to degt-1)

- WT: Crowded *** vs Isolated
- asic-1: ***
- mec-10: **
- degt-1: n.s.
- asic-1; mec-10: ***
- mec-10; degt-1: n.s.
- asic-1; degt-1: *
- asic-1; mec-10; degt-1: n.s.

Legend: ■ Crowded ■ Isolated

B

% straight 4ry branches

** **

- WT: ***
- asic-1: p=0.077
- mec-10: ***
- degt-1: **
- asic-1; mec-10: ***
- mec-10; degt-1: n.s.
- asic-1; degt-1: *
- asic-1; mec-10; degt-1: **

C

% self avoidance defects

- WT: *
- asic-1: n.s.
- mec-10: n.s.
- degt-1: n.s.
- asic-1; mec-10: n.s.
- mec-10; degt-1: n.s.
- asic-1; degt-1: n.s.
- asic-1; mec-10; degt-1: n.s.

D

	Crowded	Isolated		Crowded	Isolated
WT	Straight	Ectopic branching	asic-1; mec-10		
asic-1			mec-10; degt-1		
mec-10	Non straight		asic-1; degt-1		
degt-1			asic-1; mec-10; degt-1		

A

B

C

A

Isolated growth 72 h

Monitor side of plate contact 2-3 h

Mount and image both PVDs

B

p = 0.03

Ectopic 2°-4°/100µm

● Agar contact
● Opposing side

Isolated
(n=36)

Side of plate contact

Opposing side