Sox2 and FGF20 interact to regulate organ of Corti hair cell and supporting cell

development in a spatially-graded manner

Short title: Sox2 and FGF20 in cochlea development

Lu M. Yang[1], Kathryn S.E. Cheah[2], Sung-Ho Huh[1,3,*], David M. Ornitz[1,*]

[1]Department of Developmental Biology; Washington University School of Medicine; St. Louis,

Missouri, 63110; USA

[2]School of Biomedical Sciences; The University of Hong Kong; Pokfulam, Hong Kong, China

[3]Holland Regenerative Medicine Program, and the Department of Neurological Sciences;

University of Nebraska Medical Center; Omaha, Nebraska, 68198; USA

*Corresponding authors

Email: dornitz@wustl.edu (D.M.O.), sungho.huh@unmc.edu (S.H.)

Author contributions

18 Conceptualization, L.M.Y., S.H., and D.M.O.; Methodology, L.M.Y., S.H., and D.M.O.; Formal

19 Analysis, L.M.Y. and S.H.; Investigation: L.M.Y. and S.H.; Resources: K.S.E.C., S.H., and

20 D.M.O.; Writing – Original Draft: L.M.Y. and D.M.O.; Writing – Review & Editing: L.M.Y.,

21 K.S.E.C., S.H., and D.M.O.; Supervision: D.M.O.; Funding Acquisition: S.H. and D.M.O.

Abstract

The mouse organ of Corti develops in two steps: progenitor specification and differentiation. Fibroblast Growth Factor (FGF) signaling is important in this developmental pathway, as deletion of FGF receptor 1 (*Fgfr1*) or its ligand, *Fgf20*, leads to the loss of hair cells and supporting cells from the organ of Corti. However, whether FGF20-FGFR1 signaling is required during specification or differentiation, and how it interacts with the transcription factor Sox2, also important for hair cell and supporting cell development, has been a topic of debate. Here, we show that while FGF20-FGFR1 signaling functions during progenitor differentiation, FGFR1 has an FGF20-independent, Sox2-dependent role in specification. We also show that a combination of reduction in *Sox2* expression and *Fgf20* deletion recapitulates the *Fgfr1*-deletion phenotype. Furthermore, we uncovered a strong genetic interaction between *Sox2* and *Fgf20*, especially in regulating the development of hair cells and supporting cells towards the basal end and the outer compartment of the organ of Corti. To explain this genetic interaction and its effects on the basal end of the organ of Corti, we provide evidence that decreased *Sox2* expression delays specification, which begins at the organ of Corti apex, while *Fgf20*-deletion results in premature onset of differentiation, which begins near the organ of Corti base. Thereby, *Sox2* and *Fgf20* interact to ensure that specification occurs before differentiation towards the cochlear base. These findings reveal an intricate developmental program regulating organ of Corti development along the basal-apical axis of the cochlea.

Author summary

The mammalian cochlea contains the organ of Corti, a specialized sensory epithelium populated by hair cells and supporting cells that detect sound. Hair cells are susceptible to injury by noise,

toxins, and other insults. In mammals, hair cells cannot be regenerated after injury, resulting in

permanent hearing loss. Understanding genetic pathways that regulate hair cell development in

the mammalian organ of Corti will help in developing methods to regenerate hair cells to treat

hearing loss. Many genes are essential for hair cell and supporting cell development in the

mouse organ of Corti. Among these are *Sox2*, *Fgfr1*, and *Fgf20*. Here, we investigate the

relationship between these three genes to further define their roles in development.

Interestingly, we found that *Sox2* and *Fgf20* interact to affect hair cell and supporting cell

development in a spatially-graded manner. We found that cells toward the outer compartment

and the base of the organ of Corti are more strongly affected by the loss of *Sox2* and *Fgf20*. We

provide evidence that this spatially-graded effect can be partially explained by the roles of the

two genes in the precise timing of two sequential stages of organ of Corti development,

specification and differentation.

59
60
Introduction

The inner ear contains six sensory organs required for the senses of hearing and balance. The

cochlea, a snail-like coiled duct, is the auditory organ. It contains specialized sensory epithelia,

called the organ of Corti, composed of hair cells (HCs) and supporting cells (SCs). In mammals,

this sensory epithelium is elegantly patterned, with one row of inner hair cells (IHCs) and three

rows of outer hair cells (OHCs), separated by two rows of pillar cells forming the tunnel of Corti.

Each row of OHCs is associated with a row of supporting cells called Deiters' cells. Here, we

refer to pillar cells and Deiters' cells collectively as SCs.

Organ of Corti development has been described as occurring in two main steps: prosensory

specification and differentiation [1]. During prosensory specification, proliferative progenitors at

the floor of the developing cochlear duct are specified and exit the cell cycle to form the

postmitotic prosensory domain. Here, we define specification to be a process that makes

progenitors competent to differentiate. We also use cell cycle exit as a marker for specified cells

in the prosensory domain (prosensory cells). During differentiation, prosensory cells differentiate

into both HCs and SCs [2]. Interestingly, cell cycle exit, marking the completion of specification,

and initiation of differentiation occur in waves that travel in opposite directions along the length

of the cochlear duct. At around embryonic day 12.5 (E12.5) in the mouse, progenitors begin to

exit the cell cycle and express the cyclin-dependent kinase inhibitor CDKN1B (p27^{Kip1}) in a wave

that begins at the apex of the cochlea (the cochlear tip) and reaches the base of the cochlea by

around E14.5 [3,4]. Afterwards, the specified prosensory cells begin differentiating into HCs and

SCs in a wave that begins at the mid-base at around E13.5, and spreads quickly to the rest of

the base and to the apex over the next few days [1]. Notably, while the basal end of the

cochlear duct differentiates immediately after prosensory specification, the apical end has a

longer time between specification and differentiation, providing a larger "temporal buffer" for

apical development. The spiral ganglion, containing neurons synapsing to HCs, has been

shown to be important for this delay in apical differentiation, via Sonic Hedgehog (SHH)

signaling [5–8].

The transcription factor *Sox2* is one of the earliest markers of prosensory cells [9,10]. Mice with

specific *Sox2* hypomorphic mutations that affect inner ear expression have hearing impairment

due to decreased HC and SC number, while mice with inner ear-specific *Sox2* null mutations

are completely deaf and have no HCs or SCs [11,12]. Genetic experiments show that *Sox2* is

both necessary and sufficient for prosensory specification. Absence of *Sox2* expression leads to

the loss of *Cdkn1b* expression at E14, a marker for the prosensory domain [12], while ectopic

Sox2 expression in cochlear nonsensory epithelium can induce ectopic sensory patches [13–

15].

The Fibroblast Growth Factor (FGF) signaling pathway also plays vital roles in organ of Corti

development [16]. Studies utilizing cochlear explants showed that inhibition of FGF signaling

prior to and during stages of HC and SC differentiation results in decreased HC and SC number

[17]. Signaling through FGF receptor 1 (FGFR1), in particular, is essential during this process.

Conditional deletion of *Fgfr1* (Fgfr1-CKO) in the developing cochlear epithelium resulted in

dramatically reduced HC and SC number [18–20]. This has been attributed to decreased *Sox2*

expression in the prosensory domain of Fgfr1-CKO mice, leading to a defect in prosensory

specification [19].

FGF20 has been hypothesized to be the FGFR1 ligand during organ of Corti development. Both

in vitro inhibition of FGF20 with an anti-FGF20 antibody [17] and *in vivo* knockout of *Fgf20*

(Fgf20-KO) [21] led to decreased HC and SC number, similar to the Fgfr1-CKO phenotype.

However, the Fgf20-KO phenotype is clearly not as severe as that of Fgfr1-CKO. Almost all

OHCs and some IHCs are missing in Fgfr1-CKO mice [19], while only 2/3 of OHCs are missing

in Fgf20-KO mice, without any loss of IHCs [21]. This suggests that another FGF ligand may be

redundant with and compensating for the loss of FGF20, the identity of which is currently

unknown.

Another difference between Fgfr1-CKO and Fgf20-KO mice is the proposed mechanism

accounting for the decrease in HCs and SCs. Interestingly, unlike in Fgfr1-CKO mice, Sox2

expression in the prosensory domain is not disrupted in Fgf20-KO mice [19,21]. Rather, FGF20

seems to function during HC and SC differentiation. These differences between the Fgfr1-CKO

and Fgf20-KO phenotypes and their relationship with *Sox2* suggest that FGF20/FGFR1

signaling has a more complex and as yet unexplained role during organ of Corti development.

Here, we hypothesize that FGFR1 signaling has functions in both steps of organ of Corti

development: an earlier role in prosensory specification that involves *Sox2*, and a later role in

the initiation of differentiation. We provide evidence that FGF20 regulates differentiation but not

specification. Moreover, while *Fgfr1* functions upstream of *Sox2*, *Fgf20* is downstream of *Sox2*.

We further show that *Sox2* and *Fgf20* genetically interact during organ of Corti development.

Interestingly, downregulation of both genes leads to the loss of hair cells and supporting cells

preferentially towards the outer compartment and the basal end of the cochlear duct. To explain

the more severe basal phenotype, we provide evidence that *Sox2* regulates the timing of

prosensory specification, while *Fgf20* regulates the timing of differentiation. As these two steps

occur along a developmental pathway, we hypothesize that prosensory specification must occur

prior to differentiation. In *Sox2* hypomorphic mice, prosensory specification is delayed, while in

Fgf20-KO mice, onset of differentiation occurs prematurely. When combined, these two defects

led to differentiation attempting to initiate prior to the completion of specification towards the

basal end of the cochlear duct. These results define unique functions of and complex

interactions among FGF20, FGFR1, and Sox2 during organ of Corti development and highlight

the potential importance of the timing of specification and differentiation along different regions

of the cochlear duct.

Results

The Fgf20-KO cochlear phenotype is less severe than the Fgfr1-CKO phenotype

Previous studies showed that deletion of *Fgf20* leads to a loss of two thirds of OHCs in the

mouse organ of Corti [21], while conditional deletion of *Fgfr1* from the cochlear epithelium leads

to a loss of almost all OHCs and some IHCs [19,20]. To rule out the effect of genetic

background accounting for these differences, we generated *Fgf20* knockout (Fgf20-KO: *Fgf20$^{-/-}$*)

and *Fgfr1* conditional knockout (Fgfr1-CKO: *Foxg1$^{Cre/+}$*; *Fgfr1$^{flox/-}$*) mice along with littermate

controls (*Fgf20$^{+/-}$* for Fgf20-KO and *Fgfr1$^{flox/+}$*, *Fgfr1$^{flox/-}$*, and *Foxg1$^{Cre/+}$*;*Fgfr1$^{flox/+}$* for Fgfr1-CKO)

on a mixed C57BL/6J and 129X1/SvJ genetic background. Fgf20-KO and Fgfr1-CKO mice were

generated in separate matings; therefore, some genetic background differences could persist.

Foxg1^{Cre} targets most of the otic vesicle as early as E9.5 [22] and has been used in other

studies to conditionally delete *Fgfr1* [18–20]. In the *Fgf20^{-}* allele, exon 1 of *Fgf20* is replaced by

a sequence encoding a GFP-Cre fusion protein [18]. We also refer to this null allele as *Fgf20^{Cre}*.

We examined the cochleae at P0 (Figs 1A and 1B) and quantified the length of the cochlear

duct and the total number of IHCs, OHCs, and SCs (Figs 1C-1F), as well as the number of cells

along the basal, middle, and apical turns of the cochlear duct (S1A-S1C Figs). Refer to Fig 1G

for the positions of basal, middle, and apical turns along the cochlear duct. We identified HCs

based on Phalloidin labeling and SCs based on Prox1/Sox2 labeling. IHCs and OHCs were

distinguished based on location relative to p75NTR-labeled inner pillar cells (IHCs are neural, or

towards the center of the coiled duct; OHCs are abneural).

In both Fgf20-KO and Fgfr1-CKO cochleae, there were gaps in the sensory epithelium that

lacked HCs and SCs along the entire cochlear duct. Quantitatively, Fgf20-KO cochleae had a

6% reduction in cochlear length compared to control (*Fgf20$^{+/-}$*) cochleae, while Fgfr1-CKO

cochleae had a 28% reduction compared to control (*Fgfr1$^{Cre/+}$*;*Fgfr1$^{flox/+}$*). Fgf20-KO did not have

a significant reduction in the number of IHCs, while Fgfr1-CKO cochleae had a 40% reduction.

Fgf20-KO cochleae only had a 76% reduction in the number of OHCs, while Fgfr1-CKO

cochleae had almost a complete lack of OHCs, a 97% reduction. For SCs, Fgf20-KO cochleae

had a 59% reduction, while Fgfr1-CKO cochleae had an 84% reduction. These patterns

persisted when HC and SC numbers were normalized to cochlear length. These results were all

consistent with previous studies [19,21] and showed that the Fgfr1-CKO phenotype is more

severe than the Fgf20-KO phenotype in cochlear length and in the number of HCs and SCs. We

hypothesize that during organ of Corti development, an additional FGFR1 ligand compensates

for the loss of FGF20.

Notably, while the total number of IHCs was decreased in Fgfr1-CKO cochleae, the decrease

was only observed in the basal and middle turns of the cochlea, not in the apical turn (S1A Fig).

In addition, the number of IHCs normalized to cochlear length was slightly increased in Fgf20-

KO cochleae (Fig 1D), and this increase was only prominent in the middle and apical turns of

the cochlea, but not in the basal turn (S1A Fig). The increase in IHCs could be explained by the

shortened cochlear duct length in Fgf20-KO mice. No such basal/middle/apical turn

discrepancies existed in the number of OHCs or SCs in either genotype (S1B and S1C Figs).

Our previous studies also noted that the apical tip of Fgf20-KO cochleae has delayed

differentiation relative to control at E16.5 and P0, but catches up by P7 [21]. We confirmed this

result, finding that at P0 in control cochleae, sensory epithelium at the apical tip has begun to

differentiate, based on phalloidin and p75NTR expression, while in Fgf20-KO cochleae, there

was no sign of differentiation at the apical tip. There was a similar delay in differentiation at the

apical tip of Fgfr1-CKO cochleae relative to control (S1E Fig). Refer to S1D Fig for the location

of the apical tip.

FGFR1 but not FGF20 regulates *Sox2* expression

Next, we examined *Sox2* expression in Fgf20-KO and Fgfr1-CKO cochleae at E14.5 by RNA in

situ hybridization and immunofluorescence. In control cochleae, *Sox2* mRNA and protein were

highly expressed in the prosensory domain (Fig 2A, refer to Fig 2C). The expression of *Sox2*

was not changed in Fgf20-KO cochleae compared to control; however, it was noticeably

decreased in Fgfr1-CKO cochleae (Fig 2A), in agreement with previous findings [19–21]. This

indicates that FGFR1 has an additional role, independent of FGF20, in regulating *Sox2*, which is

required for prosensory specification [12]. Similar to Sox2, CDKN1B expression in the

prosensory domain is also regulated by FGFR1, but not by FGF20 [18,19,21]. We confirmed

these results, finding that while CDKN1B expression was not changed in Fgf20-KO cochleae at

E14.5 relative to control, it was dramatically downregulated in Fgfr1-CKO cochleae (Fig 2B).

This is consistent with the role of Sox2 in regulating CDKN1B expression [12]. We hypothesize

that a yet unidentified FGF ligand (in addition to or independent of FGF20) signaling via FGFR1

regulates Sox2 expression (and therefore CDKN1B expression) during prosensory specification,

while FGF20 signaling via FGFR1 regulates differentiation (Fig 2D).

We also wanted to confirm that FGF20 signals to epithelial FGFR1 at around the initiation of

differentiation. To do so, we examined the expression of *Etv4* (also known as *Pea3*) and *Etv5*

(also known as *Erm*), two well-established downstream effectors of FGF signaling [23], by in situ

hybridization. The expression of these two genes are downregulated with FGF signaling

inhibition in E14 cochlear explants [17]. At E14.5, there were two domains of *Etv4* and *Etv5*

expression in control cochleae: the prosensory domain and the outer sulcus (S2A Fig,

brackets). The outer sulcus is the region of the cochlear epithelium abneural to the prosensory

domain at E14.5. In Fgf20-KO cochleae, expression of both genes was not detected in the

prosensory domain. In Fgfr1-CKO cochleae, expression of both genes was similarly not

detected in the prosensory domain. Expression of *Etv4* and *Etv5* in the outer sulcus was not

affected in Fgf20-KO and Fgfr1-CKO cochleae (S2A Fig). These results confirm that FGF20

signals through epithelial FGFR1 to the prosensory domain.

Previous studies have also reported a decrease in proliferation in Kölliker's organ (neural to the

prosensory domain, S2B Fig) in Fgfr1-CKO cochleae [20]. We replicated this result by

examining EdU (5-ethynyl-2'-deoxyuridine) incorporation at E14.5. *Fgfr1*-CKO mice had a

complete lack of EdU-incorporating Kölliker's organ cells, while Fgf20-KO mice did not show a

decrease in EdU incorporation (S2B Fig). This finding is also consistent with an additional FGF

ligand signaling via FGFR1, likely at an earlier stage. We do not know whether the proliferation

defect in Kölliker's organ contributes to the reduction in HC and SC number in Fgfr1-CKO mice.

Genetic rescue of the Fgf20-KO phenotype suggests that FGF20 is required for

differentiation

We have previously shown that recombinant FGF9, which is biochemically similar to FGF20

[23,24], is able to rescue the loss of HCs and SCs in Fgf20-KO explant cochleae [21].

Interestingly, while treatment with FGF9 at E13.5 and E14.5 was able to rescue the Fgf20-KO

phenotype, treatment at E15.5 was not. This temporal rescue specificity suggests that FGF20

signaling is required for the initiation of HC and SC differentiation.

To confirm the hypothesis that FGF20 is involved in differentiation and not specification (Fig

2D), we sought to more accurately determine the temporal requirement of FGF20 signaling. To

achieve this, we developed an *in vivo* genetic rescue model of the Fgf20-KO phenotype by

ectopically expressing FGF9. We combined $Fgf20^{Cre}$ with the $Fgf20^{\beta gal}$ [21], $ROSA^{rtTA}$ [25] and

TRE-Fgf9-IRES-eGfp [26] alleles to generate Fgf20-rescue ($Fgf20^{Cre/\beta gal}$;$ROSA^{rtTA/+}$;TRE-Fgf9-

IRES-eGfp) mice along with littermate controls: Fgf20-het ($Fgf20^{Cre/+}$;$ROSA^{rtTA/+}$), Fgf9-OA

($Fgf20^{Cre/+}$;$ROSA^{rtTA/+}$;TRE-Fgf9-IRES-eGfp), and Fgf20-null ($Fgf20^{Cre/\beta gal}$;$ROSA^{rtTA/+}$). These

mice express the reverse tetracycline transactivator (rtTA) in the $Fgf20^{Cre}$ lineage, which

contains the prosensory domain and Kölliker's organ at E13.5 to E15.5 [18]. In mice expressing

TRE-Fgf9-IRES-eGfp, rtTA drives the expression of FGF9 upon doxycycline (Dox) induction.

The $Fgf20^{\beta gal}$ allele is another $Fgf20$-null allele, in which exon 1 of $Fgf20$ is replaced by a

sequence encoding β-galactosidase. We combined $Fgf20^{Cre}$ with $Fgf20^{\beta gal}$ to generate

homozygous mutant mice while maintaining a constant dosage of $Fgf20^{Cre}$ in control and

knockouts.

Initially, pregnant dams were fed a Dox diet from E13.5 to E15.5 and pups were harvested at P0

to examine HC and SC development. As expected, Dox treatment itself did not appear to affect

HC or SC development in Fgf20-het and Fgf20-null cochleae, both of which showed the

expected phenotypes (Figs 3A and 3B). Ectopic expression of FGF9 during these stages also

did not affect HC or SC development in Fgf9-OA cochleae, showing that excess FGF20/FGF9

was not sufficient to produce ectopic HCs and SCs. Importantly, ectopic expression of FGF9

resulted in a full rescue of the number and patterning of HCs and SCs in Fgf20-rescue pups.

The organ of Corti in these rescue pups had one row of IHCs, three rows of OHCs, and five

rows of SCs throughout the entire length of the cochlear duct, without any gaps (Figs 3A and

3B). This shows that FGF20/FGF9 signaling at E13.5-E15.5 is sufficient for HC and SC

differentiation. The quantified results from all of the rescue experiments are summarized in Fig

3C, where the number of OHCs and SCs are represented as a percentage of that of Fgf20-het

mice treated with the same Dox regimen. All of the quantified data are presented in S3 Fig.

To more precisely determine the timing of rescue sufficiency, we fed pregnant dams Dox for a

period of 24 hours starting at E13.5, E14.5, or E15.5. With E13.5 Dox, patterning and OHC

number in the basal turn of the cochlea were completely rescued (Fig 3A). However, OHC

number in the middle and particularly the apical turns were only partially rescued, resulting in

regions with two rows of OHCs instead of three. For instance, in the apical turn, OHC number

was restored to 81% of Fgf20-het mice, which is statistically significantly increased compared to

Fgf20-null, but also statistically significantly decreased compared to Fgf20-het, indicating partial

rescue (Fig 3C). With E14.5 Dox, patterning and OHC number in the middle and apical turns

were completely rescued. However, OHC number in the basal turn was not completely rescued,

with regions of one or two rows of OHCs, instead of three. With E15.5 Dox, patterning and OHC

number was not rescued in the basal and middle turns, as gaps still formed between islands of

HCs (Fig 3A). However, OHC number in the apical turn was partially rescued, with two or three

rows of OHCs not separated by gaps towards the tip of the apex. In all of these experiments,

the rescue of SCs followed the same pattern as that of OHCs (Fig 3B).

These rescue results show that FGF20/FGF9 is sufficient for OHC and SC differentiation in the

basal turn of the cochlea at E13.5, in the middle and apical turns at E14.5-E15.5, and in the tip

of the apical turn at E15.5. Since the initiation of HC and SC differentiation occurs in the

base/mid-base of the cochlea at E13.5 and progresses apically over the next few days, these

results strongly imply that FGF20 functions during the initiation of differentiation, rather than

prosensory specification, consistent with our model (Fig 2D).

Decrease in *Sox2* expression results in similar phenotypes as disruptions to

FGFR1 signaling

Our results and previous findings suggest that FGFR1 regulates prosensory specification via

Sox2 [19]. Mice with an inner ear-specific *Sox2* hypomorphic mutation (*Sox2$^{Ysb/Ysb}$*, see below)

have defects in prosensory specification, accounting for a small loss of HCs and SCs, whereas

mice with inner-ear specific *Sox2* null mutations have a complete lack of prosensory

specification and a complete absence of sensory epithelium [12]. To examine how much the

reduction in *Sox2* expression in Fgfr1-CKO cochlea contributes to the phenotype at P0, we

combined the *Sox2*⁻ (*Sox2* constitutive null) and *Sox2*^{Ysb} alleles to closely examine the effects of

reduction in *Sox2* expression on organ of Corti development, on a similar genetic background

as our Fgf20-KO and Fgfr1-CKO mice. We hypothesized that if *Fgfr1* acts upstream of *Sox2*,

then reducing *Sox2* expression should at least partially recapitulate the Fgfr1-CKO cochlea

phenotype. The *Sox2*^{Ysb} allele is a regulatory mutant in which transgene insertion in

chromosome 3 disrupts some otic enhancers, resulting in hypomorphic *Sox2* expression in the

inner ear [11,12].

We generated a *Sox2* allelic series of mice with the following genotypes, in order of highest to

lowest levels of *Sox2* expression: *Sox2*$^{+/+}$ (wildtype), *Sox2*$^{Ysb/+}$, *Sox2*$^{Ysb/Ysb}$, and *Sox2*$^{Ysb/-}$. In this

allelic series, decrease in *Sox2* expression had a dose-dependent effect on cochlea length at

P0 (Figs 4A-4C). *Sox2*$^{Ysb/+}$ cochleae had a 6% reduction in length compared to wildtype

(although not statistically significant), *Sox2*$^{Ysb/Ysb}$ cochleae had a 24% reduction, and *Sox2*$^{Ysb/-}$

had a 46% reduction. *Sox2*$^{Ysb/+}$ organ of Corti developed relatively normally, with three rows of

OHCs and one row of IHCs (Fig 4A). Interestingly, there were occasional ectopic IHCs medial

(neural) to the normal row of IHCs, especially in the middle and apical turns of the *Sox2*$^{Ysb/+}$

cochlea (Fig 4A, arrowheads). However, there was no significant increase in IHC number (total

or normalized to length) compared to wildtype cochleae (Fig 4D). The *Sox2*$^{Ysb/Ysb}$ cochlea

appeared much more abnormal, with gaps in the sensory epithelium that lacked HCs and SCs

in the basal turn (Figs 4A and 4B), similar to what was observed previously [12]. Moreover, at

the base, in the sensory islands between the gaps, there were often four rows of OHCs and six

rows of SCs. In the middle and apical turns, there were the normal three rows of OHCs and five

rows of SCs. There were also numerous ectopic IHCs throughout the middle and apical turns,

sometimes forming an entire second row of cells (Fig 4A), resulting in increased number of IHCs

in the middle turn compared to wildtype (S4A Fig). However, the total and length-normalized

number of IHCs in $Sox2^{Ysb/Ysb}$ cochleae did not significantly differ from that of wildtype cochleae

(Fig 4D). In terms of OHCs, $Sox2^{Ysb/Ysb}$ cochleae exhibited a 40% decrease in total number

compared to wildtype cochleae (Fig 4E). This decrease was not quite as severe when

normalized to cochlear length (21% decrease). Strikingly, $Sox2^{Ysb/-}$ cochleae lacked almost all

HCs and SCs, except in the apical turn (Figs 4A and 4B). The decrease in OHC number (93%)

in $Sox2^{Ysb/-}$ cochleae compared to wildtype was more severe than the decrease in IHC number

(75%). Notably, IHC number was significantly decreased in the basal and middle turns, but not

in the apical turn (S4A Fig). OHC number was significantly decreased throughout all three turns

(S4B Fig). In all of these genotypes, the number of SCs followed the pattern of loss of OHCs

(Fig 4F and S4C Fig). Interestingly, while $Sox2^{Ysb/-}$ cochleae almost completely lacked HCs and

SCs in the basal and middle turns, in 7 of 11 $Sox2^{Ysb/-}$ cochleae examined, one or two small

islands of HCs or SCs were found at the basal tip (S4D Fig).

Overall, these results showed that the basal end of the cochlea is more sensitive to the loss of

Sox2 expression than the apical end. Furthermore, while both IHCs and OHCs were affected,

OHCs were more sensitive to decrease in *Sox2* expression than IHCs. Importantly, both of

these features were found in Fgfr1-CKO cochleae, where the decrease in IHCs was only found

in the basal and middle turns and there were almost no OHCs along the entire cochlear duct

(S1A and S1B Fig). Therefore, we conclude that decrease in *Sox2* expression, leading to

defects in prosensory specification, could account for the Fgfr1-CKO phenotype. Furthermore,

the decrease in *Sox2* expression could also account for the difference in severity between the

Fgf20-KO and Fgfr1-CKO phenotypes, since Fgf20-KO cochleae, which had normal *Sox2*

expression, did not have a decrease in the number of IHCs, unlike Fgfr1-CKO and $Sox2^{Ysb/-}$

cochleae.

Decrease in levels of Sox2 expression delays prosensory specification

We sought to determine why a decrease in *Sox2* expression more severely affected the basal

end of the cochlear duct. Initially, we examined *Sox2* expression at E14.5. As expected, Sox2

expression was almost completely absent in *Sox2$^{Ysb/-}$* cochleae (S5A Fig). This decrease in

expression was not more severe at the basal turn of the cochlear duct, relative to the middle

and apical turns, suggesting that the more severe basal phenotype in *Sox2$^{Ysb/-}$* cochleae cannot

be explained by differential Sox2 expression. Similarly, CDKN1B expression was downregulated

in the prosensory domain of *Sox2$^{Ysb/-}$* cochleae, consistent with previous studies [12].

Interestingly, the decrease in expression was also not more severe at the basal turn relative to

the middle and apical turns (S5B Fig). Using CDKN1B as a marker of prosensory specification,

this suggests that the more severe basal phenotype also cannot be explained by differential

regulation of prosensory specification along the length of the cochlea.

As described in the introduction, waves of cell cycle exit (marking the completion of prosensory

specification) and initiation of differentiation travel in opposite directions along the cochlear duct

during development, resulting in the basal end of the cochlear duct differentiating immediately

after specification. The apical end, meanwhile, exhibits a delay in differentiation, resulting in a

longer temporal buffer between specification and differentiation. In this developmental pathway,

specification must be completed prior to the initiation of differentiation. We reasoned, therefore,

that disruptions to the timing of prosensory specification will preferentially interfere with basal

sensory epithelia development, potentially accounting for the more severe basal phenotype in

Sox2 hypomorphs.

To test this hypothesis, we examined cell cycle exit in the prosensory domain via Ki67

expression, as a marker of the status of prosensory specification. Ki67 is expressed by cycling

cells, but not cells in G_0 [27]. In the developing cochlea at around E12.5 to E15.5, cells of the

prosensory domain, sometimes referred to as the zone of non-proliferation, have turned off or

are beginning to turn off Ki67 [3]. At E14.5 in $Sox2^{Ysb/+}$ cochleae, the prosensory domain along

most of the cochlear duct (serial sections 2-6) has turned off Ki67 expression, except at the very

base (serial section 1; Fig 5A, brackets). See graphical summary below Fig 5A; also see S5C

Fig for serial "mid-modiolar" sections through the cochlea. This indicates that the wave of cell

cycle exit, which starts at the apex, has reached the very base of the cochlear duct. However, in

$Sox2^{Ysb/-}$ cochleae, only the prosensory domain at the apical turn of the cochlear duct (serial

section 6) has turned off Ki67, not at the mid-basal or basal turns (serial sections 1-3); the

middle turns (serial sections 4 and 5), meanwhile, were just starting to turn off Ki67 (Fig 5A,

brackets). This indicates that cell cycle exit has only reached the middle turn, suggesting a

delay in prosensory specification. In addition, the nuclei of prosensory domain cells shift away

from the luminal surface of the cochlear epithelium upon specification [28]. This basal shift of

nuclei localization within the cell leaves a blank space between DAPI-stained nuclei and the

luminal surface of the cochlear duct, which can be visualized in all six serial sections in $Sox2^{Ysb/+}$

cochleae at E14.5 (Fig 5A, asterisks). However, in $Sox2^{Ysb/-}$ cochleae, cells of the prosensory

domain mostly did not exhibit this nuclei shift at E14.5.

At E15.5, the prosensory domain along the entire length of the cochlear duct has turned off Ki67

expression in both $Sox2^{Ysb/+}$ and $Sox2^{Ysb/-}$ cochleae (Fig 5A, brackets), indicating that

prosensory specification in $Sox2^{Ysb/-}$ cochleae has caught up by this stage. Prosensory nuclei

localization has also begun to catch up at E15.5 in $Sox2^{Ysb/-}$ cochleae (Fig 5A, asterisks).

Overall, these results suggest that prosensory specification is delayed in $Sox2^{Ysb/-}$ cochleae, but

not permanently disrupted.

By definition, prosensory specification must occur prior to differentiation to generate HCs and

SCs. Therefore, the period of time in between cell cycle exit and the initiation of differentiation

represents a temporal buffer (Fig 5B, green shading) preventing differentiation from initiating

prior to specification. As differentiation begins in the basal/mid-basal cochlear turns shortly after

specification, the delay in specification in $Sox2^{Ysb/-}$ cochleae leads to progenitors not having

been specified in time for differentiation at the basal end of the cochlear duct (Fig 5B,

crosshatch pattern). We propose that this at least partially explains why the basal end of the

cochlea is more sensitive to decreases in the level of $Sox2$ expression. Moreover, since

differentiation begins in the mid-base and spreads to the rest of the base and apex, progenitors

at the basal tip in $Sox2^{Ysb/-}$ cochleae may still undergo specification prior to differentiation. This

may explain why small islands of HCs and SCs are sometimes seen in the basal tip of $Sox2^{Ysb/-}$

cochleae (S4D Fig).

$Sox2$ is upstream of $Fgf20$

While the delay in prosensory specification can explain the preferential loss of sensory

epithelium from the basal end of $Sox2$ hypomorph cochleae, it does not readily explain the

preferential loss of OHCs, relative to IHCs. Since this preference for OHC loss is reminiscent of

the $Fgf20$/$Fgfr1$ deletion phenotypes, we investigated the possibility that $Sox2$ may be upstream

of FGF20-FGFR1 signaling. Interestingly, both $Etv4$ and $Etv5$ were dramatically downregulated

in the prosensory domain of $Sox2^{Ysb/-}$ cochleae compared to control (Fig 6A). This shows that

FGF20-FGFR1 signaling was disrupted in the $Sox2$ hypomorph cochleae. Examination of $Fgfr1$

and $Fgf20$ expression by in situ hybridization revealed that while $Fgfr1$ expression did not

appear to be affected in $Sox2^{Ysb/-}$ cochleae at E14.5, $Fgf20$ expression was absent (Fig 6B).

This suggests that while $Fgfr1$ functions upstream of $Sox2$, $Fgf20$ is downstream of $Sox2$. This

model predicts that *Fgf20* expression would be downregulated in Fgfr1-CKO cochleae, which

was confirmed by in situ hybridization (Fig 6C).

The above results indicate that the loss of *Fgf20* could partially account for the *Sox2*$^{Ysb/-}$

phenotype. To determine whether loss of *Fgf20* also causes delayed prosensory specification,

we examined Ki67 expression in Fgf20-KO cochleae. At E14.5, there was no detectable delay in

cell cycle exit in Fgf20-KO cochleae, as loss of Ki67 expression reached the base (serial section

1) in both control and Fgf20-KO cochleae (Fig 6D, brackets). See S6A Fig for serial "mid-

modiolar" sections through the cochlea. There was also no detectable delay in prosensory basal

nuclei shift in Fgf20-KO cochleae (Fig 6D, asterisks). These results were expected as the

Fgf20-KO phenotype is not more severe at the basal end of the cochlear duct. This is also

consistent with *Fgf20* being required during differentiation rather than prosensory specification

(Fig 2D). However, these results do not answer whether and how the loss of *Fgf20* contributes

to the *Sox2* hypomorph phenotype.

We also asked whether decrease in *Sox2* expression can account for the absence of

proliferation in Kölliker's organ of Fgfr1-CKO cochleae. Interestingly, EdU-incorporation was

decreased in Kölliker's organ in *Sox2*$^{Ysb/-}$ cochleae at E14.5, especially in the region adjacent to

the prosensory domain (S6B Fig, bracket). However, EdU-incorporation was not completely

absent from Kölliker's organ, unlike in Fgfr1-CKO cochleae. This suggests that loss of *Sox2* in

combination with other factors contributes to Kölliker's organ phenotype in Fgfr1-CKO cochleae.

***Sox2* and *Fgf20* interact during cochlea development**

To explore how the loss of *Fgf20* contributes to the *Sox2* hypomorph phenotype, we combined

the *Fgf20*⁻ and *Sox2*^{Ysb} alleles to generate *Fgf20* and *Sox2* compound mutants. We also

hypothesized that reducing *Sox2* expression in Fgf20-KO mice would recapitulate (or

phenocopy) the more severe Fgfr1-CKO phenotype. We interbred F1 mice from the same

parents to generate nine different F2 genotypes encompassing all possible combinations of the

Fgf20⁻ and *Sox2^Ysb^* alleles: *Fgf20^+/+^;Sox2^+/+^*, *Fgf20^+/+^;Sox2^Ysb/+^*, *Fgf20^+/-^;Sox2^+/+^*,

*Fgf20^+/-^;Sox2^Ysb/+^*, *Fgf20^+/+^;Sox2^Ysb/Ysb^*, *Fgf20^+/-^;Sox2^Ysb/Ysb^*, *Fgf20^-/-^;Sox2^+/+^*, *Fgf20^-/-^;Sox2^Ysb/+^*,

and *Fgf20^-/-^;Sox2^Ysb/Ysb^* (Figs 7A and 7B and data not shown). At P0, an overview of HCs and

SCs showed that the *Fgf20^+/-^;Sox2^Ysb/+^* phenotype mostly resembled that of *Fgf20^+/+^;Sox2^+/+^*,

*Fgf20^+/+^;Sox2^Ysb/+^*, and *Fgf20^+/-^;Sox2^+/+^* cochleae, except for the prevalence of ectopic IHCs (Fig

7A, arrowheads). The *Fgf20^+/-^;Sox2^Ysb/Ysb^* phenotype mostly resembled that of

*Fgf20^+/+^;Sox2^Ysb/Ysb^* cochleae, but with more gaps in the basal cochlear turn and two rows of

IHCs throughout the length of the cochlear duct, except where there were gaps. The *Fgf20^-^*

*^/-^;Sox2^Ysb/+^* phenotype mostly resembled that of *Fgf20^-/-^;Sox2^+/+^* cochleae, but with smaller

sensory islands in between gaps. The *Fgf20^-/-^;Sox2^Ysb/Ysb^* phenotype appeared by far the most

severe, with almost a complete absence of IHCs, OHCs, and SCs from the basal turn, and tiny

sensory islands in the middle turn; however, the apical turn appeared similar to that of *Fgf20^-^*

*^/-^;Sox2^Ysb/+^* and *Fgf20^-/-^;Sox2^+/+^* cochleae (Figs 7A and 7B).

Quantification of the phenotypes are presented in Figs 8B-8E and S7B-S7D Figs. We analyzed

the quantified P0 phenotype via two-way ANOVA with the two factors being gene dosage of

Fgf20 (levels: *Fgf20^+/+^*, *Fgf20^+/-^*, *Fgf20^-/-^*) and *Sox2* (levels: *Sox2^+/+^*, *Sox2^Ysb/+^*, *Sox2^Ysb/Ysb^*).

Results from the two-way ANOVA and post-hoc Tukey's HSD are presented in Figs 8A, 8F, and

S7A and S8 Figs. Cochlear length and the total number of IHCs, OHCs, and SCs were all

significantly affected by both the *Fgf20* dosage and the *Sox2* dosage, as well as an interaction

between the two factors (Figs 8A-8E). The statistically significant interaction between *Fgf20* and

Sox2 dosages suggests that *Fgf20* and *Sox2* have a genetic interaction in regulating cochlear

length as well as the number of IHCs, OHCs, and SCs (Fig 8A). Notably, *Fgf20^+/-^;Sox2^Ysb/Ysb^*

cochleae had significantly fewer OHCs and SCs than $Fgf20^{+/+};Sox2^{Ysb/Ysb}$ cochleae, and $Fgf20^{-}$

$^{/-};Sox2^{Ysb/+}$ cochleae had significantly fewer OHCs than $Fgf20^{-/-};Sox2^{+/+}$ cochleae (Fig 8F).

Importantly, $Fgf20^{-/-};Sox2^{Ysb/Ysb}$ cochleae had decreased total and length-normalized number of

IHCs, which was not observed in any of the other genotypes, strongly supporting a genetic

interaction between $Fgf20$ and $Sox2$ ($Fgf20^{+/+};Sox2^{Ysb/Ysb}$ cochleae did have a slight decrease in

the total number IHCs, but not in the length-normalized number of IHCs).

Interestingly, while the total number of IHCs was decreased in $Fgf20^{-/-};Sox2^{Ysb/Ysb}$ cochleae

relative to all other genotypes, this decrease was only found in the basal and middle turns, but

not the apical turn (S7B and S8 Figs). No such basal/middle/apical turn discrepancies existed in

the number of OHCs or SCs (S7C, S7D, and S8 Figs). This is reminiscent of the Fgfr1-CKO and

$Sox2^{Ysb/-}$ phenotypes.

To ensure that the $Fgf20^{-}$ and $Sox2^{Ysb}$ interaction is not purely an artifact of the $Sox2^{Ysb}$ allele,

we generated $Fgf20^{+/+};Sox2^{+/+}$ (wildtype), $Fgf20^{+/-};Sox2^{+/+}$ (Fgf20-het), $Fgf20^{+/+};Sox2^{+/-}$ (Sox2-

het), and $Fgf20^{+/-};Sox2^{+/-}$ (double het) mice to look for an interaction between the $Fgf20^{-}$ and

$Sox2^{-}$ alleles (S9A Fig). At P0, cochlear length did not significantly differ among the four

genotypes (S9B Fig). HC quantification showed that neither $Fgf20$ nor $Sox2$ exhibited

haploinsufficiency for total or length-normalized number of IHCs or OHCs (S9C and S9D Figs).

However, in Fgf20-het and much more so in Sox2-het cochleae, occasional ectopic IHCs can be

found in the middle and apical turns of the cochlear duct (S9A Fig, arrowheads). Interestingly, in

double het cochleae, many more ectopic IHCs were found, even in the basal turn. These

ectopic IHCs led to an increase in the total and length-normalized number of IHCs in double het

cochleae, compared to wildtype (S9C Fig). Notably, a significant increase in IHCs was only

found in the basal turn, not the middle or apical turns (S9E Fig). In the basal turn, IHC number

was significantly increased in double het cochleae compared to wildtype, Fgf20-het, and Sox2-

het cochleae. Double het cochleae also had a significant decrease in total and length-

normalized number of OHCs compared to wildtype (S9D Fig). Again, a significant decrease in

OHCs was only found in the basal turn, not the middle or apical turns (S9F Fig). These results

confirm a genetic interaction between *Fgf20* and *Sox2*.

Loss of *Fgf20* does not further delay prosensory specification in *Sox2*

hypomorph cochleae

We propose that the $Fgf20^{-/-}$;$Sox2^{Ysb/Ysb}$ phenotype lies in between that of Fgfr1-CKO and

$Sox2^{Ysb/-}$ in terms of severity of reductions in cochlear length and in the number of HCs and

SCs. We further hypothesize that these three phenotypes form a continuum with the Fgf20-KO

phenotype (Fig 9A). Along this continuum, all four genotypes lack FGF20 signaling, but vary in

the level of *Sox2* expression and phenotype severity in the basal end of the cochlear duct and

the outer compartment (outer rows of OHCs and SCs). From this, and from the $Fgf20^{-}$ and

$Sox2^{Ysb}$ series of alleles, we conclude that the basal end of the cochlear duct and the outer

compartment are more sensitive to the loss of *Fgf20* and *Sox2*, relative to the apical end and

inner compartment, respectively.

To determine the mechanism underlying the *Sox2* and *Fgf20* interaction, we asked whether the

similarity between the $Fgf20^{-/-}$;$Sox2^{Ysb/Ysb}$ and $Sox2^{Ysb/-}$ phenotypes could be explained by a

further decrease in *Sox2* levels in $Fgf20^{-/-}$;$Sox2^{Ysb/Ysb}$ cochleae from $Sox2^{Ysb/Ysb}$ levels. In other

words, we asked whether loss of *Fgf20* further reduces *Sox2* expression on a *Sox2*

hypomorphic background. Examination of prosensory domain *Sox2* expression at E14.5

revealed, as expected, that $Fgf20^{-/-}$;$Sox2^{Ysb/+}$ cochleae did not have a decrease in Sox2

expression compared to $Fgf20^{+/-}$;$Sox2^{Ysb/+}$ (S10A Fig). $Fgf20^{-/-}$;$Sox2^{Ysb/Ysb}$ cochleae also did not

have a further decrease in Sox2 expression compared to $Fgf20^{+/-};Sox2^{Ysb/Ysb}$ cochleae.

Moreover, despite the loss of sensory epithelium in most of the basal turn, Sox2 expression was

not further decreased in the basal turn at E14.5 relative to the rest of the $Fgf20^{-/-};Sox2^{Ysb/Ysb}$

cochlea (S10A Fig). These data confirm that *Fgf20* does not regulate *Sox2* expression. A similar

pattern of expression was observed for CDKN1B across the different genotypes (S10B Fig).

Loss of *Fgf20* did not contribute to a further decrease in CDKN1B expression on a $Sox2^{Ysb/Ysb}$

background, nor was there a basal-apical difference in CDKN1B expression in $Fgf20^{-}$

$^{/-};Sox2^{Ysb/Ysb}$ cochleae at E14.5.

Next, we asked whether *Sox2* and *Fgf20* interact to delay prosensory specification. We showed

that Fgf20-KO cochleae do not exhibit a delay in prosensory specification (Fig 6D). However,

this does not rule out the possibility that the loss of *Fgf20* on a *Sox2* hypomorphic background

may contribute to a delay. We examined Ki67 expression at E14.5 and found that in

$Fgf20^{+/-};Sox2^{Ysb/+}$ cochleae, prosensory domain cell cycle exit has reached the end of the base

(serial section 1; Fig 9B, brackets). See S10D Fig for serial "mid-modiolar" sections through the

cochlea. Similarly, cell cycle exit in $Fgf20^{-/-};Sox2^{Ysb/+}$ cochleae also reached the very base. As

expected, $Fgf20^{+/-};Sox2^{Ysb/Ysb}$ cochleae exhibited a slight delay in prosensory specification; cell

cycle exit has reached the base (serial section 2), but has not yet reached the end of the base

(serial section 1). Importantly, $Fgf20^{-/-};Sox2^{Ysb/Ysb}$ cochleae did not show a further delay relative

to $Fgf20^{+/-};Sox2^{Ysb/Ysb}$. There was also no detectable delay in basal nuclei shift in $Fgf20^{-}$

$^{/-};Sox2^{Ysb/+}$ or $Fgf20^{-/-};Sox2^{Ysb/Ysb}$ cochleae (Fig 9B, asterisks). These results suggest that the

loss of *Fgf20* does not contribute to delayed specification and that the severity of the $Fgf20^{-}$

$^{/-};Sox2^{Ysb/Ysb}$ basal phenotype cannot be completely attributed to delayed specification.

Lastly, we examined proliferation in the $Fgf20^{-}$ and $Sox2^{Ysb}$ E14.5 cochleae. Interestingly, there

was a noticeable decrease in the number of EdU-incorporating cells in Kölliker's organ in $Fgf20^{-}$

$^{/-}$;*Sox2$^{Ysb/Ysb}$* cochleae, compared to *Fgf20$^{+/-}$;Sox2$^{Ysb/+}$*, *Fgf20$^{-/-}$;Sox2$^{Ysb/+}$*, and *Fgf20$^{+/-}$;Sox2$^{Ysb/Ysb}$*

cochleae (S10C Fig). This phenotype is similar to that of *Sox2$^{Ysb/-}$* cochleae and is less severe

than that of Fgfr1-CKO cochleae. This suggests that *Fgf20* and *Sox2* interact to regulate

proliferation in Kölliker's organ, although other factors downstream of *Fgfr1* also contribute.

Fgf20-KO organ of Corti exhibits premature differentiation

We showed that *Fgf20* likely plays a role during the initiation of differentiation. Previous studies

showed that deletion of both transcription factors *Hey1* and *Hey2* results in premature

differentiation in the organ of Corti [29]. Furthermore, it has been suggested that FGF signaling,

in particular FGF20, regulates *Hey1* and *Hey2* expression during this process [8,29]. To test

whether *Fgf20* is upstream of *Hey1* and *Hey2*, we looked at the expression of the two

transcription factors via in situ hybridization. In Fgf20-KO cochleae at E14.5, *Hey1* expression is

downregulated while *Hey2* is almost completely absent compared to control (Fig 10A). To test

whether FGF20 loss leads to premature differentiation, we examined myosin VI (Myo6)

expression, a marker of differentiated HCs [29]. At E14.5, the cochleae of 3 of 12 control

embryos examined contained Myo6-expressing HCs, while the cochleae of 18 of 19 littermate

Fgf20-KO embryos contained Myo6-expressing HCs (p < 0.001, Fisher's exact test; Fig 10B). If

present, the Myo6-expressing HCs at this stage were always found in the basal and mid-basal

turns of the cochlea. These results show that there is premature onset of differentiation in

Fgf20-KO cochleae, which begins in the basal/mid-basal turns. This result is surprising given

our previous finding of delayed differentiation in the apical end of Fgf20-KO cochleae at later

stages, which we confirm here (S1E Fig). These findings suggest that apical progression of

differentiation may be slower in Fgf20-KO cochleae.

Next, we asked whether ectopic activation of FGF signaling via overexpression of FGF9 will

delay the onset of differentiation. We generated Fgf20-het ($Fgf20^{Cre/+};ROSA^{rtTA/+}$), Fgf20-null

($Fgf20^{Cre/βgal};ROSA^{rtTA/+}$), Fgf9-OA ($Fgf20^{Cre/+};ROSA^{rtTA/+}$;TRE-Fgf9-IRES-eGfp), Fgf20-rescue

($Fgf20^{Cre/βgal};ROSA^{rtTA/+}$;TRE-Fgf9-IRES-eGfp) mice as before and started Dox induction at

E13.5 until E15.0 (Fig 3). At E15.0, all of the Fgf20-het (4/4) and Fgf20-null (4/4) cochleae

contained Myo6-expressing HCs, while none of the Fgf9-OA (0/4) and Fgf20-rescue (0/4)

cochleae contained Myo6-expressing HCs (Fig 10C). This suggests that ectopic expression of

FGF9 was able to delay the onset of differentiation, even with the lack of endogenous FGF20.

Despite this delay in onset of differentiation, by P0, differentiation has apparently caught up in

both Fgf9-OA and Fgf20-rescue cochleae (Fig 3A).

Similar to a delay in prosensory specification, premature onset of differentiation narrows the

temporal buffer between the completion of specification and initiation of differentiation towards

the cochlear base. In the context of a slight delay in specification due to decreased *Sox2* levels,

premature differentiation from the loss of *Fgf20* can lead to an attempt at differentiation before

specification in the basal end of the cochlea. We propose that *Sox2* and *Fgf20* interact to

regulate the boundaries of the temporal buffer, helping to ensure that differentiation begins after

the completion of specification (Fig 11).

Discussion

***Fgfr1* is involved in prosensory specification and differentiation, while *Fgf20* is**

only involved in differentiation

Fgf20 and *Fgfr1* are required for HC and SC development. Based on similarities in the

phenotype caused by the loss of FGF20 and loss of FGFR1 signaling, FGF20 has been

hypothesized as the FGFR1 ligand during organ of Corti development [17–21]. However, the

exact role of FGF20/FGFR1 during organ of Corti development has been a topic of debate. We

previously reported that Fgf20-KO mice do not have defects in prosensory specification, and

have a normally formed prosensory domain [21]. We further showed that FGF20 signaling is

important during the initiation stage of differentiation, and that Fgf20-KO cochleae have gaps in

the differentiated sensory epithelium filled with undifferentiated prosensory progenitors.

However, other studies have shown *in vitro* that FGF20 regulates prosensory specification via

Sox2 [33] and *in vivo* that FGFR1 is required for prosensory specification via Sox2 [19]. Here,

using an *in vivo* rescue model, we show that ectopic FGF9 signaling is sufficient to rescue the

Fgf20-KO phenotype in a spatiotemporal pattern that matched the timing of initiation of

differentiation along the length of the cochlear duct. We conclude, therefore, that FGF20 is

involved in differentiation and not necessary for prosensory specification.

Notably, the Fgf20-KO phenotype, in which two-thirds of OHCs fail to develop, is not as severe

as the Fgfr1-CKO phenotype, which lacks almost all OHCs as well as half of IHCs. Potential

explanations for this include differences in mouse genetic background, and the existence of a

redundant FGF ligand(s). To rule out the former, we examine here Fgf20-KO and Fgfr1-CKO

mice on a similar genetic background, and replicated the difference in phenotype severity. We

also replicated the decrease in *Sox2* expression in the prosensory domain previously reported

in Fgfr1-CKO mice [19]. We further reaffirmed that *Sox2* expression in the prosensory domain is

not affected by the loss of *Fgf20*. This suggests that another FGF ligand signaling through

FGFR1 is required to maintain *Sox2* expression during prosensory specification. The identity of

this ligand is currently unknown.

Foxg1^{Cre} has been used in several studies to target the otic epithelium, including to conditionally

delete *Fgfr1* [18–20]. One concern with *Foxg1^{Cre}* is that it is a null allele [22]. *Foxg1*-null mice

have shortened cochlear length, although HC and SC differentiation did not appear to be

directly affected [34]. Previous work [35] and our results here showed that *Foxg1* is not

haploinsufficient during cochlea development, as *Foxg1$^{Cre/+}$;Fgfr1$^{flox/+}$* cochleae had very similar

phenotypes to *Fgfr1$^{+/-}$* cochleae. Moreover, the use of the Six1enh21-Cre transgene, which

targets the otic epithelium in a similar spatiotemporal pattern as *Foxg1^{Cre}*, to conditionally delete

Fgfr1 resulted in the same phenotype as *Foxg1$^{Cre/+}$;Fgfr1$^{flox/-}$* cochleae [19]. This included the

loss of almost all OHCs, loss of IHCs, and decreased prosensory *Sox2* expression. Therefore,

the increased severity of *Foxg1$^{Cre/+}$;Fgfr1$^{flox/-}$* cochleae relative to *Fgf20$^{-/-}$* cochleae is likely not

attributable *Foxg1* haploinsufficiency.

We hypothesized that the severity of the Fgfr1-CKO phenotype is due to the loss of FGF20

signaling during differentiation and decreased *Sox2* expression causing disrupted prosensory

specification. Consistent with this hypothesis, the combination of *Fgf20$^{-/-}$* and *Sox2$^{Ysb/Ysb}$*

mutations phenocopied Fgfr1-CKO cochleae. The similarities in phenotype include

approximately a 30% reduction in cochlear length and almost a complete loss of OHCs and SCs

and approximately a 50% loss of IHCs. Interestingly, the *Fgf20$^{-/-}$;Sox2$^{Ysb/Ysb}$* phenotype is also

similar to the *Sox2$^{Ysb/-}$* phenotype. We conclude that the Fgfr1-CKO, *Fgf20$^{-/-}$;Sox2$^{Ysb/Ysb}$*, and

Sox2$^{Ysb/-}$ phenotypes likely lie along the same continuum, as these three genotypes all exhibited

a lack of *Fgf20* expression or signaling and varying levels of *Sox2* expression (Fig 9A). Fgf20-

KO cochleae, in which *Sox2* expression was not affected, lies at the mild end of this continuum.

Interestingly, this continuum shows that in the absence of *Fgf20* expression or signaling,

reductions in the level of *Sox2* most severely affected sensory epithelium development of the

cochlear base and the outer compartment. Moving from the Fgf20-KO (mild) end of the

spectrum towards the $Sox2^{Ysb/-}$ (severe) end, increasing numbers of HCs and SCs are lost,

preferentially form the cochlear base and the outer compartment.

***Sox2* and *Fgf20* interact to affect development towards the basal end of the**

cochlea

We show here conclusive evidence that *Sox2* and *Fgf20* genetically interact during cochlea

development. Interestingly, HC and SC development towards the basal end of the cochlea is

more severely affected by the loss of *Sox2* and *Fgf20* and their interaction. While we

hypothesize that *Sox2* and *Fgf20* are involved in distinct steps during organ of Corti

development (prosensory specification and differentiation, respectively), there is nevertheless

potential for a strong interaction. We propose that the timing of specification and differentiation

define a temporal buffer that normally prevents differentiation from initiating prior to the

completion of specification, and that *Sox2* and *Fgf20* modulate the borders of this buffer. In a

developmental pathway, the upstream event (specification) must occur prior to the downstream

event (differentiation). Therefore, loss of *Sox2* and *Fgf20* leading to delayed specification and

premature differentiation onset, respectively, disrupts the temporal buffer, especially towards the

cochlear base (Fig 11).

Specification must occur prior to the onset of differentiation. However, cell cycle exit does not

need to occur prior to the onset of differentiation, as mice lacking *Cdkn1b*, which is required for

cell cycle exit, still produce HCs and SCs (Chen & Segil, 1999; Kanzaki et al., 2006). Here, we

use cell cycle exit in the prosensory domain (also known as the zone of non-proliferation) as a

marker for the completion of specification (Chen & Segil, 1999). We hypothesize that

prosensory cells become specified and primed for differentiation upon withdrawal from the cell

cycle. Previous studies showed that prosensory cells are indeed capable of differentiating into

HCs and SCs directly after cell cycle exit, even in the apex. When *Shh* was deleted from the

spiral ganglion, differentiation began in the apex shortly after cell cycle exit and progressed

towards the base [5]. This suggests that specification occurs in an apex-to-base direction. We

cannot rule out, however, that specification occurs in the same direction as differentiation (base-

to-apex), independently of cell cycle exit. Such a scenario would still be consistent with our

model that a combination of delayed specification and premature onset of differentiation

accounts for the more severe basal phenotype in *Fgf20/Sox2* mutants.

The effect of loss of *Fgf20* on the timing of differentiation is small. We estimate that the onset of

differentiation in Fgf20-KO cochleae is advanced by only around 0.5 days. By itself, this effect

does not lead to a more severe mid-basal or basal phenotype in Fgf20-KO cochleae. However,

we present evidence that on a sensitized genetic background of delayed specification, this small

change in the timing of differentiation leads to a large defect in HC and SC production towards

the basal end of the cochleae. We propose that this at least partially explains the interaction

between *Sox2* and *Fgf20*. Furthermore, the relative sparing of development towards the apical

end of $Sox2^{Ysb/Ysb}$;$Fgf20^{-/-}$ cochleae, especially of IHCs, can be further explained by a delay in

differentiation at the apical end due to the loss of *Fgf20*. We do not know why an apical-basal

difference in timing of differentiation exists in Fgf20-KO cochleae. Perhaps there is a delay in

the apical progression of differentiation, or perhaps other factors contribute to the differentiation

of the apical end of the cochlea. Consistent with the latter, by P7 in Fgf20-KO cochleae, the

apical tip contains a full complement of IHCs and OHCs [21].

Notably, while we show the potential for a *Sox2* and *Fgf20* interaction in modulating the

temporal buffer between specification and differentiation, *Sox2* also has known roles during HC

and SC differentiation [13,15,37,38]. Therefore, the genetic interaction may occur during

differentiation as well. While interaction at this stage may explain the preferential loss of outer

compartment cells in *Sox2* and *Fgf20* mutants, it does not explain the selective loss of basal

cochlear HCs and SCs. Therefore, we conclude that the *Sox2* and *Fgf20* interaction regulates

the temporal buffer, with potential further interactions during differentiation.

The Notch ligand Jagged1 (Jag1) is thought to be important for cochlear prosensory

specification via lateral induction [39–45]. Interestingly, Notch signaling has also been shown to

be upstream of both *Fgf20* and *Sox2* in the developing cochlea [33]. Conditional deletion of

Jag1 or *Rbpj*, the major transcriptional effector of canonical Notch signaling, resulted in the loss

of HCs and SCs, particularly from the basal end of the cochlear duct, similarly to *Fgf20*/*Sox2*

mutants. Unlike *Fgf20*/*Sox2* mutants, however, deletion of *Jag1* or *Rbpj* led to preferential loss

of Sox2 and CDKN1B expression from the prosensory domain at the basal end of the cochlea

[40,42,46]. This suggests that Jag1-Notch signaling is required for prosensory specification,

especially towards the cochlear base. This likely accounts for the more severe basal phenotype

of *Jag1* or *Rbpj* mutants. This same mechanism likely does not explain the more severe basal

phenotype of *Fgf20/Sox2* mutants, as Sox2 and CDKN1B expression was not more severely

reduced or absent in the cochlear base in these mice. Notably, not all studies agree that *Jag1* or

Rbpj is required for Sox2 and CDKN1B expression or for prosensory specification [47]. Further

studies are required to elucidate the functional relationship between *Jag1/Notch*, *Fgf20*, and

Sox2 during cochlea development.

Other genes that potentially interact with *Fgf20* and *Sox2* during cochlea development include

Mycn (*N-Myc*) and *Mycl* (*L-Myc*). Interestingly, deletion of *Mycn* and *Mycl* from the cochlear

epithelium results in accelerated cell cycle exit and delayed initiation of differentiation [48],

opposite to the effects of loss of *Sox2* and *Fgf20*. Addressing potential interactions between

Sox2, *Fgf20*, *Mycn*, and *Mycl* is another topic for future studies.

Outer compartment of the cochlear sensory epithelium is more sensitive to the

loss of *Fgfr1, Fgf20*, and *Sox2* than the inner compartment.

In all of the genotypes we observed in this study, the loss of outer compartment cells (i.e.

OHCs) was predominant. Only in the most severe cases in which almost all OHCs were

missing, as seen in Fgfr1-CKO, $Fgf20^{-/-};Sox2^{Ysb/Ysb}$, and $Sox2^{Ysb/-}$ cochleae, were IHCs also lost.

Similarly, reduction in SC number always preferentially affected the outermost cells. This

suggests that the organ of Corti outer compartment is more sensitive to the loss of *Fgfr1*, *Fgf20,*

and *Sox2* than the inner compartment. The combination of $Fgf20^{-}$ and $Sox2^{Ysb}$ alleles elegantly

demonstrates this: as the number of $Fgf20^{-}$ and $Sox2^{Ysb}$ alleles increased, the number of OHCs

progressively decreased. In the double homozygous mutants, the number of IHCs decreased as

well.

Previous studies noted that the dosage of *Fgfr1* affects the degree of organ of Corti outer

compartment loss. In *Fgfr1* hypomorphs with 80% reduction in transcription, only the third row of

OHCs were missing, while 90% hypomorphs had a slightly more severe phenotype [20].

Therefore, *Fgfr1* loss preferentially affects the outermost HCs. Other studies suggested that the

timing of *Fgfr1* deletion is important in determining the degree of outer compartment loss and

level of *Sox2* expression. When an earlier-expressed Cre driver (Six1enh21-Cre) was used to

conditionally delete *Fgfr1*, almost all OHCs and some IHCs were lost, with a 66% reduction in

Sox2 expression at E14.5 [19]. When a later-expressed Cre driver ($Emx2^{Cre}$) was used, many

more OHCs and IHCs remained, with only a 12% reduction in *Sox2* expression. Our results are

consistent with both of these studies. We show that FGF20-independent FGFR1 signaling and

Sox2 are required early, affecting both IHC and OHC development, while FGF20-FGFR1

signaling is important during later stages, affecting only OHC development.

Differentiation in the organ of Corti not only occurs in a basal-to-apical gradient, but also occurs

in an orthogonal inner-to-outer gradient. That is, IHCs differentiate first, followed by each

sequential row of OHCs [49]. This wave of differentiation suggests that perhaps outer

compartment HCs and SCs require a longer temporal buffer between specification and

differentiation. The genetic interaction between *Sox2* and *Fgf20* in modulating this temporal

buffer, therefore, could also account for the loss of outer compartment HCs and SCs. We

hypothesize that the requirement for a longer temporal buffer may also be involved in

determining OHC fate. In *Fgf20*$^{+/-}$;*Sox2*$^{+/-}$ cochleae, there was a slight decrease in OHCs that

was compensated for by ectopic IHCs, suggesting a fate switch from OHCs into IHCs. Here, we

confirmed previous suggestions that *Fgf20* regulates *Hey1* and *Hey2* to prevent premature

differentiation in the developing organ of Corti [8,29]. Interestingly, in *Hey1*/*Hey2* double

knockout cochleae, there was a similar slight decrease in OHCs compensated for by ectopic

IHCs [29]. Furthermore, inner ear-specific deletion of either *Smoothened* or *Neurod1*, which led

to premature differentiation in the apical cochlear turn, also led to loss of OHCs and the

presence of ectopic IHCs at the apex [6,8]. These findings further support a model where timing

of specification and differentiation affect IHC versus OHC fate, an interesting and important

topic for future studies.

Previously, we hypothesized that *Fgf20* is strictly required for the differentiation of an outer

compartment progenitor [21]. However, data we present here show that *Fgf20*, on a sensitized,

Sox2 hypomorphic background, is also required for inner compartment differentiation. We

conclude that inner and outer compartment progenitors likely are not distinct populations.

Rather, all prosensory progenitors giving rise to the organ of Corti exist on an inner-to-outer

continuum. FGF20 signaling, in combination with other factors including Sox2, are required for

the proper development of all of these cells, though with varying sensitivities.

The relationship between *Fgf20* and *Hey1/Hey2* in regulating differentiation is

complex

We show *in vivo* that *Fgf20* is upstream of *Hey1* and *Hey2*. Supporting this result, *Fgfr1* has

also been shown *in vivo* to be upstream of *Hey2* [19]. Interestingly, in explant studies, inhibition

of FGF signaling alone did not result in decreased *Hey1/Hey2* expression or premature

differentiation [29]. However, FGF inhibition was able to rescue the overexpression of

Hey1/*Hey2* and the delay in differentiation induced by SHH signaling overactivation [8,29].

These discrepancies suggest that the relationship between *Fgf20* and *Hey1*/*Hey2* is more

complicated than we currently understand. Notably, *Hey1*/*Hey2* double knockout cochleae do

not exhibit a loss of OHCs to the extent of Fgf20-KO cochleae, suggesting that other genes

downstream of *Fgf20* are important in organ of Corti development. Moreover, deletion of *Fgf20*

only led to premature differentiation at the basal and mid-basal turns. *Fgf20* deletion actually

delayed differentiation in the apical end of the cochlea. Deletion of *Hey1*/*Hey2*, contrarily, led to

premature differentiation along the entire length of the cochlear duct, although it is unclear how

Hey1/Hey2 loss affects the timing of apical differentiation beyond E15.0 [29]. This suggests that

other factors downstream of *Fgf20* interact with *Hey1*/*Hey2* to regulate the timing of

differentiation. Perhaps these same genes contribute to the loss of OHCs in Fgf20-KO

cochleae. *Mekk4*, which has been shown to be downstream of *Fgf20* and necessary for OHC

differentiation [50] could be one of these genes. Identifying other factors downstream of *Fgf20*

will be a topic of future studies.

Materials and methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (*M. musculus*)	*Fgf20^{Cre}*	Huh et al., 2015	MGI:5751785	
Genetic reagent (*M. musculus*)	*Fgf20^{βgal}*	Huh et al., 2012	RRID:MGI:5425887	
Genetic reagent (*M. musculus*)	*Foxg1^{Cre}*	Hébert and McConnell, 2000	RRID:IMSR_JAX:00 4337	
Genetic reagent (*M. musculus*)	*Fgfr1^{flox}*	Trokovic et al., 2003	RRID:IMSR_CMMR: 0268	
Genetic reagent (*M. musculus*)	Sox2^{Cre}	Hayashi et al., 2002	RRID:IMSR_JAX:00 4783	
Genetic reagent (*M. musculus*)	*Sox2^{Ysb}*	Dong et al., 2002	RRID:IMSR_EM:050 15	
Genetic reagent (*M. musculus*)	*Sox2^{flox}*	Shaham et al., 2009	RRID:IMSR_JAX:01 3093	
Genetic reagent (*M. musculus*)	*ROSA^{rtTA}*	Belteki et al., 2005	RRID:IMSR_JAX:00 5670	

Genetic reagent (*M. musculus*)	TRE-Fgf9-IRES-eGfp	White et al., 2006	MGI:5538516	
Recombinant DNA reagent	*pBluescriptKS-Fgfr1TM*	K. Peters		
Recombinant DNA reagent	*pGEMT-Fgf20*			
Recombinant DNA reagent	*pBluescriptSK-Sox2*	R. Lovell-Badge		Gift of A. Kiernan
Recombinant DNA reagent	*pGEM-Etv4*	G. Martin		
Recombinant DNA reagent	*pBluescriptSK-Etv5*	G. Martin		
Recombinant DNA reagent	*pT7T3D-Hey1*	IMAGE clone	#478014	Gift of S. Rentschler
Recombinant DNA reagent	*pCMVSPORT6-Hey2*	IMAGE clone	#5374813	Gift of S. Rentschler
Antibody	Sheep polyclonal anti-Digoxigenin-AP	Sigma-Aldrich	11093274910	1:1000
Antibody	Rabbit polyclonal anti-P75NTR	EMD Millipore	AB1554	1:300
Antibody	Rabbit polyclonal anti-Prox1	EMD Millipore	ABN278	1:1000
Antibody	Goat polyclonal anti-Sox2	Santa Cruz Biotechnology	sc-17320	1:200

Antibody	Rabbit polyclonal anti-p27Kip1	Neomarkers	RB-9019-P	1:50
Antibody	Rabbit polyclonal anti-Ki67	Abcam	ab15580	1:200
Antibody	Rabbit polyclonal anti-Myo6	Santa Cruz Biotechnology	sc-50461	1:100
Antibody	Donkey polyclonal anti-Rabbit IgG, Alexa Fluor 488	Thermo-Fisher Scientific	A-21206	1:500
Antibody	Donkey polyclonal anti-Goat IgG, Alexa Fluor 555	Thermo-Fisher Scientific	A-21432	1:500
Antibody	Goat polyclonal anti-Rabbit IgG, Flexa Fluor 555	Thermo-Fisher Scientific	A-21428	1:500
Chemical compound, drug	Alexa Fluor 488-conjugated Phalloidin	Invitrogen	A12379	1:50
Chemical compound, drug	Dox Diet, Grain-Based Doxycycline	Bio-Serv	S3888	
Chemical compound, drug	EdU (5-ethynyl-2'-deoxyuridine)	Thermo-Fisher Scientific	E10187	
Commercial assay or kit	Click-iT EdU Alexa Fluor 488 kit	Thermo-Fisher Scientific	C10337	

Mice

Mice were group housed with littermates, in breeding pairs, or in a breeding harem (2 females to

1 male), with food and water provided ad libitum.

For timed-pregnancy experiments, embryonic day 0.5 (E0.5) was assigned as noon of the day

the vaginal plug was found. For postnatal experiments, postnatal day 0 (P0) was determined as

the day of birth.

Mice were of mixed sexes and maintained on a mixed C57BL/6J x 129X1/SvJ genetic

background. All mice were backcrossed at least three generations onto this background. The

following mouse lines were used:

- $Fgf20^{Cre}$ ($Fgf20^-$): knockin allele containing a sequence encoding a GFP-Cre fusion

protein replacing exon 1 of $Fgf20$, resulting in a null mutation [18].

- $Fgf20^{\beta gal}$: knockin allele containing a sequence encoding β-galactosidase (βgal)

replacing exon 1 of $Fgf20$, resulting in a null mutation [21].

- $Foxg1^{Cre}$: knockin allele containing a sequence encoding Cre fused in-frame downstream

of the first 13 codons, resulting in a null mutation [22].

- $Fgfr1^{flox}$: allele containing loxP sequences flanking exons 8 through 15 of $Fgfr1$. Upon

Cre-mediated recombination, produces a null mutation [51]

- $Fgfr1^-$: null allele generated by combining $Fgfr1^{flox}$ with $Sox2^{Cre}$ [52] to delete $Fgfr1$ from

the epiblast.

- $ROSA^{rtTA}$: knockin allele containing a loxP-Stop-loxP sequence followed by a sequence

encoding rtTA-IRES-eGFP, targeted to the ubiquitously expressed $ROSA26$ locus. Upon

Cre-mediated recombination, reverse tetracycline transactivator (rtTA) and eGFP are

expressed [25].

• TRE-Fgf9-IRES-eGfp: transgene containing seven tetracycline-inducible regulatory elements

driving the expression of FGF9-IRES-eGFP [26].

• $Sox2^{Ysb}$: Inner ear specific $Sox2$ hypomorphic allele resulting from a random insertion of

a transgene in chromosome 3, likely interfering with tissue-specific $Sox2$ regulatory

elements [11].

• $Sox2^{-}$: null allele generated by combining $Sox2^{flox}$ [53] with $Sox2^{Cre}$ to delete $Sox2$ from

the epiblast.

All studies performed were in accordance with the Institutional Animal Care and Use Committee

at Washington University in St. Louis (protocol #20160113) and University of Nebraska Medical

Center (protocol #16-004-02 and 16-005-02).

Doxycycline induction

Pregnant dams were starved overnight the night before initiation of Dox induction and fed Dox

Diet, Grain-Based Doxycycline, 200 mg/kg (S3888, Bio-Serv, Flemington, NJ) ad libitum starting

at noon on the start date of Dox induction. On the stop date of Dox induction, Dox Diet was

replaced with regular mouse chow at noon.

Sample preparation and sectioning

For whole mount cochleae, inner ears were dissected out of P0 pups and fixed in 4% PFA in

PBS overnight at 4°C with gentle agitation. Samples were then washed x3 in PBS. Cochleae

were dissected away from the vestibule, otic capsule, and periotic mesenchyme with Dumont

#55 Forceps (RS-5010, Roboz, Gaithersburg, MD). The roof of the cochlear duct was opened

up by dissecting away the stria vascularis and Reissner's membrane; tectorial membrane was

removed to expose hair and supporting cells.

For sectioning, heads from E14.5 embryos were fixed in 4% PFA in PBS overnight at 4°C with

gentle agitation. Samples were then washed x3 in PBS and cryoprotected in 15% sucrose in

PBS overnight and then in 30% sucrose in PBS overnight. Samples were embedded in Tissue-

Tek O.C.T. compound (4583, VWR International, Radnor, PA) and frozen on dry ice. Serial

horizontal sections through base of the head were cut at 12 μm with a cryostat, dried at room

temperature, and stored at -80°C until use.

RNA in situ hybridization

Probe preparation: mouse cDNA plasmids containing the following inserts were used to make

RNA in situ probes, and were cut and transcribed with the indicated restriction enzyme (New

England Biolabs, Ipswich, MA) and RNA polymerase (New England Biolabs, Ipswich, MA):

Fgfr1 transmembrane domain (325 bp, HincII, T7, gift of K. Peters), *Fgf20* (653 bp, NcoI, Sp6),

Sox2 (750 bp, AccI, T3, gift of A. Kiernan), *Etv4* (~2300 bp, ApaI, Sp6, gift of G. Martin), *Etv5*

(~4000 bp, HindIII, T3, gift of G. Martin), *Hey1* (343 bp, EcoRI, T3, gift of S. Rentschler), *Hey2*

(819 bp, EcoRI, T7, gift of S. Rentschler). Restriction digest and *in vitro* transcription were done

according to manufacturer's instructions, with DIG RNA Labeling Mix (11277073910, Sigma-

Aldrich, St. Louis, MO). After treatment with RNase-free DNase I (04716728001, Sigma-Aldrich,

St. Louis, MO) for 15 min at 37°C, probes were hydrolyzed in hydrolysis buffer (40 mM

$NaHCO_3$, 60 mM Na_2CO_3) at 60°C for up to 30 min, depending on probe size.

Frozen section in situ hybridization: frozen slides were warmed for 20 min at room temperature

and then 5 min at 50°C on a slide warmer. Sections were fixed in 4% PFA in PBS for 20 min at

room temperature, washed x2 in PBS and treated with pre-warmed 10 µg/ml Proteinase K

(03115828001, Sigma-Aldrich, St. Louis, MO) in PBS for 7 min at 37°C. Sections were then

fixed in 4% PFA in PBS for 15 min at room temperature, washed x2 in PBS, acetylated in 0.25%

acetic anhydride in 0.1M Triethanolamine, pH 8.0, for 10 min, and washed again in PBS.

Sections were then placed in pre-warmed hybridization buffer (50% formamide, 5x SSC buffer,

5 mM EDTA, 50 µg/ml yeast tRNA) for 3 h at 60°C in humidified chamber for prehybridization.

Sections were then hybridized in 10 µg/ml probe/hybridization buffer overnight (12-16 h) at

60°C. The next day, sections were washed in 1x SSC for 10 min at 60°C, followed by 1.5x SSC

for 10 min at 60°C, 2x SSC for 20 min at 37°C x2, and 0.2x SSC for 30 min at 60°C x2. Sections

were then washed in KTBT (0.1 M Tris, pH 7.5, 0.15 M NaCl, 5 mM KCl, 0.1% Triton X-100) at

room temperature and blocked in KTBT + 20% sheep serum + 2% Blocking Reagent

(11096176001, Sigma-Aldrich, St. Louis, MO) for 4 h. Blocking Reagent was dissolved in 100

mM Maleic acid, 150 mM NaCl, pH 7.5. Sections were then incubated in sheep anti-

Digoxigenin-AP, Fab fragments (1:1000, 11093274910, Sigma-Aldrich, St. Louis, MO) in KTBT

+ 20% sheep serum + 2% Blocking Reagent overnight at 4°C. Sections were then washed x3 in

KTBT for 30 min at room temperature, and then washed x2 in NTMT (0.1 M Tris, pH 9.5, 0.1 M

NaCl, 50 mM MgCl$_2$, 0.1% Tween 20) for 15 min. Sections were next incubated in NTMT +

1:200 NBT/BCIP Stock Solution (11681451001, Sigma-Aldrich, St. Louis, MO) in the dark at

room temperature until color appeared. Sections were then washed in PBS, post-fixed in 4%

PFA in PBS for 15 min and washed x2 in PBS. Finally, sections were dehydrated in 30% and

then 70% methanol, 5 min each, followed by 100% methanol for 15 min. Sections were then

rehydrated in 70% and 30% methanol and then PBS, 5 min each, and mounted in 95% glycerol.

Immunofluorescence

Whole mount: cochleae were incubated in PBS + 0.5% Tween-20 (PBSTw) for 1 h to

permeabilize. Cochleae were then blocked using PBSTw + 5% donkey serum for 1 h and then

incubated in PBSTw + 1% donkey serum with the primary antibody overnight at 4°C. Cochleae

were then washed x3 in PBS and incubated in PBS + 1% Tween-20 with the secondary

antibody. After wash in PBS x3, cochleae were mounted in 95% glycerol with the sensory

epithelium facing up.

Frozen slides were warmed for 30 min at room temperature and washed in PBS before

incubating in PBS + 0.5% Triton X-100 (PBST) for 1 h to permeabilize the tissue. Sections were

then blocked using in PBST + 5% donkey serum for 1 h and then incubated in PBST + 1%

donkey serum with the primary antibody overnight at 4°C in a humidified chamber. Sections

were then washed x3 in PBS and incubated in PBS + 1% Triton X-100 with the secondary

antibody. After wash in PBS x3, slides were mounted in VectaShield antifade mounting medium

with DAPI (H-1200, Vector Labs, Burlingame, CA).

Cell proliferation assay

EdU (E10187, Thermo-Fisher Scientific, Waltham, MA) was injected i.p. into pregnant dams at

100 µg per gram body weight. Embryos were harvested at 1 h after injection. EdU was detected

using the Click-iT EdU Alexa Fluor 488 kit (C10337, Thermo-Fisher Scientific, Waltham, MA)

according to manufacturer's instructions.

Imaging

Brightfield microscopy was done using a Hamamatsu NanoZoomer slide scanning system with

a 20x objective. Images were processed with the NanoZoomer Digital Pathology (NDP.view2)

software.

Fluorescent microscopy was done using a Zeiss LSM 700 confocal or Zeiss Axio Imager Z1 with

Apotome 2, with z-stack step-size determined based on objective lens type (10x or 20x), as

recommended by the ZEN software (around 1 µm). Fluorescent images shown are maximum

projections. Low magnification fluorescent images shown of the whole cochlear duct required

stitching together, by hand, several images. Images were processed with ImageJ

(imagej.nih.gov).

Quantification

Measurements and cell quantification (using the Cell Counter plugin by Kurt De Vos) were done

using ImageJ. Total cochlear duct length was defined as the length from the very base of the

cochlea to the very tip of the apex, along the tunnel of Corti. Hair cells were identified via

Phalloidin, which binds to F-actin [54]. Inner pillar cells were labeled via p75NTR [55], and

supporting cells (SCs, including pillar cells and Deiters' cells) were labeled with a combination of

Prox1 [56] and Sox2 [10]. Inner hair cells (IHCs) were differentiated from outer hair cells (OHCs)

based on their neural/abneural location, respectively, relative to p75NTR-expressing inner pillar

cells. For total cell counts, IHCs, OHCs, and SCs were counted along the entire length of the

cochlea. Total cell counts were also normalized to cochlear length and presented as cell count

per 100 µm of cochlea (e.g. IHCs/100 µm). For cell quantification at the basal, middle, and

apical turns of the cochlea, the cochlear duct was evenly divided into thirds, and total IHCs,

OHCs, and SCs were quantified for each third and normalized to length. For the Fgf20-rescue

experiments in Fig 3, IHCs, OHCs, and SCs from at least 300 μm regions of the basal (10%),

middle (40%), and apical (70%) turns of the cochleae were counted and normalized to 100 μm

along the length of the cochlear duct.

In Sox2$^{Ysb/-}$ cochleae, p75NTR expression was mostly absent, resulting in sensory islands

without p75NTR-expressing inner pillar cells. In these cochleae, HCs not associated with inner

pillar cells were presumed to be IHCs during quantification. When a curved line was drawn

connecting the p75NTR islands along the organ of Corti, these presumed IHCs were always

medial (neural) to that line.

Statistical analysis and plotting

All figures were made in Canvas X (ACD systems). Data analysis was performed using the

Python programming language (python.org) in Jupyter Notebook (jupyter.org) with the following

libraries: Pandas (pandas.pydata.org), NumPy (numpy.org) and SciPy (scipy.org). Plotting was

done using the Matplotlib library (matplotlib.org). Statistics (t-test, one-way ANOVA, two-way

ANOVA, and Fisher's exact test) were performed using the SciPy module Stats; Tukey's HSD

was performed using the Statsmodels package (statsmodels.org). All comparisons of two

means were performed using two-tailed, unpaired Student's t-test. For comparisons of more

than two means, one-way ANOVA was used, except in Fig 8 and S7 Fig, where two-way

ANOVA was used, with the factors being *Fgf20* (levels: *Fgf20$^{+/+}$*, *Fgf20$^{+/-}$*, *Fgf20$^{-/-}$*) and *Sox2*

(levels: *Sox2$^{+/+}$*, *Sox2$^{Ysb/+}$*, *Sox2$^{Ysb/Ysb}$*) gene dosage. For significant ANOVA results at α = 0.05,

Tukey's HSD was performed for post-hoc pair-wise analysis. In all cases, $p < 0.05$ was

considered statistically significant. All statistical details can be found in the figures and figure

legends. In all cases, each sample (each data point in graphs) represents one animal. Based on

similar previous studies, a sample size of 3-5 was determined to be appropriate. Error bars

represent mean ± standard deviation (SD). For qualitative comparisons (comparing expression

via immunofluorescence or RNA in situ hybridization), at least three samples were examined

per genotype. All images shown are representative.

Evaluation of onset of Myo6-expressing cells (Figs 10B and 10C): 3 or 4 serial sections through

the entire cochleae were immunostained for Myo6 and evaluated, blinded to genotype, for the

presence of Myo6-expressing cells. E14.5 embryos were further stage-matched based on

interdigital webbing of the hindlimb (at E14.5, roughly half of the hindlimb interdigital webbing is

still present). Of the 34 embryos at E14.5, 3 were removed from analysis due to lack of or

minimal hindlimb interdigital webbing (too old relative to the other embryos).

1003

1004   Acknowledgements

1005   We would like to thank Drew Hagan, Kel Vin Woo, and Yongjun Yin for their critical reading of

1006   the manuscript.

1007

1008   References

1009

1.   Basch ML, Brown RM, Jen H-I, Groves AK. Where hearing starts: the development of the
mammalian cochlea. J Anat. 2016 Feb;228(2):233–54.

2.   Fekete DM, Muthukumar S, Karagogeos D. Hair Cells and Supporting Cells Share a
Common Progenitor in the Avian Inner Ear. J Neurosci. 1998 Oct 1;18(19):7811–21.

3.   Chen P, Segil N. p27(Kip1) links cell proliferation to morphogenesis in the developing
organ of Corti. Development. 1999 Apr;126(8):1581–90.

4.   Lee Y-S, Liu F, Segil N. A morphogenetic wave of p27Kip1 transcription directs cell cycle
exit during organ of Corti development. Development. 2006 Aug 1;133(15):2817–26.

5.   Bok J, Zenczak C, Hwang CH, Wu DK. Auditory ganglion source of Sonic hedgehog
regulates timing of cell cycle exit and differentiation of mammalian cochlear hair cells. Proc
Natl Acad Sci U S A. 2013 Aug 20;110(34):13869–74.

6.   Jahan I, Pan N, Kersigo J, Fritzsch B. Neurod1 Suppresses Hair Cell Differentiation in Ear
Ganglia and Regulates Hair Cell Subtype Development in the Cochlea. Riley B, editor.
PLOS ONE. 2010 Jul 22;5(7):e11661.

7.   Matei V, Pauley S, Kaing S, Rowitch D, Beisel K w., Morris K, et al. Smaller inner ear
sensory epithelia in Neurog1 null mice are related to earlier hair cell cycle exit. Dev Dyn.
2005 Nov 1;234(3):633–50.

8.   Tateya T, Imayoshi I, Tateya I, Hamaguchi K, Torii H, Ito J, et al. Hedgehog signaling
regulates prosensory cell properties during the basal-to-apical wave of hair cell
differentiation in the mammalian cochlea. Development. 2013 Sep 15;140(18):3848–57.

9.   Gu R, Brown RM, Hsu C-W, Cai T, Crowder AL, Piazza VG, et al. Lineage tracing of Sox2-
expressing progenitor cells in the mouse inner ear reveals a broad contribution to non-
sensory tissues and insights into the origin of the organ of Corti. Dev Biol. 2016
01;414(1):72–84.

10.   Mak ACY, Szeto IYY, Fritzsch B, Cheah KSE. Differential and overlapping expression
pattern of SOX2 and SOX9 in inner ear development. Gene Expr Patterns. 2009 Sep
1;9(6):444–53.

11.   Dong S, Leung KKH, Pelling AL, Lee PYT, Tang ASP, Heng HHQ, et al. Circling,
Deafness, and Yellow Coat Displayed by Yellow Submarine (Ysb) and Light Coat and
Circling (Lcc) Mice with Mutations on Chromosome 3. Genomics. 2002 Jun;79(6):777–84.

12.   Kiernan AE, Pelling AL, Leung KKH, Tang ASP, Bell DM, Tease C, et al. Sox2 is required
for sensory organ development in the mammalian inner ear. Nature. 2005 Apr
21;434(7036):1031–5.

13.   Dabdoub A, Puligilla C, Jones JM, Fritzsch B, Cheah KSE, Pevny LH, et al. Sox2 signaling
in prosensory domain specification and subsequent hair cell differentiation in the
developing cochlea. Proc Natl Acad Sci U S A. 2008 Nov 25;105(47):18396–401.

14.   Pan W, Jin Y, Chen J, Rottier RJ, Steel KP, Kiernan AE. Ectopic Expression of Activated
Notch or SOX2 Reveals Similar and Unique Roles in the Development of the Sensory Cell
Progenitors in the Mammalian Inner Ear. J Neurosci. 2013 Oct 9;33(41):16146–57.

15.   Puligilla C, Kelley MW. Dual role for Sox2 in specification of sensory competence and
regulation of Atoh1 function. Dev Neurobiol. 2016;77(1):3–13.

16.   Ebeid M, Huh S-H. FGF signaling: diverse roles during cochlear development. BMB Rep.
2017 Oct 31;50(10):487–95.

17.   Hayashi T, Ray CA, Bermingham-McDonogh O. Fgf20 Is Required for Sensory Epithelial
Specification in the Developing Cochlea. J Neurosci. 2008 Jun 4;28(23):5991–9.

18.   Huh S-H, Warchol ME, Ornitz DM. Cochlear progenitor number is controlled through
1056   mesenchymal FGF receptor signaling. eLife. 2015 Apr 27;4:e05921.

1057   19.   Ono K, Kita T, Sato S, O'Neill P, Mak S-S, Paschaki M, et al. FGFR1-Frs2/3 Signalling
1058   Maintains Sensory Progenitors during Inner Ear Hair Cell Formation. Cheah KSE, editor.
1059   PLOS Genet. 2014 Jan 23;10(1):e1004118.

1060   20.   Pirvola U, Ylikoski J, Trokovic R, Hébert JM, McConnell SK, Partanen J. FGFR1 is
1061   required for the development of the auditory sensory epithelium. Neuron. 2002 Aug
1062   15;35(4):671–80.

1063   21.   Huh S-H, Jones J, Warchol ME, Ornitz DM. Differentiation of the Lateral Compartment of
1064   the Cochlea Requires a Temporally Restricted FGF20 Signal. Groves A, editor. PLOS Biol.
1065   2012 Jan 3;10(1):e1001231.

1066   22.   Hébert JM, McConnell SK. Targeting of cre to the Foxg1 (BF-1) Locus Mediates loxP
1067   Recombination in the Telencephalon and Other Developing Head Structures. Dev Biol.
1068   2000 Jun 15;222(2):296–306.

1069   23.   Ornitz DM, Itoh N. The Fibroblast Growth Factor signaling pathway. Wiley Interdiscip Rev
1070   Dev Biol. 2015 May 1;4(3):215–66.

1071   24.   Zhang X, Ibrahimi OA, Olsen SK, Umemori H, Mohammadi M, Ornitz DM. Receptor
1072   Specificity of the Fibroblast Growth Factor Family: THE COMPLETE MAMMALIAN FGF
1073   FAMILY. J Biol Chem. 2006 Jun 9;281(23):15694–700.

1074   25.   Belteki G, Haigh J, Kabacs N, Haigh K, Sison K, Costantini F, et al. Conditional and
1075   inducible transgene expression in mice through the combinatorial use of Cre-mediated
1076   recombination and tetracycline induction. Nucleic Acids Res. 2005 Jan 1;33(5):e51–e51.

1077   26.   White AC, Xu J, Yin Y, Smith C, Schmid G, Ornitz DM. FGF9 and SHH signaling
1078   coordinate lung growth and development through regulation of distinct mesenchymal
1079   domains. Development. 2006 Apr 15;133(8):1507–17.

1080   27.   Scholzen T, Gerdes J. The Ki-67 protein: From the known and the unknown. J Cell
1081   Physiol. 2000 Mar 1;182(3):311–22.

1082   28.   Kelley MW. Cellular commitment and differentiation in the organ of Corti. Int J Dev Biol.
1083   2007;51(6–7):571–83.

1084   29.   Benito-Gonzalez A, Doetzlhofer A. Hey1 and Hey2 Control the Spatial and Temporal
1085   Pattern of Mammalian Auditory Hair Cell Differentiation Downstream of Hedgehog
1086   Signaling. J Neurosci. 2014 Sep 17;34(38):12865–76.

1087   30.   Bermingham NA, Hassan BA, Price SD, Vollrath MA, Ben-Arie N, Eatock RA, et al. Math1:
1088   an essential gene for the generation of inner ear hair cells. Science. 1999 Jun
1089   11;284(5421):1837–41.

1090   31.   Cai T, Seymour ML, Zhang H, Pereira FA, Groves AK. Conditional Deletion of Atoh1
1091   Reveals Distinct Critical Periods for Survival and Function of Hair Cells in the Organ of
1092   Corti. J Neurosci. 2013 Jun 12;33(24):10110–22.

32.   Woods C, Montcouquiol M, Kelley MW. Math1 regulates development of the sensory
epithelium in the mammalian cochlea. Nat Neurosci. 2004 Dec;7(12):1310–8.

33.   Munnamalai V, Hayashi T, Bermingham-McDonogh O. Notch Prosensory Effects in the
Mammalian Cochlea Are Partially Mediated by Fgf20. J Neurosci. 2012 Sep
12;32(37):12876–84.

34.   Pauley S, Lai E, Fritzsch B. Foxg1 is required for morphogenesis and histogenesis of the
mammalian inner ear. Dev Dyn. 2006 Sep;235(9):2470–82.

35.   Brown AS, Epstein DJ. Otic ablation of smoothened reveals direct and indirect
requirements for Hedgehog signaling in inner ear development. Development. 2011 Sep
15;138(18):3967–76.

36.   Kanzaki S, Beyer LA, Swiderski DL, Izumikawa M, Stöver T, Kawamoto K, et al. p27Kip1
deficiency causes organ of Corti pathology and hearing loss. Hear Res. 2006 Apr
1;214(1):28–36.

37.   Ahmed M, Wong EYM, Sun J, Xu J, Wang F, Xu P-X. Eya1-Six1 Interaction Is Sufficient to
Induce Hair Cell Fate in the Cochlea by Activating Atoh1 Expression in Cooperation with
Sox2. Dev Cell. 2012 Feb;22(2):377–90.

38.   Kempfle JS, Turban JL, Edge ASB. Sox2 in the differentiation of cochlear progenitor cells.
Sci Rep. 2016;6:23293.

39.   Bosman EA, Quint E, Fuchs H, Hrabé de Angelis M, Steel KP. Catweasel mice: a novel
role for Six1 in sensory patch development and a model for branchio-oto-renal syndrome.
Dev Biol. 2009 Apr 15;328(2):285–96.

40.   Brooker R. Notch ligands with contrasting functions: Jagged1 and Delta1 in the mouse
inner ear. Development. 2006 Apr 1;133(7):1277–86.

41.   Kiernan AE, Ahituv N, Fuchs H, Balling R, Avraham KB, Steel KP, et al. The Notch ligand
Jagged1 is required for inner ear sensory development. Proc Natl Acad Sci U S A.
2001;98(7):3873–3878.

42.   Kiernan AE, Xu J, Gridley T. The Notch ligand JAG1 is required for sensory progenitor
development in the mammalian inner ear. PLOS Genet. 2006;2(1):e4.

43.   Kiernan AE, Li R, Hawes NL, Churchill GA, Gridley T. Genetic background modifies inner
ear and eye phenotypes of jag1 heterozygous mice. Genetics. 2007 Sep;177(1):307–11.

44.   Moayedi Y, Basch ML, Pacheco NL, Gao SS, Wang R, Harrison W, et al. The Candidate
Splicing Factor Sfswap Regulates Growth and Patterning of Inner Ear Sensory Organs.
Wu DK, editor. PLOS Genet. 2014 Jan 2;10(1):e1004055.

45.   Tsai H, Hardisty RE, Rhodes C, Kiernan AE, Roby P, Tymowska-Lalanne Z, et al. The
mouse slalom mutant demonstrates a role for Jagged1 in neuroepithelial patterning in the
organ of Corti. Hum Mol Genet. 2001 Mar 1;10(5):507–12.

46. Yamamoto N, Chang W, Kelley MW. Rbpj regulates development of prosensory cells in the
mammalian inner ear. Dev Biol. 2011 May;353(2):367–79.

47. Basch ML, Ohyama T, Segil N, Groves AK. Canonical Notch Signaling Is Not Necessary
for Prosensory Induction in the Mouse Cochlea: Insights from a Conditional Mutant of
RBPj. J Neurosci. 2011 Jun 1;31(22):8046–58.

48. Kopecky BJ, Jahan I, Fritzsch B. Correct Timing of Proliferation and Differentiation is
Necessary for Normal Inner Ear Development and Auditory Hair Cell Viability: Mycs' Role
in Ear Development. Dev Dyn. 2013 Feb;242(2):132–47.

49. Chen P, Johnson JE, Zoghbi HY, Segil N. The role of Math1 in inner ear development:
Uncoupling the establishment of the sensory primordium from hair cell fate determination.
Development. 2002 May;129(10):2495–505.

50. Haque K, Pandey AK, Zheng H-W, Riazuddin S, Sha S-H, Puligilla C. MEKK4 Signaling
Regulates Sensory Cell Development and Function in the Mouse Inner Ear. J Neurosci.
2016 Jan 27;36(4):1347–61.

51. Trokovic N, Trokovic R, Mai P, Partanen J. Fgfr1 regulates patterning of the pharyngeal
region. Genes Dev. 2003 Jan 1;17(1):141–53.

52. Hayashi S, Lewis P, Pevny L, McMahon AP. Efficient gene modulation in mouse epiblast
using a Sox2Cre transgenic mouse strain. Mech Dev. 2002 Dec;119 Suppl 1:S97–101.

53. Shaham O, Smith AN, Robinson ML, Taketo MM, Lang RA, Ashery-Padan R. Pax6 is
essential for lens fiber cell differentiation. Development. 2009 Aug 1;136(15):2567–78.

54. Avinash GB, Nuttall AL, Raphael Y. 3-D analysis of F-actin in stereocilia of cochlear hair
cells after loud noise exposure. Hear Res. 1993 May 1;67(1):139–46.

55. Mueller KL, Jacques BE, Kelley MW. Fibroblast Growth Factor Signaling Regulates Pillar
Cell Development in the Organ of Corti. J Neurosci. 2002 Nov 1;22(21):9368–77.

56. Bermingham-McDonogh O, Oesterle EC, Stone JS, Hume CR, Huynh HM, Hayashi T.
Expression of Prox1 during mouse cochlear development. J Comp Neurol. 2006 May
10;496(2):172–86.

1156

Figure titles and legends

1158

Fig 1. The Fgf20-KO cochlear phenotype is less severe than the Fgfr1-CKO

phenotype

(A, B) Whole mount cochlea from P0 $Fgf20^{+/-}$, $Fgf20^{-/-}$, $Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and

$Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ mice showing (A) inner and outer hair cells (IHC and OHC,

phalloidin, green) separated by inner pillar cells (p75NTR, red) and (B) supporting cells

(Prox1 and Sox2, green/yellow). Magnifications show the basal, middle, and apical turns

of the cochlea. Scale bar, 100 µm (magnifications), 1 mm (whole).

(C-F) Quantification of (C) cochlear duct length, (D) total IHCs and IHCs per 100 µm of the

cochlear duct, (E) total OHCs and OHCs per 100 µm, and (F) total supporting cells

(SCs) and SCs per 100 µm at P0. $Fgf20^{+/-}$ and $Fgf20^{-/-}$ cochleae results were analyzed

by unpaired Student's t test; $Fgfr1^{flox/+}$, $Fgfr1^{flox/-}$, $Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and

$Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ cochleae results were analyzed by one-way ANOVA. P values

shown are from the t test and ANOVA. * indicates $p < 0.05$ from Student's t test or

Tukey's HSD (ANOVA post-hoc); n.s., not significant. Error bars, mean ± SD.

(G) Schematic showing the positions of basal, middle, and apical turns along the cochlear

duct.

See also S1 Fig.

1176

Fig 2. FGFR1 but not FGF20 regulates *Sox2* expression

(A) Sections through the middle turn of E14.5 cochlear ducts from $Fgf20^{+/-}$, $Fgf20^{-/-}$,

$Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and $Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ mice. RNA in situ hybridization (top) and

immunofluorescence for *Sox2* (red, bottom), which is expressed in the prosensory

domain at this stage. Refer to schematic in (C).

(B)　Immunofluorescence for CDKN1B (green) in sections through the basal, middle, and

apical turns of E14.5 $Fgf20^{+/-}$, $Fgf20^{-/-}$, $Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and $Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$

cochleae.

(C)　Schematic of a cross section through the middle turn of the E14.5 cochlear duct,

showing the location of the prosensory domain (PD). Neural indicates the side of the

duct towards the spiral ganglion cells; abneural indicates away.

(D)　A model of genetic pathways during organ of Corti development. Ligand X/FGFR1

signaling regulates $Sox2$ expression during prosensory specification; FGF20/FGFR1

signaling regulates differentiation. Ligand X may include FGF20, along with another

functionally redundant ligand.

DAPI, nuclei (blue). Scale bar, 100 μm. See also S2 Fig.

1193

Fig 3. Genetic rescue of the Fgf20-KO phenotype suggests that FGF20 is required

for differentiation

(A, B)　Whole mount cochlea from P0 $Fgf20^{+/-}$;$ROSA^{rtTA}$ (Fgf20-het), $Fgf20^{+/-}$;$ROSA^{rtTA}$; TRE-

Fgf9-IRES-eGfp (Fgf9-OA), $Fgf20^{-/-}$;$ROSA^{rtTA}$ (Fgf20-null), and $Fgf20^{-/-}$;$ROSA^{rtTA}$; TRE-

Fgf9-IRES-eGfp (Fgf20-rescue) mice showing (A) inner and outer hair cells (phalloidin,

green) separated by inner pillar cells (p75NTR, red) and (B) supporting cells (Prox1 and

Sox2, green/yellow). Images from basal, middle, and apical turns of the cochlea are

shown. Fgf20-rescue cochleae from four different doxycycline chow (Dox) regimens are

shown (E13.5-E15.5, E13.5, E14.5, and E15.5). Fgf20-het, Fgf9-OA, and Fgf20-null

cochleae shown are from the E13.5-E15.5 Dox regimen. Scale bar, 100 μm.

(C)　Quantification of outer hair cells (OHC) and supporting cells (SC) from P0 Fgf9-OA (Dox

regimen: E13.5-E15.5), Fgf20-null (Dox regimen: E13.5-E15.5), and Fgf20-rescue (all

four Dox regimens) cochleae, presented as a percentage of the number of cells in

Fgf20-het cochleae from the same Dox regimen. * indicates $p < 0.05$ compared to Fgf20-

null cochleae from the same Dox regimen; ^ indicates $p < 0.05$ compared to Fgf20-het

cochleae from the same Dox regimen; Tukey's HSD (one-way ANOVA post-hoc).

See also S3 Fig.

1211

Fig 4. Decrease in Sox2 expression results in similar phenotypes as disruptions

to FGFR1 signaling

(A, B) Whole mount cochlea from P0 mice from the *Sox2* allelic series (in order of highest to

lowest levels of *Sox2* expression: $Sox2^{+/+}$, $Sox2^{Ysb/+}$, $Sox2^{Ysb/Ysb}$, and $Sox2^{Ysb/-}$) showing

(A) inner and outer hair cells (phalloidin, green) separated by inner pillar cells (p75NTR,

red) and (B) supporting cells (Prox1 and Sox2, green/yellow). Magnifications show the

basal, middle, and apical turns of the cochlea. Arrowheads indicate ectopic inner hair

cells. Scale bar, 100 µm (magnifications), 1 mm (whole).

(C-F)   Quantification of (C) cochlear duct length, (D) total inner hair cells (IHCs) and IHCs per

100 µm of the cochlear duct, (E) total outer hair cells (OHCs) and OHCs per 100 µm,

and (F) total supporting cells (SCs) and SCs per 100 µm at P0. P values shown are from

one-way ANOVA. * indicates $p < 0.05$ from Tukey's HSD (ANOVA post-hoc). Error bars,

mean ± SD.

See also S4 Fig.

1226

Fig 5. Decrease in levels of Sox2 expression delays prosensory specification

(A)   Serial sections (1-6) through the duct of E14.5 and E15.5 $Sox2^{Ysb/+}$ and $Sox2^{Ysb/-}$

cochleae. Immunofluorescence for Ki67 (red) and DAPI (nuclei, cyan). Cochlear

epithelium is outlined. Bracket indicates prosensory domain. * indicates shift of

prosensory nuclei away from the luminal surface of the epithelium. N, neural side. Scale

bar, 100 µm. Whole mount cochlear duct schematics show relative positions of the serial

sections and progression of cell cycle exit (green arrow).

(B)   A model of organ of Corti development showing embryonic staging (x-axis) and location

along the cochlear duct (basal, middle, and apical turns, y-axis). Development occurs in

two stages: unspecified progenitors (tan shading) undergo specification and cell cycle

exit to become prosensory cells (green shading), which then differentiate into hair cells

and supporting cells (HCs & SCs; red shading). In wildtype cochleae, cell cycle exit

(indicating completion of specification) begins at the apex of the cochlea and proceeds

basally. Afterwards, differentiation initiates at the mid-base of the cochlea and proceeds

basally and apically. Temporal buffer (green shading) refers to the time between cell

cycle exit and initiation of differentiation. In *Sox2* hypomorph cochleae, specification and

cell cycle exit are delayed, resulting in failure to complete specification before initiation of

differentiation towards the basal end of the cochlea (crosshatch pattern).

See also S5 Fig.

1246

Fig 6. *Sox2* is upstream of *Fgf20*

(A-C)   Sections through the middle turn of E14.5 cochleae.

(A)   RNA in situ hybridization for *Etv4* and *Etv5* in *Sox2*$^{Ysb/+}$ and *Sox2*$^{Ysb/-}$ cochleae. The two

brackets indicate *Etv4/5* expression in the outer sulcus (OS, left) and prosensory domain

(PD, right). Refer to schematic at the bottom right of the figure.

(B)   RNA in situ hybridization for *Fgfr1* and *Fgf20* in *Sox2*$^{Ysb/+}$ and *Sox2*$^{Ysb/-}$ cochleae. Bracket

indicates *Fgfr1*/*Fgf20* expression in the prosensory domain.

(C)   RNA in situ hybridization for *Fgf20* in *Foxg1*$^{Cre/+}$;*Fgfr1*$^{flox/+}$ and *Foxg1*$^{Cre/+}$;*Fgfr1*$^{flox/-}$

cochleae. Bracket indicates *Fgf20* expression in the prosensory domain.

(D)   Serial sections (1-6) through the duct of E14.5 *Fgf20*$^{+/-}$ and *Fgf20*$^{-/-}$ cochleae.

Immunofluorescence for Ki67 (red) and DAPI (nuclei, cyan). Cochlear epithelium is

outlined. Bracket indicates prosensory domain. * indicates shift of prosensory nuclei

away from the luminal surface of the epithelium. N, neural side. Scale bar, 100 μm.

Whole mount cochlear duct schematics show relative positions of the serial sections and

progression of cell cycle exit (green arrow).

OS, outer sulcus; PD, prosensory domain; KO, Kölliker's organ. Scale bar, 100 μm. See also S6

Fig.

1264

Fig 7. *Sox2* and *Fgf20* interact during cochlea development

(A, B) Whole mount cochlea from P0 *Fgf20*$^{+/-}$;*Sox2*$^{Ysb/+}$, *Fgf20*$^{+/-}$;*Sox2*$^{Ysb/Ysb}$, *Fgf20*$^{-/-}$;*Sox2*$^{Ysb/+}$,

and *Fgf20*$^{-/-}$;*Sox2*$^{Ysb/Ysb}$ mice showing (A) inner and outer hair cells (phalloidin, green)

separated by inner pillar cells (p75NTR, red) and (B) supporting cells (Prox1 and Sox2,

green/yellow). Magnifications show the basal, middle, and apical turns of the cochlea.

Arrowheads indicate ectopic inner hair cells. Scale bar, 100 μm (magnifications), 1 mm

(whole).

1272

Fig 8. *Sox2* and *Fgf20* interact during cochlea development—quantitative analysis

(A)   P values from two-way ANOVA analyzing the quantification results in (B-E). The two

factors analyzed are *Fgf20* (*Fgf20*$^{+/+}$, *Fgf20*$^{+/-}$, *Fgf20*$^{-/-}$) and *Sox2* (*Sox2*$^{+/+}$, *Sox2*$^{Ysb/+}$,

Sox2$^{Ysb/Ysb}$) gene dosage. A p value < 0.05 (yellow highlight) for *Fgf20* or *Sox2* indicates

that the particular factor (independent variable) has a statistically significant effect on the

measurement (dependent variable). Whereas a p value < 0.05 (orange highlight) for

Interaction indicates a statistically significant interaction between the effects of the two

factors on the measurement.

(B-E) Quantification of (B) cochlear duct length, (C) total inner hair cells (IHCs) and IHCs per

100 μm of the cochlear duct, (D) total outer hair cells (OHCs) and OHCs per 100 μm,

and (E) total supporting cells (SCs) and SCs per 100 µm at P0 in $Fgf20^{+/+}$;$Sox2^{+/+}$,

$Fgf20^{+/+}$;$Sox2^{Ysb/+}$, $Fgf20^{+/-}$;$Sox2^{+/+}$, $Fgf20^{+/-}$;$Sox2^{Ysb/+}$, $Fgf20^{+/+}$;$Sox2^{Ysb/Ysb}$,

$Fgf20^{+/-}$;$Sox2^{Ysb/Ysb}$, $Fgf20^{-/-}$;$Sox2^{+/+}$, $Fgf20^{-/-}$;$Sox2^{Ysb/+}$, and $Fgf20^{-/-}$;$Sox2^{Ysb/Ysb}$ cochleae.

Error bars, mean ± SD.

(F)   Results from post-hoc Tukey's HSD analyzing the quantification results in (B-E). Letters

(L, I, J, O, P, S, T; representing each measurement in panels B-E) indicate a statistically

significant decrease ($p < 0.05$) when comparing the row genotype against the column

genotype. L, cochlear length; I, total IHCs; J, IHCs/100 µm; O, total OHCs; P, OHCs/100

µm; S, total SCs; T, SCs/100 µm.

See also S7 and S8 Figs.

1293

Fig 9. Loss of Fgf20 does not further delay prosensory specification in Sox2

hypomorph cochleae

(A)   The $Fgf20^{-/-}$, $Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$, $Fgf20^{-/-}$;$Sox2^{Ysb/Ysb}$, and $Sox2^{Ysb/-}$ cochleae phenotypes

lie along a continuum, preferentially affecting outer and basal cochlear sensory

epithelium, likely attributable to varying *Sox2* levels on an *Fgf20*-null background.

(B)   Serial sections (1-6) through the duct of E14.5 $Fgf20^{+/-}$;$Sox2^{Ysb/+}$, $Fgf20^{+/-}$;$Sox2^{Ysb/Ysb}$,

$Fgf20^{-/-}$;$Sox2^{Ysb/+}$, and $Fgf20^{-/-}$;$Sox2^{Ysb/Ysb}$ cochleae. Immunofluorescence for Ki67 (red)

and DAPI (nuclei, cyan). Cochlear epithelium is outlined. Bracket indicates prosensory

domain. * indicates shift of prosensory nuclei away from the luminal surface of the

epithelium. N, neural side. Whole mount cochlear duct schematics show relative

positions of the serial sections and progression of cell cycle exit (green arrow). Note:

unlike in Fig 7, we have switched the placement of images from $Fgf20^{-/-}$;$Sox2^{Ysb/+}$ and

$Fgf20^{+/-}$;$Sox2^{Ysb/Ysb}$ cochleae to facilitate comparison.

Scale bar, 100 µm. See also S10 Fig.

1308

Fig 10. Fgf20-KO organ of Corti exhibits premature differentiation

(A)   RNA in situ hybridization for *Hey1* and *Hey2* on sections through the middle turn of

E14.5 *Fgf20*$^{+/-}$ and *Fgf20*$^{-/-}$ cochleae*.*

(B, C) Immunofluorescence for Myo6 (red) on "mid-modiolar" sections through the (B) E14.5

Fgf20$^{+/-}$ and *Fgf20*$^{-/-}$ cochleae, and (C) E15.0 Fgf20-het (*Fgf20*$^{Cre/+}$;*ROSA*$^{rtTA/+}$), Fgf20-null

(*Fgf20*$^{Cre/βgal}$;*ROSA*$^{rtTA/+}$), Fgf9-OA (*Fgf20*$^{Cre/+}$;*ROSA*$^{rtTA/+}$;TRE-Fgf9-IRES-eGfp)*,* Fgf20-

rescue (*Fgf20*$^{Cre/βgal}$;*ROSA*$^{rtTA/+}$;TRE-Fgf9-IRES-eGfp) cochleae (Dox from E13.5 to

E15.0), with magnification*.* The number of cochleae containing Myo6-expressing cells

out of the total number of cochleae examined for each genotype are shown below each

panel. Arrows indicate Myo6-expressing hair cells. DAPI, nuclei (blue).

OS, outer sulcus; PD, prosensory domain; KO, Kölliker's organ. Scale bar, 100 μm.

1320

Fig 11. *Sox2* and *Fgf20* interact to modulate a temporal buffer between

specification and differentiation

Model of the roles of *Sox2* and *Fgf20* in organ of Corti development, which occurs in two stages:

unspecified progenitors (tan shading) undergo specification and cell cycle exit to become

prosensory cells (green shading), which then differentiate into hair cells and supporting cells

(HCs & SCs; red shading). In wildtype cochleae, cell cycle exit (indicating completion of

specification) begins at the apex of the cochlea and proceeds basally. Afterwards, differentiation

initiates at the mid-base of the cochlea and proceeds basally and apically. The prosensory cells

exist within a temporal buffer (green shading), defined as the time between cell cycle exit and

initiation of differentiation. In *Sox2*/*Fgf20* mutant cochleae, decrease in levels of *Sox2*

expression in the developing cochlea leads to delayed prosensory specification and cell cycle

exit (arrow 1), while loss of *Fgf20* leads to premature onset of differentiation at the basal and

mid-basal cochlear turns (arrow 2) as well as delayed differentiation at the apical turn (arrow 3).

Loss of both *Sox2* and *Fgf20* leads to loss of the temporal buffer between specification and

differentiation towards the base of the cochlear duct, disrupting the development of HCs and

SCs in the basal region (crosshatch pattern).

1337

1338

1339 **Supplemental figure titles and legends**

1340

S1 Fig.

(A-C)   Quantification of length-normalized number of (A) inner hair cells (IHCs/100 µm), (B)

outer hair cells (OHCs/100 µm), and (C) supporting cells (SCs/100 µm) in the basal,

middle, and apical turns of P0 cochleae from $Fgf20^{+/-}$, $Fgf20^{-/-}$ $Fgfr1^{flox/+}$, $Fgfr1^{flox/-}$,

$Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and $Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ mice. $Fgf20^{+/-}$ and $Fgf20^{-/-}$ cochleae were

analyzed by unpaired Student's t test; $Fgfr1^{flox/+}$, $Fgfr1^{flox/-}$, $Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and

$Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ cochleae were analyzed by one-way ANOVA. P values shown are

from the t test and ANOVA. * indicates $p < 0.05$ from Student's t test or Tukey's HSD

(ANOVA post-hoc); n.s., not significant. Error bars, mean ± SD.

(D)   Schematic showing the positions of basal, middle, and apical turns along the cochlear

duct. Apical tip refers to the apical end of the cochlea.

(E)   Whole mount cochlea from P0 $Fgf20^{+/-}$, $Fgf20^{-/-}$, $Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and

$Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ mice showing immunofluorescence for phalloidin (green) and

p75NTR (red) at the apical tip of the cochlea. T, towards the tip. Scale bar, 100 µm.

1355

S2 Fig.

(A, B) Sections through the middle turn of E14.5 cochlear ducts from $Fgf20^{+/-}$, $Fgf20^{-/-}$,

$Foxg1^{Cre/+}$;$Fgfr1^{flox/+}$, and $Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ mice. Scale bar, 100 µm. Refer to

schematic below. OS, outer sulcus; PD, prosensory domain; KO, Kölliker's organ.

(A)   RNA in situ hybridization for *Etv4* and *Etv5.* The two brackets indicate *Etv4/5* expression

in the outer sulcus (OS, left) and prosensory domain (PD, right; lost in $Fgf20^{-/-}$ and

$Foxg1^{Cre/+}$;$Fgfr1^{flox/-}$ cochleae).

(B)   EdU-incorporation (green). Dashed region indicates Kölliker's organ (KO). DAPI, nuclei

(blue).

1365

S3 Fig.

Quantification of length-normalized number of inner hair cells (IHCs/100 µm), outer hair cells

(OHCs/100 µm), and supporting cells (SCs/100 µm) overall (along the entire cochlea; top three

graphs) and in the basal, middle, and apical turns (bottom three graphs) of P0 cochleae from

$Fgf20^{+/-}$;$ROSA^{rtTA}$ (Fgf20-het), $Fgf20^{+/-}$;$ROSA^{rtTA}$; TRE-Fgf9-IRES-eGfp (Fgf9-OA), $Fgf20^{-}$

$^{/-}$;$ROSA^{rtTA}$ (Fgf20-null), and $Fgf20^{-/-}$;$ROSA^{rtTA}$; TRE-Fgf9-IRES-eGfp (Fgf20-rescue) mice. Dox

regimens: E13.5-E15.5, E13.5, E14.5, or E15.5. P values shown are from one-way ANOVA. *

indicates p < 0.05 from Tukey's HSD (ANOVA post-hoc); n.s., not significant. Error bars, mean ±

SD. Summarized in Fig 3C.

1375

S4 Fig.

(A-C)   Quantification of length-normalized number of (A) inner hair cells (IHCs/100 µm), (B)

outer hair cells (OHCs/100 µm), and (C) supporting cells (SCs/100 µm) in the basal,

middle, and apical turns of P0 cochleae from $Sox2^{+/+}$, $Sox2^{Ysb/+}$, $Sox2^{Ysb/Ysb}$, and $Sox2^{Ysb/-}$

mice. P values shown are from one-way ANOVA. * indicates p < 0.05 from Tukey's HSD

(ANOVA post-hoc); n.s., not significant. Error bars, mean ± SD.

(D)   Whole mount cochlea from P0 *Sox2$^{Ysb/-}$* mice showing presence of inner and outer hair

cells (phalloidin/p75NTR) and supporting cells (Prox1/Sox2, in a different cochlea) at the

basal tip. Schematic shows the location of sensory epithelium at the apical turn and

basal tip of *Sox2$^{Ysb/-}$* cochleae. Scale bar, 1 mm (whole), 100 µm (basal tip).

1386

S5 Fig.

(A, B) Immunofluorescence for (A) Sox2 (red) and (B) CKDN1B (green) in sections through the

basal, middle, and apical turns of E14.5 *Sox2$^{Ysb/+}$* and *Sox2$^{Ysb/-}$* cochleae.

(C)   Immunofluorescence for Ki67 (red) on serial "mid-modiolar" sections through the E14.5

and E15.5 *Sox2$^{Ysb/+}$* and *Sox2$^{Ysb/-}$* cochleae*. Brackets indicate prosensory domain. Nine

sections through the length of the cochlear duct are labeled. See whole mount cochlear

duct schematic (lower left) for relative positions of the sections.

DAPI, nuclei (blue). Scale bar, 100 µm.

1395

S6 Fig.

(A)   Immunofluorescence for Ki67 (red) on serial "mid-modiolar" sections through the E14.5

Fgf20$^{+/-}$ and *Fgf20$^{-/-}$* cochleae*. Brackets indicate prosensory domain. Nine sections

through the length of the cochlear duct are labeled. See whole mount cochlear duct

schematic (right) for relative positions of the sections.

(B)   EdU-incorporation (green) in sections through the middle turn of E14.5 *Sox2$^{Ysb/+}$* and

Sox2$^{Ysb/-}$ cochleae. Dashed region indicates Kölliker's organ (KO). Bracket indicates part

of Kölliker's organ without EdU-incorporating cells in *Sox2$^{Ysb/-}$* cochleae.

OS, outer sulcus; PD, prosensory domain; KO, Kölliker's organ. DAPI, nuclei (blue). Scale bar,

100 µm.

1406

1407   **S7 Fig.**

1408   (A)   P values from two-way ANOVA analyzing the quantification in (B-D). The two factors

1409   analyzed are *Fgf20* (*Fgf20$^{+/+}$*, *Fgf20$^{+/-}$*, *Fgf20$^{-/-}$*) and *Sox2* (*Sox2$^{+/+}$*, *Sox2$^{Ysb/+}$*, *Sox2$^{Ysb/Ysb}$*)

1410   gene dosage. A p value < 0.05 (yellow highlight) for *Fgf20* or *Sox2* indicates that the

1411   particular factor (independent variable) has a statistically significant effect on the

1412   measurement (dependent variable). Whereas a p value < 0.05 for Interaction indicates a

1413   statistically significant interaction between the effects of the two factors on the

1414   measurement.

1415   (B-D) Quantification of length-normalized number of (B) inner hair cells (IHCs/100 µm), (C)

1416   outer hair cells (OHCs/100 µm), and (D) supporting cells (SCs/100 µm) in the basal,

1417   middle, and apical turns of P0 cochleae from *Fgf20$^{+/+}$;Sox2$^{+/+}$*, *Fgf20$^{+/+}$;Sox2$^{Ysb/+}$*,

1418   *Fgf20$^{+/-}$;Sox2$^{+/+}$*, *Fgf20$^{+/-}$;Sox2$^{Ysb/+}$*, *Fgf20$^{+/+}$;Sox2$^{Ysb/Ysb}$*, *Fgf20$^{+/-}$;Sox2$^{Ysb/Ysb}$*, *Fgf20^{-}*

1419   *$^{/-}$;Sox2$^{+/+}$*, *Fgf20$^{-/-}$;Sox2$^{Ysb/+}$*, and *Fgf20$^{-/-}$;Sox2$^{Ysb/Ysb}$* mice. Error bars, mean ± SD.

1420

1421   **S8 Fig.**

1422   Results from post-hoc Tukey's HSD analyzing the quantification results in (B-D). Letters (L, I, J,

1423   O, P, S, T; representing each measurement in S7B-S7D Figs) indicate a statistically significant

1424   decrease (p < 0.05) when comparing the row genotype against the column genotype. L,

1425   cochlear length; I, IHCs/100 µm; O, OHCs/100 µm; S, SCs/100 µm.

1426

S9 Fig.

(A)   Whole mount cochlea from P0 $Fgf20^{+/+};Sox2^{+/+}$, $Fgf20^{+/-};Sox2^{+/+}$, $Fgf20^{+/+};Sox2^{Ysb/+}$, and

$Fgf20^{+/-};Sox2^{Ysb/+}$ mice showing inner and outer hair cells (phalloidin, green) separated

by inner pillar cells (p75NTR, red). Magnifications show the basal, middle, and apical

turns of the cochlea. Scale bar, 100 μm (magnifications), 1 mm (whole); arrowheads

indicate ectopic inner hair cells.

(B-F) Quantification of (B) cochlear duct length, (C) total inner hair cells (IHCs) and IHCs per

100 μm of the cochlear duct, (D) total outer hair cells (OHCs) and OHCs per 100 μm,

and (E) IHCs/100 μm and (F) OHCs/100 μm in the basal, middle, and apical turns at P0.

P values shown are from one-way ANOVA. * indicates $p < 0.05$ from Tukey's HSD

(ANOVA post-hoc); n.s., not significant. Error bars, mean ± SD.

1438

S10 Fig.

(A, B) Immunofluorescence for (A) Sox2 (red) and (B) CDKN1B (green) in sections through the

basal, middle, and apical turns of E14.5 $Fgf20^{+/-};Sox2^{Ysb/+}$, $Fgf20^{-/-};Sox2^{Ysb/+}$,

$Fgf20^{+/-};Sox2^{Ysb/Ysb}$, and $Fgf20^{-/-};Sox2^{Ysb/Ysb}$ cochleae.

(C)   EdU-incorporation (green) in sections through the middle turn of E14.5 $Fgf20^{+/-};Sox2^{Ysb/+}$,

$Fgf20^{-/-};Sox2^{Ysb/+}$, $Fgf20^{+/-};Sox2^{Ysb/Ysb}$, and $Fgf20^{-/-};Sox2^{Ysb/Ysb}$ cochleae. Dashed region

indicates Kölliker's organ (KO).

(D)   Immunofluorescence for Ki67 (red) on serial "mid-modiolar" sections through the E14.5

$Fgf20^{+/-};Sox2^{Ysb/+}$, $Fgf20^{-/-};Sox2^{Ysb/+}$, $Fgf20^{+/-};Sox2^{Ysb/Ysb}$, and $Fgf20^{-/-};Sox2^{Ysb/Ysb}$

cochleae. Brackets indicate prosensory domain. Nine sections through the length of the

cochlear duct are labeled. See whole mount cochlear duct schematic (upper right) for

relative positions of the sections.

Note: unlike in Fig 7, we have switched the placement of images from $Fgf20^{-/-};Sox2^{Ysb/+}$ and

$Fgf20^{+/-};Sox2^{Ysb/Ysb}$ cochleae to facilitate comparison. OS, outer sulcus; PD, prosensory domain;

KO, Kölliker's organ. DAPI, nuclei (blue). Scale bar, 100 μm.

E14.5

Fig 3

A — Fgf20-het, Fgf9-OA, Fgf20-null, Fgf20-rescue (E13.5-15.5, E13.5, E14.5, E15.5 Dox); basal, middle, apical; P0

Phalloidin p75NTR

B — basal, middle, apical; P0

Prox1 Sox2

C

		Fgf9-OA	Fgf20-null	Fgf20-rescue E13.5-15.5	E13.5	E14.5	E15.5 Dox
OHC	basal	95.3%*	48.6%^	105.8%*	102.4%*	76.2%*	41.7%^
	middle	99.6%*	32.4%^	108.2%*	89.0%*	94.4%*	44.7%^
	apical	97.1%*	29.0%^	101.3%*	81.2%*^	94.8%*	75.7%*^
SC	basal	99.0%*	44.1%^	103.4%*	85.6%*^	66.9%^	47.1%^
	middle	97.6%*	40.3%^	102.8%*	88.7%*	80.7%*	49.5%^
	apical	88.0%*	28.4%^	94.0%*	83.3%	74.7%	70.9%*^

<60.0%	60.0-69.9%	70.0-79.9%	80.0-89.9%	>90.0%

A

E14.5

Sox2^{Ysb/+}   *Sox2^{Ysb/-}*

E15.5

Sox2^{Ysb/+}   *Sox2^{Ysb/-}*

serial sections

Ki67   DAPI

E14.5

Sox2^{Ysb/+}   *Sox2^{Ysb/-}*

E15.5

Sox2^{Ysb/+}   *Sox2^{Ysb/-}*

→ cell cycle exit

B

Wildtype

Sox2 hypomorph

basal / middle / apical

Unspecified progenitors

HCs & SCs

Cell cycle exit

Initiation of differentiation

Prosensory cells (Temporal buffer)

Unspecified progenitors

HCs & SCs

Cell cycle exit

Initiation of differentiation

Prosensory cells

Prosensory specification

Differentiation

Unspecified progenitors

Prosensory cells

HCs & SCs

A

Sox2^(Ysb/+)   *Sox2^(Ysb/-)*

Etv4

OS   PD

Etv5

B

Fgfr1

Fgf20

E14.5

C

Foxg1^(Cre/+)   *Foxg1^(Cre/+)*
Fgfr1^(flox/+)   *Fgfr1^(flox/-)*

Fgf20

E14.5

D   *Fgf20^(+/-)*   E14.5   *Fgf20^(-/-)*

serial sections

Ki67   DAPI

Fgf20^(+/-)   *Fgf20^(-/-)*

apical   basal
cell cycle exit

Abneural   Neural
OS   PD   KO

Fig 7

Fig 8

L   cochlear length   P   OHCs/100 μm
I   total IHCs   S   total SCs
J   IHCs/100 μm   T   SCs/100 μm
O   total OHCs

A

Organ of Corti phenotype

mild   severe

$Fgf20^{-/-}$ $Foxg1^{Cre/+};Fgfr1^{flox/-}$ $Fgf20^{-/-};Sox2^{Ysb/Ysb}$ $Sox2^{Ysb/-}$

Sox2 levels

Basal sensory epithelium

Outer compartment

Inner compartment

B   E14.5

$Fgf20^{+/-};Sox2^{Ysb/+}$ $Fgf20^{-/-};Sox2^{Ysb/+}$ $Fgf20^{+/-};Sox2^{Ysb/Ysb}$ $Fgf20^{-/-};Sox2^{Ysb/Ysb}$

serial sections

1 2 3 4 5 6

Ki67   DAPI

$Fgf20^{+/-};Sox2^{Ysb/+}$ $Fgf20^{-/-};Sox2^{Ysb/+}$ $Fgf20^{+/-};Sox2^{Ysb/Ysb}$ $Fgf20^{-/-};Sox2^{Ysb/Ysb}$

apical   basal

cell cycle exit

Ligand X    FGF20

FGFR1    FGFR1

Sox2     HEY1/2

Prosensory  Premature onset of
specification   differentiation

Unspecified  Prosensory  HCs & SCs
progenitors   cells