# Black or red: A sex-linked colour polymorphism in a songbird is maintained by balancing selection

Kang-Wook Kim[1][§][*], Benjamin C. Jackson[1][§], Hanyuan Zhang[1], David P. L. Toews[2,3], Scott A. Taylor[2,3,4], Emma I. Greig[2], Irby J. Lovette[2,3], Mengning M. Liu[5], Angus Davison[5], Simon C. Griffith[6], Kai Zeng[1][*], Terry Burke[1][*]

1. Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, United Kingdom.

2. Fuller Evolutionary Biology Program, Cornell Lab of Ornithology, Cornell University, 159 Sapsucker Woods Road, Ithaca, NY 14850, USA

3. Department of Ecology and Evolutionary Biology, Cornell University, Corson Hall, Ithaca, NY 14853, USA

4. Department of Ecology and Evolutionary Biology, University of Colorado - Boulder, 1900 Pleasant Street, Boulder, CO, USA

5. School of Biology, University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom

6. Department of Biological Sciences, Macquarie University, Sydney, NSW 2109, Australia.

[§] Equal contribution

[*] Correspondence to: k.kim@sheffield.ac.uk, k.zeng@sheffield.ac.uk, t.a.burke@sheffield.ac.uk;

Tel: +44 114 222 0096 (T.B.)

Word count - Abstract: 153, Main text: 3,372
Figures: 4
References - Main: 70; Methods & Supplementary Materials: 63
Supplementary Informations: Supplementary Figures: 16, Tables: 14, Text: 1
Supplementary Dataset: One Excel file summarising gene expression analysis.

**Colour polymorphisms play a key role in sexual selection and speciation, yet the mechanisms**

**that generate and maintain them are not fully understood. We therefore used genomic and**

**transcriptomic tools to identify the precise genetic architecture and evolutionary history of a**

**sex-linked colour polymorphism in the Gouldian finch *Erythrura gouldiae* that is also**

**accompanied by remarkable differences in behaviour and physiology. We found that differences**

**in colour are associated with an ~72-kbp region of the Z chromosome in a putative regulatory**

**region for *follistatin*, an antagonist of the *TGF-β* superfamily genes. The region is highly**

**differentiated between morphs, unlike the rest of the genome, yet we found no evidence that an**

**inversion is involved in maintaining the distinct haplotypes. Coalescent simulations confirm that**

**that there is elevated nucleotide diversity and an excess of intermediate frequency alleles at this**

**locus. We conclude that this pleiotropic colour polymorphism is most probably maintained by**

**balancing selection.**

Colour-polymorphic species are observed across diverse taxa, but are relatively rare and

younger than monomorphic species[1,2]. This is probably because any difference in fitness between

alternative morphs should result in either the fixation of one of the causative alleles under natural or

sexual selection, or lead to sympatric speciation under disruptive selection – potentially increasing the

rate of speciation and promoting the transition from polymorphism to monomorphism[2-4]. Even if there

are no fitness differences, genetic drift is expected to remove alternative morphs from a small

population[3]. Therefore, the evolutionary processes that maintain colour polymorphisms in some

species over long periods are not well understood. This is partly because the genetic basis of

pigmentation, especially for non-melanic pigments like carotenoids and psittacofulvins, has only

recently been elucidated[5-10].

Gouldian finch (*Erythrura gouldiae*) head plumage is a classic example of a Mendelian

colour polymorphism in which the morphs co-occur in sympatry. The striking head colour morphs of

the Gouldian finch reflect variation in both carotenoid and melanin pigmentation[11-13]; a pigmentation

locus on the Z chromosome (hereafter the *Red* locus, with a recessive $Z^r$ allele and a less frequent

dominant $Z^R$ allele) produces melanin-based black ($Z^rZ^r$ in male, $Z^r$ in female) or carotenoid-based red

($Z^R Z^R$, $Z^R Z^r$ in male, $Z^R$ in female) head morphs in both sexes, as well as differences in feather

structure[11,12]. Rarely in the wild (<1% of birds), an epistatic interaction of the *Red* locus with an

autosomal locus produces a yellow morph[14].

With regards to the black and red morphs, there are remarkable inter-morph differences in

stress hormone levels, personalities, social dominance, sperm-length plasticity, and sex allocation

associated with the colour polymorphism (see Supplementary Table 1 for summary and references).

Moreover, pre- and postzygotic incompatibilities have been documented between these morphs in a

captive population (Supplementary Table 1), but the same phenomena have not been fully

investigated in wild populations[15].

Despite these differences, red and black morphs coexist in natural populations at stable

phenotypic frequencies[16], suggesting that there are mechanisms maintaining this multi-functional

colour polymorphism in nature. Given that multiple phenotypes are associated with head colour, one

hypothesis is that the *Red* locus is a supergene, defined as 'a group of genes acting as a mechanical

unit in particular allelic combinations'[17], which is often facilitated by a chromosomal inversion, as

recently demonstrated in several avian species[18-21]. Alternatively, the *Red* locus might be a single

pleiotropic locus that can regulate multiple genes[22]. To test these hypotheses, we identified the

genomic location of the *Red* locus and undertook a detailed analysis of the region using genomic and

transcriptomic tools. In addition, we examined the evolutionary mechanism behind the maintenance

of the colour polymorphism using coalescent simulations.

**Results and discussion**

**Identification of the *Red* locus**

By using high-depth / low SNP density RADSeq[23] and low-depth / high SNP density whole-genome

sequencing (see Methods), we identified a region on the Z chromosome strongly associated with red–

black head colouration (1,000 permutations, $P < 0.05$, Fig. 1a, Supplementary Figs. 1–2) within the

candidate interval identified in a previous linkage analysis[24]. The genotypes of the wild birds ($n_{male}$ =

89, $n_{female}$ = 72) from an allele-specific PCR in this region showed a near-perfect association with head

colour, with one exception, in a female (Fisher's exact test, male: $P < 10^{-15}$; female: $P = 7.52 \times 10^{-10}$,

Fig. 1b, Supplementary Fig. 2). The frequencies of the black-linked allele in this population were

similar in females (0.88) and males (0.85) (Fisher's exact test, $P = 0.69$). The genotypic frequencies in

males did not depart from Hardy–Weinberg Equilibrium ($P_{permutation} = 0.24$, Supplementary Table 3).

The low frequency of homozygous red males is not significantly different from expectation for a sex-

linked locus under random mating (Supplementary Table 3) and a prediction based on the phenotypic

frequency[11].

**High linkage disequilibrium but no genomic rearrangements around the *Red* locus**

Next, we examined if the multi-functional polymorphism is caused by a supergene. Previously-

described supergenes have large genomic blocks of elevated linkage disequilibrium (LD) that harbour

multiple genes. We measured the extent of LD around the *Red* locus, defined as the region with the

most significant association with the colour polymorphism and its flanking regions (Fig. 1c), using

Sanger sequencing of multiple loci. Intriguingly, the LD block significantly associated with the colour

polymorphism was limited to a single intergenic region (where no known protein-coding genes are

annotated in zebra finch) that spans approximately 72 kilobase pairs (kbp) and lies between the coding

regions of the genes *molybdenum cofactor synthesis 2* (*MOCS2*) and *follistatin* (*FST*) (Fig. 1c). In this

region, the density of fixed differences between the $Z^R$ and $Z^r$ alleles was high (6.8–27.8 per kbp;

Supplementary Tables 4-5; this region is referred to as the "core" region hereafter).

Centromeres and genomic rearrangements, particularly chromosomal inversions, may reduce

recombination rates and generate significant LD blocks between haplotypes[25-27]. However, our

previous work showed that the *Red* locus is not located near the centromere or in a large genomic

rearrangement between morphs[24]. We confirmed this finding with a fine-scale test for inversion

breakpoints using long-range PCR: our genomic assembly of the *Red* locus region for each genotype

contained contiguous homologous sequence, consistent with the absence of an inversion

differentiating the $Z^R$ and $Z^r$ alleles (Fig. 1c, Supplementary Table 7).

**The *Red* locus as a candidate regulatory region of pleiotropic *FST***

Gouldian finch head colour is associated with multiple phenotypic traits (see above). As such, the intergenic location of the *Red* locus suggests that this locus acts as a regulatory region that controls a pleiotropic gene. From a small-scale transcriptomic analysis of regenerating feather follicles from the scalp, we identified a list of differentially expressed genes belonging to functionally correlated gene ontology (GO) networks with potential roles in feather development and pigmentation (Supplementary Fig. 3, Supplementary Dataset 1). Although neither *MOCS2* or *FST* were differentially expressed between morphs, some genes in this network are known to interact with *FST* (Supplementary Fig. 3, Supplementary Dataset 1).

The multiple differences in gene expression suggest that the fates of feather precursors are determined by gene(s) that act upstream of the regulatory network at an earlier developmental stage. Previous studies indicate that *FST* is a strong candidate for regulating the colour polymorphism. Follistatin is known to regulate hair and feather development through its antagonistic action on genes of the *TGF-β* superfamily[28,29]. The genes in this family (e.g. *activin*, *TGF-β* and *BMP*) may control melanocyte differentiation and melanogenesis[30-34]. *FST* and its downstream genes might also be responsible for the structural differences in the feathers found in the Gouldian finch polymorphism as *BMP* and its alternative antagonist, *noggin*, are important determinants of the morphogenesis of feather branching[29,35-37]. The *TGF-β* superfamily and its regulator *FST* are also involved in the development and regulation of the visual[38,39] and reproductive systems[40-43].

We found no differences in the *FST* protein-coding sequences between black- and red-headed birds, and the amino-acid sequences were conserved between the Gouldian finch and the zebra finch (Supplementary Fig. 4). However, we identified several evolutionarily conserved sites within the core of the *Red* locus located upstream of *FST* (Supplementary Fig. 5), suggesting that this region has functional importance. It is therefore likely that *cis*-regulatory region(s) in the *Red* locus determine spatio-temporal variation in the expression of *FST* that then controls multiple genes in the *TGF-β* superfamily, which in turn may be responsible for the pigmentation difference and other pleiotropic effects. We note that one of the few divergent regions that distinguish the genomes of hybridizing golden-winged (*Vermivora chrysoptera*) and blue-winged (*V. cyanoptera*) warblers has been mapped to the same intergenic region between *MOCS2* and *FST*[44], suggesting convergent evolution of

plumage colouration mechanisms and a possibly important role of the *Red* locus as a barrier to gene

flow in other taxa. However, the fact that this locus has not led to speciation in the Gouldian finch

indicates that selection may instead promote the persistence of divergent morphs in this species.

**The pattern of nucleotide diversity and divergence at the *Red* locus**

If selection has played a role in the origin and maintenance of the *Red* locus then it may have left a

signature in the pattern of nucleotide diversity in this region. We calculated point estimates of

summary statistics at the *Red* locus, as well as in sliding-windows across the *Red* locus to characterize

the pattern of nucleotide diversity and divergence within and between morphs (Figs. 2a-d;

Supplementary Table 9). We then compared the *Red* locus to 24 randomly-chosen intronic loci on the

Z chromosome, as a putatively neutrally-evolving reference set (Figs. 2e-h; Supplementary Table 9).

Notably, diversity for the combined sample of alleles ($Z^R$ and $Z^r$ together) at the *Red* locus is elevated

compared to the reference loci, as is the level of differentiation between $Z^R$ and $Z^r$ alleles.

The mean nucleotide diversity at the reference loci ($\pi_{ref}$) was 0.0033 and varied little between

the alleles of each morph (Fig. 2e). The point estimate of nucleotide diversity for the combined

sample at the *Red* locus ($\pi_{total} = 0.0092$) was above that at all 24 reference loci (Fig. 2a), while

diversity within the $Z^R$ ($\pi_{red} = 0.0008$) and the $Z^r$ ($\pi_{black} = 0.0015$) alleles was similar to that at the

reference loci.

The estimate of absolute divergence $d_{xy}$ (the mean number of pairwise differences per site[45])

between the $Z^R$ and $Z^r$ alleles at the *Red* locus ($d_{xy\ Red\ locus} = 0.0149$), was again greater than between

morphs at all 24 reference loci ($d_{xy\ ref} = 0.0033$; Fig. 2b, f) and the estimate of net divergence $d_a$ ($d_{xy}$

minus the mean of the two within-morph $\pi$ values) was not greatly different from $d_{xy}$ due to the low

nucleotide diversity within each allelic class (data not shown). In contrast, the divergence between

zebra finch and alleles of each morph was similar at both the *Red* locus and the reference loci (Fig. 2b,

f).

Tajima's $D$[46] was moderately negative at the reference loci, both within the alleles of each

morph and with both morphs' alleles combined ($D_{Taj\ black} = -0.599$, $D_{Taj\ red} = -0.304$ and $D_{Taj\ total} = -$

0.818), as well as within the alleles of each morph at the *Red* locus ($D_{Taj\ red} = -0.199$ and $D_{Taj\ black} = -$

0.911; Fig. 2g, Supplementary Table 9), consistent with population expansion (see Supplementary Text). However, $D_{Taj}$ was positive in the combined sample at the *Red* locus ($D_{Taj\ total} = 2.013$; Fig. 2c), which is significantly greater than zero under the standard neutral model ($P_{coalescent} = 0.002$) using simulations in DnaSP v5[47], and again higher than at all 24 reference loci.

In captive populations of the Gouldian finch, pronounced pre- and postzygotic incompatibilities between morphs have been described, and mixed-morph pairings resulted in reduced survival: 40% of male and 80% of female offspring died before 90 days of age[48,49]. If these incompatibilities persisted in the wild, then genome-wide differentiation between the black and red morphs might be expected[50]. To quantify this, we estimated Wright's fixation index $F_{ST}$[51] between each morph's alleles. $F_{ST}$ between alleles of each morph reached nearly maximal values ($F_{ST} = 0.935$) in the core region and was significantly greater than zero ($P_{permutation} = 0.002$). This value was greater than $F_{ST}$ between each morph's alleles at all the Z-linked reference loci (composite $F_{ST} = 0.038$, Figs. 2d, h, Supplementary Fig. 1).

In contrast, we found no genomic differentiation with regards to colour morph across the genome outside of the *Red* locus in the wild sample. First, a "coancestry matrix"[52-54] constructed for 32 wild birds ($n_{black} = 22$, $n_{red} = 10$) using autosomal RADSeq loci, designed to fully exploit the available haplotypic information, failed to identify any clusters that separated the morphs (Supplementary Fig. 6). Second, a STRUCTURE[55] analysis between morphs using nine unlinked autosomal microsatellite markers for all the wild birds ($n_{black} = 126$, $n_{red} = 35$) indicated that the most-likely number of subpopulations was one ($K = 1$; Supplementary Fig. 7). These results suggest that there is no or little incompatibility leading to any genome-wide differentiation between morphs in the wild.

**Test for neutrality**

Overall, the patterns of nucleotide diversity and divergence at the *Red* locus (i.e. elevated $\pi$, excess of intermediate frequency alleles and maximal $F_{ST}$) are consistent with balancing selection[56]. However, such patterns can be obtained even under neutral evolution if sampling is conditioned on a site associated with the phenotypes of interest, as in this case (e.g. see the simulated patterns of

polymorphism close to the focal site presented in Figure 3). Therefore, to formally test for neutrality

we carried out coalescent simulations under a neutral model that incorporated demography,

recombination, and accounted for our sampling scheme (see Methods, Supplementary Text,

Supplementary Figs. 8-13, Supplementary Tables 11-14).

Our simulations show that the observed patterns of genetic variation at the *Red* locus cannot

be explained by either neutral processes or biases caused by the sampling scheme (Fig. 3,

Supplementary Text). Assuming the $Z^R$ allele is derived (i.e., the derived allele frequency, DAF =

0.144), the observed nucleotide diversity ($\pi_{\text{total}}$) and Tajima's *D* for the combined sample of $Z^R$ and $Z^r$

alleles at the *Red* locus ($D_{\text{Taj total}}$) both lie above the 95% confidence intervals (CIs) of the simulations

for most windows, as do $F_{\text{ST}}$ and $d_{\text{xy}}$ between the $Z^R$ and $Z^r$ alleles (Figs. 3a-d). In contrast, statistics

calculated within the $Z^R$ or $Z^r$ alleles fell within the 95% CIs of the simulated distributions, with the

exception of nucleotide diversity within the $Z^R$ allele ($\pi_{\text{red}}$, Supplementary Fig. 9). The same pattern

was observed when we assumed that the $Z^r$ allele was derived (DAF = 0.856, Supplementary Fig. 10).

In particular, the observed region of greater than expected differentiation extended over an ~70-kbp

interval, with a rapid drop on either side – contrasting with neutral expectations for a single site on

which sampling is conditioned, which results in a gradual decline towards background levels over a

20-kbp interval (Fig. 3d). As we have demonstrated here, there is no evidence of an inversion across

this region (Supplementary Table 7). Moreover, we found evidence for the presence of recombination

within the *Red* locus (see Supplementary Text). Therefore, the highly diverged haplotype blocks are

not likely to be maintained by reduced recombination, but rather by selection against recombinants. A

sliding-window implementation of the HKA test[57], taking into account non-random sampling,

demography and recombination at the *Red* locus, also revealed significant departures from neutrality

for multiple windows across the core of the *Red* locus (Fig. 3e).

**The evolutionary history of the black and red haplotypes**

Balancing selection can maintain polymorphism over extended periods of time, resulting in deeply

diverged haplotypes that may be even older than the species in which they reside[56]. In a gene

genealogy of the *M1* region (see Fig. 1c) for 14 species in the family *Estrildidae,* sequence divergence

between black and red haplotypes was deeper than that between other species or even between genera

(Fig. 4, Supplementary Fig. 14). Four other *Erythrura* species formed a monophyletic group

(bootstrap support 100%), separate from the Gouldian finch, whereas none of the Gouldian finch

haplotypes clustered with the sequences of red-faced finches (*E. pealii* and *E. psittacea*) or blue-faced

finches (*E. tricolor and E. trichroa*). Despite the high divergence, the two Gouldian finch haplotypes

formed a clear monophyletic group (bootstrap support 100%), distinct from other *Erythrura* species,

suggesting that these haplotypes evolved within the Gouldian finch lineage (Fig. 4). These patterns

suggest that the divergent haplotypes at the *Red* locus are old (nearly half of the divergence between

zebra finch and the Gouldian finch), although we did not attempt to estimate the age since balancing

selection is likely to affect this estimation. Near the highly divergent *M1* region, we identified an

insertion with the characteristic feature of a long terminal repeat (LTR) retrotransposon, which is

absent in the zebra finch (Fig. 2). Transposable elements are often associated with adaptive

polymorphism[58]. Indeed, signals of selection, including the departure of the observed data from

expectations under neutrality derived from simulations as well as from the HKA test, were strongest

in this region (Fig. 3e, Supplementary Text). However, comparison of the 3' and 5' LTRs in each

haplotype suggests that the divergence between the colour-associated haplotypes did not occur until

after the insertion of the retrotransposon, which occurred some time after the divergence of Gouldian

finch and zebra finch (Supplementary Fig. 15).

In a relative rate test, the red haplotype showed disproportionately longer branch lengths than

the black haplotype across the *Red* locus (Supplementary Figs. 14, 16), suggesting that the two

haplotypes have evolved with unequal evolutionary rates. Although the driver of this asymmetry

remains unclear, one possible explanation is that the red haplotype may have undergone more rounds

of adaptive substitution, which would also explain the lower diversity of the red haplotype over a long

stretch (Supplementary Fig. 9b, 10f).

**The origin of divergence and the mode of balancing selection**

There are several hypotheses for the origin of haplotypic divergence at the *Red* locus, including

former allopatry followed by secondary contact[59] or the lateral transfer of ancestral variation[60]. For

example, in northern Australia, the historical Ord Arid Intrusion event resulted in the splitting of

many co-distributed taxa into three distinct biogeographical populations, and some became (and

remain) distinct subspecies[61].

Regardless of the initial cause of divergence, our analyses suggest that balancing selection

may be responsible for maintaining the divergent haplotypes at the *Red* locus. There are a few

possible mechanisms of balancing selection in this species. First, red males may have an advantage

under sexual selection, as females of all morphs of the Gouldian finch show a preference for red

males[48]. Moreover, the dominant status of red males in the social structure[62] may be beneficial when

competing for limited resources, such as nest sites, albeit at the expense of reduced paternal care[63].

Since the red allele is genetically dominant, its allele frequency would be expected to quickly rise to a

high level under such a selective advantage. However, the red phenotype is present at low frequency

in natural populations[16]. Therefore, opposing selection has probably prevented the spread of the red

allele in the population. Red males suffer from high stress levels and have significantly poorer

reproductive outcomes than black birds when aggressive red males are in high frequency, and are

consequently in a more competitive and stressful environment[64-66]. Thus, frequency-dependent

selection may have acted to maintain a low equilibrium frequency of the red individuals.

Alternatively, this may be an example of sexual antagonism[67], in which the fitness effects of different

colouration are not the same in both sexes, and selection against red females prevents the red

haplotype from increasing to a high frequency, despite being favoured in males. Red females fully

express red head colour in captive populations (as a result of strong artificial selection by

aviculturists). However, those in the wild population are less brightly coloured generally; the

proportion of red feathers on their heads is typically lower, and they are often indistinguishable from

black females[11], suggesting selection against redness in females in the wild.

In this study, we have shown that a single locus upstream of *FST* is associated with

colouration. This locus is intergenic, which does not support a role for a 'supergene' in the classical

sense[17] in differentiating the red and black colour morphs of Gouldian finches. But it provides an

example of a putative regulatory element with downstream effects on multiple phenotypes being

maintained in the long-term by balancing selection. This has resulted in the maintenance of markedly

divergent morphs, despite gene flow within the species, and in the absence of genomic

rearrangements.

The lack of wider genomic divergence between morphs suggests that the *Red* locus may not,

alone, be sufficient to induce the incompatibilities observed in captive populations. This, in turn,

suggests that the extended LD in captive populations (e.g. an ~8 Mb region showing zero

recombination around the *Red* locus[24]) might encompass other linked loci underlying these

incompatibilities. Therefore, further studies of the region around the *Red* locus will provide an

important opportunity to address novel questions about the evolution of colour polymorphism[68,69]. In

particular, understanding the functional role of *Red*, a pleiotropic "switch" gene associated with both

melanin and carotenoid colouration, may shed light on the mechanistic basis of sexual selection and

genetic incompatibilities between colour morphs, which are both key evolutionary factors in

speciation[70].

**Methods**

**DNA samples**: Blood samples were collected from 161 Gouldian finches (male: $n_{black} = 62$, $n_{red} = 27$; female: $n_{black} = 64$, $n_{red} = 8$) in 2008 from a single wild population in Western Australia (within a 50-km radius of Wyndham in East Kimberley). Birds were either caught in mist nets at waterholes or at their nests. The population size was estimated to be around 300 adults. Genomic DNA was extracted using a modified ammonium acetate method[71]. The sex of each sample was determined using two sex-linked markers, *Z-002D*[72] and *Z37B*[73] that amplify alleles on both the Z and W chromosomes. No power analysis was performed prior to the study. All 161 samples were used for the association test and a population structure analysis. A subset of samples was randomly selected and used for RADSeq and subsequent Sanger and MiSeq sequencing and genotyping (see below). Throughout the study, traits were measured blind to treatment and to values of other traits.

For whole-genome sequencing (WGS), blood samples were collected from 24 Gouldian finches (males: $n_{black} = 4$, $n_{red} = 4$, $n_{yellow} = 3$; females: $n_{black} = 3$, $n_{red} = 5$, $n_{yellow} = 5$) that were sampled from a single captive population of domesticated birds in Ithaca, NY, in 2016. Yellow birds have the same allele at the *Red* locus as the red morph but develop as yellow under the control of an unknown epistatic autosomal locus[13]. Genomic DNA was extracted from these blood samples using DNeasy kits (tissue protocol; Qiagen, Hilden, Germany).

**RADSeq**: We genotyped 32 birds from the wild population described above using RADSeq[23] in two separate batches. In the first batch, DNA fragment libraries for 20 individuals ($n_{black\_male} = 10$, $n_{red\_female} = 8$ and $n_{red\_male} = 2$) were constructed following restriction digestion with *Sbf*I, and the pooled libraries were sequenced using two lanes of Illumina Genome Analyzer IIx. This sampling scheme was to identify the *Red* locus efficiently with limited sample size in GWAS (see below). In the second batch, a pooled library including 12 additional black males from the same population was constructed using the same protocol but sequenced on an Illumina HiSeq2000 instrument. The quality of the short sequence reads (2 x 100 bp) was checked using FASTQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). We cleaned and deduplicated the sequence reads using the *process_radtags* and *clone_filter* utility programs from STACKS[74]. We then

mapped the sequence reads against the zebra finch (*Taeniopygia guttata*) reference genome assembly

(taeGut3.2.4) using SeqMan NGen 4.0 (DNASTAR, Madison, Wisconsin USA), allowing up to 15%

divergence between sequence reads and the reference sequence. Single-nucleotide variants were

called using SAMTOOLS v1.2[75] and filtered based on the minimum depth for each site per individual

($> 2$), minimum SNP quality ($> 250$), maximum of missing genotypes across all individuals ($< 30\%$)

and minor allele frequency filter ($> 0.05$) using custom R scripts and VCFtools v0.1.12[76]. Although

the HiSeq2000 sequencing runs had more coverage than the GAIIx runs, by applying filtering based

on missing genotypes we could effectively remove any sites that were found only in the HiSeq2000

run. Mean depth per locus in an individual was different among libraries (GAIIx: L1 = 3.2, L2 = 7.8;

HiSeq2000: L3 = 13.2). However, because we mixed equal numbers of black and red birds in L1 and

L2, any bias due to the coverage in the inference of structure between morphs should be limited.

Females were encoded to have only one allele for the genotypes of Z-linked loci using the *vcf-fix-*

*ploidy* function of VCFtools. In an $F_{ST}$ outlier analysis (Supplementary Fig. 1), we included 32

individuals ($n_{black\_male} = 22$, $n_{red\_female} = 8$ and $n_{red\_male} = 2$) whose genotypes at the *Red* locus were

either homozygous or hemizygous, except for the two heterozygous red males.

**GWAS**: SNPs linked to the causative site at the *Red* locus are expected to show a recessive

homozygous genotype in black males (*a/a*), to be hemizygous (*A*) in red females and to be dominant

homozygotes (*A/A*) or heterozygotes (*A/a*) in red males (see main text). In a genome-wide association

study (GWAS) in 32 RAD-sequenced samples, we used Fisher's exact tests with one degree of

freedom for a dominant model of penetrance in a 2 x 2 contingency table of phenotype (black or red)

by genotype (*a/a* or not *a/a*) using PLINK v1.9[77,78] (Fig. 1a). We treated the hemizygous Z-

chromosome genotypes of females as homozygotes. We corrected for genomic structure among

samples using the *--within* option and the family-wise estimate of the SNP's significance was

determined and corrected for multiple testing using the *--mperm* option (1,000 permutations).

***De-novo* low-coverage whole-genome sequencing**: Sequencing libraries for the 24 captive Gouldian

finches were generated using the Illumina TruSEQ PCR-free Library Preparation Kit following the

protocol for generating 350-bp libraries. These 24 individuals were sequenced on a single lane of an

Illumina NextSeq500 instrument to produce 150-bp paired-end reads. We used AdapterRemoval[79] to

collapse overlapping paired reads, trim sequences of "Ns" along the 5' and 3' ends of reads, and

remove sequences shorter than 20 bp. This resulted in a mean of 33.7 million total reads per

individual (range 20.8–49.3 million reads). We used BOWTIE2[80] to map each of the individual reads

to the zebra finch genome to a mean coverage of approximately 3.6×, using the "very sensitive local"

set of alignment presets. For SNP discovery and variant calling, we used the UnifiedGenotyper in

GATK[81]. We removed possible variants that had "quality by depth" (QD) of < 2 and "mapping

quality" (MQ) of < 30 (the filtering conditions were: QD < 2.0, FisherStrand > 40.0, MQ < 30.0,

HaplotypeScore > 12.0, MappingQualityRankSum < -12.5, ReadPosRankSum < -8.0). Variant

confidence is a measure of sequencing depth at a given variant site; mapping quality refers to the root-

mean-square of the mapping confidence (from BOWTIE2) of reads across all samples. We applied

additional filters using VCFtools. First, we coded genotypes with a Phred-scaled quality lower than 8

as missing data. This is a permissive filter, necessary given our low-coverage data, which allows for

genotypic calls with as few as three reads at a given site for an individual. We excluded loci with

more than 30% missing data and used a minor allele frequency filter of 5%. This resulted in 4,818,133

SNPs distributed across the genome.

To properly correct for the ploidy of genotypes identified on the Z chromosome, we used the

*vcf-fix-ploidy* function of VCFtools to assign one allele at each site for Z-linked loci in females. To

increase the chance of detecting the location of the *Red* locus in an $F_{ST}$ outlier analysis

(Supplementary Fig. 1), we only included data from 17 individuals whose genotypes were known to

be homozygous or hemizygous for alternative alleles at the *Red* locus based on the pedigree (black:

$n_{black\ male} = 4$, $n_{black\ female} = 3$; red: $n_{red\ female} = 5$ and $n_{yellow\ female} = 5$).

**Allele-specific PCR analyses to test for the association with colour phenotype**: Several

segregating sites associated with head colour were identified in the sequences around a RADSeq locus

we found to be associated with head colour (locus *Ego165*, Supplementary Table 4). To confirm the

association with plumage colour, 161 birds were genotyped for alternative two-base pair variants (TG

/ CA) using an allele-specific PCR with the Multiplex PCR Kit (QIAGEN) and a set of primers (locus

*M1*, see Supplementary Table 2). The conditions for the PCR were 95°C for 15 min, 30 cycles of

94°C for 30 s, 64°C for 90 s and 72°C for 1 min, followed by 72°C for 5 min. The PCR products were

separated on an ABI 3730 DNA Analyzer (Applied Biosystems, Foster City, California, USA) and

scored using GENEMAPPER v 3.7 (Applied Biosystems). The association of the genotypes with face

colour was tested for each sex using Fisher's exact tests in R v3.2.5 (R Core Team, 2015. Fig. 1b). In

addition to the locus *M1*, a multiplex marker set was developed including two allele-specific markers

to genotype loci at the edges of the LD block around the *Red* locus (*M2* and *M3*, Supplementary Table

2, Supplementary Fig. 2). The same condition as above was used in the PCR, except for an annealing

temperature of 60°C. We examined the deviation of the *M1* genotypes from Hardy–Weinberg

equilibrium (HWE) using Graffelman–Weir's exact test for sex-linked markers implemented in the

HardyWeinberg package and obtained the *P*-value by permuting individual genotypes 10,000

times[82,83] (Supplementary Table 3).

**The extent of linkage disequilibrium and sequencing of a candidate gene**: To determine the extent

of LD across the *Red* locus and the association of marker loci with the colour trait, the DNA

sequences of 26 short fragments in the upstream region and the whole exon of an adjacent candidate

gene (*follistatin*) were obtained (Supplementary Table 4). We sequenced an additional 15 loci on the

Z chromosome that are not close to the *Red* locus as controls (Supplementary Table 4). Genomic

DNA from 16 females ($n_{black}$ = 8, $n_{red}$ = 8) was amplified and cleaned with ExoSAP-IT (USB,

Cleveland, Ohio, USA). After Sanger sequencing with the BigDye Terminator v 3.1 cycle sequencing

kit (Applied Biosystems), and precipitation using the ethanol/EDTA method, the products were

separated on an ABI 3730 DNA Analyzer and the bases were called using CodonCode Aligner v2.0.6

(CodonCode Corporation). Association tests between phenotype and 282 SNPs from 26 fragments,

and the estimation of pairwise LD between SNPs, were performed using Haploview v4.2[84] and

visualized using LDheatmap[85] (Fig. 1c). The coding sequences of *follistatin* in 16 females were

aligned, along with the homologous assembled zebra finch sequence, to examine the differences

between species and morphs (Supplementary Fig. 4).

**Test for the presence of an inversion using long-range PCR and *de novo* sequencing of the *Red***

**locus using the Illumina MiSeq**: If there is an inversion breakpoint between two neighbouring loci,

then the inverted chromosome will not produce amplicons across the breakpoint in a PCR that

produces an amplicon in the non-inverted chromosome. To test for an inversion across the *Red* locus,

19 contiguous, overlapping fragments were amplified in 12 females ($n_{black} = 6$, $n_{red} = 6$) from the wild

population by combining each forward primer with a reverse primer designed within the adjacent

amplicon (Supplementary Table 6) and using the LongRange PCR kit (QIAGEN). When the primer

sets originally designed from zebra finch sequences failed to produce long amplicons, we designed

new primers using the Gouldian finch sequences obtained from previous sequencing efforts to ensure

the failure was not due to mismatching sequences in the primer-binding sites (Supplementary Table

6). We confirmed that the long fragments amplified the correct sites by sequencing the boundaries of

each long fragment using PCR primers and comparing with the zebra finch reference genome.

Long amplicons across the *Red* locus from the above PCR experiments and short amplicons

for the reference loci (see below) were labelled by individual, pooled and sequenced using the

Illumina MiSeq platform. All traces of adapter sequences added to amplicons during paired-end short-

read (250-bp) sequencing were removed using a custom Perl script (TagCle v0.80, available upon

request from K.-W.K.). *De novo* assemblies were performed for each individual using NGen.

Supercontigs were produced by BLAST searching the contig sequences against the zebra finch

reference sequence and comparing the positions of the contigs and the overlapping sequences. The

assembled sequences were aligned using UGENE v1.12.3[86] and manually edited using BioEdit

v7.1.11[87]. These data were used to measure DNA polymorphism across the *Red* locus and in

coalescent simulations to test for selection (see below).

**Transcriptome analysis**: We tested for differential gene expression in the feather follicles between

the Gouldian finch morphs. First, we sampled regenerating head feathers from birds held by breeders

in Australia during the annual moulting period and stored them directly in RNAlater (Qiagen). The

feathers were subsequently frozen solid with liquid nitrogen, homogenized using a pestle and mortar,

and RNA was extracted using TRIzol (Thermo Fisher Scientific, Waltham, Massachusetts, United

States) and RNeasy Mini kits (Qiagen). The quality of the RNA was measured on a Bioanalyzer

(Agilent, Santa Clara, California, United States) and high-quality samples with an RNA integrity

value > 7 were used in microarray analysis. We obtained the gene expression profiles of 6 birds

($n_{black\_male} = 3$, $n_{red\_male} = 2$, $n_{red\_female} = 1$) using the Affymetrix zebra finch gene array ST1.0 that

contains probe sets from 18,595 genes. Gene expression was normalized using a robust multi-array

average algorithm implemented in the Affymetrix Expression Console v1.3. The expression

differences were tested using one-way ANOVA implemented in Transcriptome Analysis Console

v2.0 (Affymetrix). Because of the small sample size, no gene passed the threshold required to correct

for the false discovery rate. Therefore, we used uncorrected $P$-values < 0.05 and $\log_2$-fold changes ≥

1.3 to select 304 potentially differentially expressed (DE) genes. Human orthologues of DE genes

were obtained from UniProt (www.uniprot.org). A functionally grouped annotation network for the

DE genes was constructed using the Cytoscape plugin ClueGO v2.3.5[88]. Functionally related Gene

Ontology (GO) terms for biological processes in human (23.02.2017) were grouped based on a kappa

score > 0.4 with network specificity of 3 to 5. The significance was examined using a two-sided

hypergeometric test and the false discovery rate was corrected using the Benjamini–Hochberg

method[89].

**Evolutionarily conserved elements within the *Red* locus**: We used PhastCon values[90] to examine

the presence of the evolutionarily conserved elements within the *Red* locus that might represent

regulatory regions potentially responsible for the expression of a candidate gene underlying colour

polymorphism. The position of human and mouse transcripts for *MOCS2* and *FST*, and the pre-

calculated PhastCon values across the *Red* locus, were obtained from the UCSC Genome Browser

Gateway (https://genome.ucsc.edu/cgi-bin/hgGateway). PhastCons values were based on multi-

genome alignments among three vertebrates; medium ground finch (*Geospiza fortis*), human (*Homo*

*sapiens*) and mouse (*Mus musculus*), and among five birds: medium ground finch, zebra finch,

budgerigar (*Melopsittacus undulatus*), chicken (*Gallus gallus*) and turkey (*Meleagris gallopavo*). The

records were visualized using *rtracklayer*[91] and *Gviz*[92]. Genomic positions of the *Red* locus and the

core region in the genome of the medium ground finch were determined using a BLAT search in the

UCSC.

**Genomic regions under consideration in the test for neutrality**: We obtained a 99,669-bp stretch

of sequence using the Illumina MiSeq (see above) encompassing the *Red* locus (46,503,706–

46,601,744 bp in zebra finch, Supplementary Table 6) in 12 female birds ($n_{black}$ = 6, $n_{red}$ = 6) from the

wild population. Because females are hemizygous at the *Red* locus, these sequences represent six each

of the black ($Z^r$) and red ($Z^R$) alleles. We defined the entire ~100 kbp sequenced region as the '*Red*

locus' and the central region with $F_{ST} > 0.8$ (Fig. 2d) as the 'core' region in subsequent analyses.

472           In an additional MiSeq run, we also randomly chose 24 loci (henceforth the 'reference' loci,

Supplementary Table 8) across the Z chromosome, outside of the LD block of the *Red* locus, to

measure the background level of diversity for comparison with that seen at the *Red* locus. These loci

consisted of single introns, sampled in 11–12 female individuals, 10 of which (5 each of black and

red) were also represented in our sample of 12 birds at the *Red* locus.

**Estimates of DNA polymorphism and divergence**: In order to compare diversity at the *Red* locus

(combining the $Z^r$ and $Z^R$ alleles) with that at the reference loci using the MiSeq data, we calculated $\pi$

and Tajima's *D* for the whole sample at each locus using a custom R script. We then calculated the

same statistics within each allelic class (we define two allelic classes: one each for birds having the $Z^r$

or the $Z^R$ allele at the *Red* locus, respectively) at the reference loci and the *Red* locus, in order to

compare polymorphism for the combined sample with that within each allelic class. In addition, we

calculated $F_{ST}$ and $D_{XY}$ between the two allelic classes at the reference and the *Red* loci using custom

R and Python scripts. We used Weir and Cockerham (1984)'s definition of $F_{ST}$[51], and the estimator

provided by Hudson *et al*. (1992)[93] to calculate site-by-site $F_{ST}$ values (Supplementary Fig. 1). To

obtain a composite estimate of $F_{ST}$ from multiple SNPs for each window, we used equation (6) of

Jackson *et al*. (2015; see also Bhatia *et al*. 2013)[94,95]. In the estimation of $F_{ST}$, $D_{XY}$ and $\pi$ at the *Red*

locus using sliding-window analyses (window size of 1,000 bp with a step of 500 bp; Figs. 2-3), we

retained sites with missing data to maximise the number of SNPs in each window. However, because

Tajima's *D* cannot be calculated when data are missing, we removed individuals with missing data on

a *per window* basis for this statistic.

We define statistics that refer to subsets of the data using subscripts: for example, $\pi_{red}$ refers to

the nucleotide diversity among the alleles of red birds, and $\pi_{total}$ refers to nucleotide diversity among

the alleles of both black and red birds combined. When conducting point estimates of summary

statistics *per locus* we removed sites with missing data in any individual. To test for the statistical

significance of the point estimate of Tajima's *D*, we used a coalescent simulation under the standard

neutral model implemented in DnaSP v5[47].

To examine the genetic structure with regards to colour polymorphism, we constructed a

"coancestry matrix" for samples for which we had RADSeq data that summarises the nearest-

neighbour haplotype relationship and provides a more sensitive inference on population structure than

the STRUCTURE and PCA approaches[52,53,60] (Supplementary Fig. 6). We selected autosomal

RADSeq loci with ≤ 5 SNPs (total 18,719 RADSeq loci) and individuals with ≤ 20% missing data.

We used fineRADstructure[52] to obtain co-ancestry coefficients between individuals and data

missingness for each individual. Two individuals had significantly high levels of missing data and

were excluded from calculation of the coancestry matrix (Supplementary Fig. 6).

To examine the differentiation between morphs in all wild-population individuals, we used

nine autosomal microsatellite markers from a set of previously-tested cross-species markers[73]

(Supplementary Table 10) and genotyped 161 birds ($n_{black}$ = 126, and $n_{red}$ = 35) using the ABI3730

DNA Analyzer. Deviation of microsatellite genotypes from HWE was examined using CERVUS

v3.0.3[96]. A model-based clustering method implemented in STRUCTURE v2.3.4[55] was used to infer

the number of subpopulations (*K*) and to assign individuals to populations (length of burnin period =

500,000; number of MCMC repeats after burnin = 750,000; 20 iterations for each run, Supplementary

Fig. 7).

**Considering non-random sampling at the *Red* locus in the test for selection**: It is important to take

into account non-random sampling with respect to the *Red* locus in simulations to generate

polymorphism data under the null hypothesis of neutrality because, even under neutrality, it is

possible to obtain high $F_{ST}$ values, as well as elevated $\pi$ at a focal site, if the sampling scheme is

conditioned on that site, as was the case here. In order to replicate our observed sampling scheme for

the purpose of statistical testing, we used the allele frequencies from the genotypes of 161 wild birds

at the *Red* locus for this purpose (Supplementary Table 3). As we do not know which allele ($Z^r$ or $Z^R$)

is ancestral, it was necessary to consider both scenarios in turn; i.e. $Z^r$ is derived (derived allele

frequency, DAF = 0.856); or $Z^R$ is derived (DAF = 0.144) (see Supplementary Text for details about

allele frequency estimation). This is, in part, because we cannot be sure which site(s) within the *Red*

locus is causal, and so cannot polarise that site using an outgroup sequence. In addition, we lack

outgroup sequence data for a portion of the *Red* locus that coincides with a putative TE insertion in

the Gouldian finch lineage, but which does not exist at the same location in the zebra finch genome.

As such, for the majority of the simulations and analyses below, we arbitrarily set the focal site at the

centre of the *Red* locus. However, our results are robust to different choices of this parameter (see

Supplementary Text).

**Testing for selection**: We employed two widely-used methods to test for evidence of natural

selection. First, we used coalescent simulations to generate polymorphism data under neutrality at a

locus that is equivalent to the real *Red* locus in Gouldian finches in terms of its allele frequencies in

nature (see above), the sampling scheme (6 each of red and black birds) and relevant population

genetic parameters (the population mutation rate, $\theta$ and the population recombination rate, $\rho$). By

assessing whether the observed polymorphism at the *Red* locus is compatible with these simulations,

we can accept or reject the null hypothesis of neutrality. We used the programs SelSim[97] and mbs[98] to

simulate polymorphism linked to a focal site at which a mutation has arisen and spread by genetic

drift to the frequencies we observed at the *Red* locus (0.144 or 0.856). These programs allowed us to

replicate the allele frequencies, sampling scheme and population genetic parameters (see above)

observed at the *Red* locus and obtain expectations under neutrality. In addition, mbs can model

demographic changes. We also used a modified version of the HKA test[57] to test for selection. See

Supplementary Text for more detailed methods.

**Gene genealogy and relative rate test**: To infer the evolutionary history of the haplotypes at the *Red*

locus we constructed gene genealogies and compared the level of evolutionary divergence between

the black and red haplotypes with those of orthologous sequences in other species. First, blood

samples of 14 species in the family *Estrildidae* including red-faced (*E. pealii*, *E. psittacae*) or blue-

faced (*E. tricolor and E. trichroa*) *Erythrura* species were obtained from an aviculturist. Blood

samples were collected and stored using Whatman FTA cards (GE Healthcare, Little Chalfont, UK)

and DNA was extracted following the manufacturer's instructions. Sequences of the fragments

amplifiable using the *M1–M3* primers (Supplementary Table 2) were obtained using Sanger

sequencing. For zebra finch, the sequences from the reference genome assembly were used.

We tested alternative nucleotide substitution models using the R package, *phangorn* v2.3.1[99],

which suggested GTR+Γ+I to be the best model for the sequence of the entire core region of the *Red*

locus. Maximum-likelihood trees were then constructed using this model and tested with 1,000

bootstraps using *phangorn* and visualised using *ggtree* v1.4.20[100]. To examine if the molecular clock

hypothesis applied to the core of the *Red* locus, we used Tajima's nonparametric relative rate test[101]

implemented in pegas v0.10[102].

**References**

1    Galeotti, P., Rubolini, D., Dunn, P. O. & Fasola, M. Colour polymorphism in birds: causes
and functions. *J. Evolution. Biol.* **16**, 635-646 (2003).

2    Hugall, A. F. & Stuart-Fox, D. Accelerated speciation in colour-polymorphic birds. *Nature*
**485**, 631-634 (2012).

3    Roulin, A. The evolution, maintenance and adaptive function of genetic colour polymorphism
in birds. *Biol. Rev.* **79**, 815-848 (2004).

4    Gray, S. M. & McKinnon, J. S. Linking color polymorphism maintenance and speciation.
*Trends Ecol. Evol.* **22**, 71-79 (2007).

5    Hill, G. E. & McGraw, K. J. *Bird Coloration, Vol.I. Mechanisms and Measurements*. Vol. 1
(Harvard University Press, 2006).

6    Hubbard, J. K., Uy, J. A. C., Hauber, M. E., Hoekstra, H. E. & Safran, R. J. Vertebrate
pigmentation: from underlying genes to adaptive function. *Trends Genet.* **26**, 231-239 (2010).

7    Mundy, Nicholas I. *et al.* Red carotenoid coloration in the zebra finch is controlled by a
cytochrome P450 gene cluster. *Curr. Biol.* **26**, 1435-1440 (2016).

8    Toews, D. P. L., Hofmeister, N. R. & Taylor, S. A. The evolution and genetics of carotenoid
processing in animals. *Trends Genet.* **33**, 171-182 (2017).

9    Toomey, M. B. *et al.* High-density lipoprotein receptor SCARB1 is required for carotenoid
coloration in birds. *Proc. Natl. Acad. Sci. U.S.A.* (2017).

10   Cooke, T. F. *et al.* Genetic mapping and biochemical basis of yellow feather pigmentation in
budgerigars. *Cell* **171**, 427-439.e421 (2017).

11   Southern, H. N. Polymorphism in *Poephila gouldiae* Gould. *J. Genet.* **47**, 51-57 (1945).

12   Brush, A. H. & Seifried, H. Pigmentation and feather structure in genetic variants of the
Gouldian finch, *Poephila gouldiae*. *The Auk* **85**, 416-430 (1968).

13   Cooke, F. & Buckley, P. A. *Avian genetics: A population and ecological approach*.
(Academic Press, 1987).

14   Franklin, D. C. & Dostine, P. L. A note on the frequency and genetics of head colour morphs
in the Gouldian finch. *Emu* **100**, 236-239 (2000).

15   Bolton, P. E. *et al.* The colour of paternity: extra-pair paternity in the wild Gouldian finch
does not appear to be driven by genetic incompatibility between morphs. *J. Evolution. Biol.*
**30**, 174-190 (2017).

16   Gilby, A. J., Pryke, S. R. & Griffith, S. C. The historical frequency of head-colour morphs in
the Gouldian finch (*Erythrura gouldiae*). *Emu* **109**, 222-229 (2009).

17   Darlington, C. D. & Mather, K. *Elements of Genetics*.  (George Allen & Unwin Ltd, 1949).

18   Lamichhaney, S. *et al.* Structural genomic changes underlie alternative reproductive strategies
in the ruff (Philomachus pugnax). *Nat. Genet.* **48**, 84-88 (2016).

19  Küpper, C. *et al.* A supergene determines highly divergent male reproductive morphs in the
ruff. *Nat. Genet.* **48**, 79-83 (2016).

20  Tuttle, Elaina M. *et al.* Divergence and functional degradation of a sex chromosome-like
supergene. *Curr. Biol.* **26**, 344-350 (2016).

21  Kim, K.-W. *et al.* A sex-linked supergene controls sperm morphology and swimming speed in
a songbird. *Nat. Ecol. Evol.* **1**, 1168-1176 (2017).

22  Ducrest, A.-L., Keller, L. & Roulin, A. Pleiotropy in the melanocortin system, coloration and
behavioural syndromes. *Trends Ecol. Evol.* **23**, 502-510 (2008).

23  Baird, N. A. *et al.* Rapid SNP discovery and genetic mapping using sequenced RAD markers.
*PLoS ONE* **3**, e3376 (2008).

24  Kim, K. W., Griffith, S. C. & Burke, T. Linkage mapping of a polymorphic plumage locus
associated with intermorph incompatibility in the Gouldian finch (*Erythrura gouldiae*).
*Heredity* **116**, 409-416 (2016).

25  Carneiro, M., Ferrand, N. & Nachman, M. W. Recombination and speciation: Loci near
centromeres are more differentiated than loci near telomeres between subspecies of the
European rabbit (*Oryctolagus cuniculus*). *Genetics* **181**, 593-606 (2009).

26  Wang, J. *et al.* A Y-like social chromosome causes alternative colony organization in fire
ants. *Nature* **493**, 664-668 (2013).

27  Schwander, T., Libbrecht, R. & Keller, L. Supergenes and complex phenotypes. *Curr. Biol.*
**24**, R288-R294 (2014).

28  Ohyama, A., Saito, F., Ohuchi, H. & Noji, S. Differential expression of two BMP antagonists,
*gremlin* and *Follistatin*, during development of the chick feather bud. *Mech. Develop.* **100**,
331-333 (2001).

29  McDowall, M., Edwards, N. M., Jahoda, C. A. B. & Hynd, P. I. The role of activins and
follistatins in skin and hair follicle development and function. *Cytokine Growth F. R.* **19**, 415-
426 (2008).

30  Bilodeau, M. L. *et al.* BMP-2 stimulates *tyrosinase* gene expression and melanogenesis in
differentiated melanocytes. *Pigm. Cell Res.* **14**, 328-336 (2001).

31  Sharov, A. A. *et al.* Bone morphogenetic protein (BMP) signaling controls hair pigmentation
by means of cross-talk with the melanocortin receptor-1 pathway. *Proc. Natl. Acad. Sci.*
*U.S.A.* **102**, 93-98 (2005).

32  Moustakas, A. TGF-β targets *PAX3* to control melanocyte differentiation. *Dev. Cell* **15**, 797-
799 (2008).

33  Yaar, M. *et al.* Bone Morphogenetic Protein-4, a novel modulator of melanogenesis. *J. Biol.*
*Chem.* **281**, 25307-25314 (2006).

34  Murakami, M., Matsuzaki, F. & Funaba, M. Regulation of melanin synthesis by the TGF-β
family in B16 melanoma cells. *Mol. Biol. Rep.* **36**, 1247-1250 (2009).

35 Chuong, C.-M., Chodankar, R., Widelitz, R. B. & Jiang, T.-X. Evo-Devo of feathers and
scales: building complex epithelial appendages: Commentary. *Curr. Opin. Genet. Dev.* **10**,
449-456 (2000).

36 Harris, M. P., Fallon, J. F. & Prum, R. O. *Shh-Bmp2* signaling module and the evolutionary
origin and diversification of feathers. *J. Exp. Zool.* **294**, 160-176 (2002).

37 Yu, M., Wu, P., Widelitz, R. B. & Chuong, C.-M. The morphogenesis of feathers. *Nature*
**420**, 308-312 (2002).

38 Belecky-Adams, T. L., Scheurer, D. & Adler, R. Activin family members in the developing
chick retina: expression patterns, protein distribution, and *in vitro* effects. *Dev. Biol.* **210**, 107-
123 (1999).

39 Moreira, E. F. & Adler, R. Effects of follistatin overexpression on cell differentiation in the
chick embryo retina. *Dev. Biol.* **298**, 272-284 (2006).

40 Knight, P. G. Roles of inhibins, activins, and follistatin in the female reproductive system.
*Front. Neuroendocrin.* **17**, 476-509 (1996).

41 Guo, Q. *et al.* Overexpression of mouse Follistatin causes reproductive defects in transgenic
mice. *Mol. Endocrinol.* **12**, 96-106 (1998).

42 Knight, P. G. & Glister, C. Potential local regulatory functions of inhibins, activins and
follistatin in the ovary. *Reproduction* **121**, 503-512 (2001).

43 Yao, H. H. C. *et al. Follistatin* operates downstream of *Wnt4* in mammalian ovary
organogenesis. *Dev. Dynam.* **230**, 210-215 (2004).

44 Toews, David P. L. *et al.* Plumage genes and little else distinguish the genomes of hybridizing
warblers. *Curr. Biol.* **26**, 2313-2318 (2016).

45 Nei, M. *Molecular Evolutionary Genetics*. (Columbia Univ. Press, 1987).

46 Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics* **123**, 585-595 (1989).

47 Librado, P. & Rozas, J. DnaSP v5: a software for comprehensive analysis of DNA
polymorphism data. *Bioinformatics* **25**, 1451-1452 (2009).

48 Pryke, S. R. & Griffith, S. C. The relative role of male vs. female mate choice in maintaining
assortative pairing among discrete colour morphs. *J. Evolution. Biol.* **20**, 1512-1521 (2007).

49 Pryke, S. R. & Griffith, S. C. Postzygotic genetic incompatibility between sympatric color
morphs. *Evolution* **63**, 793-798 (2009).

50 Wu, C.-I. & Ting, C.-T. Genes and speciation. *Nat. Rev. Genet.* **5**, 114-122 (2004).

51 Weir, B. S. & Cockerham, C. C. Estimating F-statistics for the analysis of population
structure. *Evolution* **38**, 1358-1370 (1984).

52 Malinsky, M., Trucchi, E., Lawson, D. J. & Falush, D. RADpainter and fineRADstructure:
Population inference from RADseq data. *Mol. Biol. Evol.* **35**, 1284-1290 (2018).

53   Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of population structure using
dense haplotype data. *PLoS Genet.* **8**, e1002453 (2012).

54   Evans, B. J., Zeng, K., Esselstyn, J. A., Charlesworth, B. & Melnick, D. J. Reduced
representation genome sequencing suggests low diversity on the sex chromosomes of
Tonkean macaque monkeys. *Mol. Biol. Evol.* **31**, 2425-2440 (2014).

55   Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using
multilocus genotype data. *Genetics* **155**, 945-959 (2000).

56   Charlesworth, D. Balancing selection and its effects on sequences in nearby genome regions.
*PLoS Genet.* **2**, e64 (2006).

57   Hudson, R. R., Kreitman, M. & Aguadé, M. A test of neutral molecular evolution based on
nucleotide data. *Genetics* **116**, 153-159 (1987).

58   Hof, A. E. v. t. *et al.* The industrial melanism mutation in British peppered moths is a
transposable element. *Nature* **534**, 102 (2016).

59   Garrigan, D., Mobasher, Z., Kingan, S. B., Wilder, J. A. & Hammer, M. F. Deep haplotype
divergence and long-range linkage disequilibrium at Xp21.1 provide evidence that humans
descend from a structured ancestral population. *Genetics* **170**, 1849-1856 (2005).

60   Stryjewski, K. F. & Sorenson, M. D. Mosaic genome evolution in a recent and rapid avian
radiation. *Nat. Ecol. Evol.* **1**, 1912–1922 (2017).

61   Bowman, D. M. J. S. *et al.* Biogeography of the Australian monsoon tropics. *J. Biogeogr.* **37**,
201-216 (2010).

62   Pryke, S. R. & Griffith, S. C. Red dominates black: agonistic signalling among head morphs
in the colour polymorphic Gouldian finch. *Proc. R. Soc. B* **273**, 949-957 (2006).

63   Brazill-Boast, J., Griffith, S. & Pryke, S. Morph-dependent resource acquisition and fitness in
a polymorphic bird. *Evol. Ecol.* **27**, 1189-1198 (2013).

64   Pryke, S. R., Astheimer, L. B., Buttemer, W. A. & Griffith, S. C. Frequency-dependent
physiological trade-offs between competing colour morphs. *Biol. Letters* **3**, 494-497 (2007).

65   Pryke, S. R. & Griffith, S. C. Socially mediated trade-offs between aggression and parental
effort in competing color morphs. *Am. Nat.* **174**, 455-464 (2009).

66   Pryke, S. R., Astheimer, L. B., Griffith, S. C. & Buttemer, W. A. Covariation in life-history
traits: differential effects of diet on condition, hormones, behavior, and reproduction in
genetic finch morphs. *Am. Nat.* **179**, 375-390 (2012).

67   Arnqvist, G. & Rowe, L. *Sexual conflict.* (Princeton University Press, 2005).

68   Wellenreuther, M., Svensson, E. I. & Hansson, B. Sexual selection and genetic colour
polymorphisms in animals. *Mol. Ecol.* **23**, 5398-5414 (2014).

69   Svensson, E. I. Back to basics: using colour polymorphisms to study evolutionary processes.
*Mol. Ecol.* **26**, 2204-2211 (2017).

70   Price, T. D. *Speciation in birds.* (Roberts and Company, 2008).

71    Bruford, M. W., Hanotte, O., Brookfield, J. F. Y. & Burke, T. Multilocus and single-locus
DNA fingerprinting in *Molecular genetic analysis of populations: A practical approach* (ed
A. R. Hoelzel) 225-269 (IRL Press, 1998).

72    Dawson, D. A. *Genomic analysis of passerine birds using conserved microsatellite loci* PhD
thesis, University of Sheffield, UK, (2007).

73    Kim, K.-W. *et al.* Four-way development of microsatellite markers for the Gouldian finch
(*Erythrura gouldiae*). *Conserv. Genet. Resour.* **7**, 899-907 (2015).

74    Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W. & Postlethwait, J. H. Stacks: building
and genotyping loci de novo from short-read sequences. *G3-Genes Genom. Genet.* **1**, 171-182
(2011).

75    Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987-
2993 (2011).

76    Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158
(2011).

77    Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-based
linkage analyses. *Am. J. Hum. Genet.* **81**, 559-575 (2007).

78    Clarke, G. M. *et al.* Basic statistical analysis in genetic case-control studies. *Nat. Protoc.* **6**,
121-133 (2011).

79    Lindgreen, S. AdapterRemoval: easy cleaning of next-generation sequencing reads. *BMC Res.*
*Notes* **5**, 337 (2012).

80    Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357 (2012).

81    McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res.* **20**, 1297-1303 (2010).

82    Graffelman, J. & Weir, B. S. Testing for Hardy–Weinberg equilibrium at biallelic genetic
markers on the X chromosome. *Heredity* **116**, 558 (2016).

83    Graffelman, J. Exploring diallelic genetic markers: The HardyWeinberg package. *Journal of*
*Statistical Software* **64**, 1-23 (2015).

84    Barrett, J. C., Fry, B., Maller, J. & Daly, M. J. Haploview: analysis and visualization of LD
and haplotype maps. *Bioinformatics* **21**, 263-265 (2005).

85    Shin, J.-H., Blay, S., McNeney, B. & Graham, J. LDheatmap: An R function for graphical
display of pairwise linkage disequilibria between single nucleotide polymorphisms. *J. Stat.*
*Softw.* **16**, 9 (2006).

86    Okonechnikov, K., Golosova, O. & Fursov, M. Unipro UGENE: a unified bioinformatics
toolkit. *Bioinformatics* **28**, 1166-1167 (2012).

87    Hall, T. A. BioEdit: a user-friendly biological sequence alignment editor and analysis
program for Windows 95/98/NT. *Nucl. Acid. S.* **41**, 95–98 (1999).

88    Bindea, G. *et al.* ClueGO: a Cytoscape plug-in to decipher functionally grouped gene
ontology and pathway annotation networks. *Bioinformatics* **25**, 1091-1093 (2009).

89    Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**, 289-300 (1995).

90    Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
genomes. *Genome Res.* **15**, 1034-1050 (2005).

91    Lawrence, M., Gentleman, R. & Carey, V. rtracklayer: an R package for interfacing with
genome browsers. *Bioinformatics* **25**, 1841-1842 (2009).

92    Hahne, F. & Ivanek, R. Visualizing genomic data using Gviz and Bioconductor in *Statistical
Genomics: Methods and Protocols* Vol. 1418    335-351 (Humana Press, 2016).

93    Hudson, R. R., Slatkin, M. & Maddison, W. P. Estimation of levels of gene flow from DNA
sequence data. *Genetics* **132**, 583-589 (1992).

94    Jackson, B. C., Campos, J. L. & Zeng, K. The effects of purifying selection on patterns of
genetic differentiation between Drosophila melanogaster populations. *Heredity* **114**, 163-174
(2015).

95    Bhatia, G., Patterson, N., Sankararaman, S. & Price, A. L. Estimating and interpreting $F_{ST}$:
The impact of rare variants. *Genome Res.* **23**, 1514-1521 (2013).

96    Kalinowski, S. T., Taper, M. L. & Marshall, T. C. Revising how the computer program
Cervus accommodates genotyping error increases success in paternity assignment. *Mol. Ecol.*
**16**, 1099-1106 (2007).

97    Spencer, C. C. A. & Coop, G. SelSim: a program to simulate population genetic data with
natural selection and recombination. *Bioinformatics* **20**, 3673-3675 (2004).

98    Teshima, K. M. & Innan, H. mbs: modifying Hudson's ms software to generate samples of
DNA sequences with a biallelic site under selection. *BMC Bioinformatics* **10**, 166 (2009).

99    Schliep, K. P. phangorn: phylogenetic analysis in R. *Bioinformatics* **27**, 592-593 (2011).

100    Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. ggtree: an r package for
visualization and annotation of phylogenetic trees with their covariates and other associated
data. *Methods Ecol. Evol.* **8**, 28-36 (2017).

101    Tajima, F. Simple methods for testing the molecular evolutionary clock hypothesis. *Genetics*
**135**, 599-607 (1993).

102    Paradis, E. pegas: an R package for population genetics with an integrated–modular approach.
*Bioinformatics* **26**, 419-420 (2010).

103    Ashique, A. M., Fu, K. & Richman, J. M. Signalling via type IA and type IB bone
morphogenetic protein receptors (BMPR) regulates intramembranous bone formation,

chondrogenesis and feather formation in the chicken embryo. *Int. J. Dev. Biol.* **46**, 243 - 253
(2002).

104   Kawano, M., Suzuki, S., Suzuki, M., Oki, J. & Imamura, T. Bulge- and basal layer-specific
expression of fibroblast growth factor-13 (FHF-2) in mouse skin. *J. Invest. Dermatol.* **122**,
1084-1090 (2004).

105   Kobielak, K., Stokes, N., de la Cruz, J., Polak, L. & Fuchs, E. Loss of a quiescent niche but
not follicle stem cells in the absence of bone morphogenetic protein signaling. *Proc. Natl.*
*Acad. Sci. U.S.A.* **104**, 10063-10068 (2007).

106   Menon, G. K. & Menon, J. Avian epidermal lipids: functional considerations and relationship
to feathering. *Am. Zool.* **40**, 540-552 (2000).

107   Ni, Y. *et al.* Astaxanthin prevents and reverses diet-induced insulin resistance and
steatohepatitis in mice: A comparison with vitamin E. *Sci. Rep.* **5**, 17192 (2015).

108   Jia, Y., Wu, C., Kim, J., Kim, B. & Lee, S.-J. Astaxanthin reduces hepatic lipid accumulations
in high-fat-fed C57BL/6J mice via activation of peroxisome proliferator-activated receptor
(PPAR) alpha and inhibition of PPAR gamma and Akt. *J. Nutr. Biochem.* **28**, 9-18 (2016).

109   Stradi, R. Pigmenti e sistematica degli uccelli in *Colori in volo: il piumaggio degli uccelli*
(eds L. Brambilla, G. Canali, & E. Mannucci) 117-146 (Universita degli Studi di Milano,
1999).

110   Völker, O. Die gelben mutanten des Rotbauchwürgers (*Laniarius atrococcineus*) und der
Gouldamadine (*Chloebia gouldiae*) in biochemischer sicht. *J. Ornithol.* **105**, 186-189 (1964).

111   Pryke, S. R. Fiery red heads: female dominance among head color morphs in the Gouldian
finch. *Behav. Ecol.* **18**, 621-627 (2007).

112   Williams, L. J., King, A. J. & Mettke-Hofmann, C. Colourful characters: head colour reflects
personality in a social bird, the Gouldian finch, *Erythrura gouldiae*. *Anim. Behav.* **84**, 159-
165 (2012).

113   Mettke-Hofmann, C. Head colour and age relate to personality traits in Gouldian finches.
*Ethology* **118**, 906-916 (2012).

114   Immler, S., Pryke, S. R., Birkhead, T. R. & Griffith, S. C. Pronounced within-individual
plasticity in sperm morphometry across social environments. *Evolution* **64**, 1634-1643 (2010).

115   Fox, S., Johnson, C. N., Brooks, R. & Lewis, M. J. Polymorphism, mate choice and sexual
selection in the Gouldian finch (*Erythrura gouldiae*). *Aust. J. Zool.* **50**, 125-134 (2002).

116   Pryke, S. R. & Griffith, S. C. Maternal adjustment of parental effort in relation to mate
compatibility affects offspring development. *Behav. Ecol.* **21**, 226-232 (2010).

117   Pryke, S. R. & Griffith, S. C. Genetic incompatibility drives sex allocation and maternal
investment in a polymorphic finch. *Science* **323**, 1605-1607 (2009).

118   Tajima, F. The effect of change in population size on DNA polymorphism. *Genetics* **123**,
597-601 (1989).

119  Zeng, K., Fu, Y.-X., Shi, S. & Wu, C.-I. Statistical tests for detecting positive selection by
utilizing high-frequency variants. *Genetics* **174**, 1431-1439 (2006).

120  Eyre-Walker, A. Evidence of selection on silent site base composition in mammals: potential
implications for the evolution of isochores and junk DNA. *Genetics* **152**, 675-683 (1999).

121  Duret, L. & Galtier, N. Biased gene conversion and the evolution of mammalian genomic
landscapes. *Annu. Rev. Genomics Hum. Genet.* **10**, 285-311 (2009).

122  Zeng, K. The application of population genetics in the study of codon usage bias in *Codon*
*evolution: mechanisms and models*  (eds G.M. Cannarozzi & A. Schneider)  245–254
(Oxford University Press, 2012).

123  Backström, N., Zhang, Q. & Edwards, S. V. Evidence from a house finch (*Haemorhous*
*mexicanus*) spleen transcriptome for adaptive evolution and biased gene conversion in
passerine birds. *Mol. Biol. Evol.* **30**, 1046-1050 (2013).

124  Weber, C. C., Boussau, B., Romiguier, J., Jarvis, E. D. & Ellegren, H. Evidence for GC-
biased gene conversion as a driver of between-lineage differences in avian base composition.
*Genome Biol.* **15**, 549 (2014).

125  Bolívar, P., Mugal, C. F., Nater, A. & Ellegren, H. Recombination rate variation modulates
gene sequence evolution mainly via GC-biased gene conversion, not Hill–Robertson
interference, in an avian system. *Mol. Biol. Evol.* **33**, 216-227 (2016).

126  Zeng, K. & Charlesworth, B. Estimating selection intensity on synonymous codon usage in a
nonequilibrium population. *Genetics* **183**, 651-662 (2009).

127  Haddrill, P. R., Zeng, K. & Charlesworth, B. Determinants of synonymous and
nonsynonymous variability in three species of Drosophila. *Mol. Biol. Evol.* **28**, 1731-1743
(2011).

128  Auton, A. & McVean, G. Recombination rate estimation in the presence of hotspots. *Genome*
*Res.* **17**, 1219-1227 (2007).

129  Chan, A. H., Jenkins, P. A. & Song, Y. S. Genome-wide fine-scale recombination rate
variation in *Drosophila melanogaster*. *PLoS Genet.* **8**, e1003090 (2012).

130  Hudson, R. R. & Kaplan, N. L. Statistical properties of the number of recombination events in
the history of a sample of DNA sequences. *Genetics* **111**, 147-164 (1985).

131  Myers, S. R. & Griffiths, R. C. Bounds on the minimum number of recombination events in a
sample history. *Genetics* **163**, 375-394 (2003).

132  Nordborg, M. Structured coalescent processes on different time scales. *Genetics* **146**, 1501-
1514 (1997).

133  Hudson, R. R. Gene genealogies and the coalescent process. *Oxford Surv. Evol. Biol.* **7**, 1-44
(1990).

**Supplementary Information**: Supplementary Figures: 16, Tables: 14, Text: 1

**Supplementary Dataset**: One Excel file summarising gene expression analysis.

**Acknowledgements**

We thank Brian and Deborah Charlesworth for comments on an earlier manuscript. Sarah Pryke and David Harris provided blood samples of the Gouldian finches and other species of *Estrildidae*. Peri Bolton assisted with the collection of cheek feathers. We thank the Natural Environment Research Council Biomolecular Analysis Facility (NBAF) at the University of Edinburgh for sequencing RAD libraries and NBAF at the University of Liverpool for gene expression microarray scanning. This study was supported by a University of Sheffield scholarship to K.-W.K., a joint University of Sheffield/ Natural Environmental Research Council (NE/H524881/1, NE/K500914/1) Ph.D. studentship to B.C.J., a UK Department for Business, Innovation and Skills/ Ministry of Education of China UK–China Scholarships for Excellence studentship to H.Z., an NSERC Banting postdoctoral fellowship to D.P.L.T., an Australian Research Council (DP130100418) grant to S.C.G. and T.B., and a Leverhulme Fellowship to T.B.

**Author Contributions**

K.-W.K., B.C.J., S.C.G., K.Z. and T.B. conceived and designed the study. S.C.G. managed the work on Gouldian finches in the wild in Australia. K.-W.K., M.M.L. and A.D. constructed the RAD library. D.P.L.T, S.A.T., E.I.G and I.J.L. performed whole-genome sequencing. K.-W.K. and H.Z. performed molecular work. K.-W.K. and B.C.J. analysed the data. K.-W.K. and B.C.J. wrote the paper with contributions from all other authors.

**Author Information**

Compliance with Ethical Standards: *Samples*: Blood sampling of birds was conducted under the approval of the Animal Ethics Committees at the University of New South Wales and Macquarie University. Correspondence and requests for materials should be addressed to K.-W.K. (k.kim@sheffield.ac.uk).

**Data availability**

PLINK files for RADSeq analysis and sequence alignment of *de novo* sequencing for the *Red* locus

can be downloaded from Dryad (**doi:XXXXXXX**). Raw Illumina reads are available at NCBI SRA

under BioProject **XXXXXXX**. Microarray data are available on the Gene Expression Omnibus (GEO)

under accession number **GSE-XXXXXX**.

**Competing interests:** The authors declare no financial competing interests.

**Figure 1 | Identification of the *Red* locus. a,** Association between RADSeq markers and head colour in Gouldian finches ($n_{black}$ = 22, $n_{red}$ = 10). A circle around a dot indicates the only significant signal in GWAS ($P$ < 0.05, 1,000 permutations). The region shaded in green on the Z chromosome represents the candidate region (~7.2 cM) based on a previous linkage mapping study[24], where the *Red* locus co-segregated perfectly with microsatellite markers. **b,** Genotypes of 161 birds from a wild population (male: $n_{black}$ = 62, $n_{red}$ = 27; female: $n_{black}$ = 64, $n_{red}$ = 8) based on two contiguous SNPs in the core of the *Red* locus, where bar colours represent head colour. These SNPs (TG/CA) were identified by RADSeq (*M1* at 46,561,597–46,561,598 bp in the zebra finch) and were typed using allele-specific PCR (Supplementary Fig. 2, Supplementary Table 2). **c,** The extent of linkage disequilibrium (LD) associated with the colour trait. In the top panel, the locations of the protein-coding regions on the zebra finch Z chromosome (Mb) are indicated by vertical bars on the horizontal axis and the horizontal arrows below the gene names indicate the direction of transcription. Horizontal blue bars show the overlapping long-range PCR products that were subsequently sequenced and the entire span (~100 kbp), which includes the core region with elevated LD (central 72 kbp region with $F_{ST}$ > 0.8 in Fig. 2) and two short flanking regions, is defined as the *Red* locus in this study (see Fig. 2, Supplementary Tables 6-7). The three vertical red arrows indicate the locations of markers (*M1–M3*) that were used to genotype the wild birds in association tests (Supplementary Fig. 2, Supplementary Table 2). The middle panel shows -$\log_{10}P$ for association tests between phenotype and 282 SNPs from 26 fragments in 16 females ($n_{black}$ = 8, $n_{red}$ = 8) typed using Sanger sequencing. The lower panel shows pairwise LD ($r^2$) between SNPs.

**Figure 2 | Diversity and divergence between morph-determining haplotypes around the *Red***
**locus and between reference loci**. Position in **a-d** represents ~100 kbp alignment of MiSeq assembly
for 12 females ($n_{black}$ = 6, $n_{red}$ = 6), which corresponds to the position between 46,503,706–
46,601,744bp in the zebra finch reference genome (the region connected by horizontal blue bars in
Fig. 1c). **a-d** Sliding-window analysis (window size of 1,000 bp with a step of 500 bp) was used to
obtain estimates of summary statistics. Green shading indicates an insertion of a putative LTR
retrotransposon in the Gouldian finch that is not found in zebra finch. **a,** Nucleotide diversity ($\pi$) of
black (black line), red (red line) morphs and all samples combined (blue line). **b,** Sequence divergence
between zebra finch and each morph (ZB: black line; ZR: red line), and between morphs (BR: blue
line). **c,** Tajima's *D* for all samples combined. **d,** $F_{ST}$ between morphs. A horizontal dashed line
indicates $F_{ST}$ = 0.8. The data in **e-h** were based on 24 intronic reference loci on the Z chromosome,
with **e-g** showing the same summary statistics as **a-c** and **h** a histogram of $F_{ST}$. We define the entire
~100 kbp sequenced region as the '*Red* locus' and the central region with $F_{ST}$ > 0.8 in **d** as the 'core'
region in subsequent analyses.

**Figure 3 | Test for neutrality using coalescent simulations. a-d**, Shown are sliding-window

comparisons of the observed polymorphism at the *Red* locus with the polymorphism simulated under

a neutral model that incorporated the sampling scheme (6 alleles for each of the red and black

morphs), demography, recombination, and a derived allele frequency (DAF) of 0.144 at the focal site

(see Supplementary Text). Solid lines represent the observed data (*cf*. Fig. 2) and the grey-shaded

regions represent the space encompassed by the 95% confidence intervals derived from the

simulations. **a,** Nucleotide diversity for both morphs' alleles combined. **b,** $d_{XY}$ between the two

morphs' alleles. **c,** Tajima's *D* for both morphs' alleles combined. **d,** $F_{ST}$ between the two morphs'

alleles across the *Red* locus. **e**, A sliding-window implementation of the modified HKA test, taking

into account the observed sampling scheme, allele frequencies, mutation rate and recombination rate.

One minus the *P*-value is plotted for each window for DAF of 0.144 (yellow lines) and 0.856 (blue

lines), using the chi-squared distribution (dashed lines) and simulations (solid lines). The red

horizontal line indicates a *P*-value threshold of 0.05. The shaded grey bar indicates an insertion of a

putative LTR retrotransposon in the Gouldian finch that is not found in zebra finch.

**Figure 4 | Gene genealogy and estimates of evolutionary divergence between sequences in the core of the *Red* locus. a,** Illustrations of species in the genus *Erythrura*. From left, *E. psittacea*, *E. trichrora*, *E. pealii*, *E. tricolor*, and black and red morph of *E. gouldiae* (paintings courtesy of the *Handbook of the Birds of the World*, Lynx Edicions, Spain). **b,** Maximum-likelihood tree (GTR+Γ+I) for the species in the family *Estrildidae* using 774-bp alignment of *M1* (Fig. 1). Node support values (%) were generated from 1,000 bootstrap replicates and values >80% are shown next to the branches.