Genome-wide disruption of DNA methylation by 5-aza-2'-deoxycytidine in

the parasitoid wasp *Nasonia vitripennis*

Nicola Cook[1,¶,*], Darren J Parker[2,3,¶], Frances Turner[4], Eran Tauber[5], Bart A Pannebakker[6] and

David M Shuker[1]

1. School of Biology, University of St Andrews, UK

2. Department of Ecology and Evolution, University of Lausanne, Switzerland

3. Swiss Institute of Bioinformatics, Lausanne, Switzerland

4. Edinburgh Genomics, School of Biological Sciences, University of Edinburgh, UK

5. Faculty of Natural Sciences, University of Haifa, Israel

6. Laboratory of Genetics, Wageningen University, The Netherlands

*Corresponding author

E-mail: nc52@st-andrews.ac.uk

¶These authors contributed equally

Abstract

DNA methylation of cytosine residues across the genome influences how genes and phenotypes are regulated in a wide range of organisms. As such, understanding the role of DNA methylation and other epigenetic mechanisms has become very much a part of mapping genotype to phenotype, a major question in evolutionary biology. Ideally, we would like to manipulate DNA methylation patterns on a genome-wide scale, to help us to elucidate the role that epigenetic modifications play in phenotypic expression. Recently, the demethylating agent 5-aza-2'-deoxycytidine (5-aza-dC; commonly used in the epigenetic treatment of certain cancers), has been deployed to explore the epigenetic regulation of a number of traits of interest to evolutionary ecologists, including facultative sex allocation in the parasitoid wasp *Nasonia vitripennis*. In a recent study, we showed that treatment with 5-aza-dC did not ablate the facultative sex allocation response in *Nasonia*, but shifted the patterns of sex allocation in a way predicted by genomic conflict theory. This was the first (albeit indirect) experimental evidence for genomic conflict over sex allocation facilitated by DNA methylation. However, that work lacked direct evidence of the effects of 5-aza-dC on DNA methylation, and indeed the effect of the chemical has since been questioned in *Nasonia*. Here, using whole-genome bisulphite sequencing of more than 4 million CpGs, across more than 11,000 genes, we demonstrate unequivocally that 5-aza-dC disrupts methylation on a large scale across the *Nasonia vitripennis* genome. We show that the disruption can lead to both hypo- and hyper-methylation, may vary across tissues and time of sampling, and that the effects of 5-aza-dC are context- and sequence specific. We conclude that 5-aza-dC does indeed have the potential to be repurposed as a tool for studying the role of DNA methylation in evolutionary ecology, whilst many details of its action remain to be discovered.

Author Summary

Shedding light on the mechanistic basis of phenotypes is a major aim in the field of evolutionary biology. If we understand how phenotypes are controlled at the molecular level, we can begin to understand how evolution has shaped that phenotype and conversely, how genetic architecture may constrain trait evolution. Epigenetic markers (such as DNA methylation) also influence phenotypic expression by regulating how and when genes are expressed. Recently, 5-aza-2'-deoxycytidine (5-aza-dC), a hypomethylating agent used in the treatment of certain cancers, has been used to explore the epigenetic regulation of traits of interest to evolutionary ecologists. Previously, we used 5-aza-dC to validate a role for DNA methylation in facultative sex allocation behaviour in the parasitoid wasp *Nasonia vitripennis*. However, the direct effects of the chemical were not examined at that point and its efficacy in insects was questioned. Here, we demonstrate that 5-aza-dC disrupts DNA methylation

on a genome-wide scale in a context- and sequence-specific manner and results in both hypo- and

hyper-methylation. Our work demonstrates that 5-aza-dC has the potential to be repurposed as a tool

for studying the role of DNA methylation in phenotypic expression.

Introduction

The genomics revolution has transformed our understanding of how genomes build phenotypes. The

extraordinary diversity of morphologies, behaviours, and physiological adaptations we see across the

natural world are not necessarily the product of very different sets of genes, or very different numbers

of genes in organisms, but rather differences in how, where and when genes are expressed during

development. The genomics revolution has therefore been as much about understanding gene

regulation as it has been about identifying allelic variants at genetic loci [1, 2]. At the forefront of our

growing knowledge of gene regulation has been the role of epigenetic modification of DNA, including

DNA methylation. In the vast majority of cases, DNA methylation occurs at the C5 position of the

cytosine ring in CpG dinucleotides and is catalysed by DNA methyltransferase enzymes (Dnmts). Three

types of Dnmt are known in eukaryotes; Dnmt3 establishes DNA methylation patterns *de novo*, Dnmt1

maintains these patterns, and Dnmt2 is involved in tRNA methylation [3]. In mammals, broadly, DNA

methylation is found throughout the genome (except at CpG islands near the promoters of genes) and

is associated with transcriptional repression [4, 5]. By contrast, in insects, DNA methylation is

concentrated in gene bodies and is associated with more stable patterns of gene expression [6].

Notably, vertebrate and invertebrate DNA methylation varies greatly in its extent throughout the

genome, with the proportion of methylated CpGs much lower in invertebrates: 0-14% as compared to

60-90% in mammals [6, 7]. The evolutionary reasons for these differences are not yet clear, although

it is thought that gene body methylation is an ancient feature of eukaryote genomes [5]. However,

DNA methylation status influences how genes (and phenotypes) are expressed in many organisms,

including reproductive vs non-reproductive worker bumblebees [8], caste differentiation [9, 10], and

learning and memory [11] in honey bees, and behavioural division of labour in ants (*Pogonomyrmex*

barbatus [12]; but see [13] for a robust critique of earlier studies of DNA methylation and caste

determination in social insects). Therefore, DNA methylation patterns are very much part of mapping

genotypes to phenotypes.

The flexible control of gene expression provided by mechanisms such as DNA methylation has

also caught the interest of biologists interested in the underlying mechanisms of phenotypic plasticity,

and indeed for evolutionary biologists interested in how such plasticity may evolve [14, 15]. Moreover,

there has also been great interest in the extent to which epigenetic modifications may be inherited

across generations [14, 16, 17]. Transmission of epigenetic state allows transmission of parent-of-

origin information, or so-called genomic imprinting, setting the scene for genomic conflict [18, 19].

Genomic conflict arises when maternally- and paternally-inherited alleles have different evolutionary

optima for traits expressed in an individual [19, 20]. A classic example is the expression of maternally-

or paternally-inherited alleles during embryonic development in mammals: paternally-inherited

alleles are selected to extract more resources from the mother *in utero* compared to maternally-

inherited alleles [21]. This conflict is now known to be, at least in part, facilitated by parent-of-origin

specific gene expression of insulin-like growth factor II (Igf2) and Igf2r. In mammals, the Igf2 gene

promotes growth and cellular differentiation during development, and also regulates the placental

supply of nutrients and the demand of nutrients by the foetus [22].

To further understand the role epigenetic modifications such as DNA methylation have in

influencing phenotypes and their evolution, we need robust ways of manipulating them. Here we

consider the efficacy of the chemical 5-aza-2'-deoxycytidine (5-aza-dC) as a way of manipulating DNA

methylation, using the parasitoid wasp *Nasonia vitripennis*. The extent and nature of DNA methylation

across the *N. vitripennis* genome has recently been characterised [23, 24], and the presence in the

genome of a full "methylation toolkit" has also been confirmed [25]. 5-aza-dC is well-known from the

cancer literature as a hypomethylating agent, and is used in the treatment of acute myeloid leukaemia

and myelodysplastic syndrome [26-28]. In cancer patients, many genes that suppress leukemogenesis

in humans are silenced by aberrant DNA methylation. 5-aza-dC acts as a demethylating agent to

reactivate these genes (see below for details of action). In recent years, 5-aza-dC has been used across

a range of species as an experimental tool to explore how DNA methylation influences phenotypes

including diapause in *Nasonia vitripennis* [24] and worker reproductive status in bumble bees [8].

Previously, we used 5-aza-dC to explore whether DNA methylation was involved in the control of

facultative sex allocation in *N. vitripennis* [29].

Like all Hymenoptera, *N. vitripennis* is haplodiploid, with males arising from unfertilised

haploid eggs and females arising from fertilised diploid eggs. Females have putative control over the

sex of their offspring, releasing sperm to produce daughters or not to produce sons. Female *Nasonia*

allocate sex broadly in line with the predictions of Local Mate Competition theory [30-35; reviewed by

36]. Females laying eggs alone on their blowfly pupae hosts produce very female-biased offspring sex

ratios to minimise competition amongst her sons for mating opportunities. As more females

(foundresses) lay eggs on the same host(s) together, females produce less female-biased sex ratios,

as LMC is reduced and the fitness gained through the production of sons and daughters moves

towards equality [30]. When treated with 5-aza-dC, the facultative sex allocation response in female

Nasonia was maintained, suggesting that this plastic behaviour in and of itself did not require certain

patterns of methylation. Rather, the sex ratios (as proportion male) produced by females were shifted

slightly upwards, with more sons produced by treated females than by controls. This subtle shift is as

predicted by genomic conflict over sex allocation theory, which requires some form of genomic

imprinting [37]. Assuming that DNA methylation is the mechanism by which parent-of-origin allelic

information is carried in *Nasonia*, our data therefore provided the first (albeit indirect) evidence for

genomic conflict influencing sex allocation [29].

Recently, Ellers *et al.* [38] challenged the effects of 5-aza-dC as a general modifier of

methylation status, considering CpGs in ten genes in *Nasonia vitripennis*. Ellers *et al.* [38] failed to find

changes in methylation status across any of the 155 CpGs in their study, questioning the action of 5-

aza-dC in *Nasonia*, and hence of course the interpretation of the effects of 5-aza-dC in our earlier study

[29]. Fully negative results are of course very hard to interpret, but Ellers *et al.* [38] nonetheless raised

valid concerns of the efficacy of the chemical. As noted above, levels of methylation in insects is

generally low, and a genome-wide study would provide greater resolution to explore the effects of

this chemical in an ecologically valuable model system. Here we consider the effects of 5-aza-dC across

the whole genome using a Bisulfite sequencing (BS-Seq) experiment. There is evidence to suggest that

the effects of 5-aza-dC are unlikely to be random across methylated CpGs in a given genome. For

instance, in mammals, genomic context can influence the extent to which 5-aza-dC is effective as a

demethylating agent. DNA sequence context, the distribution of transcription factor binding sites, and

chromatin structure are known to influence the DNA methylation patterns produced by 5-aza-dC

exposure [39-41]. Although patterns of methylation differ among eukaryotes, the epigenetic

modification itself is chemically identical. Here, we verify that 5-aza-dC influences patterns of

methylation across the genome in *N. vitripennis*. Altered methylation patterns produced by 5-aza-dC

exposure were non-random, with enrichment in genes associated with transcription-factor activity

and sequence specific binding.

Materials and Methods

Study species

Nasonia vitripennis (Hymenoptera, Chalcicoidea) is a generalist parasitoid of large dipteran pupae including species of Calliphoridae. Females oviposit between 20 and 50 eggs in an individual host, with male offspring emerging just before females (after approximately 14 d at 25 °C: [42]). Males are brachypterous and unable to fly, remaining close to the emergence site where they compete with each other for emerging females, including their sisters. Females disperse after mating to locate new hosts. The females used in this experiment were from the wild-type AsymC strain, originally isolated in 1986 by curing the wild-type strain LabII of *Wolbachia*. Wasps have been maintained on *Calliphora vomitoria* or *C. vicina* hosts at 25 °C, 16L:8D light conditions ever since, and AsymC is also the reference genome strain [25]. The quantitative genetic basis of sex ratio variation has been repeatedly quantified [43, 44], whilst whole-genome transcription studies have confirmed that facultative sex allocation takes place without discernible changes in gene expression [45, 46].

5-aza-2'-deoxycytidine

We used 5-aza-dC (Sigma Aldrich Company Ltd., Gillingham, UK) at a non-lethal concentration (10 µM). Widely used in epigenetic anticancer treatments [26], 5-aza-dC is a nucleoside analogue that works by inhibiting DNA methyltransferase activity [47-50]. When initially introduced to a cell, 5-aza-dC is essentially inert but is then converted by processes inherent in the cell to its active form 5-aza-dCTP [26]. This active form is readily incorporated into DNA during S-phase of the cell cycle in the place of cytosine. In the daughter strands arising from a replication event, DNA methyltransferase 1 (Dnmt1) will attempt to restore the patterns of methylation that were present in the parent strand. However, Dnmt1 becomes irreversibly bound to the 5-aza-dCTP residues in the daughter strands, resulting in a diminishing pool of Dnmt1. Thus, administering 5-aza-dC results in a passive loss of DNA methylation, the key effect of 5-aza-dC [for a fuller description, see 51].

Experimental design

To control for possible host and other maternal effects, experimental females were not drawn straight from stock populations. Instead, two-day-old, mated, wild-type AsymC females were isolated from the mass cultures into individual glass vials. Each female was provided with three hosts and allowed to oviposit. Experimental females were drawn from the F1 generation, two days after emergence, one

female per "grandmother". We then employed a three by three factorial design whereby females

were allocated to one of three 5-aza-dC exposure regimes and were harvested post-feeding at three

time points. Exposure regimes were a) 20% sucrose for 24h, b) 20% sucrose supplemented with 10

μM 5-aza-dC for 24h and c) 20% sucrose supplemented with 10 μM 5-aza-dC for 48h. For each of the

three regimes, females were harvested at 0, 24 or 48h post-exposure. Throughout the experiment,

females were kept individually in glass vials and incubated at 25 °C, 16L :8D. Solutions (200 μl) were

provided in the lid of a 2 ml microcentrifuge tube. For individuals exposed to 5-aza-dC for 48 h, the

solutions were topped up by 100μl half-way through the exposure period to compensate for

evaporation. All solutions were stained green with food colouring such that a subset of 10 females

from each treatment combination could be dissected to check that feeding had occurred. At the point

of harvest, females were flash-frozen in liquid nitrogen and stored at -80 °C prior to DNA extraction.

DNA extraction and bisulphite-conversion

Prior to DNA extraction, individuals were removed from -80 °C storage and heads were quickly excised

from bodies using a sterile scalpel and a cryloyser to keep samples cold. Heads (and likewise bodies)

from the same treatment combination were randomly pooled into groups of 10 to give a maximum of

3 biological replicates for each treatment combination giving a total of 54 DNA samples. Briefly, tissue

was homogenised using a micropestle in 350 μl CTAB (Hexadecyltrimethylammonium Bromide) buffer

with subsequent overnight proteinase K digestion (400 μg) at 56 °C. Samples were cooled to room

temperature prior to RNase A digestion for 1 hour at 37 °C. A chloroform: isoamyl-alcohol wash (300

μl) was performed twice before ethanol/sodium acetate precipitation of the DNA. DNA purity was

analysed using the Nanodrop spectrophotometer with all samples having a 260/280 ratio of ≥1.8. DNA

integrity was confirmed using agarose gel electrophoresis.

Bisulphite sequencing

Bisulphite conversion, library preparation and sequencing were carried out by Edinburgh Genomics.

Briefly, bisulphite conversion was carried out using the EZ DNA Methylation Kit (Zymo Research, Irvine

USA), designed to reduce template degradation during the harsh bisulphite treatment whilst ensuring

complete conversion of the DNA. Library preparation was carried out using the TruSeq DNA

methylation Kit which takes the single-stranded DNA that results from bisulphite treatment and

converts it into an Illumina sequencing library. Sequencing of bisulphite-treated DNA was carried out

on the Illumina HiSeq 2500 in high output mode with 125bp paired-end reads. Library preparation

failed for three of the samples due to low levels of input DNA, leaving only two biological replicates

for three out of 18 treatment combinations and a total of 51 Bisulphite-sequenced samples (Table 1).

Table 1. Experimental design. Female wasps were exposed to 1 of 3 5-aza-dC exposure regimes
including a control. Females were then harvested at 1 of 3 timepoints after the exposure period
heads were excised from bodies to give a degree of tissue-specificity to the analysis. Three biological
replicates were available for almost all treatment combinations with each biological replicate
consisting of tissue from three *N. vitripennis* females.

Treatment Combination	5-aza-dC exposure regime	Harvest time	Tissue type	No. replicates (1 x rep. = 1 pool of 10)
1	20% sucrose for 24 h	0h	Bodies	3
2	20% sucrose for 24 h	0h	Heads	3
3	20% sucrose for 24 h	24h	Bodies	3
4	20% sucrose for 24 h	24h	Heads	3
5	20% sucrose for 24 h	48h	Bodies	3
6	20% sucrose for 24 h	48h	Heads	3
7	10µM 5-aza-dC for 24 h	0h	Bodies	3
8	10µM 5-aza-dC for 24 h	0h	Heads	3
9	10µM 5-aza-dC for 24 h	24h	Bodies	3
10	10µM 5-aza-dC for 24 h	24h	Heads	2
11	10µM 5-aza-dC for 24 h	48h	Bodies	3
12	10µM 5-aza-dC for 24 h	48h	Heads	2
13	10µM 5-aza-dC for 48 h	0h	Bodies	3
14	10µM 5-aza-dC for 48 h	0h	Heads	2
15	10µM 5-aza-dC for 48 h	24h	Bodies	3
16	10µM 5-aza-dC for 48 h	24h	Heads	3
17	10µM 5-aza-dC for 48 h	48h	Bodies	3
18	10µM 5-aza-dC for 48 h	48h	Heads	3

Raw data processing

Initial data processing was carried out by Edinburgh Genomics. Briefly, reads were trimmed using

Cutadapt version 1.12 removing any adapter sequences and for quality using a threshold of 30. After

trimming reads were required to have a minimum length of 50 bp. *Nasonia vitripennis* genome version

2.1   and   the   associated   annotation,   available   for   download   at   Ensembl

(http://metazoa.ensembl.org/Nasonia_vitripennis/Info/Index), were used as a reference. Reads were

aligned to the reference genome using Bismark (version 0.16) with the parameter "bowtie2". PCR

duplicates were removed from the resulting BAM files using samtools with the parameter "view -F

1024". Data for each CpG was then extracted from the BAM files using the Bioconductor package

methylKit (version 0.99.3) with R (version 3.3.1; [52]). The methylation status of each CpG for all

samples was calculated using the function "processBismarkAln" using the default parameters, except

"min.cov=1 no;ap=TRUE", which excludes CpG's not covered by any reads across all samples and does

not count CpGs covered by overlapping ends of a paired read as covered twice. The 1% of bases with

the highest coverage were removed using the function "filterByCoverage" to eliminate potential PCR

bias. After initial processing as outlined above, the raw data comprised percentage methylation at

4,150,376 CpG loci (no. of reads in which the CpG is methylated/total no. of reads covering the CpG)

for each of 51 samples. CpG read counts were then summed by gene, (including 1Kb up and

downstream) using gene annotation from Ensembl (version 2.1. available from:

ftp://ftp.ensemblgenomes.org/pub/metazoa/release-40/gff3/nasonia_vitripennis).   Around   half

(2,119,643) of the CpG's in the full dataset fell within genic regions with 14,765/17,279 genes (85%)

having at least one CpG present.

Statistical analysis

The overall structure of our dataset in terms of the proportion of methylated reads per gene and per

loci was visualised with two complementary analyses: a principle component analysis (PCA), which

allows samples to be visualised in relation to the two major axes of variation in our experiment, and a

T-distributed Stochastic Neighbour Embedding (t-SNE) analysis. t-SNE is a machine-learning algorithm

for data visualisation well-suited for high-dimensional datasets [53]. The PCA was conducted using the

R base function prcomp with default values. To perform the t-SNE analysis we used the R package

Rtsne [54], with default values and perplexity set to 15. For the PCA and t-SNE analyses any genes or

loci with 0 coverage in any one sample were discarded prior to analysis.

To examine the effect of 5-aza-dC exposure on individual genes, we modelled the number of

methylated and unmethylated reads from a gene as a generalized linear model (GLM) with a binomial

distribution with the following terms: exposure, tissue-type, collection time and their interactions.

Genes with 0 coverage for all samples in any particular treatment group were excluded from our

analysis (leaving a total of 11582 genes). Significance of each term was determined by a likelihood

ratio test, which was corrected for multiple testing using Benjamini and Hochberg's algorithm [55],

with statistical significance set to 5%. Coefficients (log odds of methylation) for each term were

extracted from the full model. All statistical analyses were conducted in R (v. 3.4.1; [52]).

Functional enrichment

N.   vitripennis   GO-terms   were   downloaded   from   "hymenopteramine"

(http://hymenopteragenome.org/hymenopteramine/begin.do) on 27th September 2018. In addition,

we used hymenopteramine to firstly obtain *D. melanogaster* orthologs for each *N. vitripennis* gene,

from which we could then obtain the associated *D. melanogaster* GO terms. We used both of these

GO-term sets for enrichment analyses (below). Some *N. vitripennis* genes had multiple *D.*

melanogaster orthologs. In these cases, all GO-terms from each *D. melanogaster* ortholog were

combined together. To ensure this did not bias our results we repeated our analyses, but instead of

combining GO-terms we kept only GO-terms that were shared between all orthologs.

To examine which processes are most affected by 5-aza-dC exposure, we conducted gene set

enrichment analyses using the R package TopGO (v. 2.28.0; [56]) using the elim algorithm to account

for the GO topology. Gene set enrichment analyses identify enriched GO terms in a threshold-free

way, by finding GO-terms that are overrepresented at the top of a ranked list of genes. Here we ranked

genes by FDR for the exposure effect. GO terms were considered to be significantly enriched when p

< 0.05.

Results

Median coverage across the whole experiment (N = 51 BS-seq libraries) was 898X, but coverage was

heterogeneous between loci (Fig 1) and samples (Table S1). Mean read coverage per sample ranged

from 2.6X to 48.2X. Four samples had less than 10X mean coverage per CpG, however, these samples

were distributed across treatment groups.

Treatment with 5-aza-dC had wide-ranging effects on DNA methylation across the genome.

First, samples clustered strongly by exposure regime at the gene-level in both the t-SNE visualisation

(Fig 2) and in the principle component analyses (Fig 3), indicating that 5-aza-dC has a strong effect on

the methylation of genes. Both analyses show the strong clustering at the gene loci, and that wasps

exposed for 24 hours are more like controls than wasps treated for 48 hours. In addition, the PCA

analysis showed clear separation along PCA1 (which presumably is heavily associated with exposure

regime; Fig 3). Importantly though, this analysis also emphasises the extent of variation in response

across the three treatment regimes, which is smallest for the controls, and largest for the 48 hour-

exposed wasps (note the spread along PCA axis 2 in Fig 3). In addition, both cluster analyses placed

the controls as some way intermediate between the two 5-aza-dC exposed treatments, which we

consider further in the next two paragraphs. Both analyses also show that exposure regime dominated

the clustering, over and above of any effects of either tissue-type or collection time, suggesting that

this is the key effect. When we consider clustering at the level of individual CpG loci, clustering is much

less obvious, using both t-SNE and PCA. However, the control treatment appears to cluster within the

others in the t-SNE analyses, and the 24-hour and 48-hour treatments do separate to some extent,

e.g. with the 24h samples more the bottom and left of Fig S1 and the 48h samples more to the top
and right (see Figs S1 and S2). The differences between the individual CpG and gene-level cluster
analyses are likely to be due to a combination of low coverage and a greater stochasticity of
methylation at the individual CpG level (see discussion).

Second, a more in-depth examination of the data using a generalised linear modelling (GLM)
approach to examine differential methylation at the gene level found that 5-aza-dC exposure had a
significant effect on the methylation status of the majority of genes examined; exposure treatment
was significant as a main effect at an FDR level of 0.05 for 8,725 out of 11,582 genes, or 75.3% of genes
(see also Fig S3). Moreover, both tissue-type and collection time also influenced methylation status
across a large number of genes (significant as main effects in the GLMs for 3833 and 3566 genes
respectively). Significant interactions between factors were also present for the majority of genes
(exposure regime * collection time = 10,776 genes, exposure regime * tissue type = 10,484 genes,
collection time * tissue type = 5,364 genes, and exposure * collection time * tissue type = 10,484
genes), showing widespread evidence for significant interplay between these factors. These results
emphasise the context-dependent nature of the action of 5-aza-dC. Full results from the GLMs are
presented in Table S2.

Interestingly, it appears that methylation tends to increase initially in response to exposure to
5-aza-dC, an effect that is noticeable after 24h of exposure, before decreasing after 48h of exposure
(Fig 4A). This means that how long individuals are exposed to the chemical may well influence what
kind of methylation changes are observed; in other words, both hyper- and hypo-methylation can
result from 5-aza-dC treatment. In terms of when wasps were collected after exposure, 3,566 genes
displayed significantly altered methylation in association with collection time, with a more
pronounced decrease in methylation in samples collected 48h post-exposure relative to 24h post-
exposure (when compared to 0h post-exposure; Fig 4B). This suggests that the effect of 5-aza-dC
persists at least 48h after exposure has stopped. Finally, 3,833 genes displayed significantly altered
methylation patterns in association with tissue type (Fig 4C), with lower methylation overall in the
head. However, we re-iterate that these main effects need to be contextualised within the pattern of
widespread interactions seen in the data, with 91% of genes exhibiting a significant second-order
interaction between exposure, tissue type and collection time.

Gene-set enrichment analysis, using *N. vitripennis* GO terms, when genes were ranked by their
response to exposure regime is presented in Table S3. Broadly, we identified 169 significantly enriched
GO terms (p < 0.05), and 80 of these terms fell under the "Biological process" (BP) category, 18 under
"Cellular component" (CC), and 71 under "Molecular function" (MF). The top five enriched GO terms

were "sequence-specific DNA binding" (GO:0043565), "regulation of transcription, DNA templated" (GO:0006355), "ATP binding" (GO:0005524), "DNA binding transcription factor activity" (GO: 0003700), and "protein kinase activity" (GO:0004672) (Table S3). When the analysis was repeated using GO terms for the *D. melanogaster* orthologs, a higher number of significantly enriched GO terms was obtained (700; 538 BP, CC 71, MF 91; Table S4). The top five GO terms for the *D. melanogaster* analysis were "nucleus" (GO:0005634), "positive regulation of transcription by RNA polymerase II" (GO:0045944), "regulation of Notch signalling pathway" (GO:0008593), "imaginal disc-derived wing morphogenesis" (GO:0007476), and, making a reappearance, "sequence-specific DNA binding" (GO:0043565). This analysis combined GO-terms when *N. vitripennis* genes had multiple *D. melanogaster* orthologs, which could lead to bias, however, we retrieve very similar enriched terms when we used only the GO terms that were shared between the multiple orthologs (Table S5), suggesting any bias caused by this approach is small.

Discussion

A genome-wide analysis of CpGs across more than 11,000 genes, using 51 biological replicates across our treatment combinations, has shown that the demethylating agent 5-aza-dC has very widespread effects on patterns of DNA methylation in the *Nasonia vitripennis* genome. The methylome has been characterised in *Nasonia vitripennis* by Wang *et al.* [23] and Beeler *et al.* [57], and in common with other insect species, DNA methylation appears to be primarily associated with gene bodies (including intron-exon boundaries), and a rather low proportion of CpGs are methylated. As mentioned above, previous work that has attempted to confirm the effects of 5-aza-dC on DNA methylation have had mixed results. Pegoraro *et al.* [24] confirmed the effects of 5-aza-dC across five genes (four via a qPCR MethylQuant assay and one via cloning a fragment of the *msn* gene from treated versus untreated individuals), whilst Ellers *et al.* [38] have recently failed to show any effects of 5-aza-dC on the methylation status of 155 CpGs across ten genes (nine of which provided libraries; they used high-throughput bisulfite amplicon sequencing: BSAS; see [58] for a discussion of different methylation quantification assays). Our larger-scale study has both confirmed the action of 5-aza-dC as a disruptor of methylation in *Nasonia*, but also suggested why it might be so difficult to pin down those effects. In particular, we find that whilst we see statistically significant main effects of exposure regime on DNA methylation (which includes differences in length of exposure), we also see effects of tissue type (albeit at the crude level of heads versus bodies), and time since exposure to 5-aza-dC. Moreover, we also see many statistically significant interactions between these factors; all told, the action of 5-aza-dC is likely to be context-dependent, both in terms of the sequences more likely to be targeted but also in terms of the experimental details, including target tissues. Nonetheless, our results strongly

suggest that 5-aza-dC is not chemically inert in terms of influencing DNA methylation in *Nasonia* –

indeed, quite the opposite.

That the effects of 5-aza-dC are non-random in terms of the genes it influences are further

confirmed by our gene ontology enrichment analyses. We see enrichment for genes associated with

DNA binding, regulation of transcription, and transcription factor activity. These processes are of

interest given that we might expect the regulation of transcription to be a crucial component of the

epigenetic role of DNA methylation, given what we know from both mammals and insects, even

though the targets (throughout the genome versus exons) and outcomes (transcriptional silencing

versus stable gene expression) are typically different in the two taxa. Our analysis certainly confirms

the sequence-specific action of 5-aza-dC seen in other species.

A further important effect we see is that 5-aza-dC exposure can result in hyper-methylation

as well as hypo-methylation. This means that the characterisation of 5-aza-dC as a strictly

demethylating agent may be somewhat misleading. The complexity of outcomes of 5-aza-dC in

mammals, in particular in terms of trying to understand its action for cancer treatments, is becoming

increasingly well-appreciated [26]. As such, our data confirm in an insect what is being seen in

mammalian cells. We are not well placed to speculate on the mechanism(s) by which 5-aza-dC may

lead to hyper-methylation (see instead [26]), even though its hypomethylating effect is well

characterised chemically (see Introduction). However, if 5-aza-dC disrupts the regulation of the DNA

methylation machinery itself, for instance in terms of how the Dnmt genes are expressed and act, then

the downstream outcomes could be both hyper- and hypo-methylation. Perhaps related to this, in

their study of DNA methylation and the control of the diapause response in *Nasonia*, Pegoraro *et al.*

[24] used both 5-aza-dC and RNA interference (RNAi) to manipulate methylation, in the latter case

disrupting DNA methylation by knocking down both Dnmt1a and Dnmt3. Whilst both exposure to 5-

aza-dC and the RNAi constructs affected how diapause was influenced by exposure to short or long

day length, they did so in different ways. Whilst both manipulations abolished the day-length

response, the RNAi of Dnmt1a and Dnmt3 resulted in a change in how wasps responded to long-day

length (increasing the diapause response), whilst 5-aza-dC led to a change in response to short-day

length (decreasing the diapause response). As such, different manipulations of DNA methylation may

lead to different phenotypic outcomes, depending on the regulatory causes and consequences of the

methylation pattern itself. More generally then, it is clear that there is much that remains to be

elucidated in how 5-aza-dC shapes methylation patterns, in both insects and mammals [26].

On a more technical note, perhaps unsurprisingly we found more significant genes in the GO

enrichment analysis using the GO terms for *Drosophila melanogaster* orthologs. This is perhaps

because of the more detailed gene ontology information available for the *D. melanogaster* genome.
Nonetheless, the two analyses provide a similar story in terms of GO enrichment. In addition, the
majority of our analyses were undertaken at the level of the gene. Perhaps given the variation in
coverage at the level of individual CpGs, the clustering analyses were not by any means as clear cut as
the gene-level results. However, in the t-SNE analysis (Fig S1), the control replicates cluster together,
albeit within the broader distribution of the other two exposure treatment groups. The effect of
genomic and sequence context may be particularly relevant here: our analysis at the level of the gene
(see Methods for details) brings together CpGs that share a lot of context, by virtue of being associated
with a given genetic locus. The differences in the clustering at the individual CpG versus gene levels
may therefore represent the fact that the action of 5-aza-dC does appear to be non-random, and that
we gain clarity when we consider closely associated CpGs whose patterns of methylation, and
responses to 5-aza-dC, are more similar than randomly chosen CpGs across the genome.

In conclusion, our results suggest that 5-aza-dC remains a potential tool for evolutionary
ecologists wishing to explore how DNA methylation influences phenotypic expression. As with all
genomic approaches however, no one approach gives all the answers. 5-aza-dC may prove a very
useful, if rather blunt, tool for first assessing the range of phenotypes that are influenced by
methylation status, whilst no doubt more targeted knock-downs and genetic manipulations will be
needed to link phenotypes to causal CpGs.

Acknowledgments

We are very grateful to Mark Lammers and Jacintha Ellers for their collegiate and constructive
engagement with our previous DNA methylation work. We are also grateful to Matthew Arno and the
Edinburgh Genomics team for their sequencing and downstream support.

References

424 **References**

425 1. Ashbrook DG, Mulligan MK, Williams RW. Post-genomic behavioural genetics: From
426   revolution to routine. *Genes Brain Behav.* 2018; 17: e12441.
427

428 2. Biémont C. From genotype to phenotype. What do epigenetics and epigenomics tell us?
429   *Heredity.* 2010; 105: 1-3.
430

431 3. Lyko F. The DNA methyltransferase family: a versatile toolkit for epigenetic regulation. *Nat*
432   *Rev Genet.* 2018; 19: 81-92.
433

434 4. Feng S, Cokus SJ, Zhang X, Chen P, Bostick M *et al.* Conservation and divergence of
435   methylation patterning in plants and animals. *Proc Natl Acad Sci USA. 2010;* 107: 8689-8694.
436

437 5. Zemach A, McDaniel IE, Silva P, Zilberman D. Genome-wide evolutionary analysis of
438   eukaryotic DNA methylation. *Science.* 2010; 328: 916-919.
439

440 6. Bewick AJ, Vogel KJ, Moore AJ, Schmitz RJ. Evolution of DNA methylation across Insects. *Mol*
441   *Biol Evol.* 2017; 34: 654-665.
442

443 7. Glastad KM, Hunt BG, Yi SV, Goodisman MAD. DNA methylation in insects: on the brink of the
444   epigenomic era. *Insect Mol Biol.* 2011; 20: 553-65.
445

446 8. Amarasinghe HE, Clayton CI, Mallon EB. Methylation and worker reproduction in the bumble-
447   bee (*Bombus terrestris*). *Proc R Soc Lond B Biol Sci.* 2014; 281: 20132502.
448

449 9. Kucharski R, Maleszka J, Foret S, Maleszka R. Nutritional control of reproductive status in
450   honeybees via DNA methylation. *Science. 2008;* 319: 1827-1830.
451

452 10. Lyko F, Foret S, Kucharski R, Wolf S, Falckenhayn C *et al.* The honey bee epigenomes:
453   differential metjhylation of brain DNA in queens and workers. *PLoS Biol.* 2010; 8: e1000506.
454

455 11. Li Y, Zhang L, Yi Y, Hu W, Guo Z *et al.* Genome-wide DNA methylation changes associated with
456   olfactory learning and memory in *Apis mellifera*. *Sci Rep.* 2017; 7: 17017.
457

458 12. Smith CR, Mutti NS, Jasper WC, Naidu A, Smith CD *et al.* Patterns of DNA methylation in
459   development, division of labor and hybridisation in an ant with genetic caste determination.
460   *PLoS One* 2012; 7: e42433.
461

462 13. Libbrecht R, Oxley PR, Keller L, Kronauer DJ. Robust DNA methylation in the clonal raider ant
463   brain. *Curr Biol.* 2016; 26: 391-395.
464

465 14. Bonduriansky R, Day T. Nongenetic inheritance and its evolutionary implications. *Annu Rev*
466   *Ecol Evol Syst.* 2009; 40: 103-125.
467

468 15. Bonduriansky R, Day T. *Extended Heredity*. Princeton: Princeton University Press; 2018.
469

470 16. Lind MI, Spagopoulou F. Evolutionary consequences of epigenetic inheritance. *Heredity.* 2018;
471   121: 205-209.
472

473   17. Burggren W. Epigenetic Inheritance and its role in evolutionary biology: re-evaluation and new
474   perspectives. *Biology. 2016;* 5: 24.
475

476   18. Patten MM, Ross L, Curley JP, Queller DC, Bonduriansky R *et al.* The evolution of genomic
477   imprinting: theories, predictions and empirical tests. *Heredity.* 2014; 113: 119-128.
478

479   19. Haig D. *Genomic imprinting and kinship.* New Brunswick: Rutgers University Press; 2002.
480

481   20. Haig D. Parental antagonism, relatedness asymmetries and genomic imprinting. *Proc R Soc
482   Lond B Biol Sci.* 1997; 264: 1657-1662.
483

484   21. Moore T, Haig D. Genomic imprinting in mammalian development: a parental tug-of-war.
485   Trends Genet. 1991; 7: 45-49.
486

487   22. Constancia M, Hemberger M, Hughes J, Dean W, Fergusin-Smith A *et al.* Placental-specific IGF-
488   II is a major modulator of placental and fetal growth. *Nature.* 2002; 417(6892): 945-948.
489

490   23. Wang X, Wheeler D, Avery A, Rago A, Choi JH. Function and evolution of DNA methylation in
491   *Nasonia vitripennis*. *PLoS Genet*. 2013; 9: e1003872.
492

493   24. Pegoraro M, Bafna A, Davies NJ, Shuker DM, Tauber E. DNA methylation changes induced by
494   long and short photoperiods in Nasonia. *Genome Res.* 2016; 26: 203-210.
495

496   25. Werren JH, *et al.* Functional and evolutionary insights from the genomes of three parasitoid
497   *Nasonia* species. *Science*. 2010; 327: 343-348.
498

499   26. Seelan RS, Mukhopadhyay P, Pisano MM, Greene RM. Effects of 5-Aza-2'-deoxycytidine
500   (decitabine) on gene expression. *Drug Metab Rev.* 2018; 50: 193-207.
501

502   27. Bryan J, Kantarjian H, Garcia-Manero G, Jabbour E. Pharmacokinetic evaluation of decitabine
503   for the treatment of leukemia. *Expert Opin Drug Metab Toxicol.* 2011; 7: 661-672.
504

505   28. Momparler RL. A perspective on the comparative antileukemic activity of 5-aza-2'-
506   deoxycytidine (Decitabine) and 5-azacytidine (Vidaza). *Pharmaceuticals* (Basel). 2012; 5: 875-
507   881.
508

509   29. Cook N, Pannebakker BA, Tauber E, Shuker DM. DNA methylation and sex allocation in the
510   parasitoid wasp Nasonia vitripennis. *Am Nat. 2015;* 186: 513-518.
511

512   30. Hamilton WD. Extraordinary sex ratios. *Science.* 1967*;* 156: 477-488.
513

514   31. Taylor PD, Bulmer MG. Local Mate Competition and the sex-ratio. *J Theoret Biol.* 1980; 86:
515   409-419.
516

517   32. Werren JH. Sex ratio adaptations to local mate competition in a parasitic wasp. *Science.* 1980;
518   208: 1157-1159.
519

520   33. Werren JH. Sex ratio evolution under local mate competition in a parasitic wasp. *Evolution.*
521   1983; 37: 116-124.
522

34. Shuker DM, West SA. Information constraints and the precision of adaptation: sex ratio manipulation in wasps. *Proc Natl Acad Sci USA.* 2004*;* 101: 10363-10367.

35. Burton-Chellew MN, Koevoets T, Grillenberger BK, Sykes EM, Underwood SL *et al.* Facultative sex ratio adjustment in natural populations of wasps: cues of local mate competition. *Am Nat.* 2008; 172: 393-404.

36. West SA. Sex allocation. Princeton: Princeton University Press; 2009.

37. Wild G, West SA. Genomic imprinting and sex allocation. *Am Nat.* 2009; 173: E1-14.

38. Ellers J, Visser M, Mariën J, Kraaijeveld K, Lammers M. The importance of validating the demethylating effect of 5-aza-dC in model species. *Am Nat*. 2019; Forthcoming.

39. Ramos M, Wijetunga NA, McLellan AS, Suzuki M, Greally JM. DNA demethylation by 5-aza-2'-deoxycytidine is imprinted, targeted to euchromatin and has limited transcriptional consequences. *Epigenetics Chromatin.* 2015; 8:11.

40. Mossman D, Kim K, Scott RJ. Demethylation by 5-aza-2'-deoxycytidine in colorectal cancer cells targets genomic DNA whilst promoter CpG island methylation persists. *BMC Cancer* 2010; 10: 366.

41. Hagemann S, Heil O, Lyko F, Brueckner B. Azacytidine and decitabine induce gene-specific and non-random DNA demethylation in human cancer cell lines. *PLoS One.* 2011; 6: e17388.

42. Whiting AR. The biology of the parasitic wasp *Mormoniella vitripennis* [=*Nasonia brevicornis*] (Walker). *Q Rev Biol.* 1967; 42: 333-406.

43. Pannebakker BA, Halligan DL, Reynolds KT, Ballantyne GA, Shuker DM *et al.* Effects of spontaneous mutation accumulation on sex ratio traits. *Evolution*. 2008; 62: 1921-1935.

44. Pannebakker BA, Watt R, Knott SA, West SA, Shuker DM. The quantitative genetic basis of sex ratio variation in *Nasonia vitripennis*: a QTL study. *J Evol Biol. 2011;* 24: 12-22.

45. Cook N, Trivedi U, Pannebakker BA, Blaxter M, Ritchie MG *et al.* Oviposition but not sex allocation is associated with transcriptomic changes in females of the parasitoid wasp *Nasonia vitripennis*. *G3 (Bethesda).* 2015; 5: 2885-2892.

46. Cook N, Boulton RA, Green J, Trivedi U, Tauber E *et al.* Differential gene expression is not required for facultative sex allocation: a transcriptome analysis of brain tissue in the parasitoid wasp *Nasonia vitripennis*. *Roy Soc Open Sci*. 2018; 5: 171718.

47. Piskala, Sorm F. Nucleic acid components and their analogues. LI. Synthesis of 1-glycosyl derivatives of 5-azauracil and 5-azacytosine. *Collect Czechoslov Chem Commun.* 1964; 29: 2060-2076.

48. Jones PA, Taylor SM. Cellular differentiation, cytidine analogs and DNA methylation. *Cell. 1980;* 20: 85-93.

572   49. Momparler RL. Molecular, cellular and animal pharmacology of 5-aza-2'-deoxycytidine.
573   *Pharmacol Ther.* 1985; 30: 287-299.
574

575   50. Christman JK 5-azacytidine and 5-aza-2'-deoxycytidine as inhibitors of DNA methylation:
576   mechanistic studies and their implications for cancer therapy. *Oncogene.* 2002; 21: 5483-
577   5495.
578

579   51. Cook N, Parker DJ, Tauber E, Pannebakker BA, Shuker DM. Validating the demethylating
580   effects of 5-aza-2'-deoxycytidine in insects requires a whole-genome approach (A reply to
581   Ellers *et al.*) *Am Nat.* 2019; Forthcoming.
582

583   52. R Core Team. R: A language and environment for statistical computing. R Foundation for
584   Statistical Computing, Vienna, Austria. 2017. Available from https://www.R-project.org
585

586   53. van der Maaten LJP, Hinton GE. Visualizing high-dimensional data using t-SNE. *J Mach Learn
587   Res.* 2008; 9: 2579-2605.
588

589   54. Krijthe JH. Rtsne: T-Distributed Stochastic Neighbor Embedding using a Barnes-Hut
590   Implementation. 2015. Available from: https://github.com/jkrijthe/Rtsne
591

592   55. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful
593   approach to multiple testing. *J R Stat Soc Series B Stat Methodol.* 1995; 57: 289–300.
594

595   56. Rahnenfuhrer AAJ. *topGO: Enrichment Analysis for Gene Ontology*. R package version 2.28.0.
596   2016. Available from http://bioconductor.org/packages/release/bioc/html/topGO.html
597

598   57. Beeler SM, Wong GT, Zheng JM, Bush EC, Remnant EJ et al. Whole-genome DNA methylation
599   profile of the Jewel Wasp (*Nasonia vitripennis*). G3 (Bethesda). 2014; 4: 383-388.
600

601   58. Redshaw N, Huggett JF, Taylor MS, Carole A, Foy CA *et al.* Quantification of epigenetic
602   biomarkers: an evaluation of established and emerging methods for DNA methylation
603   analysis. *BMC Genomics*. 2014; 15: 1174.
604

Data and code availability

Raw sequence reads and processed data (proportion methylated reads per CpG) have been deposited

in the GEO hosted by NCBI (Accession: GSE125388). Scripts for the analyses in this paper are available

at: https://github.com/DarrenJParker/5-Aza-dC_methylation.

Figures

Fig 1. Read coverage by locus across all 51 BS-Seq libraries.

Fig 2. t-SNE of the proportion of methylated reads per gene. Shows the effects of (A) Exposure

regime and tissue type and (B) Exposure regime and harvest time.

Fig 3. PCA of the proportion of methylated reads per gene. Shows the effects of (A) Exposure regime and tissue type and (B) Exposure regime and harvest time.

Fig 4. Gene methylation status displayed as log odds of methylation for each main effect in the experiment. A significant effect on gene methylation status was noted in response to (A) 5-aza-dC exposure regime (relative to sucrose control), (B) harvest time after exposure (relative to zero hours) and (C) tissue-type (heads relative to bodies). N= number of genes with significantly differentially methylated CpGs in association with each main effect.