1

Title: The best of both worlds: A new lipid complex has micelle and bicelle-like properties

Running title: Characterizing a new lipid complex

Authors: Monica D. Rieth *

*corresponding author

9
10

Abbreviations: DMPC, 1,2-dimyristoyl-*sn*-glycero-3-phosphocholine; DHPC, 1,2-dihexanoyl-*sn*-glycero-3-phosphocholine; C_8E_5, n-octylpentaoxyethylene; NBD-DMPE (1,2-dimyristoyl-*sn*-glycero-3-phosphoethanolamine-N-(7-nitro-2-1,3-benzoxadiazol-4-yl) (ammonium salt)); ^{31}P-NMR, phosphorus nuclear magnetic resonance spectroscopy; D_2O, deuterium oxide; CHAPS, 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate; CHAPSO, 3-([3-Cholamidopropyl]dimethylammonio)-2-hydroxy-1-propanesulfonate; HEPES, 2-[4-(2-hydroxyethyl)piperazin-1-yl]ethanesulfonic acid; DLS, dynamic light scattering

Abstract

Bicelles have been demonstrated to be a valuable tool for studying membrane protein interactions and structure *in vitro*. They are distinguished by a distinct lipid bilayer that mimics the plasma membrane of cells making it more native-like than its detergent micelle counter-part. Bicelles are typically comprised of a long-chain phospholipid such as dimyristoylphosphatidylcholine (DMPC) and a short-chain phospholipid such as dihexanoylphosphatidylcholine (DHPC). When mixed together in solution DMPC-DHPC bicelles assume a discoidal structure comprised of a heterogeneous arrangement where the short-chain lipids gather around the rim of the disk and the long-chain lipids form the flat, planar, bilayer region. In this study, the nonionic surfactant, C_8E_5, was used to prepare mixtures with DMPC to determine if it adopts properties similar to bicelles with a $q \geq 0.5$. At $q \geq 0.5$, DMPC-DHPC bicelles are bilayered and DMPC is sequestered from the detergent micelle-like DHPC. Mixtures of DMPC and C_8E_5 were prepared at various q values, a parameter used to describe the mole ratio of DMPC to DHPC in the preparation of bicelles. Employing biophysical methods like dynamic light scattering, ^{31}P-NMR and analytical ultracentrifugation, properties of these lipid-detergent complexes are described. Interestingly they adopted a spherical-shaped micellar structure morphology and did not assume a discoidal shape typical of bicelles at $q \geq 0.5$. However, they appear to retain bilayer-like properties that may prove beneficial for *in vitro* biophysical studies of membrane proteins.

1.   Introduction

It has been well-established that 1,2-dimyristoyl-*sn*-glycero-3-phosphocholine (DMPC), a naturally derived phospholipid with a 14-carbon chain length and a choline head group, spontaneously assembles into discoidal lipid structures called bicelles when mixed with short-chain phospholipids such as 1,2-dihexanoyl-*sn*-glycero-3-phosphocholine (DHPC) - also naturally derived but containing a 6-carbon chain length with a choline head group [1, 2]. These bilayered structures can also be prepared using DMPC and bile salt detergents such as CHAPS or CHAPSO in lieu of a short-chain phospholipid [1, 3, 4]. DMPC-DHPC bicelles are among the most extensively characterized bicellar structures, and they have been used successfully as membrane mimics in solution NMR studies of membrane protein structure [4 – 7]. These lipid complexes can be tailored to different applications by altering the properties of the lipids used in their preparation. For example, the lipid tail length and the ratio of DMPC to DHPC can affect the resulting physical properties of bicelles. Anionic lipids can also be included to alter the charge at the surface [8]. Collectively, these properties affect the thickness and the planar length of the bicellar structure both of which can influence membrane protein structure when they are incorporated and ultimately, their stability [9]. Bicelles are of unique interest for membrane protein studies because unlike their detergent counterpart, they contain a bilayered planar region, which captures a true membrane-like environment (Figure 1) [4, 5]. In membrane protein studies, choosing the optimal detergent or lipid system usually necessitates a screening of potential candidates because not all systems are universally adaptable to all types of membrane proteins [10 – 13]. In this study, a novel lipid-detergent complex was prepared with DMPC, the long-chain phospholipid, and a nonionic detergent, pentaethylene glycol n-octyl ether or C_8E_5 (Figure 1). Physical properties of the resulting complexes were characterized to assess whether bicelle-like properties were retained. Two parameters commonly used to describe bicelles were investigated. The *q*-value, which describes the ratio of the long-chain lipid to that of the short-chain lipid and

the total lipid concentration (% w/w), which is defined as the total concentration of DMPC + C_8E_5.

Our analysis shows the lipid complexes assume a shape that is closer to spherical than discoidal.

Further, at various q-values, DMPC-C_8E_5 complexes show remarkable changes in solution

viscosity that are not typically observed for DMPC-DHPC bicelles at $q \geq 0.5$. Because of this, the

q-value for DMPC-C_8E_5 samples was kept at a $q \leq 0.5$, a range we refer to as the "solution

dynamic range." All q-values reported were determined on a theoretical basis and do not account

for any free DHPC and C_8E_5 detergent in solution. It should be acknowledged that accounting for

free detergent in solution based on the critical micelle concentration results in a larger observed

q, q_{eff} [2]. For example, $q = 0.7$ bicelles, results in a $q_{eff} = 1.0$ determined from the CBC of DHPC

[2, 14]. This accounts for the DHPC associated with bicelles, which is in equilibrium with free

DHPC in solution.

In addition to characterizing important parameters of these complexes, we also show that

they can be adapted to studies of membrane protein self-association (oligomerization) using

sedimentation equilibrium in the analytical ultracentrifuge. A method that has been used

successfully to investigate membrane protein oligomerization in both micellar and bicellar

environments [2, 15]. With a partial specific volume similar to water, 0.983 cm^3 / g, C_8E_5 micelles

require little need for density matching [16]. However, in lipid complexes, only a small quantity of

the density modifier deuterium oxide, D_2O, was required for matching.

Materials and methods

2.1   Chemicals and reagents

DMPC (1,2-dimyristoyl-*sn*-glycero-3-phosphocholine) and NBD-DMPE (1,2-dimyristoyl-*sn*-

glycero-3-phosphoethanolamine-N-(7-nitro-2-1,3-benzoxadiazol-4-yl) (ammonium salt)) were

purchased from Avanti Polar Lipids (Alabaster, AL, USA). C_8E_5 (n-octylpentaoxyethylene) was

purchased from Bachem (King of Prussia, PA, USA). All lipids and detergents were used without

further purification. NBD-DMPE was incorporated in small molar quantities so that absorption

optics could be used to analyze the lipid complexes in the analytical ultracentrifuge. HEPES (2-

[4-(2-hydroxyethyl)piperazin-1-yl] ethanesulfonic acid) was purchased from EMD Millipore

(Billerica, MA USA) and NaCl (sodium chloride) was purchased from Avantor (Center Valley, PA,

USA). D_2O (deuterium oxide) was purchased from Cambridge Isotopes (Tewksbury, MA, USA).

Milli-Q water (Millipore) was used throughout all experiments.

2.2   Preparation of DMPC-C_8E_5 mixtures

2.1.1   Samples for analytical ultracentrifugation

DMPC-C_8E_5 lipid solutions were prepared and evaluated at q = 0.25 and a total lipid

composition of 5% (w/w). A small amount of NBD-labeled DMPE was incorporated into DMPC-

C_8E_5 samples to allow the lipid complexes to be monitored using absorption optics. The ratio of

NBD-labeled lipid to unlabeled lipid was kept sufficiently low so that the label did not influence the

physical properties of the complexes (1:500 mole ratio of NBD-DMPE:DMPC). To a 1.5 mL

eppendorf tube a sufficient quantity of 50 mg / mL DMPC in chloroform was added to give the

appropriate q value. An appropriate quantity of 66 µg / µL of NBD-DMPE chloroform solution was

immediately added to yield a 1:500 mole ratio of fluorescent-labeled lipid to unlabeled lipid. The

samples were briefly vortexed and spun in a microfuge. 500 µL of Millipore water was added

immediately before freezing the samples in liquid nitrogen. The samples were lyophilized to a

powder. To each lyophilized sample an appropriate amount of Millipore water was added followed

by a 40X concentrated stock solution of buffer, which yielded a final concentration of 10 mM

HEPES, 100 mM NaCl pH 7.4. The samples were vortexed to a milky white suspension and C_8E_5

detergent was added to each sample in quantities that would assume a q-value of 0.25. Samples

were vortexed for several seconds to clarity. D_2O was added to each sample ranging from 0%

(by volume) up to 25 % for a total of nine samples. Reference samples (buffer only) were

prepared to a final volume of 150 µL according to the following: To a 1.5 mL eppendorf tube was

added 146.0 μL of Millipore water (less D₂O) followed by 40X concentrated buffer solution to final

composition of 10 mM HEPES 100 mM NaCl pH 7.4.

Samples were loaded into a 6-channel equilibrium charcoal-filled epon centerpiece with a

pathlength of 1.2 cm. 120 μL of each sample and reference was loaded in the appropriate

channels. All sedimentation equilibrium studies were carried out on a Beckman XL-A analytical

ultracentrifuge at 25 °C using a four-hole An-Ti 60 rotor. Absorbance spectroscopy was used to

monitor the sedimentation of the lipid-detergent complexes down the solution column at a

wavelength of 464 nm, which corresponds to the absorbance of the NBD-labeled lipid. Samples

were initially equilibrated at 10,000 rpm (8,050 x g) for 16 hours to allow equilibrium to be reached.

The point of equilibrium was evaluated at each speed using Match in the program, *HeteroAnalysis*

(University of Connecticut, Storrs CT). Thereafter speeds were increased by 5,000 rpm up to

30,000 rpm (72,446 x g), and samples were left to equilibrate for 6 hours at each speed before

acquiring data. Data was collected at rotor speeds of 15,000 rpm (18,112 x g), 20,000 (32,198 x

g), 25,000 rpm (50,310 x g), and 30,000 rpm (72,446 x g). The sedimentation equilibrium profiles

were analyzed and the data were fitted to the Lamm equation, Eq. 1, to extract the buoyant MW,

M_{eff}, of the lipid-detergent complexes in each of the three samples:

147
$$\ln(c_r) = \frac{M(1 - v\rho)\omega^2}{2RT} r^2$$
Eq. 1

where c_r is the concentration of NBD-DMPE at radius r; M is the calculated molecular weight of

the lipid-detergent complex, v is the partial specific volume of the lipid-detergent complex which

was measured using a Kyoto density meter, ρ is the density of the HEPES buffer containing NaCl

pH 7.4, which was also measured using the Kyoto density meter. ω is the angular velocity of the

rotor, r is the radius from the center axis of rotation, R is the Boltzmann constant and T is the

temperature in Kelvin.

2.2.2 ^{31}P-NMR sample preparation

Samples with a 25% (w/w) total lipid composition were prepared for ^{31}P-phosphorus NMR

experiments. Each sample was prepared at a **q** = 0.5 on a 990 µL scale. To two separate 1.5

mL eppendorf tubes, DMPC was added to a final concentration of 158 mM. To both samples

81.45 µL of Millipore water was added followed by 99.0 µL of D$_2$O (10% (v/v)) and 6.6 µL 3.0 M

sodium acetate pH 5.6 buffer. The samples were vortexed to a milky, white suspension. Next,

555.44 µL of a 25% (w/w) DHPC solution was added to one sample and 222.48 µL of a 50% C$_8$E$_5$

(w/w) solution was added to the second sample. The samples were vortexed to clarity. Last,

praseodymium (III) chloride powder was added to a final concentration of 197 mM as per Glover

et. al [4]. This concentration was previously determined to saturate the lanthanide-phospholipid

interaction making the DMPC and DHPC ^{31}P-phosphorus signals distinct from one another in the

NMR [17].

NMR experiments were carried out on a Bruker DRX500 spectrometer equipped with a BBI

probe. One dimensional ^{31}P-phosphorus NMR spectra were recorded at 25 °C and 37 °C using

a proton-decoupled single-pulse experiment with a minimum of 16 scans and a 100 ppm sweep

width. The experiments were processed in TopSpin v. 1.3 (Bruker Corporation).

2.2.3 Samples for DLS

Mixtures of DMPC-C$_8$E$_5$ were prepared based on the bicelle **q**–value. **q** is determined by the

mole ratio of DMPC to DHPC according to the following equation:

174
$$q = \frac{[DMPC]}{[DHPC]}$$
Eq. 2

DMPC-C$_8$E$_5$ samples with the following **q**-values were prepared: 0.05, 0.10, 0.15, 0.20, 0.30, 0.35,

0.40, 0.45, and 0.50 with a total lipid-detergent concentration kept constant at 5% (w/w). Samples

were prepared on a 12 mL volume scale. To nine 15-mL centrifugal tubes, DMPC was added to

achieve the bicelle **q**-values described above. Next, 11.10 mL of Millipore water was added

followed by 300 µL of 400 mM HEPES, 4.0 M NaCl pH 7.4 buffer. The DMPC mixture was

vortexed for several seconds to create a milky, white suspension. To each of the samples, C_8E_5

was added in quantities that yielded q-values of 0.05, 0.10, 0.15, 0.20, 0.25, 0.30, 0.35, 0.40,

0.45, and 0.50, respectively. The samples were vortexed after adding C_8E_5 and turned clear.

To determine the size dependence of DMPC-C_8E_5 lipid complexes on the total lipid-detergent

concentration six samples were prepared on a 12 mL volume scale with a total lipid concentration

of 1, 5, 10, 15, 20 and 25% (w/w). The q-value was held constant at 0.25 for all six samples: To

six 15 mL conical tubes DMPC was added to produce a final lipid-detergent concentration of 1, 5,

10, 15, 20, 25% (w/w) followed by the appropriate quantity of Millipore water. To each of the six

samples 300 µL of 40X HEPES; NaCl pH 7.4 buffer was added to a final concentration of 10 mM

HEPES; 100 mM NaCl pH 7.4 was also added to each sample. The samples were vortexed to a

milky, white suspension and C_8E_5 was added in the proper amounts for a final lipid-detergent

concentration of 1, 5, 10, 15, 20, and 25% (w/w) and a q-value of 0.25. The samples were

vortexed briefly until clear.

Viscosity measurements were carried out on a Cannon Fenske viscometer at a temperature

of 20 °C and dynamic light scattering experiments were carried out on an ALV-CGS3 compact

goniometer equipped with a 22 mW HeNe laser at a wavelength of 632.8 nm. The scattered light

intensity was measured at a detector angle of 90 degrees for eight samples. Diffusion constants

for the lipid-detergent complexes were acquired from the decay of the autocorrelation function,

and the hydrodynamic radii were determined by the Stokes-Einstein equation for a spherical

particle [18].   Results are summarized in Tables 1 and 2.

2.3   Curve fitting and shape analysis

The shape of the lipid-detergent complexes was deduced following the detailed method of

Mazer and co-workers to characterize mixed micelle formation in bile salt-lecithin solutions. This

method was used by Glover and co-workers to deduce the discoidal shape of DMPC-DHPC

bicelles [4, 19]. The lipid-detergent solutions were treated as monodisperse and non-interacting

where the mean scattering intensity, I, is given by the following equation:

$$I = \bar{C}MP$$ Eq. 3

where C is the concentration of the lipid-detergent *(w/w)*, M is the molecular weight of the lipid-

detergent complex, and P describes the scattering form factor. The quantity I/C is proportional to

MP. Using this approach, the product MP will have different values depending on the shape of

the lipid detergent complex. It follows that the quantity I/C, which is equivalent to MP, can be

measured experimentally for DMPC-C_8E_5 complexes and the shape deduced from a semi-log plot

of the normalized I/C versus R_h. The dependence of I/C on R_h was fitted to one of three curves

that were generated for complexes with a spherical shape (Figure 4 - solid line), disk shape

(Figure 4 - dashed line), and a rod shape (Figure 4 - dotted line).

2.   Results and Discussion

3.1   Sedimentation equilibrium

Density matching of DMPC-C_8E_5 complexes

All sedimentation equilibrium experiments were performed at 25 °C using a Beckman XL-A

analytical ultracentrifuge. To follow the sedimentation of the lipid complexes using absorbance

optics, an NBD-labeled phospholipid probe was incorporated in a trace amount, 1:10,000 mole

ratio of NBD-PE to DMPC.

Sedimentation equilibrium data was collected for DMPC-C_8E_5 lipid-detergent complexes at a

q-value of 0.25 and a total lipid concentration of 5% (w/w). A total of nine samples were prepared

at D_2O concentrations of 0.00, 3.25, 6.50, 9.75, 13.00, 16.25, 19.50, 22.75, and 26.00% D_2O (v/v).

Sedimentation profiles were generated for all nine samples at rotor speeds of 15,000 rpm (18,112

x g), 20,000 rpm (32,198 x g), 25,000 rpm (50,310 x g) and 30,000 rpm (72,446 x g). The

sedimentation profiles were mathematically treated using the Lamm equation (Eq. 1) and the

effective molecular weight, M_{eff}, was calculated for each sample at each D_2O concentration. M_{eff}

was plotted against the percentage D_2O concentration to determine how much was required to

density match the DMPC-C_8E_5 or reach M_{eff} = 0. Figure 5 shows that approximately 18.4% D_2O

(v/v) is required to be matched, much lower than the 71.7% required to match DMPC-DHPC

bicelles [2]. High quantities of density modifiers introduce the risk of generating a concentration

gradient of the density modifier at high rotor speeds, which are commonly required when

sedimenting small proteins. Thus, DMPC-C_8E_5 lipid-detergent complexes are advantageous over

DMPC-DHPC bicelles for this reason.

3.2   Analysis of DMPC in bicelles and C_8E_5 complexes

To probe the lipid environment of DMPC in both DMPC-DHPC bicelles and C_8E_5 complexes,

[31]P-phosphorus NMR was used following the method of Glover and co-workers [4]. We reasoned

that if DMPC assumed a similar arrangement in DMPC-C_8E_5 complexes then it follows that the

chemical shifts would be the same or very similar to that of DMPC-DHPC bicelles. Figure 6 shows

an overlay of both spectra at 25 °C and 37 °C (Figures 6A and 6B, respectively). We can see

from these spectra that the chemical shift of DMPC is very similar to its chemical shift in DMPC-

DHPC bicelles with only a slight shift. These experiments were run at both 25 °C and 37 °C to

determine if there were any temperature-related effects, and although at higher temperatures this

shift in the signal becomes more pronounced, there is little indication that temperature largely

influences the lipid environment of DMPC in either case. We would predict DMPC to have a

largely different chemical shift compared to DMPC-DHPC bicelles were it not associated with

C_8E_5 in a bicelle-like environment, though it is unclear what the magnitude of that shift would be.

The close similarity of the chemical shifts suggests that DMPC occupies a bilayer region in both

bicelles and C_8E_5 complexes at *q* = 0.5.

 3.3 Dynamic light scattering to evaluate complex size and shape

To investigate the shape morphology of DMPC-C_8E_5 complexes, light scattering studies were

undertaken using the method of Mazer and Glover [4, 19]. Samples were prepared at a ***q***-value

of 0.25 at various % total lipid-detergent compositions ranging from 1% up to 25% (w/w). At a

total lipid-detergent concentration of 25% (w/w), dynamic light scattering data could not be

acquired because the lower limit of size detection had been reached. This trend in decreasing

complex size could be seen across the different samples (Figure 3). One plausible explanation

for this is that the concentration of freely available C_8E_5 in solution increases as the total lipid-

detergent concentration goes up. When this happens, the resulting size of the complexes

decreases because there is more freely available C_8E_5 in solution that can associate with DMPC

to form smaller complexes.

From the Stoke's-Einstein equation and the decay of the autocorrelation function, the

hydrodynamic radius, R_h, of the DMPC-C_8E_5 complexes was determined for each lipid-detergent

sample and plotted versus ***q***-value. Figure 2 shows the size of the complexes, R_h, increases with

respect to ***q***. A similar increase in the hydrodynamic radius was reported for DMPC-DHPC

bicelles, which initially supported our hypothesis that DMPC-C_8E_5 lipid-detergent structures

assume a bicellar-like arrangement. To further investigate the shape of these structures we

followed the method of Mazer and co-workers which was later applied by Glover et al. in detail [4,

19]. Using this approach, three curves were generated. Each describes the behavior of model

spherical complexes (solid line), discoidal complexes (dashed line), and rod-like complexes

(dotted line) (Figure 4). The molecular weight, M, was calculated for each type of complex along

with a unique form factor, P. The product of the molecular weight and the form factor, MP, is

equivalent to the experimentally determined scattering intensity, I, divided by the lipid-detergent

concentration, C, for the DMPC-C_8E_5 structures:

$$\frac{I}{C} = MP$$

<div align="right">Eq. 4</div>

For the curves, the product of the molecular weight and form factor, MP, were calculated and

plotted on a semi-log plot versus the hydrodynamic radius for spheres, disks, and rods. The range

for the hydrodynamic radius was determined based on the measured hydrodynamic radius, R_h,

for the DMPC-C_8E_5 samples. These R_h values were substituted into the values of R_h for each of

the models and the calculated scattering intensity was plotted in Figure 4. We chose to use

dimensions of DMPC that corresponded to the previous findings by Glover and co-workers, which

agreed with the calculations introduced by Vold and Prosser [4, 5, 20]. For the disk and rod-

shaped models, the thickness, t, and the diameter, d, were kept constant at 5 nm, governed by

the length of DMPC [4]. In these two models the radius of the disk, r, and the length, L, of the

rod-shaped complexes were adjusted to the measured values of the radius for the DMPC-C_8E_5

complexes. For the spherical model, the radius, *a*, was left to float between values of 1 and 10.

The data were normalized to the smallest value and ten points were obtained for each curve.

For DMPC-C_8E_5 the scattering intensity, I, was divided by the total lipid-detergent

concentration, C, for each **q**-value. The data were normalized to the lowest scattering intensity

and plotted against the measured hydrodynamic radius, R_h. Our analysis revealed that when the

normalized scattered light intensity was plotted on a semi-log plot versus the hydrodynamic radius

for DMPC-C_8E_5, the resulting curve overlays remarkably well with the curve for a spherical micellar

complex. A result that is more consistent with a micellar-like complex and not a disk-shaped

complex. All C_8E_5 lipid samples were kept at a **q** ≤ 0.5 due to observed phase changes and

reported changes in solution viscosity at higher **q** (Table 1). Viscosity also increased substantially

when **q** was held constant at 0.25 and the total lipid concentration, [DMPC + C_8E_5], was varied

from 1 to 25% (w/w) (Table 2).

DHPC and C_8E_5 have very different reported CMC values at approximately 16 mM and 8 mM,

respectively [14, 21, 22]. The differences in CMC may account for the viscosity differences

observed and measured for DMPC-C_8E_5 at higher q. The lower CMC of C_8E_5 means it more

readily forms micelles in solution. This means less free C_8E_5 is available to associate with DMPC

potentially leading to a much higher q_{eff} for the DMPC-C_8E_5 lipid complexes. For bicelles at high

q, ≥ 3.0, they assume more gel-like physical characteristics, which can be directly attributed to a

lower availability of free DHPC in solution (see Eq. 2) [23]. All samples prepared with C_8E_5 in this

study were carried out at concentrations above the CMC.

3.   Conclusions

In this study a new lipid bicelle system was prepared using the phospholipid, DMPC and

detergent, C_8E_5. Two important parameters that are used to describe bicelles were varied and

the biophysical properties of the resulting complexes were evaluated; the q-value and the total

lipid concentration (% w/w) defined as the sum of the concentrations of DMPC and C_8E_5, [DMPC

+ C_8E_5]. The properties of the lipid-detergent complexes were characterized and found to assume

a more sphere-like structure based on modeling studies using light scattering rather than disc-

shaped complexes adopted by DMPC-DHPC bicelles. Further, the hydrodynamic radius, R_h, of

the C_8E_5 complexes at q = 0.5 was reported to be approximately 5.25 nm, which is more

consistent with what we expect for detergent micelle complexes rather than bicelles, which tend

to be larger at a similar q.   DMPC-C_8E_5 complexes were density matched in the analytical

ultracentrifuge suggesting that they can be used for sedimentation equilibrium analysis aimed at

studying membrane protein oligomerization. The significant increase in solution viscosity at q-

values greater than 0.5 suggests a "solution dynamic range" that is below that for DMPC-DHPC

bicelles. Though we have used parameters typically reserved to describe the properties of DMPC-

DHPC bicelles, it is important to note that this new system may not behave in the same way and

necessitates further characterization of its physical properties expanding into a broader range of

q-values. Additionally, an investigation into their adaptability to NMR analysis of membrane

proteins will be beneficial for structural studies.

4.   **Acknowledgements**

I would like to thank those who supported this work by providing access to instrumentation and

data collection. I would also like to thank the reviewers, past and present, who took the time to

critically read versions of the manuscript and offer valuable feedback. Lastly, thank you to the

Chemistry Department at Southern Illinois University, Edwardsville.

Table 1. Viscosity changes as a function of *q*. Viscosity increases with *q*.

q value, 5% lipid (w/w)	Dynamic Viscosity (Pa * s)	Viscosity (cP)
0.05	0.00128	1.27659
0.10	0.00130	1.29977
0.15	0.00129	1.29047
0.20	0.00130	1.29959
0.25	0.00129	1.29459
0.30	0.00131	1.31296
0.35	0.00135	1.35020
0.40	0.00132	1.32339
0.45	0.00150	1.50336
0.50	0.00151	1.51056

Table 2. Viscosity and R_h changes as a function of total lipid concentration.

q = 0.25, % lipid (w/w)	Dynamic Viscosity (Pa*s)	Viscosity (cP)	R_h (nm)
1	0.00109	1.09144	8.69
5	0.00129	1.28765	4.84
10	0.00182	1.82314	2.22
15	0.00286	2.85859	1.33
20	0.00441	4.41406	0.82
25	0.00674	6.73514	N/A

Figure 1. Schematic diagram of a $q \geq 0.5$ bicelle containing DMPC and DHPC. Bicelles with a $q \geq 0.5$ have an appreciable bilayer region. C_8E_5 lipid complex samples at $q \leq 0.5$ contain the C_8E_5 detergent on the rim instead of DHPC.

Figure 2. Dynamic light scattering data measuring R_h as a function of $\boldsymbol{q}$. Increasing $\boldsymbol{q}$ corresponds to an increase in the hydrodynamic radius of the particle for DMPC-C_8E_5 complexes. All $\boldsymbol{q}$ values were kept below 0.5.

358   **Figure 3.** Dynamic light scattering data measuring R_h as a function of % total lipid (w/w). All samples were prepared at a ***q*** = 0.25. Particle size decreases as total lipid concentration increases.

Figure 4. Semi-log plot of normalized scattering intensity versus radius of hydration, R_h. The solid line represents the model of a spherical particle, the dashed line represents the model of a disk, the dotted line represents the model of a rod. The dotted-square line represents the DLS data. The DLS data fit the curve for the sphere indicating the shape of the DMPC- C_8E_5 complexes favor spheres over discoidal-shaped complexes at $q \leq 0.5$.

Figure 5. M_{eff} versus D_2O concentration. Approximately 18.4% D_2O is required to density match DMPC-C_8E_5 lipid-detergent complexes prepared at a **q** = 0.25.

Figure 6. ^{31}P-phosphorus NMR spectra of DMPC in C_8E_5 lipid complexes (blue) and bicelles (red) at 25 ºC (A) and 37 ºC (B). DMPC-DHPC bicelles and DMPC-C_8E_5 lipid complexes were prepared at a *q* of 0.5. The chemical shift of DHPC is approximately 35.1 ppm in bicelles at both 25 and 37 ºC. The chemical shift of DMPC appears at 41.0 ppm at 37 ºC and 42.0 ppm at 25 ºC in bicelles. In C_8E_5 complexes DMPC appears at 42.5 ppm at both 37 and 25 ºC. The inset in each panels shows a schematic visualization of how small bicelles, *q* ≤ 0.5 (top), might look with a spherical morphology and a narrow bilayered region compared with larger bicelles, *q* ≥ 0.5 (bottom), with a broader planar region.

References

[1] Sanders CR, Prosser RS. 1998. Bicelles: a model membrane system for all seasons? Structure. 6:1227–34. doi: 10.1016/S0969-2126(98)00123-3.

[2] Rieth MD, Root KT, and Glover KJ. 2020. "Reconstitution of full-length human caveolin-1 into phospholipid bicelles: Validation by analytical ultracentrifugation." *Biophys. Chem.* 259: 106339. doi: 10.1016/j.bpc.2020.106339.

[3] Sanders CR and Prestegard JH. 1990. "Magnetically orientable phospholipid bilayers containing small amounts of a bile salt analogue, CHAPSO." *Biophys. J.* 58: 447–460. doi: 10.1016/S0006-3495(90)82390-0.

[4] Glover KJ, Whiles JA, Wu G, Yu N, Deems R, Struppe JO, Stark RE, Komives EA, Vold RR. 2001. "Structural evaluation of phospholipid bicelles for solution-state studies of membrane-associated biomolecules." *Biophys. J.* 81: 2163–2171. doi: 10.1016/S0006-3495(01)75864-X.

[5] Struppe J, Whiles JA, and Vold RR. 2000. "Acidic phospholipid bicelles: a versatile model membrane system." *Biophys. J.* 78: 281–289. doi: 10.1016/S0006-3495(00)76591-X.

[6] Sanders CR. 2006. Development and Application of Bicelles for Use in Biological NMR and Other Biophysical Studies. In: Webb G.A. (eds) Modern Magnetic Resonance. Springer, Dordrecht. https://doi.org/10.1007/1-4020-3910-7_28

[7] Lu Z, Van Horn WD, Chen J, Mathew S, Zent R, and Sanders CR. 2012. "Bicelles at low concentrations." *Mol. Pharm.* 9(4): 752–761. doi: 10.1021/mp2004687.

[8] Semchyschyn DJ and Macdonald PM. 2004. "Conformational response of the phosphatidylcholine headgroup to bilayer surface charge: Torsion angle constraints from dipolar and quadrupolar couplings in bicelles." *Magn. Reson. Chem.* 42(2): 89–104. doi: 10.1002/mrc.1333.

[9] Yang Z, Wang C, Zhou Q, An J, Hildebrandt E, Aleksandrov LA, Kappes JC, DeLucas LJ, Riordan JR, Urbatsch IL, Hunt JF, Brouillette CG. 2014. "Membrane protein stability can be compromised by detergent interactions with the extramembranous soluble domains." *Protein Sci.* 23(6): 769–789. doi: 10.1002/pro.2460.

[10] Agharkar A, Rzadkowolski J, McBroom M, and Gonzales EB. 2014. "Detergent screening of the human voltage-gated proton channel using fluorescence-detection size-exclusion chromatography." *Protein Sci.* 23(8): 1136–1147. doi: 10.1002/pro.2492.

418 [11] Arachea BT, Sun Z, Potente N, Malik R, Isailovic D, and Viola RE. 2012. "Detergent
419   selection for enhanced extraction of membrane proteins." *Protein Expr. Purif.* 86(1): 12–20.
420   doi: 10.1016/j.pep.2012.08.016.
421

422 [12] Tzitzilonis C, Eichmann C, Maslennikov I, Choe S, and Riek R. 2013. "Detergent/Nanodisc
423   Screening for High-Resolution NMR Studies of an Integral Membrane Protein Containing a
424   Cytoplasmic Domain." *PLoS ONE.* 8(1): e54378. doi: 10.1371/journal.pone.0054378.
425

426 [13] Chen L, Lai C, Lai J, and Tian C. 2011. "Expression, purification, detergent screening and
427   solution NMR backbone assignment of the human potassium channel accessory subunit
428   MiRP1," *Protein Expr. Purif.*, 76(2): 205–210. doi: 10.1016/j.pep.2010.11.007.
429

430 [14] Beaugrand M, Arnold AA, Hénin J, Warschawski DE, Williamson PTF, and Marcotte I.
431   2014. "Lipid Concentration and Molar Ratio Boundaries for the Use of Isotropic Bicelles,"
432   *Langmuir*, 30(21): 6162–6170. doi: 10.1021/la5004353.
433

434 [15] Rieth MD, Lee J, and Glover KJ. 2012. "Probing the caveolin-1 P132L mutant: Critical
435   insights into its oligomeric behavior and structure." *Biochemistry*, 51(18): 3911-18.   doi:
436   10.1021/bi3001853.
437

438 [16] Burgess NK, Stanley AM, and Fleming KG. 2008. "Determination of membrane protein
439   molecular weights and association equilibrium constants using sedimentation equilibrium
440   and sedimentation velocity." *Methods Cell Biol*, 84: 181–211.
441

442 [17] Kumar VV and Baumann WJ. 1991. "Lanthanide-induced phosphorus-31 NMR downfield
443   chemical shifts of lysophosphatidylcholines are sensitive to lysophospholipid critical micelle
444   concentration," *Biophys. J.* 59(1): 103–107. doi: 10.1016/S0006-3495(91)82202-0.
445

446 [18] Chu B. 1970. "Laser Light Scattering." *Annu. Rev. Phys. Chem.*, 21(1): 145–174. doi:
447   10.1146/annurev.pc.21.100170.001045.
448

449 [19] Mazer NA, Benedek GB, and Carey MC. 1980. "Quasi-Elastic Light-Scattering-Studies of
450   Aqueous Biliary Lipid Systems - Mixed Micelle Formation in Bile-Salt Lecithin Solutions,"
451   *Biochemistry.* (19)4: 601–615. doi: 10.1021/bi00545a001.
452
453 [20] Vold RR and Prosser RS. 1996. "Magnetically oriented phospholipid bilayered micelles for
454   structural studies of polypeptides. Does the ideal bicelle exist?" *J. Magn. Reson. - Ser. B.*
455   113(3): 267–271. doi: 10.1006/jmrb.1996.0187.
456

457 [21] Marsh D. *Handbook of lipid bilayers.* CRC press, 2019.
458

459   [22] D'Errico G, Ortona GO, Paduano L, Tedeschi A, and Vitagliano V. 2002. "Mixed micellar
460   aggregates of cationic and nonionic surfactants with short hydrophobic tails. An
461   intradiffusion study," *Phys. Chem. Chem. Phys.* 4(21): 5317–5324. doi:
462   10.1039/B202731C.
463

464   [23] van Dam L, Karlsson G, and Edwards K. 2004. "Direct observation and characterization of
465   DMPC/DHPC aggregates under conditions relevant for biological solution NMR." *Biochim.*
466   *Biophys. Acta BBA - Biomembr.* 1664(2): 241–256. doi: 10.1016/j.bbamem.2004.06.005.
467