Title:

Inferring the heritability of large-scale functional networks with a multivariate ACE

modeling approach

Authors:

Fernanda L. Ribeiro[1,2*], Felipe R. C. dos Santos[3,4*], João R. Sato[1], Walter H. L. Pinaya[1,5†],

Claudinei E. Biazoli Jr.[1†]

1 – Center of Mathematics, Computing and Cognition, Universidade Federal do ABC;

São Bernardo do Campo 09606-045; Brazil

2 – School of Psychology, The University of Queensland; Brisbane QLD 4072; Australia

3 – Centro de Oncologia Molecular, Hospital Sírio-Libanês; São Paulo 01308-060; Brazil

4 – Programa Interunidades em Bioinformatica, Universidade de São Paulo; São Paulo

05508-220; Brazil

5 – Department of Biomedical Engineering; School of Biomedical Engineering &

Imaging Sciences; King's College London; London WC2R 2LS; United Kingdom

*These authors contributed equally to this work.

†These authors contributed equally to this work.

Corresponding author:

Fernanda L. Ribeiro

E-mail address: fernanda.ribeiro@uq.edu.au

Postal address: School of Psychology, The University of Queensland; Brisbane QLD
4072; Australia

Abstract

Recent evidence suggests that the functional connectome is stable at different time scales

and unique. These characteristics posit the functional connectome not only as an

individual marker but also as a powerful discriminatory measure characterized by the

high intersubject variability. Among distinct sources of intersubject variability, the long-

term sources include functional patterns that emerge from genetic factors. Here, we

sought to investigate the contribution of genetic factors to the variability of functional

networks by determining the heritability of the connectivity strength in a multivariate

fashion. First, we reproduced and extended the connectome fingerprinting analysis to the

identification of twin pairs. Then, we estimated the heritability of functional networks by

a multivariate ACE modeling approach with bootstrapping. We found that a visual (0.41)

and the medial frontal (0.35) functional networks were the most heritable, while the

subcortical-cerebellum (28.6%) and the medial frontal (21.1%) networks were the most

accurate on twin pair identification. Taken together, our findings suggest that twin

identification accuracy does not necessarily relate to the heritability of a given functional

network, indicating that heritability estimation and connectome fingerprinting are both

required to study the influence of genetic factors on the functional organization of human

brain at the level of large-scale networks.

Keywords: Connectome fingerprint; Multivariate modeling; Twin study

Introduction

In the past few years, fMRI research has been living a paradigm shift, moving from population inferences to the study of individual differences[1,2]. Previous studies have paved the way for the study of individual variability in functional connectivity patterns of the human brain[3–5]. In this context, resting-state fMRI (rs-fMRI) showed to be particularly powerful in determining underlying differences in the wiring patterns of functional connectome (FC) profiles. Indeed, connectome-based individual predictions achieved identification accuracies as high as 99% when comparing functional connectivity matrices[4]. Hence, the endeavor to identify and to characterize the individual functional connectivity architecture has been shown to have an imperative place in the study of individual differences.

Recent and mounting evidence suggests that FC profiles are stable at different time scales[6–9]. This characteristic posits the FC not only as an individual marker due to the comparably low intrasubject variability but also as a powerful discriminatory measure characterized by the high intersubject variability. Gratton et al. (2018) showed that despite functional networks displaying common organizational features at the group-level, the similarity between functional networks substantially increased at the individual level when evaluating the same participant in different tasks and sessions. This evidence supports the fact that individual stable patterns are crucial to explain the intersubject variability of functional networks. Therefore, these findings suggest that sources of intersubject variability are stable over time, acting as individual signatures or 'fingerprints'.

Seghier and Price (2018) refer to the presence of distinct sources of intersubject variability that differ in their timescale. In the lower bound, there are sources of variability due to

mood states and context. The medium to long-term sources of intersubject variability include functional patterns built from the intimate interaction of an individual with the environment and genetic factors[1], respectively. Interestingly, functional networks show distinct levels of intersubject variability. Networks comprising higher-order associative cortical areas seem to remarkably contribute to the FC distinctiveness [3–6,9,10], which, in turn, might be due to a high intersubject[5,8] and low intrasubject variability[11,12]. On the other hand, functional connectivity within networks that comprises primary sensory and motor regions showed high intrasubject and low intersubject variability[5,8,11,12]. The importance of genetic factors to these different levels of intersubject variability, however, is yet to be further investigated.

Recent reports suggest that genetic factors crucially influence the intersubject variability in the functional connectome[6,13–17]. Connectome-based identification analyses were extended to the identification of twin pairs suggesting that part of the intersubject variability is due to genetic factors[6,17]. Accordingly, studies indicate that the average heritability of the connectivity strength of the whole brain connectome is between 15% to 25% within the Human Connectome Project dataset[13,16,18]. On the other hand, the heritability of the connectivity strength within some functional networks seems to be much higher[14,19] than the whole brain connectome.

However, exclusive univariate approaches, in association with substantial differences in brain parcellation schemas[20–22], undermines the effort to determine the relationship between heritability and the different levels of intersubject variability. Crucially, such univariate approaches may bias heritability estimates of large-scale functional networks, considering that the reliability of the connectivity strength of individual edges is low[23,24]. Here, we sought to overcome this caveat by quantifying the heritability of functional networks using a multivariate ACE modeling approach. To this end, we (1) reproduced

90 and extended the identification analysis introduced by Finn et al. (2015) to determine the

91 functional networks that best uncovered individual uniqueness and intersubject similarity

92 among matched twin pairs, and (2) we investigated how the different levels of intersubject

93 variability of functional networks relate to their heritability.

Results

Functional connectivity based identification analyses

Individual identification. Whole-brain functional connectivity matrices were determined by using two distinct parcellation schemas: "Shen"[25] (268 nodes, 71,824 edges) and "Gordon"[26] (333 nodes, 110,889 edges). For brevity, we only report the results using "Shen" parcels with appropriate reference to equivalent results using "Gordon" parcels in the supplementary material. Connectivity-based identifications were performed comparing pairs of resting-state functional connectivity matrices[4]. Resting-state data were acquired in two different days for every participant included in this study, resulting in two distinct functional connectivity matrices per participant. These pairs of connectivity matrices were separated into a 'target' and a 'database' set. Individual identification was determined by computing the Pearson's correlation score of a target connectivity matrix from the 'target' set (n=380) with all connectivity matrices from the 'database' set (n=380). Following that, the maximum correlation score among all comparisons between the target matrix and each of the FC matrices from the 'database' set should correspond to the correlation of the functional connectivity matrices of the same participant in different sessions. This process was repeated for all functional connectivity matrices within the 'target' set (Figure 1A). The accuracy of the method was defined by the proportion of correct predicted participants.

Individual identification analyses were determined with whole-brain functional connectome and individual functional networks (Supplementary Table 1). The resulting accuracy of whole-brain connectome based individual predictions was 97.8% (SD = 0.4%), in agreement with previous studies[4,23]. We also investigated the relevance of individual functional networks for individual predictions by sectioning the whole-brain

functional connectome into sub-matrices of single networks. From the 8 functional

networks previously defined[4], the most successful networks were the medial frontal (93.9

$\pm 0.5\%$), and frontoparietal ($91.8 \pm 0.3\%$) networks (Figure 1B and Supplementary Table

1). Note that the visual networks and the default mode network were the ones with the

worst individual identification accuracy.

Figure 1 - Connectome-based identifications. A) Functional connectivity matrices from
different sessions were grouped into two datasets, which could be either the 'target' set or the
'database'. Following that, we computed the Pearson's correlation of each individual connectivity
matrix from a 'target' set with each connectivity matrix from the 'database'. Therefore, each row
within the individuals vs. individuals matrix contains the correlation scores between a target's FC
and all functional connectivity matrices of the database. **B)** Mean identification accuracies for
individual and twin identification analyses for all functional networks (whole-brain included).
Mean identification for individual prediction was determined from two combinations of
'database' and 'target' sets (RESTX × RESTY, where X and Y ∈ {1, 2} and X≠Y), while the
mean twin identification was determined from four combinations (RESTX × RESTY, where X
and Y ∈ {1, 2}). Error bars represent the standard deviation. All, whole-brain; MF, medial frontal;
FP, frontoparietal; SC, subcortical-cerebellum; M, motor; VI, visual I; DMN, default mode
network; VA, visual association; VII, visual II. We also present the number of nodes in each
network.

Twin identification. Previous studies indicate that functional connectivity among higher-

order associative brain regions greatly varies across individuals[5,8], even though they are

comparably more stable within an individual across sessions[12,13]. Thus, we hypothesized

that genetic factors governed sources of high intersubject and low intrasubject variability

in the functional connectome. In order to test this hypothesis, we sought to determine

whether the FC profiles from pairs of twins were more similar compared to the ones from

pairs of unrelated individuals by using connectome-based predictions.

First, we evaluated monozygotic twin identification by computing the correlation

coefficients of the functional connectivity matrices of monozygotic individuals (n=246)

within the 'target' set with all matrices in the database (246x380=93,480 comparisons).

Our prediction was based on the selection of the highest correlation score (excluding the

correlation scores between functional connectivity matrices of the same individual) for

each 'target' participant vs. 'database' iteration. The mean whole-brain based prediction

accuracy was 57.2% (SD = 2.6%). This result indicates that the idiosyncratic FC profiles

might be genetically determined and they are sufficiently stable so one could identify

monozygotic twins well above chance. Indeed, we have performed a permutation test, by

exchanging twin pairs' identities 1,000 times, such that for each identification iteration,

a new twin pair identity was assigned. The maximum identification accuracy found

through these 1,000 permutations was 1.6%, indicating that the whole-brain based

identification performance is significantly different from the chance level (p-value <

0.001).

Later on, we investigated the ability of specific functional networks in discriminating a

twin pair from pairs of unrelated individuals (Figure 1B). At this stage, the most

successful functional networks were the subcortical-cerebellum ($28.6 \pm 1.5\%$) and medial

frontal ($21.1 \pm 2.2\%$) networks. Noteworthy, the most successful functional networks on

twin identification were amongst the ones that best performed on individual

identifications. Nonetheless, a substantial decrease in the successful twin identification

rates was observed for functional networks when compared to the whole-brain

connectome, and these results were particularly affected by the number of nodes within

each network. The least successful functional networks on twin identification were the

ones with the least number of nodes, while the networks with a larger number of nodes

tended to present higher accuracies. The Pearson correlation score between the number

of nodes of each network and its ability to correctly identify monozygotic twins (Figure

2) was r = 0.95 (p-value = 6.3E-5; Supplementary Table 2), as opposed to a nonsignificant

correlation between the number of nodes and individual identification accuracy (r = 0.52,

p-value = 0.15). This implies that the ability of *a priori* defined functional networks to

capture similarities in the FC profiles of monozygotic twins differentially relies on the

amount of information provided (i.e., by the number of nodes).

Finally, we performed all the previous analyses for the identification of dizygotic twins.

At this time, we selected only the dizygotic individuals (n=134) within the 'target' set,

giving 134x380=50,920 comparisons. For the whole-brain based identification, the mean

prediction accuracy was 8.9% (SD = 2.3%; p-value <0.001). This abrupt change in twin

identification accuracy indicates that the functional connectivity patterns of monozygotic

twins are strictly more similar in comparison to dizygotic twins, which indicates the

relevance of shared genetic background. At the level of individual functional networks,

identification accuracies dropped even further (Figure 1B), and they were also correlated

with the number of nodes of the networks (r = 0.92, p-value < 0.001).

Intra and intersubject variability in the functional connectome

In order to characterize the intra and intersubject variabilities (i.e. among unrelated

individuals, monozygotic and dizygotic twin pairs) for the whole-brain connectome and

each functional network, we arranged the correlation coefficients in four groups

according to their relationship: 1) same individual - SI (n=380); 2) monozygotic twins -

MZ (n=246); 3) dizygotic twins - DZ (n=134) and 4) unrelated individuals - UN

(n=143,640). The distributions of correlations across all these pairs for the whole-brain

and functional networks are illustrated in Figure 2 (Supplementary Figure 1).

Figure 2 - Distribution of correlation coefficients between pairs of functional connectivity matrices for the whole-brain and individual functional networks. Pearson's correlation scores were determined from pairs of connectivity matrices (REST1 × REST2), and they were grouped based on individuals' genetic relationship. Hence, violin plots show the distribution of the correlation scores between pairs of matrices of the same individual (SI), monozygotic twin (MZ), dizygotic twin (DZ) and unrelated individuals (UN).

As one could expect, the mean of the distributions of correlation scores from the SI group

is notably higher than the ones from the remaining groups. This is observed not only for

the whole-brain connectome but also for most of the functional networks, especially for

the medial frontal and frontoparietal functional networks. In order to characterize the

importance of the distance between these distributions - that is, the effect size - to

identification analyses, we determined identification accuracy as a function of effect size,

Cliff's delta[24] (Figure 3, Supplementary Figure 2 and Supplementary Table 3). In Figure

3, we observe that high prediction accuracy is associated with high effect size, while low

prediction accuracy was associated with low effect size. This suggests that high

intersubject variability (which is related to low correlation between unrelated individuals'

connectivity matrices) and low intrasubject variability (high correlation between the

connectivity matrices of the same individual in a different session) are crucial for high

prediction accuracy. Additionally, the higher similarity between monozygotic twins in

comparison to unrelated individuals (medium to high effect sizes) suggests that a portion

of this intersubject variability is heritable and differs across functional networks.

Figure 3 - Dependence of connectome-based predictions on effect size. Mean prediction
accuracies from all functional networks (whole-brain included) as a function of the effect size of
the difference between the group of interest (same individual – SI, monozygotic twins – MZ, or
dizygotic twins – DZ) and unrelated individuals.

Heritability of the functional connectome

To further investigate these functional networks, we performed a heritability analysis

using a multivariate ACE modeling approach with bootstrapping. High dimensionality is

a common hurdle when multivariate processing is considered for regression or inference

methods. Hence, univariate analyses are usually preferred to avoid the necessity of

increasing computational resources and time due to high dimensional multivariate

analyses trade-off, despite the fact that multivariate analyses tend to be more suitable for

complex data that includes several thousand of covariates. In neuroscience, the

heritability of functional networks is usually determined as the average heritability of

individual functional connections (edges) over their constituent brain regions

(nodes)[13,14,16]. In order to overcome such a challenge, we fitted a lower-dimensional

multivariate ACE model several times to obtain a distribution of heritability estimates for

each functional network (Figure 4).

Figure 4 – Multivariate ACE model with bootstrapping. A) The lower triangles of mean
functional connectivity matrices were vectorized, and the effect of age and sex were regressed out
from each edge. In an iterative process, 10 edges were randomly selected and used as covariates
to fit an ACE model. This procedure was repeated with reposition for 8,000 times for whole-brain
(or 1,000 times for each functional network). **B)** This approach provides a distribution of
heritability estimates for each functional network by considering the pattern of covariation among

edges. **C)** Null distributions were similarly obtained by randomly shuffling monozygotic and
dizygotic twin statuses at each iteration.

This multivariate approach involved the random selection of 10 edges (within the

functional network of interest) that were used as covariates to fit a multivariate ACE

model (Figure 4A). This procedure was repeated with reposition for 8,000 times for

whole-brain and 1,000 times for each functional network, and it provided a distribution

of heritability estimates, "A" - in addition to "C", shared environment, and "E",

measurement error, or external sources of variability - for each functional network (Figure

4B). Finally, null distributions were similarly obtained by randomly shuffling

monozygotic and dizygotic twin statuses at each iteration (Figure 4C). The heritability

distributions with their respective null distributions for all functional networks are

illustrated in Figure 5A (Supplementary Figure 3). As expected, the mean heritability of

all null distributions was virtually equal to zero. Apart from that, all heritability estimates

distributions were significantly different from their respective null distributions

(independent t-test, $p<.001$). Among all functional networks, the visual II has shown to

be the most heritable with mean heritability of 0.41 (41% of the variance of the phenotype

is attributed to additive shared genetics; Supplementary Table 4), while the subcortical-

cerebellum was the least heritable with mean heritability of 0.22 (Figure 5B and

Supplementary Table 4). The heritability distributions of the whole-brain, visual I, and

visual association networks were not significantly distinct from one another

(Supplementary Table 5 and 6).

Figure 5 – Heritability distributions for each functional network. A) Heritability estimates and null distributions for each functional network. **B)** Heritability estimates distributions are plotted from the most heritable (visual II) to the least heritable (subcortical-cerebellum).

Discussion

Here, we found that the functional connectivity profiles of twin pairs were more similar

than of unrelated individuals, although the degree of similarity varied across functional

networks. Indeed, we demonstrated that functional networks have distinct discriminatory

power in connectome fingerprinting analyses, in both individual and twin identifications.

However, we found that high intersubject variability (i.e. variability of a trait between

individuals) is crucial for connectome fingerprinting. Additionally, our multivariate ACE

modeling approach suggests that the heritability of functional networks are consistent

throughout the brain, although the visual II and medial frontal functional networks were

the most heritable. Altogether, we were able to establish the influence of genetic factors

to intersubject variability of functional networks by leveraging a multivariate ACE model

in addition to the multivariate connectome fingerprinting approach.

Intra and intersubject variability trade-off in connectome fingerprinting

Evidence suggests that the different levels of inter and intrasubject variability in

functional networks contribute to their distinctiveness, such that high intersubject[5,8] and

low intrasubject[11,12] variability in higher-order associative networks are often related to

their high discriminability[3–6,9,10] and the opposite pattern to the low discriminability of

primary sensory and motor networks[5,8,11,12]. We confirmed that higher-order associative

networks were the most discriminatory, while visual networks were the least

discriminatory, although they showed similar levels of intrasubject variability. This

finding was similarly seen in twin pair identifications, although in the latter the prediction

accuracy was positively correlated with the number of nodes defining each functional

network. To further investigate the inter and intrasubject variability trade-off in

connectome fingerprinting, we determined the prediction accuracy as a function of the

difference between the similarity scores of functional networks derived from the same

individual - in different resting-state sessions - and unrelated individuals. We found that

high identification accuracy requires high intersubject variability, suggesting that

although the stability of idiosyncratic functional connectivity patterns is relevant and seen

across all functional networks, fingerprinting seems to rely prominently on high

intersubject variability.

Genetic influence on functional networks

To investigate the impact of genetic factors in determining stable patterns of intersubject variability, we performed an alternative approach to the univariate ACE model. The low reliability of individual edges' connectivity strength[23,24] suggests that a multivariate approach to estimate the heritability of functional networks is required. In our multivariate ACE model, rather than determining the heritability of the connectivity strength of individual edges and taking the average across edges within a functional network, we obtain a distribution of heritability estimates by randomly selecting a fixed number of edges to be used in our model. The contribution of this method is, thus, twofold: it takes into consideration the covariation pattern of edges within a functional network and, by doing so, is likely more reliable than estimating the heritability of individual edges.

We found that the functional networks that were the most heritable were not the ones that best performed on twin identifications. This is particularly prominent for the visual II and subcortical-cerebellum functional networks. The first has shown to be the most heritable, but it had a poor performance on monozygotic twin identification and individual identification. This indicates that the intersubject variability was low, thus being difficult to discriminate between pairs of connectomes from UN/Twin/SI groups. However, a great portion of this low intersubject variability is due to genetic factors, thus the high heritability. On the other hand, the subcortical-cerebellum network has shown low heritability but the best performance on twin identification (after whole-brain network), which means that the high intersubject variability allowed a better discrimination between unrelated individuals *versus* twin pairs, even though a smaller portion of its intersubject variability was due to genetic factors. This suggests that twin identification accuracy does not necessarily relate to the heritability of a given functional network.

Parcellation schema

The individual and twin identification analyses resulted in high prediction accuracy using both parcellation schemas, "Shen" and "Gordon". Notably, individual identification accuracies using "Shen" parcellation schema is about the same as in previous studies[4,27], even though we have a more homogenous sample. At the network-level, higher-order associative networks were particularly better at discriminations. This result further supports that associative networks accommodate higher intersubject variability in comparison to sensorimotor networks[8]. Despite that, we observed that the default mode network (DMN) defined by both parcellation schemas differed in performance during identification analyses. For "Gordon" parcels, the DMN figured among the most distinctive networks, similarly to other associative networks. However, this pattern was not observed using "Shen" parcels, in which the defined DMN figured among the worst functional networks on individual predictions. This distinction could be due to the different number of nodes attributed to DMN in both schemas. Additionally, the heritability level of functional networks differed between parcellations, although the mean heritability of the whole-brain functional network was 0.20 using "Gordon" parcels and 0.26 using "Shen" parcels (Supplementary Table 4). This suggests that different brain areas definition greatly impact on heritability estimates, which is a potential topic for further investigation.

Using "Gordon" parcellation, we found that the cingulo parietal and retrosplenial temporal networks were the most heritable networks, whilst the somato-sensory mouth and salience networks were the least heritable. On the other hand, Miranda-Dominguez et al. (2018) found that the retrosplenial and somato-sensory mouth were the most heritable, and the visual and salience networks the least heritable. Additionally, their heritability estimates ranged from 0.11 to 0.14, with the heritability of the whole-brain

network being 0.20[6]; while our estimates ranged from 0.52 to 0.10. These differences are

likely due to differences in heritability estimation approaches; whilst we used the

conventional ACE modeling approach, they used three-way repeated-measures

ANOVAs. Although the heritability estimates we obtained using "Shen" parcels were

more homogeneous, we were still able to capture the different levels of heritability of

functional networks, suggesting that our approach is more suitable for capturing such

differences. Additionally, using a similar methodology, Colclough et al. (2017) found that

the heritability of the connectivity strength averaged over parcels was 0.17 for the whole-

brain network, and Elliot et al. (2019) found a value of about 0.20. This suggests that,

although heritability estimates of functional networks vary depending on the parcellation

being used, the whole-brain functional network heritability seems to be reasonably

consistent across studies using different methodologies and parcellations.

Future directions

Our multivariate ACE model suggests that part of the intersubject variability seen in

functional networks is due to genetic factors. Transcriptomics and genomics approaches

have indicated that many brain disorders are, at least partly, determined by the genetic

background[28–31]. Additionally, disruptions in the human functional and structural

connectomes have been associated with neurological conditions, such as amyotrophic

lateral sclerosis (ALS)[32], Parkinson's disease[33,34], and epilepsy[35]. Specifically,

neurotoxic accumulation of amyloid plaques in Alzheimer's disease has been located in

areas consistent with cortical hubs, indicating that while cortical hubs are fundamental

for information processing, they also bring vulnerability to the human brain[36]. Also, many

compelling studies have linked psychiatric disorders to fundamental connectome

disruptions[37]. Despite their unique functional and structural connectivity patterns, these

conditions also exhibit some shared patterns that differ from healthy connectomes. The

common features of many of these disorders make it difficult to diagnose them and to

determine the mechanisms behind their onset, particularly for psychiatric disorders. Thus,

detailed scrutiny of the human connectome and genome may lead to a promising new era

for precision medicine in psychiatry and neurology. Connectome fingerprinting in

addition to heritability analyses may allow for the search of connectome features that

bring general and specific vulnerabilities to the human brain, which may be highly

heritable, and are central factors among brain disorders[37].

Materials and Methods

Database and participant information

In this study, we used the dataset from the "1200 subjects data release" of the Human

Connectome Project - HCP[38]. We restricted our analysis to monozygotic (MZ) and

dizygotic (DZ) individuals. So, we initially selected all MZ and DZ individuals from the

original sample. From this subsample, we excluded the participants who did not have both

resting-state fMRI sessions (ICA-FIX versions) available, and who did not have the twin

within the group. Therefore, our final sample size was n=380. Table 1 summarizes the

demographic data.

Table 1 - Demographic information.

	Monozygotic (n=246)	Dizygotic (n=134)
Age, y		
Mean ± SD	29.4 ± 3.3	29.1 ± 3.5
Range (min-max)	22 – 36	22 – 35
Sex, n (%)		
Female	144 (58.5)	78 (58.2)
Male	102 (41.5)	56 (41.8)

Data acquisition

The acquisition protocol has been previously described[38]. In summary, functional and

structural data were acquired in a 3T Siemens Skyra scanner using a 32-channel head coil.

Resting-state data were collected in two separated sessions (REST1 and REST2) in

different days, each session containing two runs of 15 minutes. In this protocol,

participants had to keep their eyes open with a relaxed fixation on a projected bright cross-

hair in a dark background. Each run within a session is distinguished by the oblique axial

acquisition, of which one run used phase encoding in a right-to-left (RL) direction and

the other used phase encoding in a left-to-right (LR) direction.

Data pre-processing

Pre-processing pipeline

For this study, we used the spatial and temporal pre-processed rs-fMRI timecourses[39,40],

which have undergone the steps of artifact removal, motion correction, and registration

to standard space. Furthermore, we applied additional pre-processing steps by using the

CONN toolbox (v.17.f)[41], which included: structural segmentation, functional outlier

detection (intermediate setting: 5 for z-score scan-to-scan global signal changes and 0.9

mm for scan-to-scan head-motion composite changes), and functional smoothing.

Following that, a component-based noise correction method (CompCor)[42] and a temporal

band-pass filtering (preserving frequencies between 0.01 and 0.10Hz) were applied. For

spatial smoothing, a Gaussian with the full width at half maximum (FWHM) equal to

6mm was used. We also included participant movement parameters as first-level

covariates to regress out their linear components from the BOLD time series.

Parcellations and functional networks

Timecourses were calculated as the mean signal within the regions of interest (ROIs)

defined by different parcellation schemas used: "Gordon"[26] and "Shen"[25]. Both "Gordon"

and "Shen" schemas are data-driven parcellation schemas. The first defines 333 ROIs

clustered in 12 functional networks (Supplementary Table 1), in addition to 47 ROIs not

assigned to any specific network. The latter defines 268 ROIs clustered in 8 networks

(Supplementary Table 1).

Functional connectivity matrices

Finally, for the two resting-state sessions, data from both the left-right (LR) and right-left

(RL) phase-encoding runs were used to calculate the connectivity matrices. To obtain the

connectivity matrices, ROI-to-ROI bivariate correlation connectivity measures were

computed for all ROIs defined by both parcellation methods, obtaining two symmetric

connectivity matrices for each session for each participant.

Individual identification

The identification analysis was based on previous work[4] with few alterations. Initially,

two databases were created containing the functional connectivity matrices for each

session (REST1 and REST2). The individual identification was determined by computing

the Pearson's correlation of each individual connectivity matrix from one database with

all the other connectivity matrices from the second database (RESTX $\times$ RESTY, where

X and Y $\in$ {1, 2} and X$\neq$Y). For a pair of functional connectivity matrices linearly

transformed in a column vector (vectorization), T_i and D_n, where T_i is the connectivity

matrix of a target participant i, and D_n is the connectivity matrix of a participant (n=1, ...,

380) from the other database, the Pearson's correlation coefficient r is:

433
$$r_{i,N} = \frac{\sum_{j=1}^{e} (T_{i_j} - \underline{T}_i)(D_{N_j} - \underline{D}_N)}{\sqrt{\sum_{j=1}^{e} (T_{i_j} - \underline{T}_i)^2} \sqrt{\sum_{j=1}^{e} (D_{N_j} - \underline{D}_N)^2}}$$
Equation 1

where e is the number of edges. In order to predict the identity of the target participant,

the maximal Pearson's correlation coefficient was selected (Figure 1A). Additionally, we

also investigated the contribution of single networks to identification accuracy by sub-

sectioning the functional connectivity matrices into sub-matrices of single networks. To

perform this, we selected only connection within a specified network. Then, we calculated

the Pearson's correlation coefficients, similarly to the previous approach. Results are

reported as mean ± SD.

Twin identification

The twin pair identification algorithm was based on the previous individual identification

analysis. At this stage, we removed the correlations corresponding to the same individual

in different sessions, that is the diagonal of individuals × individuals matrices, and then

performed a new set of identification analyses. In this condition, if the chosen maximum

correlation value belonged to the target subject's twin, the prediction was considered

correct. Monozygotic and dizygotic twins were analyzed separately, and all conditions

(RESTX × RESTY, where X and Y ∈ {1, 2}) were tested. Results are reported as mean

± SD.

Statistical significance assessment

To assess the statistical significance of twin identification analyses, we performed a

permutation testing. To ensure the independence of the dataset, we permuted the twin

pairs' identities, such that for each row of the 'individuals vs. individuals' matrix (Figure

1A) a new twin pair identity was assigned. The permutation process was repeated 1,000

times for each functional network.

Effect size

The distribution of correlation scores between pairs of connectivity matrices (i.e.

correlation among the vectorized form of the connectivity matrices) was determined by

grouping these scores based on familial relationship: 1) same individual - SI; 2)

monozygotic twins - MZ; 3) dizygotic twins - DZ and 4) unrelated individuals - UN.

Following that, the effect size of the differences between the distributions of correlation

values was measured through the calculation of Cliff's delta. This a non-parametric effect

size measure based on all pairwise differences[24], which gives how often values from one

distribution are larger than the ones from a second distribution (Equation 2).

$$Delta\ (d) = \frac{Sum(x_1 > x_2) - Sum(x_1 < x_2)}{n_1 n_2} \qquad Equation\ 2$$

Therefore, the number of times that values from one group are higher than the ones from

a second group is calculated for all possible combinations of values between the two

groups ($n_1 n_2$, where n_1 and n_2 are the number of values within the distribution 1 and 2,

respectively). The final Cliff's delta value is the difference between the previous

calculations divided by all possible combinations. Thus, a positive and high value of d

($d_{maximum} = 1$) mean that values within distribution 1 are mostly higher than the ones within

distribution 2, a negative and high absolute value of d ($d_{minimum} = -1$) means the other way

round, that values within distribution 1 are mostly lower than the ones within distribution

2, and d = 0 means that the distribution 1 and 2 are equal.

Heritability analyses

Functional connectivity measures from two different days (REST1 and REST2) were

averaged, giving a functional connectivity matrix per participant. As mentioned before,

whole-brain functional connectivity matrices were determined by using two distinct

parcellation schemas: "Shen"[25] (268 nodes, 71,824 edges) and "Gordon"[26] (333 nodes,

110,889 edges). The first step involved the vectorization of functional connectivity

matrices' lower triangle ("Shen" - 35,778 edges; "Gordon" – 55,278 edges). The

heritability analyses were performed using the umx package[43], after regressing out the

effect of age and sex using 'umx_residualize'.

Multivariate ACE models were fitted with bootstrapping. The Cholesky's ACE model[43]

used in this framework allocates observed phenotypic variability into three factors: A

(heritability – h^2), C (shared environment – c^2) and E (measurement error or external

sources of variability – e^2) in a multivariate fashion. In our iterative multivariate approach,

10 edges were randomly selected and used as covariates to fit an ACE model (umxACEv).

This procedure was repeated with reposition for 8,000 times (or 12,000 times when

"Gordon" parcels was used) for whole-brain, and 1,000 times for each functional network.

The number of iterations was determined such that every edge would be selected at least

twice (i.e. 8,000 iterations * 10 edges = 80,000). This method approach provides a

distribution of heritability estimates for each functional network by considering the

pattern of covariation among edges. Finally, null distributions were similarly obtained by

randomly shuffling monozygotic and dizygotic twin statuses at each iteration[13].

Independent t-student tests were performed separately to evaluate whether each

functional network's heritability distribution significantly differed from their respective

null distribution.

Code availability

All source codes will be available at

https://github.com/felenitaribeiro/fingerprinting_twinStudy and

https://github.com/frcsantos/heritability upon publication of this manuscript.

Citation diversity statement

Recent work in neuroscience and other fields identified a bias in citation practices such

that papers from women and other minorities are under-cited relative to the number of

such papers in the field[44–48]. Here we sought to proactively consider choosing references

that reflect the diversity of the field in thought, form of contribution, gender, and other

factors. Gender of the first and last author of each reference was predicted by using

databases that store the probability of a name being carried by a man or a woman[46]. By

this measure (and excluding self-citations to the first and last authors of our current

paper), our references contain 12.8% woman(first)/woman(last), 19.1% man/woman,

25.5% woman/man, and 42.6% man/man. We look forward to future work that could help

us to better understand how to support equitable practices in science.

Acknowledgments

This work was supported by the Universidade Federal do ABC (UFABC) and

Coordination of Improvement of Higher Education Personnel (CAPES). Data were

provided by the Human Connectome Project, WU-Minn Consortium (Principal

Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16

NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research;

and by the McDonnell Center for Systems Neuroscience at Washington University.

Author contributions

F.L.R., F.R.C.S and C.E.B. conceptualized the study. F.L.R. and F.R.C.S. designed and

performed the analyses with support from W.H.L.P. F.L.R. wrote the original draft. All

authors revised and edited the manuscript. J.R.S., W.H.L.P., and C.E.B provided support

and guidance with data interpretation.

Competing Interests

The authors declare no competing interests.

References

1. Seghier, M. L. & Price, C. J. Interpreting and Utilising Intersubject Variability in Brain Function. *Trends Cogn. Sci.* **22**, 517–530 (2018).

2. Dubois, J. & Adolphs, R. Building a science of individual differences from fMRI. *Trends Cogn. Sci.* **20**, 425–443 (2016).

3. Miranda-Dominguez, O. *et al.* Connectotyping: Model based fingerprinting of the functional connectome. *PLoS One* **9**, (2014).

4. Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nat. Neurosci.* **18**, 1664–1671 (2015).

5. Mueller, S. *et al.* Individual Variability in Functional Connectivity Architecture of the Human Brain. *Neuron* **77**, 586–595 (2013).

6. Miranda-Dominguez, O. *et al.* Heritability of the human connectome: a connectotyping study. *Netw. Neurosci.* **2**, 175–199 (2018).

7. Biazoli, C. E. *et al.* Commentary: Functional connectome fingerprint: identifying individuals using patterns of brain connectivity. *Front. Hum. Neurosci.* **11**, 47 (2017).

8. Gratton, C. *et al.* Functional Brain Networks Are Dominated by Stable Group and Individual Factors, Not Cognitive or Daily Variation. *Neuron* 439–452 (2018). doi:10.1016/j.neuron.2018.03.035

9.   Jalbrzikowski, M. *et al.* Functional connectome fingerprinting accuracy in youths

and adults is similar when examined on the same day and 1.5-years apart. *Hum.*

Brain Mapp. 1–13 (2020). doi:10.1002/hbm.25118

10. Kaufmann, T. *et al.* Delayed stabilization and individualization in connectome

development are related to psychiatric disorders. *Nat. Neurosci.* **20**, 513–515

(2017).

11. Laumann, T. O. *et al.* Functional System and Areal Organization of a Highly

Sampled Individual Human Brain. *Neuron* **87**, 658–671 (2015).

12. Poldrack, R. A. *et al.* Long-term neural and physiological phenotyping of a single

human. *Nat. Commun.* **6**, (2015).

13. Colclough, G. L. *et al.* The heritability of multi-modal connectivity in human brain

activity. *Elife* **6**, 1–19 (2017).

14. Ge, T., Holmes, A. J., Buckner, R. L., Smoller, J. W. & Sabuncu, M. R. Heritability

analysis with repeat measurements and its application to resting-state functional

connectivity. *Proc. Natl. Acad. Sci.* **114**, 5521–5526 (2017).

15. Yang, Z. *et al.* Genetic and Environmental Contributions to Functional

Connectivity Architecture of the Human Brain. *Cereb. Cortex* **26**, 2341–2352

(2016).

16. Elliott, M. L. *et al.* General functional connectivity: Shared features of resting-state

and task fMRI drive reliable and heritable individual differences in functional brain

networks. *Neuroimage* **189**, 516–532 (2019).

17. Demeter, D. V. *et al.* Functional Connectivity Fingerprints at Rest Are Similar

across Youths and Adults and Vary with Genetic Similarity. *iScience* **23**, 100801

(2020).

18.   Adhikari, B. M. *et al.* Comparison of heritability estimates on resting state fMRI

connectivity phenotypes using the ENIGMA analysis pipeline. *Hum. Brain Mapp.*

1–10 (2018). doi:10.1002/hbm.24331

19.   Teeuw, J. *et al.* Genetic and environmental influences on functional connectivity

within and between canonical cortical resting-state networks throughout

adolescent development in boys and girls. *Neuroimage* **202**, (2019).

20.   Arslan, S. *et al.* Human brain mapping: A systematic comparison of parcellation

methods for the human cerebral cortex. *Neuroimage* **170**, 5–30 (2018).

21.   Salehi, M. *et al.* There is no single functional atlas even for a single individual:

Functional parcel definitions change with task. *Neuroimage* **208**, 116366 (2020).

22.   Eickhoff, S. B., Yeo, B. T. T. & Genon, S. Imaging-based parcellations of the

human brain. *Nat. Rev. Neurosci.* **19**, 672–686 (2018).

23.   Noble, S. *et al.* Influences on the Test-Retest Reliability of Functional Connectivity

MRI and its Relationship with Behavioral Utility. *Cereb. Cortex* **27**, 5415–5429

(2017).

24.   Noble, S., Scheinost, D. & Constable, R. T. A decade of test-retest reliability of

functional connectivity: A systematic review and meta-analysis. *Neuroimage* **203**,

116157 (2019).

25.   Shen, X., Tokoglu, F., Papademetris, X. & Constable, R. T. Groupwise whole-

brain parcellation from resting-state fMRI data for network node identification.

Neuroimage **82**, 403–415 (2013).

26.   Gordon, E. M. *et al.* Generation and Evaluation of a Cortical Area Parcellation

595  from Resting-State Correlations. *Cereb. Cortex* **26**, 288–303 (2014).

596 27. Waller, L. *et al.* Evaluating the replicability, specificity, and generalizability of

597  connectome fingerprints. *Neuroimage* **158**, 371–377 (2017).

598 28. Gandal, M. J. *et al.* Shared molecular neuropathology across major psychiatric

599  disorders parallels polygenic overlap. *Science (80-.).* **359**, 693–697 (2018).

600 29. Prata, D. P., Costa-Neves, B., Cosme, G. & Vassos, E. Unravelling the genetic

601  basis of schizophrenia and bipolar disorder with GWAS: A systematic review. *J.*

602  *Psychiatr. Res.* **114**, 178–207 (2019).

603 30. Kasten, M. *et al.* Genotype-Phenotype Relations for the Parkinson's Disease Genes

604  Parkin, PINK1, DJ1: MDSGene Systematic Review. *Mov. Disord.* **33**, 730–741

605  (2018).

606 31. Sims, R., Hill, M. & Williams, J. The multiplex model of the genetics of

607  Alzheimer's disease. *Nat. Neurosci.* **23**, 311–322 (2020).

608 32. Chenji, S. *et al.* Investigating default mode and sensorimotor network connectivity

609  in amyotrophic lateral sclerosis. *PLoS One* **11**, 1–14 (2016).

610 33. Gratton, C. *et al.* Emergent Functional Network Effects in Parkinson Disease.

611  *Cereb. Cortex* **29**, 2509–2523 (2019).

612 34. Hall, J. M. *et al.* Changes in structural network topology correlate with severity of

613  hallucinatory behavior in Parkinson's disease. *Netw. Neurosci.* **3**, 521–538 (2019).

614 35. Lee, K. *et al.* Disruption, emergence and lateralization of brain network hubs in

615  mesial temporal lobe epilepsy. *NeuroImage Clin.* **20**, 71–84 (2018).

616 36. Buckner, R. L. *et al.* Cortical hubs revealed by intrinsic functional connectivity:

Mapping, assessment of stability, and relation to Alzheimer's disease. *J. Neurosci.*

29, 1860–1873 (2009).

37.   van den Heuvel, M. P. & Sporns, O. A cross-disorder connectome landscape of

brain dysconnectivity. *Nat. Rev. Neurosci.* **20**, 435–446 (2019).

38.   Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: An overview.

Neuroimage **80**, 62–79 (2013).

39.   Smith, S. M. *et al.* Resting-state fMRI in the Human Connectome Project.

Neuroimage **80**, 144–168 (2013).

40.   Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human

Connectome Project. *Neuroimage* **80**, 105–124 (2013).

41.   Whitfield-Gabrieli, S. & Nieto-Castanon, A. *Conn* : A Functional Connectivity

Toolbox for Correlated and Anticorrelated Brain Networks. *Brain Connect.* **2**,

125–141 (2012).

42.   Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction

method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage* **37**, 90–101

(2007).

43.   Bates, T. C., Maes, H. & Neale, M. C. Umx: Twin and path-based structural

equation modeling in R. *Twin Res. Hum. Genet.* **22**, 27–41 (2019).

44.   Dion, M. L., Sumner, J. L. & Mitchell, S. M. Gendered Citation Patterns across

Political Science and Social Science Methodology Fields. *Polit. Anal.* **26**, 312–327

(2018).

45.   Mitchell, S. M., Lange, S. & Brus, H. Gendered Citation Patterns in International

Relations Journals. *Int. Stud. Perspect.* **14**, 485–492 (2013).

46.   Dworkin, J. D. *et al.* The extent and drivers of gender imbalance in neuroscience

641   reference lists. *Nat. Neurosci.* **23**, 918–926 (2020).

642   47.   Caplar, N., Tacchella, S. & Birrer, S. Quantitative evaluation of gender bias in

643   astronomical publications from citation counts. *Nat. Astron.* **1**, (2017).

644   48.   Maliniak, D., Powers, R. & Walter, B. F. *The gender citation gap in international*

645   *relations*. *International Organization* **67**, (2013).