Histone acetyltransferase inhibition rescues differentiation of emerin-deficient

myogenic progenitors

Katherine A. Bossone, B.S.[1,2], Joseph A. Ellis, B.S.[2] and James M. Holaska, Ph.D.[1,2]*

[1]Department of Biomedical Sciences, Cooper Medical School of Rowan University

[2]Department of Pharmaceutical Sciences, University of the Sciences

*Correspondence: holaska@rowan.edu; Tel: 856-956-2746

Author contributions: Conceptualization, methodology, and validation: J.M.H.; Formal

analysis: J.M.H. and K.A.B.; Investigation: K.A.B. and J.A.E.; Resources: J.M.H.; Data

curation: J.M.H., K.A.B., and J.A.E.; Writing- original draft: J.M.H.; Writing- reviewing and

editing: J.M.H., K.A.B., and J.A.E.; Visualization: K.A.B. and J.M.H.; Supervision: J.M.H.;

Project administration: J.M.H., K.A.B., and J.A.E. Funding acquisition: J.M.H.

Acknowledgements: We thank the members of the Holaska laboratory for the many

helpful discussions regarding these studies and preparation of this manuscript. This

study was supported by the National Institute of Arthritis and Musculoskeletal and Skin

Diseases of the National Institutes of Health under Award Number R15AR069935 (to

J.M.H.). The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health.

Number of words in abstract: 147

Number of words in manuscript (excluding abstract, references, table titles and

figure legends): 3,073

Corresponding Author: James M. Holaska, Department of Biomedical Sciences,

Cooper Medical School of Rowan University, MEB 534, 401 South Broadway, Camden,

NJ 08103. Email: holaska@rowan.edu

Running title: H4K5ac dynamics in differentiation

Ethical Publication Statement: We confirm that we have read the Journal's position on

issues involved in ethical publication and affirm that this report is consistent with those

guidelines.

Conflicts of interests: The authors declare no competing or financial interests

Histone acetyltransferase inhibition rescues differentiation of emerin-deficient myogenic progenitors

ABSTRACT

Introduction: Emery-Dreifuss Muscular Dystrophy (EDMD) is a disease characterized by skeletal muscle wasting, major tendon contractures, and cardiac conduction defects. Mutations in the gene encoding emerin cause EDMD1. Our previous studies suggested emerin activation of Histone Deacetylase 3 (HDAC3) to reduce Histone 4-Lysine 5 (H4K5) acetylation (ac) is important for myogenic differentiation. **Methods:** Pharmacological inhibitors (Nu9056, L002) of histone acetyltransferases targeting acetylated H4K5 were used to test if increased acetylated H4K5 was responsible for the impaired differentiation seen in emerin deficient myogenic progenitors. **Results:** Nu9056 and L002 rescued impaired differentiation in emerin deficiency. SRT1720, which inhibits the NAD^+-dependent deacetylase Sirtuin 1 (SIRT1), failed to rescue myotube formation. **Discussion:** We conclude emerin regulation of HDAC3 activity to affect H4K5 acetylation dynamics is important for myogenic differentiation. Targeting H4K5ac dynamics represents a new strategy for ameliorating the skeletal muscle wasting seen in EDMD1.

Keywords: Cell signaling, Emerin, Emery-Dreifuss Muscular Dystrophy, Myogenic differentiation

INTRODUCTION

The nuclear envelope is composed of two lipid bilayers, the outer nuclear membrane,

which is contiguous with the endoplasmic reticulum, and the inner nuclear membrane[1].

Although the outer and inner nuclear membranes arise from a common membrane, they

are functionally distinct membranes. Underlying the inner nuclear membrane is a

network of Type V intermediate filament proteins named lamins that provide nuclear

rigidity and elasticity[2]. The inner nuclear membrane contains a large number of unique

integral inner nuclear membrane proteins[3], many of which show cell-type-specific

expression[4-11]. Inner nuclear membrane proteins function in diverse roles, including

nuclear structure, genomic organization, chromatin architecture, gene expression, cell

cycle regulation, and cytoskeletal organization[12,1]. The nuclear lamins and its associated

inner nuclear membrane proteins define the nuclear lamina.

Emerin is a lamin-binding, integral inner nuclear membrane protein. Mutations in the

gene encoding emerin cause X-linked Emery-Dreifuss muscular dystrophy (EDMD1), an

inherited disorder causing progressive skeletal muscle wasting, irregular heart rhythms,

and contractures of major tendons[13-16]. Evidence suggests the skeletal muscle wasting

seen in EDMD is not caused by increased damage to the myofiber, but by impaired

differentiation of skeletal muscle stem cells. Supporting this hypothesis, skeletal muscle

necrosis and increased skeletal muscle fiber permeability are rarely seen in EDMD

patients[17]. Further, emerin knockout mice (also commonly referred to as emerin-null or

emerin-deficient mice) exhibit delayed skeletal muscle regeneration and repair, motor

coordination defects, and mild atrioventricular conduction defects[18,19]. Skeletal muscle

from EDMD1 and EDMD2 patients and emerin-deficient mice both showed altered

expression of muscle regeneration pathway components[20,18]. Emerin-deficient myogenic

progenitors and emerin-downregulated C2C12 myoblasts exhibit impaired differentiation

and myotube formation[21-23] due to aberrant temporal activation of myogenic

differentiation genes[24] and disruption of key signaling pathways[25], suggesting defective

muscle regeneration contributes to the EDMD skeletal muscle phenotype[22,21,18].

The coordinated temporal expression of *MyoD, Myf5, Pax3* and *Pax7*, which are

important for proper differentiation, was disrupted in emerin-deficient myogenic

progenitors[26] due to the inability of the genome to properly reorganize during

differentiation[20,18,25]. This supports the hypothesis that emerin-deficient myogenic

progenitors fail to undergo the transcriptional reprogramming required for myogenic

differentiation. Furthermore, emerin was shown to bind directly to Histone Deacetylase 3

(HDAC3) and activate its deacetylase activity[27]. HDAC3 activity is required for proper

dynamic reorganization of *MyoD, Myf5, Pax3* and *Pax7*[26]. Thus, regulation of HDAC3

activity by emerin is critical for transcriptional reprogramming during myogenic

differentiation.

We used histone acetyltransferase (HAT) inhibitors targeting HATs mediating H4K5

acetylation (e.g., Tip60/KAT5) to further test the hypothesis that acetylation dynamics on

lysine 5 of Histone 4 (H4K5) were important for in myogenic differentiation. Here we

show increased H4K5 acetylation (H4K5ac) contributes to the impaired differentiation of

emerin-deficient myogenic progenitors. Targeting H4K5ac dynamics represents a

potential new strategy for ameliorating the skeletal muscle wasting seen in EDMD1.

METHODS

Pharmacological treatments

We previously showed emerin-deficient myogenic progenitors had impaired

differentiation and was rescued by activation of HDAC3[23]. To independently test if

altered H4K5 acetylation dynamics was responsible for the impaired differentiation of

emerin-deficient progenitors we chose to inhibit HATs. HAT inhibitors (HATi) selected for

these studies were chosen because they preferentially inhibit acetylation of lysine

residues targeted by HDAC3 (e.g., H4K5)[34]. Cell cycle withdrawal, myosin heavy chain

(MyHC) expression and myotube formation were analyzed 36 hours post-differentiation

induction, as previously described[23]. HAT inhibitor Nu9056 was selected because it is a

highly specific inhibitor of histone acetyltransferase Tip60/KAT5[28]. Tip60/KAT5 mediates

the acetylation of H4K5, H4K8, H4K12 and H4K16ac (Table 1). A second HAT inhibitor,

L002, was used to test whether inhibition of H4K5 acetylation rescued myogenic

differentiation of emerin-deficient progenitors. L002 inhibits H4 acetylation in cells at low

micromolar concentrations (Table 1)[29]. A Sirtuin 1 (SIRT1) activator (SRT1720) was

used to confirm the rescue of emerin-deficient progenitor differentiation was due to

changes in acetylation states of HDAC3 target residues (e.g., H4K5ac). Unlike HDAC3,

SIRT1 is an NAD+-dependent protein deacetylase[35]. SIRT1 deacetylates H3K9ac, but

does not affect H4K5, H4K8 or H4K12 acetylation[36]. 1.5 µM SRT1720 was added to

wildtype or emerin-deficient myogenic progenitors upon differentiation induction and

differentiation was analyzed after 36 hours (Figure 1A).

Concentrations of compounds used in this study were based upon the reported half-

maximal inhibitory concentration (IC50) for each compound[28-31] and per manufacturer

(EMD Millipore) instructions. 0.25 µM to 30 µM or each compound was added to

wildtype and emerin-deficient myogenic progenitors and tested for viability and impaired

proliferation. The concentrations used in this study failed to inhibit cell proliferation and

had no effect on cell viability. A 1.0 mM stock solution of L002 in DMSO was added to a

final concentration of 0.5 µM in differentiation medium. A 1.0 mM stock solution of

Nu9056 in DMSO was added to a final concentration of 0.5 µM in differentiation medium.

A 3.0 mM stock solution of SRT1720 in DMSO was added to a final concentration of 1.5

µM in differentiation medium. Differentiation media containing each inhibitor or DMSO

was added to induce differentiation of wildtype or emerin-deficient myogenic progenitors.

Cell culture

Wildtype and Emerin-null H2K mouse myogenic progenitors were a generous gift from

Tatiana Cohen and Terence Partridge (Children's National Medical Center, Washington,

DC). Emerin-null H2K mice were generated by Tatiana Cohen and Terence Partridge by

breeding emerin-null (C57Bl/6) and H-2K[b]tsA58 mice to create emerin-deficient mice in

the H-2K[b]tsA58 background[32,33]. Myogenic progenitors were isolated and maintained as

previously described[24,33,25]. Briefly, extensor digitorum longus (EDL) muscles were

isolated and placed into 2 mg/ml collagenase (Sigma, product #C0130) in DMEM

(Invitrogen, product #11995-065) for 1–2 hours at 35°C. Individual fibers were then

isolated and each fiber was transferred serially through 2–4 petri dishes containing

DMEM to select for undamaged fibers. Fibers were placed into matrigel-coated petri

dishes containing DMEM, 10% horse serum (Invitrogen product #16050-098), 0.5%

chick embryo extract (Accurate Chemical, product #CE6507), 2% L-Glutamine

(Invitrogen, product #25030-081) and 1% penicillin/streptomycin (Invitrogen, product

#15140-122) for 3–4 hours at 37°C. Myogenic progenitors were isolated from individual

fibers by transferring each fiber into one matrigel-coated well of a 24-well plate

containing proliferation media consisting of DMEM, 20% Heat inactivated FBS

(Invitrogen, product #10082-147), 2% chick embryo extract, 2% L-glutamine, 1% pen-

strep and 20 ng/ml γ-Interferon (Millipore, product #IF005). The fibers were incubated for

24–48 hours at 33°C, 10% CO_2. Upon attachment of a single myogenic progenitor to the

well, the fiber was removed and the myogenic progenitor was incubated in proliferation

media for another 48 hours at 33°C, 10% CO_2. Approximately 200 cells are expected

after 48 hours and these were split and proliferated until enough cells were obtained for

our analyses. H2K myogenic progenitors were maintained in proliferation media at 33°C

and 10% CO_2. Cells between passages 4–10 were used for these studies.

Cell culture of proliferation and differentiation of H2Ks were done as previously

described[23]. Briefly, for proliferation, wildtype and emerin-deficient H2K myogenic

progenitors were seeded onto tissue culture plates (Falcon cat no. 353046 and

3530003) and maintained at 33℃ and 10% CO_2 in proliferation medium (high glucose

DMEM supplemented with 20% heat-inactivated fetal bovine serum, 2% L-glutamine, 2%

chick embryo extract, 1% penicillin/streptomycin, sodium pyruvate, 20 units/ml γ-

interferon, ThermoFisher Scientific). The plates were coated with 0.01% gelatin (Sigma-

Aldrich) prior to seeding.

Wildtype and emerin-deficient H2K myogenic progenitors were seeded onto 12 well

tissue culture plates coated with 0.01% gelatin (Sigma-Aldrich) for differentiation

induction. Cells were seeded at 23,500 cells/cm^2 in proliferation media for 24h at 33℃

and 10% CO_2. Differentiation was stimulated by replacing the proliferation medium with

differentiation medium (high glucose DMEM with sodium pyruvate, 5% horse serum, 2%

L-glutamine, ThermoFisher Scientific). The cells were maintained at 37℃ and 5% CO_2

throughout differentiation.

EdU assays and immunofluorescence microscopy

Cells were treated with 10 µM EdU (ThermoFisher Scientific) in DMSO 2h prior to fixing,

while incubating at 37℃ and 5% CO_2. Cells were then fixed with 3.7% formaldehyde for

15 min and washed three times with PBS. Fixed cells were then stored at 4°C with 0.1%

sodium azide in PBS. The cells were permeabilized with 0.5% triton X-100 in PBS for 20

minutes, washed twice with 3% BSA in PBS for five minutes per wash and treated with

the Click-IT EdU reaction cocktail for 25 minutes. Cells were washed with PBS and

blocked for 1 h at room temperature with 3% BSA with 0.1% Triton X-100. Myosin heavy

chain (MyHC) antibodies (1:20, Santa Cruz Biotechnologies, H-300 for L002 and Nu506

experiments; 1:50, Santa Cruz Biotechnologies, B-5 for SRT1720 treatments) were

added and the cells were incubated at room temperature for 1 h. The cells were washed

with PBS three times and treated with Alexa Fluor 594 secondary antibodies (1:200,

C10637; A11032, ThermoFisher Scientific) at room temperature for 1 hour, washed with

PBS and incubated with DAPI for 5 minutes.

Images were taken using the EVOS-FL imaging system (ThermoFisher LifeSciences) for

experiments with L002. The remainder of the images were taken with the EVOS-FL Auto

(ThermoFisher LifeSciences). All images were obtained using a long working distance

40x objective. At least three replicates, with each replicate containing three culture wells

per group, were done for each drug treatment. Images from five different sections from

each well were taken, with each section containing approximately 50-200 cells. The total

number of cells analyzed for each experiment ranged between 500-1500.

The cell counter plugin on ImageJ was used to count proliferating cells. The percent of

cells still in the cycle was determined by dividing the number of EdU positive nuclei by

the total number of nuclei. The DAPI and MyHC images were superimposed to calculate

the percentage of cells expressing MyHC. Myotube formation was determined by

superimposing the phase contrast image, which allowed for monitoring nuclei within a

shared cytoplasm, with DAPI and MyHC images. Myotube formation, or the

differentiation index, was determined by counting the number of nuclei in MyHC positive

cells that contained three or more nuclei divided by the total number of nuclei in the field.

Western Blotting

Differentiating H2K cells were resuspended directly in sample buffer and 50,000 to

100,000 cell equivalents were separated by SDS-PAGE and transferred to a

nitrocellulose membrane. The membranes were blocked either at room temperature for

2 h or overnight at 4°C in 3% BSA in PBST (PBS with 0.1% Tween 20). Antibodies

against H4 (1:50,000; Millipore, 05-858), H4K5ac (1:1,000; Millipore, 07-327), H3K9ac

(1:10,000; Abcam, ab4441), H3K18ac (1:1,000; Abcam, ab1191), H3K27ac (1:1,000;

Abcam, ab4729), H4K16ac (1:2,000; Abcam, ab109463), and MyHC (1:1,000; Santa

Cruz, B-5) were then incubated either at room temperature for 2 h or overnight at 4°C.

The membranes were washed three times in PBS and incubated with Goat Anti-Rabbit

HRP or Goat Anti-Mouse HRP secondary antibody (1:10,000; ThermoFisher Scientific)

in PBST either at room temperature for 2 h or overnight at 4°C. The membranes were

treated with ECL chemiluminescence detection reagent (GE healthcare, product #

RPN2106V1 and RPN2106V20 and imaged using the Bio-Rad Chemidoc system (Bio-

Rad Laboratories). Densitometry was done using ImageLab software (Bio-Rad

Laboratories) as per the manufacturer's instructions.

RESULTS

Wildtype and emerin-deficient myogenic progenitors were differentiated for 36 hours in

the presence of the HAT inhibitor Nu9056 to independently confirm HAT inhibition

rescued emerin-deficient myogenic differentiation. 0.5 µM Nu9056 in DMSO or DMSO

alone were incubated with wildtype or emerin-deficient myogenic progenitors upon

differentiation induction (Figure 1A). Nu9056 treatment had no effect on cell cycle

withdrawal of wildtype or emerin-deficient myogenic progenitors, (Figure 1C, G, J).

Nu9056 treatment failed to rescue myoblast commitment, as the number of MyHC-

expressing cells was similar in Nu9056-treated (51.0%) and untreated emerin-deficient

myogenic progenitors (50.3%; Figure 1H, K). Myotube formation in emerin-deficient

progenitors was rescued by Nu9056 treatment, as 15.1% of Nu9056-treated emerin-

deficient progenitors fused to form myotubes, compared to 10.8% of DMSO-treated

emerin-deficient progenitors (Figure 1I, L). Myotube formation in Nu9056-treated emerin-

deficient progenitors was statistically similar to wildtype progenitors (p=0.11).

A second HAT inhibitor, L002, was used to test whether inhibition of H4K5 acetylation

rescued myogenic differentiation of emerin-deficient progenitors. Wildtype and emerin-

deficient myogenic progenitors were treated with 0.5 µM L002 upon differentiation

induction (Figure 1A). L002-treated wildtype progenitors exited the cell cycle normally

(Figure 2B', I). 2.7% of emerin-deficient progenitors failed to exit the cell cycle after 36

hours, as expected (Figure 2F, I). Emerin-deficient progenitors treated with L002 showed

a trend toward reducing the number of emerin-deficient cells in the cell cycle (2.1%;

p=0.06; Figure 2F', I). L002 treatment significantly increased the percentage of

differentiating emerin-deficient progenitors expressing MyHC (46%, Figure 2G, J;

p=0.015). The number of MyHC-positive cells in L002-treated differentiating emerin-

deficient progenitors is statistically similar to untreated wildtype progenitors (47.8% in

wildtype, p=0.35; Figure 2C, G', J), indicating rescue of myoblast commitment. L002

treatment increased myotube formation 1.8-fold in differentiating emerin-deficient

progenitors (Figure 2H, K) completely rescuing myotube formation to wildtype levels

(p=0.97 for L002-treated emerin-deficient cells vs. wildtype cells; Figure 2D, H', K).

A Sirtuin 1 (SIRT1) activator (SRT1720) was used to confirm the rescue of emerin-

deficient progenitor differentiation was due to changes in acetylation states of HDAC3

target residues (e.g., H4K5ac). Activation of SIRT1 by treatment with 1.5 µM SRT1720

failed to rescue cell cycle withdrawal of differentiating emerin-deficient progenitors, as

7.0% of DMSO-treated and 5.4% of SRT1720-treated cells were cycling (Figure 3F, I;

p=0.09). 41.1% of SRT1720-treated differentiating emerin-deficient progenitors

expressed MyHC compared to 42.7% of DMSO-treated emerin-deficient progenitors

(Figure 3G, J; p=0.49). SRT1720 treatment also failed to rescue myotube formation in

emerin-deficient progenitors (Figure 3H, K; p=0.44).

Treatment of emerin-deficient progenitors with L002 during differentiation reduced

H4K5ac 3.8-fold (Figure 4A, B), comparable to H4K5ac levels in wildtype progenitors.

Nu9056 treatment decreased H4K5ac 3.2-fold in emerin-deficient myogenic progenitors

(Figure 4A, B). Decreased H4K5ac seen in emerin-deficient myogenic progenitors

treated with Nu9056 is similar to the H4K5ac levels seen in differentiating wildtype

progenitors. Western blotting confirmed the levels of H4K5ac were unchanged by

treatment with SRT1720 (Figure 4C, D).

Acetylation of H3K9, H3K18, H3K27 and H4K16 were monitored during impaired

differentiation in emerin deficiency in the presence of Nu9056, L002 and SRT1720.

H3K9ac, H3K18ac and H3K27ac were all increased in emerin-deficient myogenic

progenitors and during emerin-deficient myogenic differentiation (Figure 5), including

during the transition to myoblast commitment. Treatment of emerin-deficient progenitors

with Nu9056 had no significant effect on acetylation of H3K9, H3K18, H3K27 or H4K16

(Figure 5). L002 treatment had a small effect on H3K18 and H3K27 acetylation.

Treatment with the SIRT 1 activator, SRT1720, reduced H3K9ac activity, as expected,

since SIRT1 deacetylates H3K9[35,36].

DISCUSSION

The studies presented here used a cell-based system to follow differentiation, in which

myotubes are formed by myoblast-to-myoblast fusion or myoblast-to-myotube fusion.

Upon stimulation of myogenic progenitors to differentiate, transcriptional reprogramming

is initiated, leading to cell cycle exit. This reprogramming activates the myogenic

differentiation program and represses the proliferative program, thereby leading to

myoblast commitment followed by fusion to form myotubes[37]. Transcriptional

reprogramming is compromised in emerin-deficient progenitors[38]. The failure of emerin-

deficient progenitors to coordinate the temporal reorganization of their genome during

differentiation is predicted to cause this defective transcriptional reprogramming.

Emerin binds directly to HDAC3, the catalytic component of the Nuclear Co-Repressor

(NCoR) complex[27,39] and activates its activity. Emerin-binding recruits HDAC3 to the

nuclear envelope. The functional interaction between emerin and HDAC3 coordinates

the spatiotemporal nuclear envelope localization of genomic regions containing

important transcription factors that control the temporal expression of differentiation

genes[27,26]. Loss of emerin disrupts this genomic reorganization resulting in impaired

myogenic differentiation. Activation of HDAC3 rescues emerin-deficient myotube

formation[26,23]. Nuclear envelope localization of HDAC3 is also important in

cardiomyocyte differentiation[40]. In the absence of nuclear envelope-localized HDAC3,

repressed genomic loci were aberrantly localized to the nuclear interior resulting in

precocious differentiation. Thus, controlling HDAC3 nuclear envelope localization and

activation is an important regulatory mechanism used for differentiation.

The results presented here support the role of emerin in controlling histone acetylation

dynamics by regulating HDAC3 activity. Using HATi specifically targeting acetylation of

residues deacetylated by HDAC3 (e.g., H4K5ac), we found HAT inhibition rescued

impaired differentiation in emerin deficiency. This recapitulated the rescue seen by

treatment of emerin-deficient progenitors with an HDAC3 activator. Thus, H4K5

acetylation dynamics are predicted to be important for ensuring proper transcriptional

reprogramming upon differentiation induction (Figure 6). Similar to HDAC3 activation,

HDAC inhibition primarily affected later differentiation transitions[23], suggesting emerin

regulation of HDAC3 activity controls the temporal expression of these later genes. This

may result from failure to completely reprogram the transcriptome upon differentiation

induction or by specifically regulating the latter steps of the gene expression program.

Nu9056 exhibits high specificity, as the half-maximal inhibitory concentration (IC50) of

Nu9056 for Tip60/KAT5 is 20-40-fold lower than for the histone acetyltransferase p300

or the histone acetyltransferase pCAF/GCN5 (Table 1). L002 was identified as a p300-

specific inhibitor[29]. p300 acetylates H3K18 and H3K27; it has also been reported to

acetylate H4K5, H4K8, H4K12 and H4K16[41,42]. L002 was previously reported to

decrease H4 acetylation[29], but this study was the first to show L002 inhibits acetylation

of H4K5. Although H3K9ac, H3K18ac and H3K27ac were increased in emerin-deficient

myogenic progenitors, treatment with Nu9056 had no significant effect on acetylation of

H3K9, H3K18 or H3K27; L002 treatment had a small effect on H3K18 and H3K27

acetylation; SRT1720 only affected H3K9 acetylation (Figure 5). Collectively, the distinct

specificities of these compounds at the concentrations used in this study demonstrates

that changes in acetylation of H3K9, H3K18, H3K27 and H4K16 cannot be responsible

for the rescued myotube formation. Rather, these results suggest rescue by L002 and

Nu9056 occurs primarily through rescuing H4K5 acetylation dynamics. This is consistent

with our previous studies using HDAC3 inhibitors and activators[23,26]. It is possible L002

may act to rescue impaired myogenic commitment by p300-mediated acetylation of

H3K18 or H3K27 (Figure 6). Alternatively, L002 may act through an unknown

mechanism or act on other HATs to rescue myoblast commitment, as L002 appears to

be more promiscuous at lower concentrations. Targeting these dynamic epigenetic

changes represents a new potential strategy for ameliorating the skeletal muscle wasting

seen in EDMD1.

It is important to elucidate how emerin regulates the dynamic epigenetic changes

occurring during myogenic differentiation to control the transcriptional programs needed

for passage through specific transition points. HAT inhibition and HDAC3 activation

successfully rescued the latter steps of emerin-deficient myogenic differentiation (this

study)[23,26], suggesting emerin regulation of H4K5ac dynamics during transcriptional

reprogramming likely impairs the expression of genes acting at later stages of

differentiation (e.g., myotube formation). Consistent with these results, HDAC3 inhibition

by RGFP966 blocked MyHC expression and myotube fusion in both differentiating

wildtype and emerin-deficient myogenic progenitors[23]. We propose H4K5 acetylation

levels are tightly regulated and that increases or decreases in H4K5 acetylation levels

impairs the transition from committed, differentiating myoblasts, to myotubes by altering

transcription reprogramming upon differentiation induction. Collectively, our results

support pharmacological targeting of H4K5 acetylation as a potential therapeutic

strategy for rescuing muscle regeneration in EDMD.

Abbreviations:

BSA: Bovine Serum Albumin

DAPI: 4',6-diamidino-2-phenylindole

DMEM: Dulbecco's Modified Eagle's medium

DMSO: Dimethyl Sulfoxide

ECL: Enhanced Chemiluminescence

EDL: Extensor Digitorum Longus

EDMD: Emery-Dreifuss Muscular Dystrophy

EdU: 5-Ethynyl-2'-deoxyuridine

FBS: Fetal Bovine Serum

H3K9: Histone 3 lysine 9

H3K9ac: Histone 3 acetylated on lysine 9

H3K18: Histone 3 lysine 18

H3K18ac: Histone 3 acetylated on lysine 18

H3K27: Histone 3 lysine 27

H3K27ac: Histone 3 acetylated on lysine 27

H4: Histone 4

H4K5: Histone 4 lysine 5

H4K5ac: Histone 4 acetylated on lysine 5

H4K16: Histone 4 lysine 16

H4K16ac: Histone 4 acetylated on lysine 16

HAT: Histone Acetyltransferase

HATi: Histone Acetyltransferase inhibitor

HDAC: Histone Deacetylase

HDACi: Histone Deacetylase Inhibitor

HRP: Horseradish Peroxidase

IC50: Half-maximal Inhibitory Concentration

NCoR: Nuclear Co-Repressor

MyHC: Myosin Heavy Chain

PBS: Phosphate Buffered Saline

PBST: Phosphate Buffered Saline with Tween

SIRT1: Sirtuin 1

References

1.   Holaska JM. Diseases of the Nucleoskeleton. Compr Physiol 2016;6(4):1655-1674.

2.   Tatli M, Medalia O. Insight into the functional organization of nuclear lamins in health and disease. Curr Opin Cell Biol 2018;54:72-79.

3.   Gruenbaum Y, Foisner R. Lamins: nuclear intermediate filament proteins with fundamental functions in nuclear mechanics and genome regulation. Annu Rev Biochem 2015;84:131-164.

4.   Gonzalez Y, Saito A, Sazer S. Fission yeast Lem2 and Man1 perform fundamental functions of the animal cell nuclear lamina. Nucleus 2012;3(1):60-76.

5.   Korfali N, Wilkie GS, Swanson SK, Srsen V, Batrakou DG, Fairley EA, Malik P, Zuleger N, Goncharevich A, de Las Heras J, Kelly DA, Kerr AR, Florens L, Schirmer EC. The leukocyte nuclear envelope proteome varies with cell activation and contains novel transmembrane proteins that affect genome architecture. Mol Cell Proteomics 2010;9(12):2571-2585.

6.   Schirmer EC, Florens L, Guan T, Yates JR, 3rd, Gerace L. Nuclear membrane proteins with potential disease links found by subtractive proteomics. Science 2003;301(5638):1380-1382.

7.   Worman HJ, Schirmer EC. Nuclear membrane diversity: underlying tissue-specific pathologies in disease? Curr Opin Cell Biol 2015;34:101-112.

8.   de Las Heras JI, Meinke P, Batrakou DG, Srsen V, Zuleger N, Kerr AR, Schirmer EC. Tissue specificity in the nuclear envelope supports its functional complexity. Nucleus 2013;4(6):460-477.

9.   Wilkie GS, Korfali N, Swanson SK, Malik P, Srsen V, Batrakou DG, de las Heras J, Zuleger N, Kerr AR, Florens L, Schirmer EC. Several novel nuclear envelope

transmembrane proteins identified in skeletal muscle have cytoskeletal associations. Mol

Cell Proteomics 2011;10(1):M110 003129.

10.   Malik P, Korfali N, Srsen V, Lazou V, Batrakou DG, Zuleger N, Kavanagh DM,

Wilkie GS, Goldberg MW, Schirmer EC. Cell-specific and lamin-dependent targeting of

novel transmembrane proteins in the nuclear envelope. Cell Mol Life Sci

2010;67(8):1353-1369.

11.   Korfali N, Wilkie GS, Swanson SK, Srsen V, de Las Heras J, Batrakou DG, Malik

P, Zuleger N, Kerr AR, Florens L, Schirmer EC. The nuclear envelope proteome differs

notably between tissues. Nucleus 2012;3(6):552-564.

12.   Barton LJ, Soshnev AA, Geyer PK. Networking in the nucleus: a spotlight on

LEM-domain proteins. Curr Opin Cell Biol 2015;34:1-8.

13.   Bione S, Maestrini E, Rivella S, Mancini M, Regis S, Romeo G, Toniolo D.

Identification of a novel X-linked gene responsible for Emery-Dreifuss muscular

dystrophy. Nat Genet 1994;8:323-327.

14.   Mendez-Lopez I, Worman HJ. Inner nuclear membrane proteins: impact on

human disease. Chromosoma 2012;121(2):153-167.

15.   Vlcek S, Foisner R. Lamins and lamin-associated proteins in aging and disease.

Curr Opin Cell Biol 2007;19(3):298-304.

16.   Worman HJ. Nuclear lamins and laminopathies. J Pathol 2012;226(2):316-325.

17.   Bonne G, Leturcq F, Ben Yaou R. Emery-Dreifuss Muscular Dystrophy. In:

Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, Bird TD,

Ledbetter N, Mefford HC, Smith RJH, Stephens K, editors. GeneReviews(R). Seattle

(WA)1993.

18.   Melcon G, Kozlov S, Cutler DA, Sullivan T, Hernandez L, Zhao P, Mitchell S,

Nader G, Bakay M, Rottman JN, Hoffman EP, Stewart CL. Loss of emerin at the nuclear

envelope disrupts the Rb1/E2F and MyoD pathways during muscle regeneration. Hum

Mol Genet 2006;15(4):637-651.

19.   Ozawa R, Hayashi YK, Ogawa M, Kurokawa R, Matsumoto H, Noguchi S,

Nonaka I, Nishino I. Emerin-lacking mice show minimal motor and cardiac dysfunctions

with nuclear-associated vacuoles. Am J Pathol 2006;168(3):907-917.

20.   Bakay M, Wang Z, Melcon G, Schiltz L, Xuan J, Zhao P, Sartorelli V, Seo J,

Pegoraro E, Angelini C, Shneiderman B, Escolar D, Chen YW, Winokur ST, Pachman

LM, Fan C, Mandler R, Nevo Y, Gordon E, Zhu Y, Dong Y, Wang Y, Hoffman EP.

Nuclear envelope dystrophies show a transcriptional fingerprint suggesting disruption of

Rb-MyoD pathways in muscle regeneration. Brain 2006;129(Pt 4):996-1013.

21.   Frock RL, Kudlow BA, Evans AM, Jameson SA, Hauschka SD, Kennedy BK.

Lamin A/C and emerin are critical for skeletal muscle satellite cell differentiation. Genes

Dev 2006;20(4):486-500.

22.   Huber MD, Guan T, Gerace L. Overlapping functions of nuclear envelope

proteins NET25 (Lem2) and emerin in regulation of extracellular signal-regulated kinase

signaling in myoblast differentiation. Mol Cell Biol 2009;29(21):5718-5728.

23.   Collins CM, Ellis JA, Holaska JM. MAPK signaling pathways and HDAC3 activity

are disrupted during differentiation of emerin-null myogenic progenitor cells. Dis Model

Mech 2017;10(4):385-397.

24.   Dedeic Z, Cetera M, Cohen TV, Holaska JM. Emerin inhibits Lmo7 binding to the

Pax3 and MyoD promoters and expression of myoblast proliferation genes. J Cell Sci

2011;124(Pt 10):1691-1702.

25.   Koch AJ, Holaska JM. Loss of Emerin Alters Myogenic Signaling and miRNA

Expression in Mouse Myogenic Progenitors. PLoS One 2012;7(5):e37262.

26.   Demmerle J, Koch AJ, Holaska JM. Emerin and histone deacetylase 3 (HDAC3)

cooperatively regulate expression and nuclear positions of MyoD, Myf5, and Pax7 genes

during myogenesis. Chromosome Res 2013;21(8):765-779.

27.   Demmerle J, Koch AJ, Holaska JM. The Nuclear Envelope Protein Emerin Binds

Directly to Histone Deacetylase 3 (HDAC3) and Activates HDAC3 Activity. J Biol Chem

2012;287(26):22080-22088.

28.   Coffey K, Blackburn TJ, Cook S, Golding BT, Griffin RJ, Hardcastle IR, Hewitt L,

Huberman K, McNeill HV, Newell DR, Roche C, Ryan-Munden CA, Watson A, Robson

CN. Characterisation of a Tip60 specific inhibitor, NU9056, in prostate cancer. PLoS

One 2012;7(10):e45539.

29.   Yang H, Pinello CE, Luo J, Li D, Wang Y, Zhao LY, Jahn SC, Saldanha SA,

Chase P, Planck J, Geary KR, Ma H, Law BK, Roush WR, Hodder P, Liao D. Small-

molecule inhibitors of acetyltransferase p300 identified by high-throughput screening are

potent anticancer agents. Mol Cancer Ther 2013;12(5):610-620.

30.   Milne JC, Lambert PD, Schenk S, Carney DP, Smith JJ, Gagne DJ, Jin L, Boss

O, Perni RB, Vu CB, Bemis JE, Xie R, Disch JS, Ng PY, Nunes JJ, Lynch AV, Yang H,

Galonek H, Israelian K, Choy W, Iffland A, Lavu S, Medvedik O, Sinclair DA, Olefsky JM,

Jirousek MR, Elliott PJ, Westphal CH. Small molecule activators of SIRT1 as

therapeutics for the treatment of type 2 diabetes. Nature 2007;450(7170):712-716.

31.   Feige JN, Lagouge M, Canto C, Strehle A, Houten SM, Milne JC, Lambert PD,

Mataki C, Elliott PJ, Auwerx J. Specific SIRT1 activation mimics low energy levels and

protects against diet-induced metabolic disorders by enhancing fat oxidation. Cell Metab

2008;8(5):347-358.

32.   Jat PS, Noble MD, Ataliotis P, Tanaka Y, Yannoutsos N, Larsen L, Kioussis D.

Direct derivation of conditionally immortal cell lines from an H-2Kb-tsA58 transgenic

mouse. Proc Natl Acad Sci U S A 1991;88(12):5096-5100.

33.   Morgan JE, Beauchamp JR, Pagel CN, Peckham M, Ataliotis P, Jat PS, Noble

MD, Farmer K, Partridge TA. Myogenic cell lines derived from transgenic mice carrying a

thermolabile T antigen: a model system for the derivation of tissue-specific and mutation-

specific cell lines. Dev Biol 1994;162(2):486-498.

34.   Anamika K, Krebs AR, Thompson J, Poch O, Devys D, Tora L. Lessons from

genome-wide studies: an integrated definition of the coactivator function of histone

acetyl transferases. Epigenetics Chromatin 2010;3(1):18.

35.   Bao J, Sack MN. Protein deacetylation by sirtuins: delineating a post-translational

regulatory program responsive to nutrient and redox stressors. Cell Mol Life Sci

2010;67(18):3073-3087.

36.   Chalkiadaki A, Guarente L. The multifaceted functions of sirtuins in cancer. Nat

Rev Cancer 2015;15(10):608-624.

37.   Mashinchian O, Pisconti A, Le Moal E, Bentzinger CF. The Muscle Stem Cell

Niche in Health and Disease. Curr Top Dev Biol 2018;126:23-65.

38.   Iyer A, Koch AJ, Holaska JM. Expression Profiling of Differentiating Emerin-Null

Myogenic Progenitor Identifies Molecular Pathways Implicated in Their Impaired

Differentiation. Cells 2017;6(4).

39.   Holaska JM, Wilson KL. An emerin "proteome": purification of distinct emerin-

containing complexes from HeLa cells suggests molecular basis for diverse roles

including gene regulation, mRNA splicing, signaling, mechanosensing, and nuclear

architecture. Biochemistry 2007;46(30):8897-8908.

40.   Poleshko A, Shah PP, Gupta M, Babu A, Morley MP, Manderfield LJ, Ifkovits JL,

Calderon D, Aghajanian H, Sierra-Pagan JE, Sun Z, Wang Q, Li L, Dubois NC, Morrisey

EE, Lazar MA, Smith CL, Epstein JA, Jain R. Genome-Nuclear Lamina Interactions

Regulate Cardiac Stem Cell Lineage Restriction. Cell 2017;171(3):573-587 e514.

41.   Henry RA, Kuo YM, Bhattacharjee V, Yen TJ, Andrews AJ. Changing the

selectivity of p300 by acetyl-CoA modulation of histone acetylation. ACS Chem Biol

2015;10(1):146-156.

42.   Henry RA, Kuo YM, Andrews AJ. Differences in specificity and selectivity

between CBP and p300 acetylation of histone H3 and H3/H4. Biochemistry

2013;52(34):5746-5759.

FIGURE LEGENDS

Figure 1. Inhibition of HAT activity with Nu9056 treatment rescues myotube

formation in emerin-deficient myogenic progenitors. (A) Timeline showing the

time point Nu9056 was added and whole cell lysate collection for western blot

analysis. Representative images at 40X magnification of vehicle-treated wildtype (B-

E) or emerin-deficient (F-I) and Nu9056-treated wildtype (B′-E′) or emerin-deficient

(F′-I′) cells 36 h after initiating differentiation. Arrows mark myotubes (e.g., 3

myotubes in I' vs 1 myotube in I). (J-L) Quantification of >500 nuclei for each

experimental treatment (n=3) was carried out to determine the percentage of

myogenic progenitors in the cell cycle (J), expressing MyHC (K) and formed tubes

(L) 36 h after inducing differentiation. Results are mean ± s.d. of n=3; N.S., not

significant; *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; ****, $P<0.0001$ using paired, two-

tailed t-tests. Scale bars are 50 µm.

Figure 2. Inhibition of HAT activity with L002 treatment rescues myotube

formation and myosin heavy chain expression in emerin-deficient myogenic

progenitors. (A-D) or emerin-deficient (E-H) and L002-treated wildtype (A′-D′) or

emerin-deficient (E′-H′) cells 36 h after initiating differentiation. Arrows mark

myotubes (e.g., 4 myotubes in H' vs 1 myotube in H). (I-K) Quantification of >500

nuclei for each experimental treatment (n=4) was carried out to determine the

percentage of myogenic progenitors in the cell cycle (I), expressing MyHC (J) and

formed tubes (K) 36 h after inducing differentiation. Results are mean ± s.d. of n=4;

N.S., not significant; *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; ****, $P<0.0001$ using paired,

two-tailed t-tests. Scale bars are 50 µm.

Figure 3. SIRT1 activation with SRT1720 treatment does not change cell cycle

exit, myosin heavy chain expression, or myotube formation in emerin-

deficient myogenic progenitors. (A-D) or emerin-deficient (E-H) and SRT1720-

treated wildtype (A′-D′) or emerin-deficient (E′-H′) cells 36 h after initiating

differentiation. Arrows mark myotubes (e.g., 1 myotube in H' and H). (I-K)

Quantification of >500 nuclei for each experimental treatment (n=3) was carried out

to determine the percentage of myogenic progenitors in the cell cycle (I), expressing

MyHC (J) and formed tubes (K) 36 h after inducing differentiation. Results are mean

±s.d. of n=3; N.S., not significant; *, P<0.05 using paired; **, P<0.01 using paired,

two-tailed t-tests. Scale bars are 50 μm.

Figure 4. H4K5 acetylation is decreased by treatment with L002 and Nu9056.

Western blotting of whole cell lysates treated with (A) Nu9056, L002, or (B)

SRT1720 to analyze H4K5 acetylation during differentiation of wildtype and emerin-

deficient progenitors. DMSO-only treatment was the control. Three biological

replicates are shown for each treatment. (B,D) Densitometry was performed and

acetylated H4K5 in each sample was normalized to total H4 protein in each sample.

Levels of acetylated H4K5 for each condition were normalized to DMSO-treated

cells. Results are mean ± s.d. of n=3 for each condition; N.S., not significant using

paired, two-tailed t-tests.

Figure 5. Acetylation of H4K16, H3K9, H3K18 or H3K27 in emerin-deficient

myogenic progenitors upon treatment with L002, Nu9056 or SRT1720. Wildtype

or emerin-deficient myogenic progenitors were treated with DMSO, L002, Nu9056 or

SRT1720 and whole cell lysates were obtained after 36 hours. (A) Western blotting

was done with the indicated antibodies to monitor histone acetylation. (B)

Quantitation of histone acetylation normalized to γ-tubulin and plotted as fold-

change in emerin-deficient cells, as compared to wildtype cells. Results are mean ±

s.d. for each condition. N.S., not significant; *P≤0.05 using paired two-tailed t-tests.

Figure 6. Effects of altered H4K5 acetylation dynamics on myogenic

differentiation. (A) Wildtype myogenic differentiation. (B) Lack of emerin results in

impaired differentiation with loss of Myf5 localization and increased H4K5 and H3

acetylation states. (C, D) Treatment with HAT inhibitors (C) and HDAC3 activators

(D) restore the H4K5 and H4K5ac equilibrium and rescue myotube formation with

no effect on cell cycle exit. (C) HAT inhibitor L002 partially rescues myoblast

commitment by targeting unknown histone modifications to alter chromatin

architecture at the nuclear envelope. (D) Treatment with HDAC3 activators induces

Myf5 nuclear envelope localization. Red arrows indicate impaired differentiation

programming, solid dark green arrows indicate normal differentiation programming,

and dashed light green arrow signifies partially impaired differentiation.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5