A tagging method for very small fish

Tessa K. Solomon-Lane[1,2*] & Hans A. Hofmann[1-4]

[1]Department of Integrative Biology, [2]Institute for Neuroscience, [3]Center for Computational

Biology and Bioinformatics, [4]Institute for Cellular and Molecular Biology, The University of

Texas at Austin, Austin, TX 78712

*Corresponding author:

Tessa Solomon-Lane

University of Texas at Austin

Department of Integrative Biology

1 University Station #C0930

Austin, TX 78712

tksolomonlane@utexas.edu

512-475-7318

KEY WORDS: mark, identification, *Astatotilapia burtoni*, fish, juvenile, development,

longitudinal study

ABSTRACT

The ability to reliably identify individuals over time and across contexts is essential in numerous areas of science. There are a variety of well-established methods for uniquely marking individuals, such as using paint or dye, visible implant elastomer tags, numbers or barcodes glued to the animal, passive integrated transponders, and more. For some species, life history stages, and/or experiments, however, these existing tagging methods are not sufficient. Here, we describe the method we developed for tagging juveniles of the African cichlid fish, *Astatotilapia burtoni*, which are too small for the methods used to tag adults. We used fishing line threaded through the needle of an insulin syringe to tie a loop of line through the dorsal muscle of juveniles as small as 10 mm standard length. Unique color patterns on the line can be used to distinguish among individuals. The tag is compatible with normal locomotion and social behavior, discernible to the eye and on camera, durable enough to last at least months, and the juvenile can grow with the tag. For *A. burtoni*, which is a model system in social neuroscience, the lack of an appropriate tagging method for very small juveniles likely contributes to the relative lack of early-life studies, and the same may be true for other small species. We expect this method to be useful in a variety of species and will facilitate the integration of organismal and behavioral development into more research programs.

INTRODUCTION

The ability to uniquely identify individuals is essential to many scientific endeavors. Although natural patterns are sufficiently distinctive in some species, such as humpback whales (Katona et al., 1979), the African cichlid fish *Neolamprologus pulcher* (Balzarini et al., 2017; Kohda et al., 2015), the guppy *Poecilia reticulata* (Kemp et al., 2009), and the bluebanded goby *Lythrypnus dalli* (Reavis and Grober, 1999), it is usually necessary to mark individuals in order to reliably identify them over time and across contexts. The ideal tag allows for easy and unambiguous identification, lasts the duration of the experiment(s), and interferes minimally with the animal and experimental conditions (Malone et al., 1999). The method also must be legal and in compliance with the local regulations for the care and use of experimental animals. There is a variety of useful and well-established methods for marking individuals. For example, paint or dye is used in diverse species, from insects to mammals (e.g., Jones et al., 2008; Williamson et al., 2016); combinations of colored and metal bands are placed on bird legs (Frazier, 2015); fish fins can be uniquely clipped (e.g., Hammer and Blankenship, 2001; Thompson et al., 2005); visible implant elastomer tags can be injected under the skin or scales of amphibians, reptiles, and fish (e.g., Campbell Grant, 2008; Malone et al., 1999; Thompson et al., 2005); unique identifiers (e.g., numbers, barcodes) can be glued to the animal (e.g., Formica et al., 2012); passive integrated transponders (PIT) can be implanted and detected by radio signal (e.g., Jørgensen et al., 2017; Kraus et al., 2017); and more (e.g., Hasler and Faber, 2018; Volk et al., 1999). New automated tracking technology also influences marking methods (e.g., Lewejohann et al., 2009; Ohayon et al., 2013; Weissbrod et al., 2013). Certain methods will be more appropriate than others depending on the spatial and temporal scales involved and the requirements and limitations of the study species and experiment.

Despite the range of existing options, suitable marking solutions are still lacking for some

species, life history stages, and/or experiments. Here, we describe the tagging method we

developed for juveniles of the African cichlid fish, *Astatotilapia burtoni*, a model system in

social neuroscience (Fernald and Maruska, 2012; Hofmann, 2003). The adults of this species are

routinely tagged using a visible implant elastomer tag or a colored bead secured with a plastic tag

through the dorsal muscle (as in Trainor and Hofmann, 2006). In juveniles, however, there is not

sufficient dorsal muscle into which to inject elastomer, and the bead and tag are too large and

heavy. We used fishing line threaded through the needle of an insulin syringe to tie a loop of line

through the dorsal muscle of fish as small as 10 mm standard length (SL). This tag does not

appear to impede the locomotion or social behavior of freely interacting fish, it is visible to the

naked eye and on camera for video analysis, it can remain in place as the animal grows, and it is

sufficiently long-lasting for the duration of our experiments (months). We expect this method

will be useful to other researchers in a variety of species and will make it feasible to incorporate

studies on organismal and behavioral development into more research programs that work with

small animals.

METHODS

Making the tag

The needle attached to the fishing line that we use is modeled on an eyeless suture, which

is expensive to purchase ($6-$24 per suture). Our alternative can be easily made and with

inexpensive materials. We used Berkley Nanofil Fishing Line, 0.006 in (0.15 mm) average

diameter, in the color clear mist (150 yards for $20) and BD Ultra-Fine™ Short Needle (8 mm,

31G) Insulin Syringes (3/10 mL) (100 for $75). Different fishing line and needle combinations

could also be used, as long as the average diameter of the line is smaller than the inner diameter

of the needle. We used the smallest, thinnest combination of materials that were available. The

procedure for making the tag is as follows:

1) To attach the fishing line to the needle, remove the plunger from the syringe and

thread the line through the sharp end of the needle. It is easiest to do under a

dissecting microscope, and it helps to work with a freshly cut end. Use a sharp, new

razor to avoid a messy cut, which can flatten the line and widen the diameter. If the

fishing line will clearly not thread into the needle, try discarding a length of line and

starting at a new place in the spool. The diameter of the line varies slightly

throughout, and it is easiest to work at an average or narrower section.

2) Once started, keep threading line through the needle until it has gone beyond the

length of the needle and is visible in the barrel of the syringe (Figure 1A).

3) Take a pair of forceps and grip the base of the needle, being careful not to dislodge

the line. Rock the needle back and forth with the forceps until it detaches from the

syringe (Figure 1B).

4) From the blunt side of the needle, pull the line through until it is sufficiently long. We

typically work with a length of line between 30 cm and 50 cm, which can be used to

tag ~4-10 fish.

5) Cut the length of line from the spool using a razor blade and pull the last bit of line

through the needle until is no longer sticking out beyond the sharp end. We do not

find it necessary to crimp the blunt end of the needle to keep the line in place.

Astatotilapia burtoni **juveniles**

We used *A. burtoni* juveniles from a laboratory population descended from a wild-caught

stock. The individuals that bred the juveniles were housed in naturalistic social groups of males

and females. Dominant males court gravid females that then lay eggs in his territory. The female

then scoops up the eggs into her mouth, where the male fertilizes them. The mother orally

incubates the larvae as they develop (Fernald and Hirata, 1979; Renn et al., 2009). When the

larvae are fully developed and ready to leave the mother's mouth 10-13 days after fertilization

(Fernald and Hirata, 1979; Renn et al., 2009), the average SL is 8.31 ± 0.039 mm (n=356, max

SL: 10.18 mm, min SL: 4.6 mm). The smallest juveniles we have successfully tagged were 10

mm SL (Figure 2A); therefore, even with this technique, it is not yet possible to tag juveniles

immediately upon release by the mother.

Tagging the fish

We anesthetized juveniles in tricaine methanesulfonate (MS-222, Sigma Aldrich) at a

dose of 0.0006 g / mL aquarium water, buffered with sodium bicarbonate to pH 7-7.5. We

removed fish from the MS-222 immediately after they stopped responding to touch, which

occurred after losing equilibrium and ceasing ventilation (opercular movement). Juveniles were

then positioned on a wet paper towel flat on their side and tagged as follows:

1) Holding the needle with forceps perpendicular to the fish, push the needle through the

dorsal muscle (Figure 2B).

2) Pull the needle, followed by the line, through the dorsal muscle with the forceps until 3-4

cm of line remains (Figure 2C). Use a razor blade to cut the line on the needle side,

leaving another 3-4 cm.

3) Using forceps, tie a square knot in the line (Figure 2D, E), and trim the excess using a

razor blade. The knot can also be tied by hand, although this method uses more line.

4) Make the loop large enough for the fish to raise its dorsal fin and to grow over the course

of the experiment. For longer experiments, place a drop of super glue on the knot to

ensure it stays tied.

5) As soon as possible, place the fish in water to recover. The glue should dry sufficiently in

less than a minute, and during this time (and throughout), the fish's gills / body can be

kept wet. Once in the water, if the fish does not start ventilating on its own, we use a

transfer pipette to gently push water over the gills until the opercula move regularly.

Anesthetizing and tagging one fish takes 2 min. The tag can be removed quickly, and

without anesthesia, by cutting the line with a sharp razor blade.

Uniquely identify individuals

To distinguish between multiple tagged fish in the same group or enclosure, we use

permanent markers to uniquely color the white fishing line (before the line is threaded through

the fish). For a short experiment (<1week), the color(s) remain vibrant under our aquarium

conditions, but over time, the colors fade. For long-term experiments, we color the line and then

add a small drop of super glue to different places along the loop to create a unique pattern. The

color under the super glue "bead" lasts at least two months. It may also be possible to thread a

seed bead onto the loop for unique identification. Adding a seed bead to the loop may also be

useful as a unique identifier. Although we have not yet tried this, seed beads are very small,

readily available in an array of colors, and inexpensive.

CONCLUSIONS

The tagging method described here has made it possible to study the behavioral

development and the underlying neuromolecular mechanisms of juvenile *A. burtoni*, a species

that is well studied in adulthood but strikingly understudied during development. The lack of an

appropriate method for identifying small juveniles likely contributes to the relative paucity of

early-life studies, in this and other small species. In social neuroscience, behavioral ecology,

animal behavior, and related fields, experiments that follow individuals through development are

critical to uncovering the emergence of individual phenotypic variation, as well as the underlying

mechanisms and fitness consequences (Taborsky, 2016). We expect this tagging method for very

small fish will be broadly useful and hope that other researchers will continue to improve upon

the technique.

ACKNOWLEDGEMENTS

We thank Isaac Miller-Crews and Gabriel Rocha for the size measurements of juvenile *A.*

burtoni and the Hofmann Lab for collecting the size data. This work was supported by NSF grant

IOS-1354942 to HAH and the BEACON Center for the Study of Evolution in Action awards

#947 (2016) and #1081 (2017, 2018) to TKSL and HAH.

REFERENCES

Balzarini, V., Taborsky, M., Villa, F., Frommen, J.G., 2017. Computer animations of color markings reveal the function of visual threat signals in Neolamprologus pulcher. Curr. Zool. 63, 45–54. https://doi.org/10.1093/cz/zow086

Campbell Grant, E.H., 2008. Visual Implant Elastomer Mark Retention through Metamorphosis in Amphibian Larvae. Source J. Wildl. Manag. 72, 1247–1252. https://doi.org/10.2193/2007-183

Fernald, R.D., Hirata, N.R., 1979. The Ontogeny of Social behavior and body coloration in the african cichlid fish haplochromis burtoni. Zeitschrif für Tierpsychologie. https://doi.org/10.1111/j.1439-0310.1979.tb01025.x

Fernald, R.D., Maruska, K.P., 2012. Social information changes the brain. Proc. Natl. Acad. Sci. 109, 17194–17199. https://doi.org/10.1073/pnas.1202552109

Formica, V., Wood, C.W., Larsen, W.B., Butterfield, R.E., Augat, M.E., Hougen, H.Y., Brodie, E.D., 2012. Fitness consequences of social network position in a wild population of forked fungus beetles (Bolitotherus cornutus). J. Evol. Biol. 25, 130–137. https://doi.org/10.1111/j.1420-9101.2011.02411.x

Frazier, K., 2015. The History of Bird Banding. Audubon (Website).

Hammer, S.A., Blankenship, H.L., 2001. Cost Comparison of Marks, Tags, and Mark-with-Tag Combinations Used in Salmonid Research. N. Am. J. Aquac. 63, 171–178. https://doi.org/10.1577/1548-8454(2001)063<0171:CCOMTA>2.0.CO;2

Hasler, A.D., Faber, W.M., 1941. A Tagging Method for Small Fish 1941, 162–165.

Hofmann, H.A., 2003. Functional genomics of neural and behavioral plasticity. J. Neurobiol. 54, 272–282. https://doi.org/10.1002/neu.10172

Jones, T.H., Godfray, H.C.J., Hassell, M.P., 2008. Relative Movement Patterns of a Tephritid Fly and Its Parasitoid Wasps Published by: Springer in cooperation with International Association for Ecology Stable URL: http://www.jstor.org/stable/4221264 Your use of the JSTOR archive indicates your acceptan 106, 317–324.

Jørgensen, M.G.P., Deurs, M. van, Butts, I.A.E., Jørgensen, K., Behrens, J.W., 2017. PIT-tagging method for small fishes: A case study using sandeel (Ammodytes tobianus). Fish. Res. 193, 95–103. https://doi.org/10.1016/j.fishres.2017.04.002

Katona, S., Baxter, B., Brazier, O., Kraus, S., Perkins, J., Whitehead, H., 1979. Identification of Humpback Whales by Fluke Photographs, in: Behavior of Marine Animals. Springer US, Boston, MA, pp. 33–44. https://doi.org/10.1007/978-1-4684-2985-5_2

Kemp, D.J., Reznick, D.N., Grether, G.F., Endler, J.A., 2009. Predicting the direction of ornament evolution in Trinidadian guppies (Poecilia reticulata). Proc. R. Soc. B Biol. Sci. 276, 4335–4343. https://doi.org/10.1098/rspb.2009.1226

Kohda, M., Jordan, L.A., Hotta, T., Kosaka, N., Karino, K., Tanaka, H., Taniyama, M., Takeyama, T., 2015. Facial recognition in a group-living cichlid fish. PLoS One 10, 1–12. https://doi.org/10.1371/journal.pone.0142552

Kraus, B.T., McCallen, E.B., Williams, R.N., 2017. Evaluating the Survival of Translocated Adult and Captive-reared, Juvenile Eastern Hellbenders (Cryptobranchus alleganiensis alleganiensis). Herpetologica 73, 271–276.

Lewejohann, L., Hoppmann, A.M., Kegel, P., Kritzler, M., Krüger, A., Sachser, N., 2009. Behavioral phenotyping of a murine model of Alzheimer's disease in a seminaturalistic environment using RFID tracking. Behav. Res. Methods 41, 850–856.

https://doi.org/10.3758/BRM.41.3.850

Malone, J.C., Forrester, G.E., Steele, M.A., 1999. Effects of subcutaneous microtags on the
growth, survival, and vulnerability to predation of small reef fishes. J. Exp. Mar. Bio. Ecol.
237, 243–253. https://doi.org/10.1016/S0022-0981(99)00003-9

Ohayon, S., Avni, O., Taylor, A.L., Perona, P., Roian Egnor, S.E., 2013. Automated multi-day
tracking of marked mice for the analysis of social behaviour. J. Neurosci. Methods 219, 10–
19. https://doi.org/10.1016/j.jneumeth.2013.05.013

Reavis, R.H., Grober, M.S., 1999. An integrative approach to sex change: social, behavioural
and neurochemical changes in Lythrypnus dali (Pisces) 2, 51–60.

Renn, S., Carleton, J.B., Magee, H., Nguyen, M.L.T., Tanner, A.C.W., 2009. Maternal care and
altered social phenotype in a recently collected stock of Astatotilapia burtoni cichlid fish.
Integr. Comp. Biol. 49, 660–673. https://doi.org/10.1093/icb/icp085

Taborsky, B., 2016. Opening the Black Box of Developmental Experiments: Behavioural
Mechanisms Underlying Long-Term Effects of Early Social Experience. Ethology 122,
267–283. https://doi.org/10.1111/eth.12473

Thompson, J.M., Hirethota, P.S., Eggold, B.T., 2005. A Comparison of Elastomer Marks and Fin
Clips as Marking Techniques for Walleye; A Comparison of Elastomer Marks and Fin Clips
as Marking Techniques for Walleye. North Am. J. Fish. Manag. 25, 308–315.
https://doi.org/10.1577/M03-237.1

Trainor, B.C., Hofmann, H.A., 2006. Somatostatin Regulates aggressive behavior in an african
cichlid fish. Endocrinology 147, 5119–5125. https://doi.org/10.1210/en.2006-0511

Volk, E.C., Schroder, S.L., Grimm, J.J., 1999. Otolith thermal marking. Fish. Res. 43, 205–219.

Weissbrod, A., Shapiro, A., Vasserman, G., Edry, L., Dayan, M., Yitzhaky, A., Hertzberg, L.,
Feinerman, O., Kimchi, T., 2013. Automated long-term tracking and social behavioural
phenotyping of animal colonies within a semi-natural environment.
https://doi.org/10.1038/ncomms3018

Williamson, C.M., Lee, W., Curley, J.P., 2016. Temporal dynamics of social hierarchy formation
and maintenance in male mice. Anim. Behav. 115, 259–272.
https://doi.org/10.1016/j.anbehav.2016.03.004

FIGURES

Figure 1: Making the tag. A) Thread the fishing line through the needle of an insulin syringe

until it is visible in the barrel of the syringe. B) Use forceps to grasp the base of the needle. Rock

back and forth (arrows) until the needle breaks off from the syringe. Not shown: From the blunt

side of the needle, pull the line through, until it no longer sticks out beyond the needle tip. Use a

razor blade to cut off excess line.

Figure 2: Tagging fish. A) Image of a tagged juvenile *A. burtoni* (10.5 mm standard length) on

a ruler (inches top, mm bottom). B) Pierce the dorsal muscle with the needle. C) Pull the needle

and line through the dorsal muscle. D) Make a loop in the line and tie the first half of a square

knot. Adjust the loop to the desired size. E) Finish the square knot, and cut the excess line using

a razor blade. Note: The same individual is pictured in each photograph.