Methods for tagging very small fish

Deijah D. Bradley[1], Elianna J. Schimke[2], Alyssa P. Alvey[1], Hans A. Hofmann[3,4], Tessa K.

Solomon-Lane[1-3,5*]

[1]Pitzer College, Claremont, CA 91711

[2]Scripps College, Claremont, CA 91711

[3]Department of Integrative Biology, [4]Institute for Neuroscience, The University of Texas at

Austin, Austin, TX 78712

[5]Claremont McKenna College, Claremont, CA 91711

*Corresponding author:

Dr. Tessa Solomon-Lane

Department of Natural Sciences / W.M. Keck Science Department

925 N. Mills Ave

Claremont, CA 91711

TSolomonLane@NatSci.Claremont.edu

909-607-7857

ABSTRACT

Identifying individuals over time and across contexts is essential in many scientific fields. There

are a variety of well-established methods for uniquely marking individuals (e.g., visible implant

elastomer, barcodes, paint). However, for some species, life history stages, and/or experiments,

existing methods are not sufficient. Here, we describe two methods – a tattoo method and a

piercing method – for tagging juveniles of the African cichlid fish, *Astatotilapia burtoni*, which

are too small for the methods used with adults, or other known methods. The tattoo method is

appropriate for the youngest, smallest *A. burtoni* juveniles and injects tattoo ink into the dorsal

muscle. Different colors and injection locations can be used to distinguish among individuals

over a period of weeks (up to 4 weeks, average 2.5-3 weeks under our conditions). Because fish

this young and small are sensitive to handling and injection, we also include physiological data

showing fish recover well from anesthetization and tagging. The piercing method is appropriate

for fish of all sizes, including the youngest, smallest juveniles. It uses very thin fishing line

threaded through the dorsal muscle and tied into a barbell or loop. Unique colors and patterns

can be used to distinguish among individuals over a period of months. We also include open

field exploration behavior data because a physical tag might impede normal movement in very

small fish. We expect these effective and inexpensive methods to be useful for a variety of small

species and will facilitate early-life, developmental, and longitudinal research.

KEY WORDS: *Astatotilapia burtoni*, development, fry, identification, juvenile, longitudinal,

mark

INTRODUCTION

The ability to uniquely identify individuals is essential to many scientific endeavors. In some species, the natural, external patterns on the body are sufficiently distinctive for identification, such as humpback whales, *Megaptera novaeangliae* (Katona et al., 1979), the African cichlid fish, *Neolamprologus pulcher* (Kohda et al., 2015; Balzarini et al., 2017), the guppy, *Poecilia reticulata* (Kemp et al., 2009), and the bluebanded goby, *Lythrypnus dalli* (Reavis and Grober, 1999). For most species, however, it is necessary to mark individuals to reliably identify them over time and across contexts. The ideal tag allows for easy and unambiguous identification, lasts the duration of the experiment(s), and interferes minimally with the animal and experimental conditions (Malone et al., 1999). The method also must be legal and in compliance with the local regulations for the care and use of experimental animals. There are a variety of useful and well-established methods for marking individuals. For example, paint or dye is used in diverse species, from insects to mammals (e.g., Jones et al., 1996; Williamson et al., 2016); combinations of colored and metal bands are placed on bird legs (Frazier, 2015); fish fins can be uniquely clipped (e.g., Hammer and Lee Blankenship, 2001; Thompson et al., 2005); visible implant elastomer tags can be injected under the skin or scales of amphibians, reptiles, and fish (e.g.,(Malone et al., 1999; Thompson et al., 2005; Grant, 2008); unique identifiers (e.g., numbers, barcodes) can be glued to the animal (e.g., Formica et al., 2012); passive integrated transponders (PIT) can be implanted and detected by radio signal (e.g., Jørgensen et al., 2017; Kraus et al., 2017); surgical steel coded wire implant tags (e.g., Martin and Wainwright, 2013; Martin and Gould, 2020), and more (e.g., Hasler and Faber, 1941; Volk et al., 1999). Developments in automated tracking technology have also influenced identification methods, such as radio frequency identification (RFID) (e.g., Lewejohann et al., 2009;

Weissbrod et al., 2013) and computer-vision systems (e.g., bleach-marks on mouse fur, Ohayon

et al., 2013). The spatial and temporal scales involved, and the requirements and limitations of

the study species and experiment, together determine the most appropriate marking method.

Despite the variety of existing options, suitable marking solutions are still lacking for

some species, life history stages, and/or experiments. Here, we describe two tagging methods – a

tattoo method and a piercing method – we developed for juveniles of the African cichlid fish,

Burton's Mouthbrooder (*Astatotilapia burtoni*), a model system in social neuroscience

(Hofmann, 2003; Maruska and Fernald, 2018). Adults of this species are routinely tagged using

visible implant elastomer tags, a colorful liquid that dries as a pliable solid (Northwest Marine

Technology, Inc.), or a colored bead secured with a plastic tag through the dorsal muscle (as in

Trainor and Hofmann, 2006). However, the smallest, youngest juvenile *A. burtoni* (average size:

8.31 ± 0.039 mm standard length; 0.022 ± 0.0005 g mass, Fig 1A) do not have sufficient dorsal

muscle into which to inject elastomer (Fig 1B-E), which uses a 30-gauge needle because of the

viscosity of the substance. The bead (often cut in half to be lighter) and plastic tag (can also be

used alone) are too large and heavy for the juveniles.

The tattoo method injects nanoliter quantities (~100 nL) of tattoo ink into the dorsal

muscle using a pulled glass capillary tube needle. We can use this method with the youngest and

smallest fish, including larva that are still developing. This method is moderately long-lasting: up

to 4 weeks, with an average of 2.5-3 weeks under our laboratory conditions. This is similar to

injection methods in other species (Thresher and Gronell, 1978). This tag does not appear to

interfere with locomotion, and it is visible to the naked eye. Because fish of this early

developmental stage are especially sensitive to handling and prone to injury or death, we also

present physiological data on how anesthetized and injected juveniles recover from the procedure

compared to anesthetized juveniles that were not injected (Sneddon, 2012).

The piercing method threads very thin fishing line through the dorsal muscle of the fish

using a pin or needle appropriate to the fish's size. The line is then tied into a barbell or loop and

remains in place. These tags are visible to the naked eye, visible on camera for video analysis

(depending on the size of the tag and cameras), it can remain in place as the juvenile grows, and

it is sufficiently long-lasting for the duration of our experiments (months). Because a physical

tag might impede locomotion in the smallest fish, we also include data from an open field

exploration behavior test using 1-day old juveniles. We chose these young fish because they are

most likely to be physically impeded. Anesthesia recovery data are not included for this method.

Overall, we hope these methods will be useful to other researchers working with a variety

of species and that they will continue to be developed, adapted, and improved. The ability to

identify individuals over time, starting in early life, will make it more feasible to carry out

developmental and longitudinal research when working with very small, young fish.

Tattoo ink injection

METHODS

Astatotilapia burtoni juveniles

We used *A. burtoni* juveniles from a laboratory population descended from a wild-caught

stock (see lab strain from Hofmann and Renn: Pauquet et al., 2018). The individuals that bred the

juveniles were housed in naturalistic social groups of males and females. Dominant males court

gravid females that then lay eggs in his territory. The female then scoops up the eggs into her

mouth, where the male fertilizes them. The mother orally incubates the larvae as they develop.

When the larvae are fully developed and ready to leave the mother's buccal cavity 10-13 days

after fertilization (Fernald and Hirata, 1979; Renn et al., 2009), the average standard length (SL)

in our laboratory populations is 8.31 ± 0.039 mm (n=356, max SL: 10.18 mm, min SL: 4.6 mm,

Fig 1A). All procedures were done in compliance with the Institutional Animal Care and Use

Committee (IACUC) of the W.M. Keck Science Department of Scripps, Pitzer, and Claremont

McKenna Colleges (protocol #19-001) and the IACUC of The University of Texas at Austin.

Tattoo ink and Nanoject injection set-up

We developed this method using commercially available sets of tattoo ink, including

seven color sets of Millennium Mom's Tattoo Ink (Amazon.com, $40 USD at the time of the

work) and HAWINK Tattoo Ink (Amazon.com, $20 USD at the time of the work). These sets

included vibrant colors such as red, yellow, green, and blue (Supplemental Fig 1), as well as

neutral colors such as white, gray, and black. We also used Keep It Wet (Eternal Tattoo Supply,

$5 USD at the time of the work) to keep the ink from drying as quickly, without affecting the

color.

Tattoo injections were performed under a dissection microscope using the Nanoject II

Auto-Nanoliter Injector (Drummond Scientific Company). The Nanoject was positioned on a

micromanipulator attached to a magnetic base, which was secured to the benchtop on a magnetic

board. See Supplemental Materials for detailed set-up information. Briefly, we used a puller

(World Precision Instruments PUL-1) to make glass capillary tube needles (1.2 mm x 68 mm, A-

M Systems). The very tip was broken off using forceps, we backfilled the needle with mineral

oil, and then we secured the needle onto the Nanoject (Figure 2A, Supplemental Fig 2). Using

the Nanoject controls, we emptied a portion of the mineral oil, and then filled the needle with

tattoo ink. We mixed the ink in a weigh boat or on Parafilm. Different ink colors and brands had

different consistencies and drying times in the needle, so we used a ratio ranging from 1:1 to 1:4

ink drops to water drops (or Keep It Wet additive). We performed multiple test injections

(Supplemental Fig 2F) at the start and between fish to ensure the ink was not dried, clogging the

needle.

Injection of tattoo ink

We removed the fish from the anesthetic (see below) and placed it on the injection

platform under the dissection scope on a paper towel wet with aquarium water (Fig 2A). If the

Nanoject is set-up to the right of the scope (for a right-handed scientist), we positioned the fish

with its head to the left and tail to the right so the needle inserted between or under the fish's

scales. Securing the fish's head between the left thumb and forefinger, pressing firmly down on

the paper towel but without squeezing the fish, we positioned the needle at the desired location

along the rostral-caudal axis of the dorsal muscle (Fig 2B). A ventral injection in or near the

abdominal cavity will not remain visible for long, and the tattoo ink may affect internal organs.

The needle was lowered using the micromanipulator until the skin was pierced and the tip

of the needle was in the muscle. Young *A. burtoni* juveniles are semi-transparent at this

developmental stage, so we could also visibly check the needle position (Fig 2B). There may be

some resistance to piercing the skin, which can indicate a dull needle that should be replaced.

Holding the fish as still as possible, we pressed "inject" twice for a total injection volume of

~100 nL of ink. Waiting a couple seconds to remove the needle can help prevent the ink from

flowing out of the injection site. The ink is immediately visible at the injection site, as well as on

the opposite side of the fish (Fig 2C-E). Over time, the tag is usually most visible just on one

side of the body (Fig 2E). We then carefully placed the fish into a beaker of fresh aquarium

water and began the resuscitation protocol (see below). The needle was cleaned with ethanol on a

Kimwipe (Kimberly-Clark) between injections. We changed the needle approximately every 4-8

animals, as the needle dulled, if the needle clogged, or to change ink color.

For experiments that require strict standardization of the procedure, we recommend

keeping the amount of time out of water consistent across fish (e.g., 1 min for tattooing only; 2

min for tattooing and an additional procedure, such as drug administration). Otherwise, we place

the fish back in water for recovery as quickly as possible. The procedure can take less than 30 s

once the fish is anesthetized.

Anesthesia and post-tagging resuscitation

We anesthetized juveniles less than two weeks old in tricaine methanesulfonate (MS-222,

Sigma Aldrich) at a dose of 0.0003 g / mL aquarium water, buffered with sodium hydroxide to

pH 7-7.5. During anesthetization, we recorded the time it took for body movement to stop; loss

of equilibrium, when the fish's dorsal side first rolled away from an upright orientation; and

when breathing stop, indicated by movement of the opercula stopping fully. We removed fish

from the MS-222 immediately after breathing stopped. When tagging fish without collecting

anesthetization and/or recovery data, we remove fish from the MS-222 immediately after

breathing stopped or after 30-35 s, whichever happens earlier.

After tagging (see above), the fish was placed in a 200 mL plastic beaker of fresh

aquarium water to recover. We immediately began resuscitating the fish using a plastic pipette to

gently push water over the gills. Recovery from MS-222 is a highly stereotyped process

(McFarland, 1959). We recorded the time it took for the fish to first move its body; first move

the opercula; breathe independently without resuscitation; initial regaining of equilibrium, when

the dorsal side of the fish first returned to an upright position; and sustained equilibrium, when

the fish was permanently upright. We also recorded when the fish looked fully recovered, which

tended to happen suddenly. Fully recovered fish swam smoothly and breathed steadily, with no

wobbles in equilibrium.

Statistics

All statistical analyses were conducted using R Studio (R version 4.4.0) (RStudio Team,

2022). Results were considered significant at the $p<0.05$ level, and averages ± standard error of

the mean are included in the text. To compare anesthetization and recovery in control and tagged

juveniles, we used t-tests for data that met the assumptions of parametric statistics. A Wilcoxon

signed ranks test was used for data that did not meet those assumptions. Cohen's d is reported to

estimate effect size for significant pairwise differences (small effect: $0.2 < d < 0.5$; medium: 0.5

$< d < 0.8$; large: $0.8 < d$). Pearson correlations were used to test for associations among recovery

metrics in control and tagged fish.

RESULTS

Recovery from anesthesia and injection

To investigate the safety and efficacy of this procedure on very young, very small

juveniles (< 2 weeks after removal from the mother's buccal cavity), we compared the

anesthetization and recovery of fish that were both anesthetized and tagged (tagged: n=8) to fish

that were anesthetized but not injected (control: n=8). Fish in both treatment groups were

removed from the MS-222 immediately after breathing stopped and kept out of water for 1 min

on a wet paper towel. Tagged fish were injected during the time out of water.

While being anesthetized, juveniles first lost equilibrium (30.6 ± 2.9 s), after which they

stopped moving their bodies (52.5 ± 3.0 s) and stopped breathing (54.1 ± 3.5 s) in close

succession. Some individuals appeared to stop breathing first, and others appeared to stop

moving first. The treatment groups did not differ in time to lose equilibrium or time to stop

moving (Table 1, Supplemental Fig 3A, B). Fish in the tagged group (although they had not been

tagged yet) took significantly longer to stop breathing (Supplemental Fig 3C), but overall, the

groups did not differ in the total time spent in MS-222 (Table 1, Supplemental Fig 3D).

All (100%) of the fish in this cohort successfully recovered. After being placed in the

recovery beaker, opercular movement during resuscitation occurred first (40.4 ± 9.5 s), followed

by independent breathing without resuscitation (107.9 ± 18 s). First body movement ($128.4 \pm$

17.2 s), initial regained equilibrium (145.3 ± 21.9 s), and sustained regained equilibrium ($162.6 \pm$

26.8 s) occurred next, in close succession. By definition, full recovery occurred last (407.3 ± 51

s). There were no significant differences for any recovery metric between fish that were

anesthetized compared to those that were anesthetized and tagged (Table 1, Fig 3A, B). Some

recovery metrics were good predictors of later recovery (Table 2, Supplemental Fig 4).

Independent breathing was significantly and positively associated with first body movement (Fig

3C) and initial regained equilibrium for both control and tagged fish, as well as sustained

regained equilibrium in control fish. Initial body movement was significantly and positively

associated with initial regained equilibrium in control and tagged fish, as well as sustained

regained equilibrium in control fish. Initial regained equilibrium was significantly and positively

associated with sustained regained equilibrium in control and tagged fish. Initial (Fig 3D) and

sustained equilibrium were the only metrics associated with full recovery, and the associations

were significant only for control fish. Initial breathing during resuscitation did not predict any

later recovery points (Table 2).

DISCUSSION

Injecting tattoo ink into the dorsal muscle of very young, small fish immediately creates a

vibrant visual mark, allowing for individual identification. Tattoo ink is inexpensive and

commercially available in an array of colors. Mixing inks may also be useful in creating unique

colors to track more individuals within a group. Depending on the size of the fish, injection

location may also be a useful unique identifier. The main limitation of this technique in our

hands is that the colors fade over time, with lighter colors (e.g., yellow) appearing to fade more

quickly than darker colors (e.g., black, blue). Our longest lasting tattoo tags lasted 3-4 weeks

(Fig 2E). Different colored inks, and different manufactures, have different compositions

(Moseman et al., 2024), which could underlie the differences in tag visibility we observed. The

different ink components could also have differential health effects (Grant et al., 2015; Moseman

et al., 2024). In humans, tattoo ink is injected into the skin, specifically the papillary layer of the

dermis. Over time, ink particles move deeper into the reticular dermis, and the tattoo persists

because sufficient pigment stays in the skin (Grant et al., 2015; Moseman et al., 2024). Here, we

injected ink into the dorsal muscle, which resulted in a moderately long-lasting tag, but the mark

was not permanent. Pigment may disperse more readily, or fail to accumulate, in muscle

compared to skin. Muscle can also change dramatically during fish development, including in

muscle fiber type, size, and number (Patruno et al., 1998; Johnston, 1999).

Although tagging can take less than 30 s, anesthesia is necessary for the welfare and

recovery of the juvenile fish. We chose to use MS-222 (pH buffered), which is a common and

safe fish anesthetic (Wagner et al., 2003; Kiessling et al., 2009; Sneddon, 2012; Zahl et al., 2012;

Topic Popovic et al., 2012) that can be cleared from circulation in less than 300 s (Kiessling et

al., 2009). Observing individuals during anesthetization and recovery is highly informative for

establishing an effective anesthetic dose, determining the optimal procedural timing, and

ensuring full recovery (Sneddon, 2012). For example, some fish species will recovery unaided

from MS-222 (e.g., bluebanded gobies, *Lythrypnus dalli*, Solomon-Lane and Grober, 2012);

however, *A. burtoni* juveniles recover best with resuscitation and continued monitoring until full

recovery. These data can also be used to identify unintended biases. Here, we found that

juveniles in the tagged group took longer to stop breathing, even though they had not yet

experienced anything different from the control fish (i.e., no fish had yet been tagged). With this

information, we can test for downstream consequences. For example, despite this significant

difference, tagged fish did not spend more total time in MS-222.

We found no significant differences in recovery timing between anesthetized control fish

compared to anesthetized and tagged fish. This suggests that, overall, tattoo ink injections are

safe, with limited negative physiological consequences. We did identify some effects of tagging

on the temporal progression of recovery. In control fish, for example, the timing of initial and

sustained regained equilibrium was significantly associated with the timing of full recovery. In

contrast, for tagged fish, no recovery metric predicted the timing of full recovery. These

differences support our protocol of using recovery data as an indicator of how an individual was

impacted by the procedure and if it is ready to return to its home aquarium or engage in an

experimental test (e.g., a behavioral assay) (Solomon-Lane and Grober, 2012). For example, we

found that the timing of first opercular movement (initial breathing) during resuscitation was not

a good predictor of subsequent stages of recovery. In contrast, independent breathing and body

movement, which occurred in close succession, predicted when equilibrium was regained.

Additional validations may be needed, depending on the species or experiment. For example, in

our future studies related to social behavior, we will test whether tag color affects how fish

behave or are treated by other members of their social group.

We also tested the efficacy of injecting acrylic paint (Lotrich and Meredith, 1974),

thinned slightly with Liquidtex® Flow Aid^{TM} to prevent the paint from drying rapidly in the

needle. These acrylic paint tags lasted approximately two weeks, with the longest tag lasting four

weeks, under our conditions. Acrylic paint is non-toxic, inexpensive, and available in many

colors, making it a potentially good option, in addition to tattoo ink. Injections of Alcian blue

(Thermo Scientific Chemicals) lasted approximately three weeks. With no color options other

than blue, this did not meet our needs for distinguishing among multiple juveniles in a social

group. Injections of food coloring lasted only a couple of days (data not shown).

Fishing line piercing method

METHODS

Making the fishing line tag

Unique tags can be made from fishing line by coloring or dying the line, or purchasing

fishing line manufactured in different colors (options may be limited). We used permanent

markers (Sharpie) to uniquely color white fishing line (before the line is threaded through the

fish). "Beads" of different colors, sizes, or patterns can also be made along the length of the line

by tying a knot (or more than one) and using a drop of super glue to preserve the color. Nail

polish can also be used over the knot(s) for additional color combinations. Note that these beads

should be on a portion of the tag that does not get threaded through the body of the fish, as they

would cause more tissue damage.

For the juveniles that are too small for a 31-gauge insulin needle (or the smallest

commercially available needle) to pass through the dorsal muscle (Fig 1D, E), we use fishing

line in combination with a minutien pin (Fine Science Tools, 0.1mm diameter, 500 pins for $32

USD at the time of the work). We used the thinnest line available, which was Berkley NanoFil

Uni-filament fishing line (2 lb break weight, 0.051 mm average diameter, 150 yards for $20 USD

at the time of the work). For this method, the fishing line and pin are not physically attached.

For juveniles that are large enough for a 31-gauge insulin needle (body size > ~12 mm

SL, ~0.08 g mass for *A. burtoni*), we find it easiest to tag many fish at once by attaching the

fishing line to a needle. This design is modeled on an eyeless suture, which can be expensive to

purchase. Our alternative is easy and inexpensive to make. We used Berkley NanoFil Uni-

filament Fishing Line (4lb break weight, 0.1 mm average diameter, in the color clear mist; 150

yards for $20 USD at the time of the work) and BD Ultra-Fine[TM] Short Needle (8 mm, 31G)

Insulin Syringes (3/10 mL) (90 count for $21 USD at the time of the work). Any fishing line and

needle combination can be used, as long as the average diameter of the line is smaller than the

inner diameter of the needle.

To attach the fishing line to the needle, remove the plunger from the syringe and thread

the line through the sharp end of the needle. It is easiest to do under a dissection microscope, and

it helps to work with a freshly cut end. Use a sharp, new razor blade to avoid a messy cut, which

can flatten the line and widen the diameter. If the fishing line will clearly not thread into the

318 needle, try discarding a length of line and starting at a new place along the spool. The diameter

319 of the line varies slightly throughout, and it is easiest to work at an average or narrower section.

320 Once started, keep threading the line through the needle until it has gone beyond the length of the

321 needle and is visible in the barrel of the syringe (Figure 4A). Then, take a pair of forceps and

322 grip the base of the needle, being careful not to dislodge the line. Rock the needle back and forth

323 with the forceps until it detaches from the syringe (Figure 4B). From the blunt side of the needle,

324 pull the line through until it is sufficiently long. We typically work with a length of line between

325 30 cm and 50 cm, which can be used to tag ~4-10 fish. Cut the length of line from the spool

326 using a razor blade and pull the last bit of line through the needle until it is no longer sticking out

327 beyond the sharp end. We do not find it necessary to crimp the blunt end of the needle to keep

328 the line in place.

330 *Anesthesia and resuscitation*

331  We anesthetized juveniles less than 2 weeks old as described above. Older, larger

332 juveniles were anesthetized in MS-222 at a dose of 0.0006 g / mL aquarium water, buffered to

333 pH 7-7.5. We removed the larger juveniles from the MS-222 immediately after they stopped

334 responding to touch, which occurred after losing equilibrium and after breathing stopped. Fish

335 were resuscitated as above.

337 *Tagging the fish*

338  For the smallest fish (Fig 5A, B), place the fish on a wet paper towel under a dissection

339 scope. Using forceps to hold the minutien pin perpendicular to the fish, push the needle into and

340 through the dorsal muscle. With the pin still in place, reposition the fish to be upright, and pull

the pin through the other side of the fish. Using forceps to hold the fishing line, immediately

push the line through the hole made by the pin. If there are knots on the line, be careful not to

pull those through the fish. For larger fish (Fig 5C), follow a similar procedure with the needle

attached to the fishing line. With the fish lying flat on a wet paper towel, use forceps to hold the

needle perpendicular to the fish. Shifting the fish to an upright position, push the needle through

the dorsal muscle (Figure 5D). Pull the needle, followed by the line, through the dorsal muscle

with the forceps until 3-4 cm of line remains (Figure 5E). Use a razor blade to cut the line on the

needle side, leaving another 3-4 cm.

Once the fishing line is through the dorsal muscle of the fish, the fishing line can be tied

into a barbell (Fig 5B) or into a loop (Fig5 A,C) using a square knot (Fig 5F,G). If using a loop,

be sure to tie it large enough for the fish to raise its dorsal fin and to grow over the course of the

experiment. However, a loop that is too large could get tangled in the tags of other fish or tank

elements (e.g., plastic plants). The knots can be tied by hand, using forceps, or a combination. If

the knots are challenging to tie, we find that loosely pre-tying and then untying a knot in the line

prior threading it through the fish can make this step much easier, as well as gentler for the fish.

Trim the excess fishing line using a razor blade. We recommend placing a drop of super glue on

the knot(s) to ensure they stay tied. As soon as possible, place the fish in a beaker of fresh

aquarium water to recover. The glue should dry sufficiently in less than a minute, and during this

time (and throughout), the fish's gills / body can be kept wet (with aquarium water or MS-222,

as needed) using a transfer pipette. The tag can be removed quickly, and without anesthesia, by

cutting the line with a sharp razor blade.

Open field exploration

A physical tag may be more likely to affect locomotion and behavior, particularly in the

smallest juveniles. We used an open field exploration to compare between juveniles that were

anesthetized and tagged (n=7) and those that were anesthetized but not tagged (n=8). This test is

commonly used across species to assess movement, activity, and anxiety-like behavior (Prut and

Belzung, 2003; Cachat et al., 2010; Solomon-Lane and Hofmann, 2019). We tagged the juveniles

less than 2 hrs after being removed from the mother's buccal cavity, and the open field test was

carried out the next day (~20 hrs later). The time it took to complete each tagging surgery was

recorded (average 257 ± 22 s), and each control fish was kept out of water and manipulated with

forceps for the same amount of time as one of the tagged fish. Additional MS-222 (or water) was

applied to the fish during the procedure, as needed.

For the open field exploration, we placed fish individually in a small, novel aquarium

(22.9 x 15.2 x 15.2 cm) without a cover for 30 min. For analysis, aquaria were divided into 4

zones: the territory zone, which contained a small terracotta pot territory / shelter, the close zone,

the far zone, and the investigate zone (Fig 6A). Video cameras (Warrior 4.0, Security Camera

Warehouse, Asheville, NC) recorded behavior from above, and we used BORIS (Friard and

Gamba, 2016) to score the number of times fish entered each zone of the aquarium, and how

long fish spent in each zone of the tank during a 10 min observation (minutes 20-30 of the test).

Statistics

To compare locomotion and behavior in the open field exploration between control and

tagged juveniles, we used a two-way mixed ANOVAs, with tagging treatment as the independent

measure and aquarium zone (frequency entering, time in) as the repeated measure. Tukey's HSD

test was used for *post hoc* analysis of significant ANOVA results. Eta-squared is reported to

estimate effect size for significant ANOVA results (small effect: $0 < \eta^2 < 0.01$; moderate: $0.01 <$

$\eta^2 < 0.06$; large: $0.06 < \eta^2$).

RESULTS

Open field exploration

We measured the time spent in each tank zone, as well as the frequency entering each

aquarium zone, for control and tagged juveniles. For the time in each zone, we found a

significant main effect of aquarium zone ($F_{(1.46, 13.13)}=13.44$, $p=0.001$), but no main effect of

treatment ($F_{(1,9)}=1.20$, $p=0.30$), and no interaction effect ($F_{(1.46, 13.12)}=1.35$, $p=0.28$). *Post*

hoc analysis showed that juveniles spent significantly less time in the investigate zone compared

to the territory ($p=0.00026$), close ($p<0.0001$), and far zones ($p=0.00029$). The other pairwise

comparisons were not significant (close-territory: $p=0.39$, far-territory: $p=1.00$, far-close: 0.31)

(Fig 6B). Similarly for the frequency of entering each zone, we found a significant main effect of

aquarium zone ($F_{(1.13, 10.2)}=16.79$, $p=0.002$), but no main effect of treatment ($F_{(1,9)}=1.90$,

$p=0.20$), and no interaction effect ($F_{(1.13, 10.2)}=0.48$, $p=0.53$). Juveniles spent significantly less

time in the investigate zone compared to the close zone ($p=0.0004$). The other pairwise

comparisons were not significant (close-territory: $p=0.23$, far-territory: $p=1.00$, investigate-

territory: $p=0.088$, far-close: $p=0.21$, investigate-far: $p=0.082$) (Fig 6C).

DISCUSSION

Our fishing line piercing method is a highly effective way of individually marking

juvenile fish that can last months. Across multiple experiments using older juveniles (e.g.,

(Solomon-Lane et al., 2022), all fish have survived the procedure and recovered very well. An

experienced researcher can also carry out these surgeries quickly, with anesthetization and

tagging taking ~2 min for the larger juveniles. All the tagged juveniles in this sample also

survived and recovered well; however, juveniles of this small size and young age are more

sensitive than older, larger fish. As needles and fishing line (or similar, e.g., 44-gauge stainless

steel wire, Remington Industries) continue to get thinner and lighter, we expect this method can

become even less invasive for very small fish. For a short experiment (< 1 week), line colored

with permanent marker remains vibrant under our aquarium conditions. Over a period of 3-4

weeks, the colors do fade. If super glue is added to the knot or to "beads" along the length of the

tag, the color under the glue remains visible for longer (at least 8 weeks). It may also be possible

to thread a seed bead onto the line for unique identification. Although we have not tested this yet,

seed beads are small, readily available in an array of colors, and inexpensive.

A physical tag may be more likely to affect locomotion and behavior than the tattoo

method. We tested fish in an open field exploration under conditions that we considered most

likely to reveal potential negative effects on behavior, while allowing juveniles time to recover

from the surgery. Juveniles were tagged less than 2 hrs after being removed from the mother's

buccal cavity, and at the time of the open field test, they had less than 24 hrs experience as

freely-swimming fish. Tagging did not have a significant effect on where juveniles spent time in

the aquarium, or how frequently they entered the different aquarium zones. Interestingly, tagged

juveniles entered the territory zone and close zone fewer times than control fish. Remaining in or

near the territory can be associated with shyness (Harmon et al., 2024), and injured fish may be

more likely to hide in the territory. However, the effect we saw was in the opposite direction.

This behavior pattern could indicate that locomotion decreases for a period of time after tagging,

and we plan to measure locomotion as a control in future studies. Depending on the experiment,

species, and developmental stage, a longer recovery period may be needed, and testing relevant

behaviors on subsequent days post-surgery will be valuable for determining appropriate

experimental timing. Overall, we were impressed by the quick recovery of these very small, very

young juveniles.

CONCLUSIONS

The tagging methods described here have made it possible for us to study the behavioral

development and underlying neuromolecular mechanisms of juvenile *A. burtoni* (e.g., Solomon-

Lane et al., 2022). The tattoo method can be used with the smallest, youngest juveniles and it has

multiple benefits: the procedure is fast, keeping the fish out of water for ~30 s, it uses the

smallest needle (pulled glass capillary tube), and the tag is fully internal. We have not yet tested

this method with larger juvenile or adult *A. burtoni*; however, once fish have sufficient dorsal

muscle, a visible implant elastomer tag or colored beads secured with a plastic tag (e.g., Trainor

and Hofmann, 2006) may become better options. The piercing method can be used with fish of

all sizes, including the smallest and youngest juveniles. The benefits of this method include

being longer-lasting than the tattoo method, and the tags can be visible on video for behavior

analysis (visibility will depend on the size of the tag and the camera system). Overall, the lack of

an appropriate method for identifying juveniles this small and young likely has contributed to the

relative paucity of early-life studies, in this and other small fish species. In social neuroscience,

behavioral ecology, animal behavior, developmental biology, fisheries research, and related

fields, experiments that follow individuals through development are critical to uncovering the

emergence of individual phenotypic variation, as well as the underlying mechanisms and fitness

consequences (Taborsky, 2016).

ACKNOWLEDGEMENTS

We thank Isaac Miller-Crews, Gabriel Rocha, Cameron Hamson, and Caitlyn Kwun for size

measurements of juvenile *A. burtoni*. We thank Dr. Sarah Budischak for use of her camera rig

and microscope for images of tagged fish. We thank members of the Solomon-Lane Lab for

assisting with resuscitation during the development of these methods, fish care, and pictures of

tagged fish, including Veronica Britton, Alex Kasiske, Jessica Gonzalez Rodriguez, Eliyah Stern,

and Andrew Yuan. Special thanks to Dr. Mel Coleman for suggesting the use of the glass

capillary tube needle for tattooing and leading us to minutien pins. We also thank members of

the Hofmann Lab and Solomon-Lane Lab for contributing to animal care and maintenance. This

work was supported by NSF grant IOS-1354942 to HAH, the BEACON Center for the Study of

Evolution in Action awards #947 (2016) and #1081 (2017, 2018) to TKSL and HAH, NSF grant

IOS-2341006 to TKSL, W.M. Keck Science Department funding to TKSL, and the Keck

Summer Research Fellowship to DDB.

LITERATURE CITED

Balzarini, V., Taborsky, M., Villa, F., Frommen, J.G., 2017. Computer animations of color

markings reveal the function of visual threat signals in *Neolamprologus pulcher*. Curr. Zool.

63, 45–54. https://doi.org/10.1093/cz/zow086

Campbell Grant, E.H., 2008. Visual Implant Elastomer Mark Retention through Metamorphosis

in Amphibian Larvae. Source J. Wildl. Manag. 72, 1247–1252.

https://doi.org/10.2193/2007-183

Fernald, R.D., Hirata, N.R., 1979. The Ontogeny of Social behavior and body coloration in the

African cichlid fish *Haplochromis burtoni*. Zeitschrif für Tierpsychologie, 50, 180-187.

https://doi.org/10.1111/j.1439-0310.1979.tb01025.x

Fernald, R.D., Maruska, K.P., 2012. Social information changes the brain. Proc. Natl. Acad. Sci.

109, 17194–17199. https://doi.org/10.1073/pnas.1202552109

Formica, V., Wood, C.W., Larsen, W.B., Butterfield, R.E., Augat, M.E., Hougen, H.Y., Brodie,

E.D., 2012. Fitness consequences of social network position in a wild population of forked

fungus beetles (Bolitotherus cornutus). J. Evol. Biol. 25, 130–137.

https://doi.org/10.1111/j.1420-9101.2011.02411.x

Frazier, K., 2015. The History of Bird Banding. Audubon (Website).

Hammer, S.A., Blankenship, H.L., 2001. Cost Comparison of Marks, Tags, and Mark-with-Tag

Combinations Used in Salmonid Research. N. Am. J. Aquac. 63, 171–178.

https://doi.org/10.1577/1548-8454(2001)063<0171:CCOMTA>2.0.CO;2

Hasler, A.D., Faber, W.M., 1941. A Tagging Method for Small Fish 1941, 162–165.

Hofmann, H.A., 2003. Functional genomics of neural and behavioral plasticity. J. Neurobiol. 54,

493   272–282. https://doi.org/10.1002/neu.10172

494 Jones, T.H., Godfray, H.C.J., Hassell, M.P., 2008. Relative Movement Patterns of a Tephritid

495   Fly and Its Parasitoid Wasps Published by: Springer in cooperation with International

496   Association for Ecology Stable URL : http://www.jstor.org/stable/4221264 Your use of the

497   JSTOR archive indicates your acceptan 106, 317–324.

498 Jørgensen, M.G.P., Deurs, M. van, Butts, I.A.E., Jørgensen, K., Behrens, J.W., 2017. PIT-

499   tagging method for small fishes: A case study using sandeel (*Ammodytes tobianus*). Fish.

500   Res. 193, 95–103. https://doi.org/10.1016/j.fishres.2017.04.002

501 Katona, S., Baxter, B., Brazier, O., Kraus, S., Perkins, J., Whitehead, H., 1979. Identification of

502   Humpback Whales by Fluke Photographs, in: Behavior of Marine Animals. Springer US,

503   Boston, MA, pp. 33–44. https://doi.org/10.1007/978-1-4684-2985-5_2

504 Kemp, D.J., Reznick, D.N., Grether, G.F., Endler, J.A., 2009. Predicting the direction of

505   ornament evolution in Trinidadian guppies (*Poecilia reticulata*). Proc. R. Soc. B Biol. Sci.

506   276, 4335–4343. https://doi.org/10.1098/rspb.2009.1226

507 Kohda, M., Jordan, L.A., Hotta, T., Kosaka, N., Karino, K., Tanaka, H., Taniyama, M.,

508   Takeyama, T., 2015. Facial recognition in a group-living cichlid fish. PLoS One 10, 1–12.

509   https://doi.org/10.1371/journal.pone.0142552

510 Kraus, B.T., McCallen, E.B., Williams, R.N., 2017. Evaluating the Survival of Translocated

511   Adult and Captive-reared, Juvenile Eastern Hellbenders (*Cryptobranchus alleganiensis*

512   *alleganiensis*). Herpetologica 73, 271–276.

513 Lewejohann, L., Hoppmann, A.M., Kegel, P., Kritzler, M., Krüger, A., Sachser, N., 2009.

514   Behavioral phenotyping of a murine model of Alzheimer's disease in a seminaturalistic

515   environment using RFID tracking. Behav. Res. Methods 41, 850–856.

https://doi.org/10.3758/BRM.41.3.850

Malone, J.C., Forrester, G.E., Steele, M.A., 1999. Effects of subcutaneous microtags on the

growth, survival, and vulnerability to predation of small reef fishes. J. Exp. Mar. Bio. Ecol.

237, 243–253. https://doi.org/10.1016/S0022-0981(99)00003-9

Ohayon, S., Avni, O., Taylor, A.L., Perona, P., Roian Egnor, S.E., 2013. Automated multi-day

tracking of marked mice for the analysis of social behaviour. J. Neurosci. Methods 219, 10–

19. https://doi.org/10.1016/j.jneumeth.2013.05.013

Reavis, R.H., Grober, M.S., 1999. An integrative approach to sex change: social, behavioural

and neurochemical changes in *Lythrypnus dalli* (Pisces) 2, 51–60.

Renn, S., Carleton, J.B., Magee, H., Nguyen, M.L.T., Tanner, A.C.W., 2009. Maternal care and

altered social phenotype in a recently collected stock of *Astatotilapia burtoni* cichlid fish.

Integr. Comp. Biol. 49, 660–673. https://doi.org/10.1093/icb/icp085

Taborsky, B., 2016. Opening the Black Box of Developmental Experiments: Behavioural

Mechanisms Underlying Long-Term Effects of Early Social Experience. Ethology 122,

267–283. https://doi.org/10.1111/eth.12473

Thresher, R.E., Gronell, A.M., 1978. Subcutaneous Tagging of Small Reef Fishes. Copeia,

1978(2), 352. https://doi.org/10.2307/1443582

Thompson, J.M., Hirethota, P.S., Eggold, B.T., 2005. A Comparison of Elastomer Marks and Fin

Clips as Marking Techniques for Walleye; A Comparison of Elastomer Marks and Fin Clips

as Marking Techniques for Walleye. North Am. J. Fish. Manag. 25, 308–315.

https://doi.org/10.1577/M03-237.1

Trainor, B.C., Hofmann, H.A., 2006. Somatostatin Regulates aggressive behavior in an African

cichlid fish. Endocrinology 147, 5119–5125. https://doi.org/10.1210/en.2006-0511

Volk, E.C., Schroder, S.L., Grimm, J.J., 1999. Otolith thermal marking. Fish. Res. 43, 205–219.

Weissbrod, A., Shapiro, A., Vasserman, G., Edry, L., Dayan, M., Yitzhaky, A., Hertzberg, L.,

Feinerman, O., Kimchi, T., 2013. Automated long-term tracking and social behavioural

phenotyping of animal colonies within a semi-natural environment.

https://doi.org/10.1038/ncomms3018

Williamson, C.M., Lee, W., Curley, J.P., 2016. Temporal dynamics of social hierarchy formation

and maintenance in male mice. Anim. Behav. 115, 259–272.

https://doi.org/10.1016/j.anbehav.2016.03.004

TABLES

Table 1: Treatment differences in anesthetization and recovery between juveniles that were

anesthetized (control) vs. juveniles that were anesthetized and tattooed. All juveniles remained

out of the water for 1 min, whether or not they were tagged.

Behavior	DF	Test Statistic	p-value	Effect size
Anesthetization in MS-222				
Loss of equilibrium	10.963	t = -0.35	0.73	
Stop moving	9.50	t = -1.07	0.31	
Stop breathing	**9.01**	**t = -2.42**	**0.039**	**-1.3**
Total time in MS-222	10.07	t = -1.388	0.20	
Recovery				
Initial breathing during resuscitation	12.13	t = 0.51	0.62	
Independent breathing	12.9	t = 0.11	0.91	
Body movement	13.47	t = 0.90	0.38	
Initial regained equilibrium	12.82	t = 0.10	0.92	
Sustained regained equilibrium	12.57	t = -0.52	0.61	
Full recovery	1	W = 18	0.28	

553   Results of pairwise comparisons using t-tests (t) or a Mann-Whitney Wilcoxon test (W). Cohen's

554   d is reported for effect size of significant results. Significant results are in bold.

555 **Table 2:** Associations between recovery metrics following anesthesia.

	Initial breathing during resuscitation	Independent breathing	Body movement	Initial regained equilibrium	Sustained regained equilibrium	Full recovery
Initial breathing during resuscitation		p=0.60	p=0.65	p=0.57	p=0.57	p=0.95
Independent breathing	p=0.59		**p=0.005, r^2=0.72**	**p=0.0036, r^2=0.81**	**p=0.0036, r^2=0.81**	p=0.14
Body movement	p=0.57	**p=0.011, r^2=0.63**		**p<0.0001, r^2=0.96**	**p<0.0001, r^2=0.96**	p=0.14
Initial regained equilibrium	p=0.12	**p=0.046, r^2=0.43**	**p=0.015, r^2=0.60**		**p<0.0001, r^2=1.00**	**p=0.034, r^2=0.55**
Sustained regained equilibrium	p=0.052	p=0.51	p=0.31	**p=0.0082, r^2=0.67**		**p=0.034, r^2=0.55**
Full recovery	p=1.00	p=0.22	p=0.37	p=0.92	p=0.95	

557 Correlations for control fish in the upper triangular (black text). Correlations for tattooed fish in

558 the lower triangular (grey text). Significant associations in bold. Adjusted r^2 values shown for

559 significant results.

560 **FIGURES**

Figure 1: Summary of appropriate tagging methods for *Astatotilapia burtoni* of different body sizes. A) Association between standard length (mm) and mass (g) (n=729). B) Image of 1-day old juvenile (8.8 mm standard length). C) Four methods used for tagging *A. burtoni* at different body sizes and life history stages, including the two methods described here: tattooing and a fishing line piercing through the dorsal muscle. The size upper ranges for the tagging methods are approximate and may vary depending on the experiment (e.g., short- vs. long-term, live vs. video behavior scoring) and the species. D) Image of a 0.1 mm diameter minutien pin (a: for piercing the smallest juveniles), fishing line (b: for tagging the smallest juveniles), pulled glass capillary tube needle (c: for tattooing), and a 31-gauge needle (d: for piercing larger juveniles). The red lines indicating diameter are the same diameter as the white circles in (E) (D and E are at the same scale). E) Cropped image of the fish in (B). The dorsal edge of the spinal cord is edged in dashed white lines. The dorsal edge of the fish is shown in the sold white line. White circles show the diameter of the minutien pin (a), fishing line (b), capillary tube needle (c), and 31-g needle (d) shown in (D).

Figure 2: Method 1: Tattooing: A) Image of the Nanoject setup under a dissection scope, with

a (plastic) fish placed on a wet napkin on the platform, prepped for injection. The injection

button is labeled on the Nanoject controller. **B)** A juvenile fish being injected with blue ink using

a pulled glass capillary tube needle. **C)** Fish immediately after injection with tattoo ink. The

injection site is shown on the left, with the dye spreading on the right side of the fish. **D)** A

juvenile fish 2 days after injection. **E)** The same juvenile fish as in (D, at the same scale) 36 days

after injection. See Supplemental Information for pictures of additional tag colors.

Figure 3: Recovery from anesthesia following tattooing: Time until A) breathing

independently and B) full recovery from anesthetization in control and tagged juveniles. C)

Association between initial body movement and independent breathing. D) Association between

initial regained equilibrium and full recovery. All recovery times start when the fish was placed

in the recovery beaker, following anesthetization and 1 min out of water. Tagged fish were

injected with dye during the 1 min out of water.

592 **Figure 4: Making the fishing line tag.** A) Thread the fishing line through the needle of an

593 insulin syringe until it is visible in the barrel of the syringe. B) Use forceps to grasp the base of

594 the needle. Rock back and forth (arrows) until the needle breaks off from the syringe. Not

595 shown: From the blunt side of the needle, pull the line through, until it no longer sticks out

596 beyond the needle tip. Use a razor blade to cut off excess line.

Figure 5: Tagging with a fishing line piercing. A) A juvenile tagged with a blue fishing line loop 1 hr after being removed from the mother's buccal cavity (8.88 mm standard length, SL). B) A young juvenile (4 days post-removal from the mother's buccal cavity) tagged with a blue fishing line barbell (10.08 mm SL). C) An older juvenile tagged with a fishing line loop (10.5 mm SL) on a ruler (inches top, mm bottom). Procedure for tagging larger juveniles: D) Pierce the dorsal muscle with the needle. E) Pull the needle and line through the dorsal muscle. F) Make a loop in the line and tie the first half of a square knot. Adjust the loop to the desired size. G) Finish the square knot. Cut the excess line using a razor blade. The same individual fish is pictured in panels C-G.

Figure 6: Open field exploration. A) Tank layout of the open field exploration, with the

territory, close, far, and investigate zones. The territory zone has a small terracotta pot territory /

shelter. **B)** Time spent (s) in each tank zone. **C)** Frequency entering each tank zone. Tagged fish

(n=7) were anesthetized and tagged with a fishing line piercing less than 2 hrs after being

removed from the mother's buccal cavity. Control fish (n=8) were anesthetized but not tagged.

The behavior test took place ~20 hrs post-surgery. Different letters indicate significant

differences.