Excitatory-inhibitory windows shape coherent neuronal dynamics driven by optogenetic stimulation in the primate brain

Ryan A. Shewcraft[1], Heather L. Dean[1], Margaret M. Fabiszak[2], Maureen A. Hagan[3,4], Yan T. Wong[3,4,5], and Bijan Pesaran[1,6,7,*]

Lead Contact:
Bijan Pesaran, Ph.D.
4 Washington Pl. Rm 809
Center for Neural Science
New York University
New York, NY 10003 USA
Tel: 212.998.3578, Fax 212.995.4011
E-mail: bijan@nyu.edu

Word counts

Summary	144	(<150)
Introduction	612	
Results	2093	
Discussion	1529	
Methods	2942	
References	58	
Figures	6	
Tables	0	
Total words	4234	

[1] Center for Neural Science; New York University; New York, NY, 10003; United States

[2] Laboratory of Neural Systems; The Rockefeller University; New York, NY, 10065; United States

[3] Biomedicine Discovery Institute; Monash University; Clayton, VIC 3800; Australia

[4] Australian Research Council, Centre of Excellence for Integrative Brain Function; Monash University; Clayton, VIC 3800; Australia

[5] Department of Electrical and Computer Systems Engineering; Monash University; Clayton, VIC 3800; Australia

Summary

Coherent neuronal dynamics play an important role in complex cognitive functions. Optogenetic stimulation promises to provide new ways to test the functional significance of coherent neural activity. However, the mechanisms by which optogenetic stimulation drives coherent dynamics remain unclear, especially in the non-human primate brain. Here, we perform computational modeling and experiments to study the mechanisms of optogenetic-stimulation-driven coherent neuronal dynamics in non-human primates. Neural responses arise from stimulation-evoked temporal windows of excitatory and inhibitory activity. The temporal properties of the E-I windows generate coherent neuronal dynamics at varied frequencies and depend on optogenetic stimulation parameters. Experimental results agree with parameter dependent predictions from computational models. These results demonstrate that responses to optogenetic stimulation are governed by local circuit properties that alter the timing of E-I activity. Temporary imbalances in excitatory and inhibitory activity may provide a general mechanism for generating coherent neuronal dynamics.

Introduction

Spiking and local field potential (LFP) activity display coherent patterns of activity across a range of temporal frequencies (Pesaran et al., 2018). Coherent temporal structure has been proposed to play an important role in the control of higher cognitive functions (Buschman and Miller, 2007; Dean et al., 2012; Womelsdorf et al., 2006; Wong et al., 2016) and enhanced cortico-cortical communication more generally (Fries, 2005; Pesaran et al., 2018, 2008). Despite this, whether and how complex cognitive functions depend on the temporal structure of neuronal activity, as opposed to the magnitude of the activity, has been highly debated. Causal manipulations that generate frequency selective coherent neuronal dynamics are important for directly testing their functional significance. However, in order to interpret the results of causal manipulations we also need to understand how extrinsic modulations of coherent neuronal dynamics recruit intrinsic cellular and network mechanisms to generate the driven response.

In general, coherent neuronal dynamics depend on the temporal precision and synchrony of spike timing (Buzsáki and Wang, 2012), which is governed by the intrinsic dynamics of excitatory (E) and inhibitory (I) post-synaptic potentials (Brunel and Wang, 2003; Wehr and Zador, 2003). Typically, excitatory and inhibitory activity are tightly correlated, with E preceding I by several milliseconds (Okun and Lampl, 2008). Thus, inputs which drive excitatory activity are closely followed by inhibitory activity, forming a brief a window for temporally precise spiking in the postsynaptic neurons (Gabernet et al., 2005; Haider et al., 2013; Salkoff et al., 2015). The instantaneous frequency of coherent neuronal activity is regulated by E-I balance (Atallah and Scanziani, 2009) and depends on the relative timing of E and I activity (Chariker et al., 2018; Keeley et al., 2017). Consistent with this, empirically observed neural coherence is dynamic and episodic in the gamma (Burns et al., 2011, 2010; Xing et al., 2012) and beta (Feingold et al., 2015; Rule et al., 2017; Sherman et al., 2016) bands.

Optogenetic stimulation has been shown to alter neuronal dynamics extrinsically by

manipulating either excitatory or inhibitory activity alone. In some studies, driving inhibitory

activity by delivering pulse trains of optogenetic stimulation to interneurons perturbed coherent

neuronal activity (Cardin et al., 2009; Sohal et al., 2009). In other studies, driving excitatory

activity by stimulating pyramidal neurons using continuous optogenetic stimulation generated

coherent dynamics, specifically gamma activity (Lu et al., 2015; Ni et al., 2016). However,

coherent neuronal dynamics depend on the interaction between excitatory and inhibitory

activity, which is governed by the intrinsic properties of the local circuit. Since stimulating either

excitatory or inhibitory activity alone recruits both excitatory and inhibitory postsynaptic

responses in the network (Buzsáki and Wang, 2012; Gloveli et al., 2005; Phillips and

Hasenstaub, 2016; Pike et al., 2000; Tukker et al., 2007), optogenetic stimulation may generate

coherent neuronal dynamics according to the relative timing of stimulation-driven E-I responses.

Motivated by this reasoning, we tested the hypothesis that optogenetic stimulation drives

coherent neuronal dynamics by generating E-I responses that arise from intrinsic E-I balance.

We first developed a computational model of coherent neuronal dynamics in response to

optogenetic stimulation in terms of a temporal window of excitatory and inhibitory activity—the

E-I window. We experimentally tested the model predictions by expressing ChR2(H134R) in

macaque cortex delivered by an AAV2/5 viral injection and measuring spiking and local field

potential (LFP) responses to sequences of optogenetic stimulation pulses. Finally, we

developed a spiking neural network model of the observed neural dynamics and found that E-I

windows are an emergent property of the model. Together these results suggest that

optogenetic stimulation extrinsically alters excitatory and inhibitory activity by recruiting intrinsic

E-I windows to shape coherent neuronal dynamics in the beta and gamma frequency bands.

Results

Modeling optogenetic-stimulation-evoked coherent neuronal dynamics.

Coherent neuronal dynamics arise from synchronous, temporally structured neuronal activity. Spike-field coherence (SFC) measures coherent neuronal dynamics by estimating the strength of correlations between spiking activity of individual neurons and neural population activity in the form of simultaneously-recorded LFP activity. When the spike times of a neuron can be predicted from fluctuations in LFP activity at a particular frequency, the spiking is said to be coherent with LFP activity at that frequency. Neurons that fire consistently at a particular phase of LFP activity are coherent with LFP activity. As a result, neuronal coherence is defined by coherent spiking and LFP activity and can be measured by estimating SFC between spiking and nearby LFP activity (Pesaran et al., 2018, 2002).

Figure 1A presents a model of optogenetic-stimulation-evoked coherent neuronal dynamics and corresponding idealized neural responses. During spontaneous activity, synaptic inputs drive activity in a local network. We model spontaneous spiking activity as being generated from a Poisson process with a constant rate (Brown et al., 2003) and spontaneous LFP activity as being generated from a Brown noise process. Under these assumptions, spontaneous spiking is not coherent with spontaneous LFP activity (**Figure 1A**, Spontaneous activity)**.**

Direct activation of transduced neurons leads to excitatory and inhibitory synaptic currents within the local network of neurons. The resulting excitatory (E) and inhibitory (I) currents in postsynaptic neurons occur at different moments in time and the relative timing of the E and I currents determines when neurons fire spikes (Cruikshank et al., 2010; Haider et al., 2013; Higley and Contreras, 2003; Wehr and Zador, 2003). Given that excitatory currents depolarize the membrane potential and inhibitory currents hyperpolarize the membrane

potential, the cell has the greatest probability of firing when E is high and I is low. Therefore, we

model the impact of optogenetic stimulation on spiking as an excitatory- and inhibitory-

dependent temporal change in the probability of the cell firing (**Figure 1A,** Spike generation).

Thus, the timing of spiking activity depends on the net postsynaptic E and I consequences of

each stimulation pulse - the E-I window.

To simulate neuronal activity that exhibits SFC, we need to model the coupling between

spiking and LFP activity. LFP activity predominantly reflects summed postsynaptic potentials

(Buzsáki et al., 2012; Pesaran et al., 2018). Since the E and I postsynaptic activity drives

spiking, we can model the driven LFP as the sum of the driven E and I responses, thus linking

spiking and LFP activity,. We add brown noise background activity to the driven LFP activity to

generate the simulated LFP (**Figure 1A**, LFP).

Figure 1. Excitation and inhibition drive coherent neuronal dynamics. (A) To model
optogenetic stimulation of a population of transduced neurons, we simulated pulsatile responses

We refer to this model of neuronal coherence as the **E-I window model.** We model

spontaneous spiking and LFP as uncorrelated so the spontaneous activity does not exhibit SFC

(**Figure 1B**). Since optogenetic stimulation evokes E-I windows in which action potentials are

more likely to occur, changes in spike probability are coupled to LFP responses. This coupling

results in spikes that are coherent with LFP activity (**Figure 1C**). Importantly, the E-I window

model predicts that the coherence is due to the temporal dynamics, or shape, of the E-I

windows. In line with the prediction, SFC varied with the E-I window duration such that longer E-

I windows generated SFC at lower frequencies (**Figure 1D**). In our empirically measured E-I

windows, excitatory responses dominated the early stage followed by inhibitory responses. This

is consistent with sensory stimulation which typically generates an increase in excitation

followed shortly by a corresponding inhibitory response (Higley and Contreras, 2003; Wehr and

Zador, 2003). However, inhibition may be stronger and have faster rise times than excitation in

certain circuits or behaviors (Cruikshank et al., 2010; Haider et al., 2013). This regime would be

captured in the E-I window model by setting $\alpha_{inh} > \alpha_{exc}$.

Note that the neuronal coherence described by the E-I window model depends on

predictability between spikes and LFP activity. This construction is consistent with, but does not

require, invoking an oscillator or other phase-consistent process. This construction is also

consistent with the use of SFC, as opposed to other metrics of phase consistency, to estimate

coherent neuronal dynamics. This is because SFC measures linear predictability between

spikes and LFP activity. Note also that our approach contrasts with and is complementary to

previous work that has used cell-type-specific optogenetic stimulation to generate neuronal

coherence.

Testing the E-I window model with optogenetic stimulation.

Next, we tested whether optogenetic stimulation generates empirically observed neural coherence according to the predictions of the E-I window model . The model suggests that the frequency of driven coherent neuronal dynamics depends on the E-I window duration. Therefore, varying the duration of the stimulation pulse should change the E-I window duration, and the frequency content of the resulting driven coherent neuronal activity.

To test this hypothesis, we injected AAV5-hSyn-ChR2(H134R)-EYFP and recorded neuronal responses in three alert macaque monkeys quietly seated in a primate chair. We recorded in the inferior and superior parietal lobules in Monkeys H and J and lateral to the post-central dimple in Monkey B (**Figure 2A**)**.** We observed strong viral expression at injection sites with robust labelling of neurons and neuropil (**Figure 2B**). We performed simultaneous *in vivo* optogenetic stimulation and recording using a fiber optic cable placed on a surgically thinned dura and adjacent to the intracortical recording electrode (**Figure 2C**). Stimulation consisted of one second long sequences of light pulses (stimulation epoch) followed by 1-3 s of no stimulation (spontaneous epoch) (**Figure 2D**). Optogenetic stimulation reliably generated neural responses. These responses consisted of action potentials as well as large evoked field potentials visible on an individual pulse basis in the raw data (**Figure 2E,F,G).**

We measured neuronal responses to stimulation sequences composed of Poisson pulse trains with different pulse rates and pulse durations. **Figure 3A** presents example neuronal responses during a stimulation block with 5 ms wide pulses delivered with a 20 pulse/s Poisson pulse train. Individual stimulation pulses drove spiking activity and evoked LFP responses at the stimulation site (**Figure 3Ai,ii**). We estimated the E-I windows by fitting the the E-I window to the empirically measured evoked LFP (**Figure 3Aiii**). We used this fitted E-I window in the E-I window model to simulate coherent neuronal dynamics. The simulated SFC closely matched the empirically measured SFC (**Figure 3Aiv**). This suggests that the E-I window model captures the phenomenology of the coherent neuronal dynamics driven by optogenetic stimulation. In addition, we computed the power spectrum of LFP activity during the one second long stimulation epoch and for one second of the spontaneous epoch (**Figure 3Av**). The stimulation LFP power spectrum revealed a significant increase in power (Chi-squared test, p<0.01).

Figure 3. Neuronal dynamics driven by optogenetic stimulation. (A) Neuronal responses to 5 ms pulse duration, 20 Hz pulse rate, Possion pulse train. (i) Raster (top) and PSTH (bottom) of the response to single light pulses (blue). (ii) Mean evoked potential for all the stimulation pulses that were the last in a sequence and not preceded by a prior stimulation pulse within 50 ms, normalized by subtracting the value of the LFP at the onset of the light. Dotted line indicates 0 mV. (iii) Modelled E-I window (black) fit to the empirically measured E-I window (grey dotted) with excitatory (blue) and inhibitory (red) components. The fitted E-I window parameters are A=0.097, α_{exc}=1, α_{inh}=3, β_{exc}=2.568, β_{inh}=3.654.(iv) Spike-field coherence for multi-unit activity and LFP measured on the same electrode. Dotted horizontal line shows expected coherence under the null hypothesis that coherence is zero. (v) Power spectral density (PSD) of 1 s of data during the stimulation epoch (black line) and the spontaneous epoch (grey line). The dotted lines are error bars determined by the Chi-squared test for p=0.01. Solid gray lines above the PSD illustrate frequencies where the driven PSD was significantly different from the spontaneous PSD (p<0.01). **(B)** Same as (A) for 20 ms, 20 pulses/s pulse rate, Poisson pulse train. The fitted E-I window parameters for (Biii) are A=0.097, α_{exc}=2, α_{inh}=4, β_{exc}=4.548, β_{inh}=4.730. **(C)** Comparison of the peak frequency of the driven empirical and simulated SFC for all pulse widths conditions and all units. **(D)** Peak frequency for the driven SFC for each unit for the empirical data (grey) and E-I window model simulation (black). Horizontal lines show the mean for each condition.

Longer pulse durations generated longer duration changes in firing rate (**Figure 3Bi**) and longer periods of excitation and inhibition in the LFP activity (**Figure 3Bii,iii**), and have more selective changes in SFC and the LFP power spectrum (**Figure 3Bivv**). This shows that, stimulation sequences with different pulse durations can be used to selectively generate coherent neuronal dynamics, consistent with changing the E-I window duration. Thus, the E-I

window model predicted the relationship between the E-I windows and the frequency content of

the stimulation-evoked coherent neuronal dynamics.

Optogenetic stimulation generates pulse duration-dependent responses.

We computed the SFC for the population of recording sites with driven spiking activity in

response to stimulation using Poisson pulse trains with 5, 10, 15, and 20 ms duration pulses.

For each recording condition, we fit the E-I window as in Figures 3A and 3B and used the E-I

window model to generate simulated SFC. For all recordings, the simulated SFC closely

matched the empirical SFC (**Figure 3C**). In addition, for both the empirical and simulated data,

the frequency of the driven SFC varied with pulse duration (**Figure 3D**), consistent with the

example site and predictions from the E-I window model.

Importantly the center frequency did not depend on the mean pulse rate for Poisson

stimulation (**Figure 3—figure supplement 1**). The decoupling of frequency from pulse rate

means that the amount of coherence can be controlled separately from the center frequency.

These results show that coherent neuronal dynamics in either beta or gamma frequency bands

can be generated by stimulating the same population of neurons using stimulation pulses of

different durations.

241

242   **Figure 3—figure supplement 1. The response dynamics do not depend on pulse rate.**
243   Average and SEM of the standard deviation of the first fitted Gaussian for all stimulus sequences
244   with a 10 ms pulse width across all recording sites.

Model-based investigation of excitatory and inhibitory dynamics.

Since we did not have direct experimental access to excitatory and inhibitory currents, we developed a spiking neural network model to study how the frequency content of the driven coherent neuronal dynamics depended on the timing of the E and I activity. We implemented a computational model that consisted of a randomly-connected network of leaky integrate-and-fire excitatory and inhibitory neurons that captures LFP responses to sensory stimulation (Mazzoni et al., 2008). A subset of neurons also expressed ChR2 ion channels (Witt et al., 2013) (**Figure 4A,B**).

The model generated spiking activity and subthreshold potentials in response to optogenetic stimulation (**Figure 4C,D**). To test whether the model responses depended on stimulation pulse duration, we simulated data with the same pulse parameters as in **Figure 3** (5-20 ms pulse widths; 20 pulse/s Poisson). The model responses exhibit pulse duration dependence (**Figure 4E**) and frequency selectivity that varies with pulse duration (**Figure 4F**). Consistent with the empirical results, longer pulses durations resulted in a decrease in the center frequency of the driven coherent neuronal dynamics (**Figure 4F**). Therefore, the spiking model captures the key features of the empirically observed optogenetically-driven coherent neuronal dynamics. As with the empirical data and E-I window model, the frequency selectivity of the driven SFC is pulse-duration-dependent.

Figure 4. Spiking model responses and dynamics. (A) Diagram of a leaky integrate-and-fire model network with ChR2-transduced neurons. **(B)** Circuit diagram of leaky integrate-and-fire neurons with ChR2 channels that make up the spiking network model. **(C)** Raster (top) and PSTH (bottom) of the spiking model response to single light pulses for 10 ms, 10 pulses/s, Poisson stimulation. **(D)** Example broadband spiking model simulated LFP activity during a pulse sequence (10 ms, 10 pulses/s Poisson stimulation; blue lines indicate pulse times). *Inset*: Broadband simulated LFP activity at the time of a single pulse. **(E)** Mean normalized evoked potentials for the spiking model during pulse trains with 5 ms, 10 ms and 15 ms pulse durations and 10 pulses/s, Poisson stimulation. The dotted light blue line indicates the onset of the light stimulation and the subsequent dotted lines show the time when the light turned off. **(F)** Peak frequency of spiking model simulated SFC for each pulse width.

The spiking model responses depended on both channel and connectivity parameters.

The closing time constant of the ChR2 channel, τ_{OFF}, governed the duration of the response.

For longer τ_{OFF}, as in Witt *et al.* (Witt et al., 2013), the duration of the evoked spiking and LFP responses in our spiking network model extended well beyond the end of the light pulse. This was because the ChR2 channels slowly closed after the light turned off, allowing excitatory currents to continue for an extended period of time. To account for the differences in ChR2 and ChR2(H134R) (Lin et al., 2009) channel kinetics and their temperature-dependence (Feldbauer

et al., 2009) we fit an exponential model to temperature-dependent channel closing times and

computed the *in vivo* closing time constant for ChR2(H134R) as approximately 3 ms (**Figure**

4—Figure Supplement 1). Setting τ_{OFF} to 3 ms generated a response that did not have

responses that extended well beyond the offset of the light pulse (**Figure 4C-E**).

Figure 4—figure supplement 1. Fitting ChR2 channel kinetics to *in vivo* temperatures.
Approximate ChR2 channel kinetics adapted from Feldbauer et al. with an exponential fit and
extrapolation to macaque body temp (about 37^0 F).

Driven coherent neuronal dynamics depend on timing of excitatory and inhibitory

responses.

The spiking model allows us to study the dynamics of intracellular currents during

stimulation. We estimated the mean driven excitatory and inhibitory currents in the neurons in

the model by taking the stimulation pulse triggered average of AMPA-mediated currents,

$I_A = \dfrac{V_A C_m}{\tau_m}$ and GABA-mediated currents, $I_G = \dfrac{V_G C_m}{\tau_m}$, respectively (**Figure 5A**). As the

stimulation pulse duration increases, the driven excitatory and inhibitory currents persist for a

longer period of time (**Figure 5A**). It is important to note that the E-I windows were not explicitly

built into the model, but rather emerged as responses to simulated optogenetic stimulation. As

we increased stimulation pulse duration, the relative timing of both the onset and offset of the

excitatory and inhibitory currents remained the same, with excitatory currents leading inhibitory

currents (**Figures 5B,C**). Thus, the frequency content of the driven coherent neuronal dynamics

depends on the duration of the E and I responses.

Figure 5. The dynamics of E and I responses varies with pulse duration. **(A)** Stimulation pulse-
triggered excitatory (blue), inhibitory (red) and combined E-I responses in neurons in the spiking
network model in response to 5, 10, 15 and 20 ms duration simulation pulses. Examples of single
pulse evoked spiking are shown above. **(B)** Onset of normalized excitatory (solid) and inhibitory
(dashed) responses for 5,10,15 and 20 ms pulse durations. Onset delay is the difference between
the time to half maximal amplitude for excitatory and inhibitory responses. **(C)** Offset for normalized
excitatory (solid) and inhibitory (dashed) responses for different pulse durations. Offset delay is the
difference between the time to half maximal amplitude from the peak of the onset response for
excitatory and inhibitory responses.

315   Previous work has suggested that balanced inhibition is responsible for the

temporal precision of spiking in response to sensory stimulation (Wehr and Zador, 2003). In our

spiking network model, E-I balance arises from correlations between excitatory synaptic

currents and inhibitory synaptic currents during both spontaneous activity and optogenetic

stimulation and is thus an intrinsic property of the circuit (**Figures 6A-C**). Consistent with

previous experimental results, we find that inhibition lags excitation in the spiking model.

Interestingly, the lag decreases during optogenetic stimulation (2.6 ms during spontaneous

activity and 1.8 ms during stimulation) (**Figure 6B**) and coherence between E and I activity

increases (**Figure 6D**) so that the inhibitory activity more closely tracks the excitatory activity.

The spiking model shows that optogenetic stimulation drives excitatory postsynaptic

currents in neurons, which increases their probability of spiking. However, due to the E-I

balance inherent in the local network, the excitatory currents are followed closely in time by a

balanced amount of inhibitory postsynaptic currents, which suppresses the increased tendency

to fire. Therefore, E-I balance, which increases during stimulation, may be responsible for the

temporal precision of the driven spiking and thus may be a mechanism for stimulation driven

coherent neuronal dynamics.

Figure 6: Balanced excitatory and inhibitory responses. (A) Example excitatory (blue) and
inhibitory (red) synaptic currents in the spiking network model during spontaneous activity, (left),
and stimulation,(right). Currents are normalized to show overlap. **(B)** Cross-correlation between E
and I synaptic currents during spontaneous activity and the 5 ms stimulation condition. Dotted lines
show time lag for maximal cross correlation. **(C)** Instantaneous excitatory and synaptic currents for
the 5 ms pulse width condition measured in 0.2 ms intervals. Spontaneous: $\rho = 0.82$, $p = 0$;
Stimulation: $\rho = 0.73$, $p = 0$. **(D)** Coherence between excitatory and inhibitory currents during
spontaneous activity (dotted line) and stimulation using 5 ms, 10 ms and 15 ms pulse durations.

Discussion

Here, we investigate the mechanisms by which optogenetic stimulation of non-human primate (NHP) cortical neurons evokes coherent neuronal dynamics. We show that optogenetically-driven neuronal dynamics arise from how extrinsic inputs recruit excitatory and inhibitory responses that follow intrinsic circuit properties, leading to an effective temporal window for spiking – the E-I window. We develop a phenomenological model to predict how the frequency of optogenetic-stimulation-evoked coherence varies parametrically with stimulation parameters according to the timing of excitatory and inhibitory currents generated by each stimulation pulse. Experimental results from optogenetically-stimulating NHP cortex confirmed the model predictions. Furthermore, we developed a spiking neural network model of the empirical coherent neuronal dynamics and observed emergent stimulation-driven E-I windows. These results show that varying the duration of optogenetic stimulation pulses can selectively generate coherent dynamics across a range of behaviorally relevant frequencies—beta and gamma—by varying the effective E-I window.

E-I window mechanisms of driven coherence

Neurons fire spikes more often during periods of excitatory and inhibitory imbalance, called integration windows. Synchronous excitatory synaptic inputs depolarize cellular membrane potentials and drive spiking activity. Inhibition dominates cortical responses to stimulation (Haider et al., 2013) and regulates the temporal properties of integration windows through temporally lagged responses which balance excitatory inputs and repolarize membrane potentials (Gabernet et al., 2005). Spontaneous excitatory and inhibitory activity are correlated, with inhibition lagging excitation, suggesting that E-I balance may arise from the properties of the neural circuit (Okun and Lampl, 2008). Therefore, both time-lagged, correlated excitatory

and inhibitory responses may play a key role in spike timing precision in response to

optogenetic stimulation, thus generating coherent neuronal dynamics.

The E-I window model is phenomenological and offers a mechanism for how excitatory

and inhibitory activity affect spike timing and driven coherent neuronal dynamics. To gain further

mechanistic insight into the consequences of imbalances in excitatory and inhibitory activity, we

developed a spiking network model. Experimental agreement with the E-I window model

predictions reinforces the idea that driven coherent neuronal dynamics depends on a

mechanism of spike timing precision involving a temporary increase in excitatory activity that

leads to an increase in spiking. This is then followed by an increase in inhibitory activity that

returns the probability of spiking to baseline. Furthermore, pulse-duration dependent E-I

windows were an emergent property of a spiking network model of driven coherent neuronal

dynamics. Together, these results suggest that the generation of E-I windows by optogenetic

stimulation precisely times spikes to generate coherent neural activity

The E-I window model may offer a more general mechanism of how spike-field

coherence arises endogenously, in addition to being driven by exogenous inputs. For example,

sensory stimulation drives E-I responses which govern spike timing (Wehr and Zador, 2003).

Since LFP activity arises from synchronous neuronal activity that occurs due to temporally

structured fluctuations of local excitation and inhibition (Buzsáki and Wang, 2012; Csicsvari et

al., 2003; Hasenstaub et al., 2005) the LFP activity measures measures summed local

postsynaptic potentials (Buzsáki and Wang, 2012; Pesaran et al., 2018), Thus the intrinsically

generated E-I windows are likely reflected in the LFP.

Since SFC depends on coupling LFP activity to spiking. The E-I window contributes to

both the spike-timing and the LFP activity by making use of an LFP-based conditional intensity

function, coupling the measurements and resulting in coherent neuronal dynamics. Moreover,

the stimulation-evoked LFPs reveal the temporal structure of stimulation-induced postsynaptic

potentials (Okun et al., 2010; Poulet and Petersen, 2008). The agreement between the E-I

model and experimental results shows that stimulation-evoked LFP activity is effective in

capturing the stimulation-driven E-I balance, and thus the E-I window may be a general

mechanism for intrinsic coherent neuronal dynamics

Relation to cell-type specific manipulations

Previous work suggests that controlling activity in a single cell-type is necessary for

driving coherence (Adesnik and Scanziani, 2010; Cardin et al., 2009; Lu et al., 2015; Sohal et

al., 2009) However, cell-type-specific stimulation recruits synaptically-mediated responses from

other strongly connected neurons in the network (Phillips and Hasenstaub, 2016). The

frequency content of coherent neuronal dynamics is shaped by the recruited neuronal ensemble

(Kato et al., 2017). This suggests that neuronal coherence evoked by cell-type-specific

optogenetic stimulation may also drive an effective E-I window. Our work shows that pan-

neuronal optogenetic stimulation generates E-I windows that drive coherent neuronal dynamics.

Whether the E-I window model is as effective in predicting the impact of cell-type-specific

manipulations that target either E or I cells alone is an open question and will likely depend on

the strength of recurrent connectivity in the network.

Local and long-range mechanisms of coherent neuronal dynamics

Models of coherent neuronal dynamics have shown that oscillatory dynamics can arise

from the timing of interactions between excitatory and inhibitory populations of neurons (Börgers

and Kopell, 2003; Brunel and Wang, 2003; Keeley et al., 2019; Wilson and Cowan, 1972)..

Thus, these local mechanisms are consistent with the E-I window model of driven coherent

neuronal dynamics. In the beta band, coherent neuronal dynamics may be generated locally

(Jensen et al., 2005; Roopun et al., 2006) or from remote sources (Holgado et al., 2010; Lee et

al., 2013; Sherman et al., 2016). The E-I window model suggests that optogenetic stimulation

drives coherent neuronal dynamics through local, intrinsic mechanisms. This has important

implications for interpretations that seek to dissociate between local and long-range beta

mechanisms. If optogenetically modulating beta band activity affects behavior, we can interpret

that changing locally generated beta band activity at the stimulation site is sufficient for altering

behavior.

Non-oscillatory dynamics

Many computational models that offer insights into the mechanisms of coherent neuronal

dynamics have focused on simulating periodic, oscillatory activity (Brunel and Wang, 2003;

Wilson and Cowan, 1972). However, coherent neuronal activity is not simply oscillatory neural

activity, and only requires the presence of statistically significant dependencies at specific

frequency bands. Empirically observed neuronal coherence is episodic (Burns et al., 2011,

2010; Feingold et al., 2015; Rule et al., 2017; Sherman et al., 2016; Xing et al., 2012), with

phase and frequency varying over time, and can be narrow-band or broad-band. The E-I

window model expands on these prior models to describe manipulations of coherent neural

activity patterns which are not necessarily oscillatory or narrow-band. Since the E-I window

model also relies on the dynamics of intrinsic excitatory and inhibitory activity to simulate

coherent neuronal dynamics, the presence of oscillatory or episodic dynamics may be explained

by the mechanism of interactions between extrinsic inputs and intrinsic circuit properties.

Poisson stimulation for studying dynamics

Continuous and periodic pulse optogenetic stimulation sequences have previously been

used generate beta and gamma activity and study their effects on behavior (Adesnik and

Scanziani, 2010; Cardin et al., 2009; Ni et al., 2016; Sohal et al., 2009). These forms of

stimulation implicitly control the E-I window using the properties of the stimulation pulse and

stimulation sequence. However, using continuous constant and pulsatile periodic stimulation

drives responses at only particular frequencies (Adesnik and Scanziani, 2010; Cardin et al.,

2009; Lu et al., 2015), likely related to properties of the local network (Buzsáki and Wang, 2012)

and thus may vary across brain regions and subjects (Lu et al., 2015; Ni et al., 2016). The effect

of Poisson stimulation depends on the temporal properties of the stimulation-generated E-I

windows rather than the pulse sequence parameters. Thus Poisson stimulation can generate

coherent activity across a broader range of behaviorally-relevant frequencies by varying the E-I

windows using different stimulation pulse durations.

Potential bias in estimates of coherent neuronal dynamics from LFP activity

Extracellular recordings of LFP activity and spiking may be biased toward activity in

pyramidal neurons. LFP activity is most sensitive to postsynaptic potentials in neurons with

separated current sources that generate an open field geometry (Pesaran et al., 2018). Thus,

contributions from neurons with extended dendritic architectures, such as pyramidal neurons,

are more heavily weighted by LFP activity. Additionally, pyramidal neurons typically have larger

cell bodies than other cortical neurons and thus have more easily measurable action potentials.

Therefore, our measurement of both spiking activity and LFP, and thus SFC, may be biased

toward contributions from pyramidal neurons. In addition to LFP activity, alternative

decompositions of neural data, such as projections of population activity onto lower dimensional

subspaces, may be used to estimate coherence. Such estimates may be more effective at

identifying spiking that is coherent with pre- and postsynaptic activity in neuronal ensembles

containing both open and closed field neurons.

In conclusion, we systematically modeled neuronal responses to optogenetic stimulation

in NHP cortex. We highlight the role of generating specific E-I windows by selecting stimulation

parameters—pulse duration—to selectively perturb coherent neuronal activity. Importantly, we

do so in a general manner without necessarily targeting optogenetic expression to specific types

of neurons. In general, our results show that combining experimental measures with model-

based predictions can be used to study how optogenetic stimulation drives network dynamics

across a wide range of experimental designs, including using different brain regions, model

species, and stimulation protocols. These results demonstrate that optogenetic stimulation

generates dynamics in behaviorally-relevant frequency ranges in the primate brain by altering

excitatory and inhibitory activity in a manner similar to sensory stimulation. Therefore,

optogenetic control of neuronal activity can be useful for understanding how coherent neuronal

dynamics give rise to complex, flexible behaviors.

Methods:

Animals

Recordings were taken from two male rhesus macaques (*Macaca mulatta*) (Monkey J: 12 years, 6.5 kg; Monkey H: 11 years, 10.5 kg) and one male cynomolgus macaque (*Macaca fascicularis*) (Monkey B: 4 years, 5 kg). Monkeys J and H had been part of previous studies using the contralateral intraparietal sulcus and the ipsilateral frontal cortex. Monkey B had not been part of previous experiments. All animals were in normal health and were group housed with other conspecifics at the facility. All surgical and animal care procedures were approved by the New York University Animal Care and Use Committee and were performed in accordance with the National Institute of Health guidelines for care and use of laboratory animals.

Injection surgery

We performed injections of optogenetic viral vectors 6-25 weeks prior to recording. Subjects were maintained at a constant depth of anesthesia with isofluorane (1-2**%**). Monkey J and H were injected in existing craniotomies. In Monkey B, after mobilizing the scalp and other soft tissue*,* we made craniotomies over motor cortex to provide access for injections. Injections were made through 26 s gauge stainless steel cannulas (Hamilton) that were inserted through a thinned dura. For each injection, one microliter of AAV5-hSyn-ChR2(h134R)-EYFP was injected at a rate of 0.05 ∫l/min. We made injections at 2-3 depths spaced 500-750 ∫m across 2-4 cortical sites per animal. We injected at multiple sites simultaneously to increase the total number of injections sites and thus increase the probability of having successful transduction at one or more sites. We waited 20 minutes between each injection to allow for the virus to diffuse through the tissue. After injections, chambers were reclosed with the chamber cap and periodically opened for cleaning and maintenance while the viral vector was expressing.

Optical Stimulation

We illuminated the injection site with a 473 nm Laser (Shanghai Sanctity Laser)

connected to a multi-mode fiber optic cable (200 ⌈m core, BFL-200, Thor Labs). One end of the

fiber optic cable was terminated with an FC/PC connecter and connected to a collimator (PAFA-

X-4-A, Thor Labs) for focusing light from the laser. At the other end of the cable we removed a

small amount of buffer with a microstripper (T12S21, Thor Labs) and cleaved it with a diamond

knife (S90R, Thorlabs) to ensure a flat edge for minimal power-loss.

The fiber optic was placed on top of the thinned dura. The cable was held in the

microdrive and placed in the same guide tube as the recording electrode. We did not insert the

fiber into the brain in order to minimize damage to the neural tissue. Light power at the tip varied

from 16-20.5 mW. (510-653 mW/mm^2), as measured prior to each experiment using a power

meter (PM100D, Thor Labs). Evoked potentials were visually inspected to ensure that there

were no signs of artifacts from the Becquerel effect, including rapid onset or offset of neural

responses locked to the timing of the light (Cardin et al., 2010). Light based artifacts typically

have a capacitive charging and discharging profile, which has a rapid onset and offset tightly

coupled to the onset and offset of the device. We ensured that the onset of the response and

the inflection point were not aligned with the times when the light was turned on and off. The

neural response is slower than the duration of the light pulse and thus continues to increase for

a brief period after the light has been turned off. The onset is delayed because the ChR2

channel dynamics are slower than the chemical response and thus the membrane potential

does not start to change immediately. The initial response recruits network activity, which has a

slow time course. This is particularly evident for short light pulses (<=5 ms). When there is no

artifact, the response inflection point occurs after the light is turned off. This is less clear for

longer pulse widths (>15 ms) because adaptation and other properties of channel kinetics can

lead to a decrease in the response prior to the end of stimulation.

Light pulse patterns were generated by proprietary software running in LabVIEW

(National Instruments) and converted to analog signals using a DAQ card (USB-6251, National

Instruments). Stimulation trials consisted of 1 s long bursts of square wave pulses. For each

block of trials, we controlled three stimulation parameters: pulse width, pulse rate and pulse

distribution. We varied pulse widths from 5-20 ms, pulse rates from 5-20 Hz and used both

Poisson and periodic pulse distributions.

In rodents, both measurement and simulation have proposed that ~1% of light power

density remains after light has passed through approximately 1 mm of brain tissue (Aravanis et

al., 2007; Huber et al., 2008; Yizhar et al., 2011). However, primate brain tissue may have

slightly different optical properties, thus altering the light transmission. In addition, we stimulated

transdurally which may have additional effects on the light. However, we were able to record

purportedly directly driven spiking (low-latency, low-jitter) as far as 1.5mm below the cortical

surface (not shown), suggesting that we were activating neurons into layer 3.

We tested each recording site in pseudo-randomly interleaved blocks of recordings.

Each block consisted of 30-60 repetitions of 1 s long sequences of square wave pulses (bursts),

with 1-3 s of no stimulation between each stimulation burst (**Fig 2C**). We held other stimulation

parameters (pulse rate, duration and intensity) constant within each block.

Poisson and periodic stimulation pulse trains were generated *in silico* using the same

pulse width and rate parameters. Simulations consisted of 1000 trials for each combination of

parameters.

Electrophysiology

We recorded neural signals from awake, alert macaques while they were seated, head-

fixed in a primate chair (Rogue Research) in a dark, quiet room. Neural signals were recorded

on glass-coated tungsten electrodes (Alpha-Omega, impedance: 0.7-0.8 M∧ at 1 kHz) held in a

FlexMT microdrive (Alpha Omega). Neural signals were acquired at 30 kHz on an NSpike DAQ

system and recorded using proprietary data acquisition software. Neural recordings were

referenced to a metal wire placed in contact with the dura and away from the recording site. We

recorded driven LFP activity and multi-unit spiking in premotor cortex in monkey B (pre-motor

cortex) and posterior parietal cortex in Monkeys J and H.

Fitting E-I Windows

E-I windows were fit to the form

$$EI = Ae^{-At}\left(\frac{\beta_{exc}\left(At\right)^{\alpha_{exc}}}{\alpha_{exc}!} - \frac{\beta_{inh}\left(At\right)^{\alpha_{inh}}}{\alpha_{inh}!} \right)$$

551   which has an excitatory component

$$E = Ae^{-At}\left(\frac{\beta_{exc}\left(At\right)^{\alpha_{exc}}}{\alpha_{exc}!} \right)$$

553   and an inhibitory component

$$I = Ae^{-At}\left(\frac{\beta_{inh}\left(At\right)^{\alpha_{inh}}}{\alpha_{inh}!} \right).$$

For the pure simulations, A, β_{exc} and β_{inh} were set to 1. To fit empirical E-I windows, all

parameters were fit using a grid search to minimize the squared error between the modelled

window and the empirical window.

E-I Window Model

We model neuronal spiking according to a conditional intensity function, $\lambda(t)$, which

represents the instantaneous firing rate, such that the probability of a spike occurring at time t

is given by $\lambda(t)\Delta t$ (Brown et al., 2003). In the absence of optogenetic stimulation, we model

background spiking activity as a Poisson process with a constant rate, λ_0 (**Figure 1A**). We

model background LFP activity as a process with increased power at low frequencies–"brown

noise". For simplicity, we further assume that background LFP activity is uncorrelated with

spiking activity and does not affect the value of $\lambda(t)$.

To simulate LFPs using an excitation window model, we generated $\dfrac{T}{dt}$ samples of

Brownian noise as background activity, with T =1 s and dt =1 ms. We estimated an E-I window

for each recording site by fitting the stimulation pulse evoked LFP response to the following

model:

$$EI = \beta_{exc} exc - \beta_{inh} inh$$

$$exc = Ae^{-At}\left(\frac{(At)^{\alpha_{exc}}}{\alpha_{exc}!}\right)$$

$$inh = Ae^{-At}\left(\frac{(At)^{\alpha_{inh}}}{\alpha_{inh}!}\right)$$

We simulated the driven LFP by convolving the site-specific E-I window with a sequence of

delta pulses determined by the stimulation parameters. The driven LFP was and added to the

background activity to generate the total simulated LFP activity.

In the model, separate E and I responses to stimulation pulses are summed to create

the E-I window. For illustrative purposes, we show the E response preceding the I response but

this can be reversed so that I precedes E (Cruikshank et al., 2010; Haider et al., 2013). The E-I

window is then convolved with a pulse train to model the combined E-I response to stimulation.

To simulate spiking responses, the positive- and negative-rectified responses to the pulse train

are passed through a non-linear transformation and summed to generate the time varying firing

rate, $\lambda(t)$. Individual action potentials are then randomly generated at each time point from a

binomial random variable with $p(spike\ at\ time\ t) = \lambda(t)\Delta t$.

We modelled spiking activity that was coherent with the LFP using a Poisson point

process with a conditional intensity function. The conditional intensity function models stochastic

spiking governed by a time-varying instantaneous firing rate modelled as

$$\lambda(t) = (\lambda_E(t) + \lambda_I(t) + \lambda_0) \cdot dt$$
$$\lambda_E(t) = f(E(t)) = e^{E(t)}$$
$$\lambda_I(t) = g(I(t)) = e^{I(t)}$$

where λ_0 is the spontaneous firing rate, and dt is the sampling rate. $E(t)$ and $I(t)$ are the time

courses of optogenetically driven E and I response, respectively, that generate the fluctuations

in the conditional intensity function, $\lambda(t)$. $E(t)$ and $I(t)$ is determined by convolving the E-I

window with a Poisson process generated from the stimulus sequence parameters, then

positively or negatively rectifying the E-I response to get the separate E and I responses. After a

spike we used a fixed refractory period by forcing $\lambda(t) = 0$ for 3ms.

Simulations of the excitation window model were run using custom scripts in MATLAB

using a 1 ms time step. For each trial, we simulated 1 s of spontaneous activity and 1 s of

stimulation. We simulated 1000 trials for set of stimulus parameters.

Spiking network model

The spiking network model consists of a network of 4000 excitatory and 1000 inhibitory

leaky integrate-and-fire neurons, each of whose membrane potential, V, is described by the

equation:

$$\tau_m \frac{dV}{dt} = -(V - V_r) + V_A - V_G + \frac{k\tau_m I_{ChR2}}{C_m}$$

Neurons are randomly connected to each other with a connection probability $P_{e/i,e/i}$. In addition,

the neurons receive connections from external inputs (described below), which represent

synaptic input from spontaneous activity in other areas of the brain. The synaptic potentials, I_A

and I_G, taken from the Mazzoni model, represent the sum of synaptic responses of each

neuron induced by individual pre-synaptic spikes from external inputs and other excitatory

(pyramidal, AMPA) and inhibitory (interneurons, GABA) neurons in the model.

$$\tau_{dA} \frac{dV_A}{dt} = -V_A + x_A$$

$$\tau_{rA} \frac{dx_A}{dt} = -x_A + \tau_m \left(J_{pyr} \sum_{pyr} \delta\left(t - t_{pyr} - \tau_L\right) + J_{ext} \sum_{ext} \delta\left(t - t_{ext} - \tau_L\right) \right)$$

$$\tau_{dG} \frac{dV_G}{dt} = -V_G + x_G$$

$$\tau_{rG} \frac{dx_G}{dt} = -x_G + \tau_m \left(J_{int} \sum_{int} \delta\left(t - t_{int} - \tau_L\right) \right)$$

In the model, $t_{pyr/int/ext}$ is the time of a spike on a presynaptic neuron. The synaptic currents

generated by pre-synaptic spikes have latency τ_L and have dynamics governed by their

respective exponential rise times $\left(\tau_{rA}, \tau_{rG}\right)$ and decay times $\left(\tau_{dA}, \tau_{dG}\right)$. The efficacy of each

connection type is given by $J_{pyr/int/ext}$ and is uniform for each neuron type. **Table 1** shows the

parameters for the Mazzoni part of the model.

External Inputs

We modeled spontaneous external activity as inputs generated from excitatory neurons

having random Poisson spike trains with a time-varying rate, $r_{ext}(t)$, that is the same for all

neurons (Mazzoni et al., 2008). The spike trains are generated from a Poisson process with a

rate governed by an Ornstein-Uhlenbeck process:

$$r_{ext}(t) = \begin{cases} c_0 + n(t), & c_0 + n(t) \geq 0 \\ 0, & otherwise \end{cases}$$

$$\tau_n \frac{dn(t)}{dt} = -n(t) + \sigma_n \sqrt{\frac{2}{\tau_n}} \theta(t)$$

622 where $\theta(t)$ is Gaussian white noise, c_0 is a non-time-varying mean firing rate and $r_{ext}(t)$ is

623 half-wave rectified to ensure that all firing rates are non-negative.

624 *Channelrhodopsin photocurrent*

625 We modeled the channelrhodopsin photocurrent in the transduced neurons according to

626 a model from Witt *et al.* (Witt et al., 2013).

627
$$I_{ChR2}(t) = -a_{refract} g_{ChR2} F_{ChR2}(t, W_{light})(V - V_{ChR2})$$

628
$$F_{ChR2}(t, W_{light}) = A_{act}\left(1 - e^{-\frac{t - t_{ON} - d}{\tau_{act}}}\right)\left(A_{persist} + A_{inact}^{(1)} e^{-\frac{t - t_{ON} - d}{\tau_{inact}^{(1)}}} + A_{inact}^{(2)} e^{-\frac{t - t_{ON} - d}{\tau_{inact}^{(2)}}}\right)$$

629
$$d = d_A + d_B W_{light} + \frac{d_c}{W_{light}}$$

630
$$\tau_{act} = \tau_{act}^{(0)} + c_{act} e^{-k_{act} W_{light}}$$

631
$$A_{act} = a_0 + \frac{a_{min} - 1}{1 + \left(\frac{W_{0.5}}{W_{light}}\right)^2}$$

632
$$A_{inact}^{(1)} = b_0 + \frac{b_1}{b_2 + \left(W_{light} - W_{inact}\right)^2}$$

633
$$A_{inact}^{(2)} = c_{inact} e^{-k_{inact} W_{light}}$$

634
$$A_{persist} = 1 - A_{inact}^{(1)} - A_{inact}^{(2)}$$

The photocurrent, I_{ChR2}, is defined as the product of the conductance, $g_{ChR2} F_{ChR2}(t, W_{light})$, and

the driving force, $V - V_{ChR2}$, from the ChR2 channels at any time, t. When light stimulation is on,

the conductance varies over time as described by the function F_{ChR2}. When the light is off, the

conductance returns to baseline following a single exponential decay with a time constant of

τ_{OFF}. The light induced conductance change is described as the product of activation and

inactivation functions which depend on the time from light onset $t - t_{ON}$, a light intensity

dependent delay, d, and their respective time constants, τ_{act}, $\tau_{inact}^{(1)}$ and $\tau_{inact}^{(2)}$. The parameters

for equations 8-15 have been previously fitted to photoconductance measured *in vitro* human

embryonic kidney cells transduced with AAV1/2-CAG-ChR2-YFP (Witt et al., 2013). **Table 1**

shows the parameters for the ChR2 current part of the model. Each neuron of each type was

randomly assigned to have ChR2 channels according to the transduction probability for that cell

type. We used a viral vector containing the pan-neuronal hSyn promoter. Therefore, we set

equal transduction probabilities equal to 1% for both excitatory neurons and inhibitory

neurons.(Diester et al., 2011).

Previous studies have shown that transduced neurons do not have robust spiking

responses to individual light pulses at higher stimulation frequencies (Lin et al., 2009), which is

likely due to the slower kinetics of ChR2 channels. This response fidelity drop-off tends to start

around 20 Hz periodic stimulation and reaches near zero around 80-100 Hz stimulation (Cardin

et al., 2009). Therefore, light pulses that occurred less than 50 ms after the previous pulse

activated ChR2 channels with a probability that depended linearly such that

$$a_{refract} = \begin{cases} 1, & \Delta t \geq 50ms \\ 1, & P < \dfrac{\Delta t}{50ms} \\ 0, & otherwise \end{cases}$$

656   where $P \in unif(0,1)$ and is chosen individually for each close pulse occurrence and for each

657   model neuron.

$\tau_{m,exc}$	20 ms
$V_{rest,pyr}$	0 mV
$V_{thresh,pyr}$	18 mV

$V_{reset,pyr}$	11 mV
$\tau_{m,int}$	10 ms
$V_{rest,int}$	0 mV
$V_{thresh,int}$	18 mV
$V_{reset,int}$	11 mV
τ_L	1 ms
τ_{rG}	0.25 ms
$\tau_{rA,int}$	0.2 ms
$\tau_{rA,pyr}$	0.4 ms
$\tau_{dA,int}$	1 ms
$\tau_{dA,pyr}$	2 ms
$\tau_{dG,pyr}$	5 ms
$\tau_{rG,pyr}$	0.25 ms
$\tau_{dG,int}$	5 ms
$\tau_{rG,int}$	0.25 ms
τ_n	16 ms
$J_{ext,int}$	0.95 mV
$J_{ext,pyr}$	0.55 mV
N_{ext}	5000
C_m	0.5 nF
$J_{int,exc}$	1.7 mV
$J_{int,inh}$	2.7 mV
$J_{pyr,pyr}$	0.42 mV
$J_{pyr,int}$	0.7 mV
$t_{refract,pyr}$	2 ms
$t_{refract,int}$	1 ms
$P_{i,i}$	0.2
$P_{e,i}$	0.2
$P_{i,e}$	0.4
$P_{e,e}$	0.2
W_{inact}	0.11
τ_{off}	3 ms
k	1e10
d_A	0.27
d_B	-0.05
d_C	-0.0126
$\tau_{act}^{(0)}$	0.74 ms
c_{act}	12
k_{act}	25
a_0	1
a_{min}	0.4
$W_{0.5}$	0.38
$\tau_{inact}^{(1)}$	9.06 ms
b_0	0.16
b_1	0.013

b_2	0.027
$\tau_{inact}^{(2)}$	59.6 ms
c_{inact}	0.29
k_{inact}	2.4
W_{light}	0.3
g_{ChR2}	0.007 uS
V_{ChR2}	65 mV

Table 1: Spiking network model parameters.

Pre-processing in vivo data

Broad-band neural activity was split into multi-unit signals and local field potentials (LFP)

by high-pass and low-pass filtering, respectively, at 300 Hz. LFP signals were down-sampled to

1 kHz. Multi-unit activity (MUA) was computed by applying a 3.5 standard deviation threshold to

the multi-unit signals to identify putative spikes. All waveforms that exceeded this peak were

labelled as multi-unit action potentials.

Pre-processing in silico data

LFPs measure currents generated by synaptic activity within a volume near the

electrode (Logothetis, 2003). Therefore, we followed Mazzoni *et al*. (Mazzoni et al., 2008) and

modeled the LFP as a sum of synaptic currents. Since interneurons typically have dendrites that

are organized in a symmetrical, closed-field configuration, we assumed their contribution to the

LFP would be minimal in comparison with synaptic currents on the typically dipole-like dendrites

of pyramidal neurons (Buzsáki and Wang, 2012). Therefore, only synaptic currents in pyramidal

neurons contributed to our measure of the model LFP. Adding together the absolute values of

synaptic currents has been shown to accurately reproduce the power spectrum and information

content of LFP recorded in macaque visual cortex (Mazzoni et al., 2008). Simulated LFPs were

also low-pass filtered at 300 Hz and down-sampled to 1 kHz. Simulated multiunit activity

consisted of all the threshold crossings in the leaky integrate-and-fire model, as recorded by the

Brian simulator. The *in vivo* and *in silico* data were pre-processed, were treated the same after

pre-processing.

Data Analysis

We computed multi-unit peri-stimulus time histograms (PSTH) by aligning MUA to the

onset time of light pulses. Individual spikes were collected in 1 ms bins and the corresponding

histogram was smoothed by convolution with a Gaussian function with a standard deviation of 1

ms. In order to avoid corruption from nearby pulses, evoked potentials were computed by

averaging LFPs aligned to the onset time of light pulses only for pulses that occurred at the end

of a pulse train and did not have a pulse preceding it by less than 50 ms.

Spectral analysis was performed using multi-taper methods (Mitra and Pesaran, 1999).

The accuracy of the spectral quantities we used does not depend on the center frequency of the

estimate for all frequencies above the minimum resolvable frequency, given by the frequency

smoothing in Hz (Jarvis and Mitra, 2001). For each recording, power spectra were computed

using a 1 s analysis window and ± 1.5 Hz frequency smoothing. We computed power spectra

for the LFP responses during entire 1 s burst (stimulation epoch) and for 1 s prior to the burst

(spontaneous epoch). Power spectra were computed for neural signals for the spontaneous

and stimulation epochs. Recording sites with power spectra that did not exhibit a significant

difference were not included in analysis. The exclusion criteria for recording sessions did not

depend on the frequency content of the driven LFP response—the desired effect—but rather

only that there was any driven LFP response at all. Sites were excluded if there was no change

in the LFP power between spontaneous and stimulation periods for the 10ms pulse width

condition, which was likely due to the electrode being placed on the periphery of light cone and

thus failing to record driven neural activity. We explicitly confirm this in our experiments by

moving the electrode and observing driven responses. We computed the spectrogram for each

recording using a ± 250 ms analysis window, stepped 50 ms between windows, with ± 2.5 Hz

frequency smoothing aligned to the start of the burst. We computed the spike-field coherence

for each recording using a 1 s analysis window with ± 3 Hz frequency smoothing averaged

across at least 30 repetitions.

DATA AND SOFTWARE AVAILABILITY

Software is available in a GitHub repository linked from http://www.pesaranlab.org. Data are

available upon reasonable request.

Acknowledgements:

This work was supported, in part, by US National Institutes of Health (NIH) BRAIN Initiative grant R01-NS104923, NIH R01-EY024067, NIH P30-EY013079, NIH Training Grant T32-EY007136, a Scholar Award from the McKnight Endowment Fund for Neuroscience (BP) and the Alfred P Sloan Foundation (BP).

Author Contributions:

Conceptualization: R.A.S., H.L.D., Y.T.W., M.A.H., and B.P.; Methodology: R.A.S., H.L.D., Y.T.W., M.A.H., M.M.F. and B.P.; Software: R.A.S. and B.P.; Formal Analysis: R.A.S. and B.P.; Investigation: R.A.S. and B.P.; Resources: R.A.S., H.L.D., and B.P.; Writing – Original Draft: R.A.S. and B.P.; Writing – Review & Editing: R.A.S., H.L.D., Y.T.W., M.A.H., M.M.F. and B.P.; Visualization: R.A.S. and B.P.; Supervision: H.L.D., Y.T.W., M.A.H., and B.P.; Funding Acquisition: B.P.

Competing Financial Interests:

The authors declare no competing financial interests.

References

Adesnik H, Scanziani M. 2010. Lateral competition for cortical space by layer-specific horizontal circuits. *Nature* **464**:1155–1160. doi:10.1038/nature08935

Aravanis AM, Wang L-P, Zhang F, Meltzer LA, Mogri MZ, Schneider MB, Deisseroth K. 2007. An optical neural interface: in vivo control of rodent motor cortex with integrated fiberoptic and optogenetic technology. *J Neural Eng* **4**:S143-56. doi:10.1088/1741-2560/4/3/S02

Atallah B V., Scanziani M. 2009. Instantaneous Modulation of Gamma Oscillation Frequency by Balancing Excitation with Inhibition. *Neuron* **62**:566–577. doi:10.1016/j.neuron.2009.04.027

Börgers C, Kopell N. 2003. Synchronization in Networks of Excitatory and Inhibitory Neurons with Sparse, Random Connectivity. *Neural Comput* **15**:509–538. doi:10.1162/089976603321192059

Brown EN, Barbieri R, Eden UT, Frank LM. 2003. Likelihood methods for neural data analysis In: Feng J, editor. Computational Neuroscience: A Comprehensive Approach. Boca Raton: CRC Press. pp. 253–286.

Brunel N, Wang X-J. 2003. What Determines the Frequency of Fast Network Oscillations With Irregular Neural Discharges? I. Synaptic Dynamics and Excitation-Inhibition Balance. *J Neurophysiol* **90**:415–430. doi:10.1152/jn.01095.2002

Burns SP, Xing D, Shapley RM. 2011. Is gamma-band activity in the local field potential of V1 cortex a "clock" or filtered noise? *J Neurosci* **31**:9658–9664. doi:10.1523/JNEUROSCI.0660-11.2011

Burns SP, Xing D, Shelley MJ, Shapley RM. 2010. Searching for autocoherence in the cortical network with a time-frequency analysis of the local field potential. *J Neurosci* **30**:4033–4047. doi:10.1523/JNEUROSCI.5319-09.2010

Buschman TJ, Miller EK. 2007. Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. *Science* **315**:1860–2. doi:10.1126/science.1138071

752 Buzsáki G, Anastassiou CA, Koch C. 2012. The origin of extracellular fields and currents —

753  EEG, ECoG, LFP and spikes. *Nat Rev Neurosci* **13**:407–420. doi:10.1038/nrn3241

754 Buzsáki G, Wang X-J. 2012. Mechanisms of Gamma Oscillations. *Annu Rev Neurosci* **35**:203–

755  225. doi:10.1146/annurev-neuro-062111-150444

756 Cardin JA, Carlén M, Meletis K, Knoblich U, Zhang F, Deisseroth K, Tsai L-H, Moore CI. 2010.

757  Targeted optogenetic stimulation and recording of neurons in vivo using cell-type-specific

758  expression of Channelrhodopsin-2. *Nat Protoc* **5**:247–254. doi:10.1038/nprot.2009.228

759 Cardin JA, Carlén M, Meletis K, Knoblich U, Zhang F, Deisseroth K, Tsai L-H, Moore CI. 2009.

760  Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature*

761  **459**:663–7. doi:10.1038/nature08002

762 Chariker L, Shapley R, Young L-S. 2018. Rhythm and Synchrony in a Cortical Network Model. *J*

763  *Neurosci* **38**:8621–8634. doi:10.1523/JNEUROSCI.0675-18.2018

764 Cruikshank SJ, Urabe H, Nurmikko A V, Connors BW. 2010. Pathway-Specific Feedforward

765  Circuits between Thalamus and Neocortex Revealed by Selective Optical Stimulation of

766  Axons. *Neuron* **65**:230–245. doi:10.1016/j.neuron.2009.12.025

767 Csicsvari J, Jamieson B, Wise KD, Buzsáki G. 2003. Mechanisms of gamma oscillations in the

768  hippocampus of the behaving rat. *Neuron* **37**:311–322. doi:10.1016/S0896-6273(02)01169-

769  8

770 Dean HL, Hagan MA, Pesaran B. 2012. Only Coherent Spiking in Posterior Parietal Cortex

771  Coordinates Looking and Reaching. *Neuron* **73**:829–841.

772  doi:10.1016/j.neuron.2011.12.035

773 Diester I, Kaufman MT, Mogri M, Pashaie R, Goo W, Yizhar O, Ramakrishnan C, Deisseroth K,

774  Shenoy K V. 2011. An optogenetic toolbox designed for primates. *Nat Neurosci* **14**:387–

775  397. doi:10.1038/nn.2749

776 Feingold J, Gibson DJ, Depasquale B, Graybiel AM. 2015. Bursts of beta oscillation differentiate

777  postperformance activity in the striatum and motor cortex of monkeys performing

movement tasks **112**. doi:10.1073/pnas.1517629112

Feldbauer K, Zimmermann D, Pintschovius V, Spitz J, Bamann C, Bamberg E. 2009.

Channelrhodopsin-2 is a leaky proton pump. *Proc Natl Acad Sci USA* **106**:12317–22.

doi:10.1073/pnas.0905852106

Fries P. 2005. A mechanism for cognitive dynamics: Neuronal communication through neuronal

coherence. *Trends Cogn Sci* **9**:474–480. doi:10.1016/j.tics.2005.08.011

Gabernet L, Jadhav SP, Feldman DE, Carandini M, Scanziani M. 2005. Somatosensory

integration controlled by dynamic thalamocortical feed-forward inhibition. *Neuron* **48**:315–

27. doi:10.1016/j.neuron.2005.09.022

Haider B, Häusser M, Carandini M. 2013. Inhibition dominates sensory responses in the awake

cortex. *Nature* **493**:97–100. doi:10.1038/nature11665

Hasenstaub A, Shu Y, Haider B, Kraushaar U, Duque A, McCormick DA. 2005. Inhibitory

postsynaptic potentials carry synchronized frequency information in active cortical

networks. *Neuron* **47**:423–435. doi:10.1016/j.neuron.2005.06.016

Higley MJ, Contreras D. 2003. Nonlinear integration of sensory responses in the rat barrel

cortex: an intracellular study in vivo. *J Neurosci* **23**:10190–200.

doi:10.1523/JNEUROSCI.23-32-10190.2003

Holgado AJN, Terry JR, Bogacz R. 2010. Conditions for the Generation of Beta Oscillations in

the Subthalamic Nucleus-Globus Pallidus Network. *J Neurosci* **30**:12340–12352.

doi:10.1523/jneurosci.0817-10.2010

Huber D, Petreanu L, Ghitani N, Ranade S, Hromádka T, Mainen Z, Svoboda K. 2008. Sparse

optical microstimulation in barrel cortex drives learned behaviour in freely moving mice.

Nature **451**:61–4. doi:10.1038/nature06445

Jarvis MR, Mitra PP. 2001. Sampling properties of the spectrum and coherency of sequences of

action potentials. *Neural Comput* **13**:717–49. doi:10.1162/089976601300014312

Jensen O, Goel P, Kopell N, Pohja M, Hari R, Ermentrout B. 2005. On the human sensorimotor-

cortex beta rhythm: Sources and modeling. *Neuroimage* **26**:347–355.

doi:10.1016/j.neuroimage.2005.02.008

Kato HK, Asinof SK, Isaacson JS. 2017. Network-Level Control of Frequency Tuning in Auditory

Cortex. *Neuron* **95**:412-423.e4. doi:10.1016/j.neuron.2017.06.019

Keeley S, Byrne Á, Fenton AA, Rinzel J. 2019. Firing Rate Models for Gamma Oscillations. *J*

Neurophysiol 2181–2190. doi:10.1152/jn.00741.2018

Keeley S, Fenton AA, Rinzel J. 2017. Modeling fast and slow gamma oscillations with

interneurons of different subtype. *J Neurophysiol* **117**:950–965. doi:10.1152/jn.00490.2016

Lee JH, Whittington MA, Kopell NJ. 2013. Top-Down Beta Rhythms Support Selective Attention

via Interlaminar Interaction: A Model. *PLoS Comput Biol* **9**.

doi:10.1371/journal.pcbi.1003164

Lin JY, Lin MZ, Steinbach P, Tsien RY. 2009. Characterization of engineered channelrhodopsin

variants with improved properties and kinetics. *Biophys J* **96**:1803–14.

doi:10.1016/j.bpj.2008.11.034

Logothetis NK. 2003. The underpinnings of the BOLD functional magnetic resonance imaging

signal. *J Neurosci* **23**:3963–3971. doi:23/10/3963 [pii]

Lu Y, Truccolo W, Wagner FB, Vargas-Irwin CE, Ozden I, Zimmermann JB, May T, Agha NS,

Wang J, Nurmikko A V. 2015. Optogenetically induced spatiotemporal gamma oscillations

and neuronal spiking activity in primate motor cortex. *J Neurophysiol* **113**:3574–3587.

doi:10.1152/jn.00792.2014

Mazzoni A, Panzeri S, Logothetis NK, Brunel N. 2008. Encoding of naturalistic stimuli by local

field potential spectra in networks of excitatory and inhibitory neurons. *PLoS Comput Biol*

4. doi:10.1371/journal.pcbi.1000239

Mitra PP, Pesaran B. 1999. Analysis of dynamic brain imaging data. *Biophys J* **76**:691–708.

doi:10.1016/S0006-3495(99)77236-X

Ni J, Wunderle T, Lewis CM, Desimone R, Diester I, Fries P. 2016. Gamma-Rhythmic Gain

Modulation. *Neuron* **92**:240–251. doi:10.1016/j.neuron.2016.09.003

Okun M, Lampl I. 2008. Instantaneous correlation of excitation and inhibition during ongoing and

sensory-evoked activities. *Nat Neurosci* **11**:535–537. doi:10.1038/nn.2105

Okun M, Naim A, Lampl I. 2010. The Subthreshold Relation between Cortical Local Field

Potential and Neuronal Firing Unveiled by Intracellular Recordings in Awake Rats. *J*

Neurosci **30**:4440–4448. doi:10.1523/JNEUROSCI.5062-09.2010

Pesaran B, Nelson MJ, Andersen R a. 2008. Free choice activates a decision circuit between

frontal and parietal cortex. *Nature* **453**:406–9. doi:10.1038/nature06849

Pesaran B, Pezaris JS, Sahani M, Mitra PP, Andersen RA. 2002. Temporal structure in

neuronal activity during working memory in macaque parietal cortex. *Nat Neurosci* **5**:805–

811. doi:10.1038/nn890

Pesaran B, Vinck M, Einevoll GT, Sirota A, Fries P, Siegel M, Truccolo W, Schroeder CE,

Srinivasan R. 2018. Investigating large-scale brain dynamics using field potential

recordings: analysis and interpretation. *Nat Neurosci*. doi:10.1038/s41593-018-0171-8

Phillips EAK, Hasenstaub AR. 2016. Asymmetric effects of activating and inactivating cortical

interneurons. *Elife* **5**:1–22. doi:10.7554/eLife.18383

Poulet JFA, Petersen CCH. 2008. Internal brain state regulates membrane potential synchrony

in barrel cortex of behaving mice. *Nature* **454**:881–885. doi:10.1038/nature07150

Roopun AK, Middleton SJ, Cunningham MO, LeBeau FEN, Bibbig A, Whittington M a, Traub

RD. 2006. A beta2-frequency (20-30 Hz) oscillation in nonsynaptic networks of

somatosensory cortex. *Proc Natl Acad Sci* **103**:15646–15650.

doi:10.1073/pnas.0607443103

Rule ME, Vargas-Irwin CE, Donoghue JP, Truccolo W. 2017. Dissociation between sustained

single-neuron spiking and transient β-LFP oscillations in primate motor cortex. *J*

Neurophysiol **117**:1524–1543. doi:10.1152/jn.00651.2016

Salkoff DB, Zagha E, Yuzgec O, McCormick DA. 2015. Synaptic Mechanisms of Tight Spike

Synchrony at Gamma Frequency in Cerebral Cortex. *J Neurosci* **35**:10236–10251.

doi:10.1523/jneurosci.0828-15.2015

Sherman MA, Lee S, Law R, Haegens S, Thorn CA, Hämäläinen MS, Moore CI, Jones SR.

2016. Neural mechanisms of transient neocortical beta rhythms: Converging evidence from

humans, computational modeling, monkeys, and mice. *Proc Natl Acad Sci* **113**:E4885–

E4894. doi:10.1073/pnas.1604135113

Sohal VS, Zhang F, Yizhar O, Deisseroth K. 2009. Parvalbumin neurons and gamma rhythms

enhance cortical circuit performance. *Nature* **459**:698–702. doi:10.1038/nature07991

Wehr M, Zador AM. 2003. Balanced inhibition underlies tuning and sharpens spike timing in

auditory cortex. *Nature* **426**:442–446. doi:10.1038/nature02116

Wilson HR, Cowan JD. 1972. Excitatory and Inhibitory Interactions in Localized Populations of

Model Neurons. *Biophys J* **12**:1–24. doi:10.1016/S0006-3495(72)86068-5

Witt A, Palmigiano A, Neef A, El Hady A, Wolf F, Battaglia D. 2013. Controlling the oscillation

phase through precisely timed closed-loop optogenetic stimulation: a computational study.

Front Neural Circuits **7**:49. doi:10.3389/fncir.2013.00049

Womelsdorf T, Fries P, Mitra PP, Desimone R. 2006. Gamma-band synchronization in visual

cortex predicts speed of change detection. *Nature* **439**:733–736. doi:10.1038/nature04258

Wong YT, Fabiszak MM, Novikov Y, Daw ND, Pesaran B. 2016. Coherent neuronal ensembles

are rapidly recruited when making a look-reach decision. *Nat Neurosci* **19**:1–11.

doi:10.1038/nn.4210

Xing D, Shen Y, Burns S, Yeh C-I, Shapley R, Li W. 2012. Stochastic Generation of Gamma-

Band Activity in Primary Visual Cortex of Awake and Anesthetized Monkeys. *J Neurosci*

32:13873-13880a. doi:10.1523/JNEUROSCI.5644-11.2012

Yizhar O, Fenno LE, Davidson TJ, Mogri M, Deisseroth K. 2011. Optogenetics in neural

systems. *Neuron* **71**:9–34. doi:10.1016/j.neuron.2011.06.004