Functional determinants of a small protein controlling a broadly conserved bacterial sensor kinase

Srujana S Yadavalli[2, 3]*, Ted Goh[4,5], Jeffrey N Carey[6], Gabriele Malengo[1,7], Sangeevan Vellappan[8], Bryce E Nickels[3], Victor Sourjik[1,7], Mark Goulian[2,9]* and Jing Yuan[1,7]*

[1]Max Planck Institute for Terrestrial Microbiology, 35043 Marburg, Germany
[2]Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
[3]Department of Genetics and Waksman Institute of Microbiology, Rutgers University, Piscataway, NJ 08854, USA
[4]Department of Biology, Swarthmore College, Swarthmore, Pennsylvania 19081, USA
[5] Boston University School of Medicine, Boston, Massachusetts 02118, USA
[6]Biochemistry and Molecular Biophysics Graduate Group, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
[7]LOEWE Center for Synthetic Microbiology (SYNMIKRO), 35043 Marburg, Germany
[8]Molecular Biosciences Graduate Program, Rutgers University, Piscataway NJ 08854
[9]Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

*Corresponding authors. J.Y.: jing.yuan@synmikro.mpi-marburg.mpg.de, S.S.Y.: sam.yadavalli@rutgers.edu, M.G.: goulian@sas.upenn.edu

Abstract

The PhoQ/PhoP two-component system plays a vital role in the regulation of Mg^{2+} homeostasis, resistance to acid and hyperosmotic stress, cationic antimicrobial peptides, and virulence in *Escherichia coli*, Salmonella and related bacteria. Previous studies have shown that MgrB, a 47 amino acid membrane protein that is part of the PhoQ/PhoP regulon, inhibits the histidine kinase PhoQ. MgrB is part of a negative feedback loop modulating this two-component system that prevents hyperactivation of PhoQ and may also provide an entry point for additional input signals for the PhoQ/PhoP pathway. To explore the mechanism of action of MgrB, we have analyzed the effects of point mutations, C-terminal truncations and transmembrane region swaps on MgrB activity. In contrast with two other known membrane protein regulators of histidine kinases in *E. coli*, we find that the MgrB TM region is necessary for PhoQ inhibition. Our results indicate that the TM region mediates interactions with PhoQ and that W20 is a key residue for PhoQ/MgrB complex formation. Additionally, mutations of the MgrB cytosolic region suggest that the two N-terminal lysines play an important role in regulating PhoQ activity. Alanine scanning mutagenesis of the periplasmic region of MgrB further indicates that, with the exception of a few highly conserved residues, most residues are not essential for MgrB's function as a PhoQ inhibitor. Our results indicate that the regulatory function of the small protein MgrB depends on distinct contributions from multiple residues spread across the protein. Interestingly, the TM region also appears to interact with other non-cognate histidine kinases in a bacterial two-hybrid assay, suggesting a potential route for evolving new small protein modulators of histidine kinases.

Introduction

Small proteins—proteins less than 50-amino acids long that are directly translated from small open reading frames—regulate a wide variety of cellular processes but, collectively, this class of proteins remains relatively understudied (reviewed in [1, 2]). For example, hundreds to thousands of small open reading frames have been reported in *Escherichia coli* [3-5], *Mycobacterium tuberculosis* [6] and human microbiomes [7]. A fraction of them have been confirmed with detections at the protein level [3-6] and the functions of most these

proteins are unknown. In bacteria and bacteriophages, small proteins can be classified into
two groups: secreted and non-secreted [1]. The former are mainly involved in
communication or competition among organisms, whereas the latter are localized in the
cytoplasm or cell membrane and play a role in diverse aspects of cell physiology. In *E. coli*,
the list of small proteins that have been identified and verified continues to grow with
improvements in computational and experimental techniques [3, 4, 8]. Interestingly, nearly
one third have been shown or are predicted to be in the cytoplasmic membrane, suggesting
many may be involved in sensing environmental cues and mediating stress responses.
63
Bacteria thrive in a multitude of niches, often under challenging physicochemical conditions,
and have therefore evolved stress response systems to monitor the environment and
modulate their physiology accordingly. Many of these stress responses are controlled by
two-component signaling systems, one of the primary modes of bacterial signal transduction
[9]. The PhoQ/PhoP system in *E. coli*, Salmonella, and related Gram-negative bacteria is a
well-studied example of a two-component system that is critical for virulence and facilitates
adaptation to conditions of low Mg^{2+} or Ca^{2+}, low pH, osmotic upshifts and the presence of
cationic antimicrobial peptides [10-14]. In response to input signal, the sensor histidine
kinase PhoQ autophosphorylates and modulates the phosphorylation state of the response
regulator PhoP, which functions as a transcription factor [10, 15]. At least three small
proteins are regulated by PhoP: the 31-amino acid membrane protein MgtS is induced under
magnesium limitation in *E. coli* and inhibits degradation of the magnesium transporter MgtA
[16]; the 30-amino acid protein MgtR promotes the degradation of the virulence factor MgtC
by the FtsH protease in *Salmonella enterica* serovar Typhimurium [17]; and the 47-amino
acid protein MgrB directly inhibits PhoQ kinase activity, thereby mediating a negative
feedback loop [18, 19]. All of these small proteins are membrane-localized, regulate proteins
expressed in the PhoP regulon, and modify cell responses to the environment.
81
MgrB orthologs are found in numerous Enterobacteriaceae, and PhoQ inhibition has been
demonstrated explicitly for a few distinct genera. The physiological importance of this
inhibition and the associated negative feedback loop remain poorly understood. MgrB could
serve as a point of control for additional input signals that modulate PhoQ. For example,
MgrB activity is affected by the redox state of the periplasm via the protein's two conserved
periplasmic cysteines [20]. MgrB is not essential and appears to mainly act in a regulatory
role, however it can have a significant impact on bacterial physiology. In *E. coli*, recent work
has shown that strong stimulation of the PhoQ/PhoP system hinders cell division [21]. Cells
grown in low levels of Mg^{2+} in the absence of MgrB form filaments due to strong activation of
PhoQ in this condition. MgrB-mediated feedback inhibition thus appears to be important to
appropriately limit PhoQ activity and prevent hyperactivation of the system. In at least some
contexts, loss of *mgrB* can also confer a fitness advantage: mutation of *mgrB* has emerged
as one of the primary mechanisms of acquired resistance to the last resort antibiotic colistin
in clinical isolates of *Klebsiella pneumoniae* [22-25].
96
In addition to MgrB, the PhoQ histidine kinase is modulated by another membrane protein,
SafA [26], which is present exclusively in *E. coli*, and acts as a connector between the
EvgS/EvgA and PhoQ/PhoP two-component systems. Stimulation of EvgS by low pH
upregulates SafA expression [27], which activates PhoQ kinase activity and leads to
increased acid resistance. In at least some *E. coli* isolates, this connection is required for
PhoQ activation in response to acid stress [28, 29]. Another example of a connector protein
that modulates the activity of a histidine kinase is MzrA [30], which links the CpxA/CpxR and
EnvZ/OmpR two-component systems. Although not as small as MgrB, both SafA and MzrA
are relatively small for membrane proteins (65 and 127 amino acids, respectively) and, like
MgrB, are bitopic with their N-termini in the cytoplasm.
107
MgrB acts by inhibiting PhoQ kinase activity [19], however, the mechanism by which this
small protein inhibits a protein that is roughly ten times larger than itself, and the sequence

determinants responsible for this action are largely unknown. In this work, we show that the
transmembrane (TM) region of MgrB is required for the protein to form a complex with its
target kinase PhoQ, in contrast with SafA and MzrA, which do not require their TM regions
for their activity [31, 32]. Using site-directed mutagenesis combined with MgrB functional
assays and *in vivo* Förster resonance energy transfer (FRET), we identified W20 in the MgrB
TM region to be crucial for MgrB/PhoQ complex formation. Additionally, we identified 11
residues across different regions of MgrB that are important for its inhibitory function. Among
these residues, two lysines in the cytoplasmic region likely act through their charged side
chains to regulate PhoQ HAMP conformation, although they do not appear to affect
MgrB/PhoQ complex formation. Surprisingly, a large number of amino acids in the small
periplasmic region of MgrB can be replaced with alanine with only minor effects on protein
function. Together our findings suggest that PhoQ inhibition by MgrB depends on
coordinated interactions from amino acid residues spanning different regions of the protein.
We also uncovered interactions of MgrB with other histidine kinases in bacterial two-hybrid
assays, which may provide a starting point for evolving novel small membrane proteins that
modulate other histidine kinases.
126
127 Results
128
129 **Sequence analysis of MgrB**
130
MgrB was shown to be a membrane-localized small protein with N-terminal-in and C-
terminal-out topology [18]. To predict the transmembrane (TM) region in MgrB, we used
eight algorithms (PHDHTM, DAS-TMFILTER, SCAMPI, TMHMM, TMpred, TMMTOP,
PRED-TMR and MEMSAT) to analyze the entire amino acid sequence (Fig 1). The
predictions were fairly convergent on the TM cytoplasm/membrane boundary, which was
located at W6 or V7. In contrast, the predictions of the membrane/periplasm boundary were
quite different among these programs, ranging from F24 to M27. In this manuscript, we use
W6 to F24 as the predicted transmembrane region of MgrB.
139
To gain insight on potential functionally important residues in MgrB, we aligned over 100
bacterial MgrB sequences obtained from KEGG protein BLAST using *E. coli* MgrB as query
(Fig 1 and supplementary file 1). As indicated in the sequence logo of the alignment, the
amino acids were fairly conserved in the cytosolic and periplasmic regions. This high degree
of conservation might originate from limitations in the diversity of MgrB sequences currently
available in databases or suggest the functional importance of each conserved amino acid in
these regions. In contrast, more variations and insertion of one amino acid were shown in
the predicted TM region. Interestingly, two amino acids with polar side chains (C16 and Q22)
and two tryptophans (W6 and W20) in the TM region are highly conserved, suggesting a
functional importance.
150
151 **MgrB TM and periplasmic regions are necessary but not sufficient for MgrB function**
152
As a first step towards understanding how MgrB inhibits PhoQ at a structural level, we
looked at the role of the protein's TM and periplasmic regions. To test the functional role of
the MgrB TM region, we swapped the predicted MgrB TM region with either (i) a 20-amino
acid TM from the *E. coli* maltose transporter MalF or (ii) a 21-amino acid variant of human
glycophorin A (GpA) TM [33] (Fig 2A). The MgrB variants described in this section are
derivatives of a functional N-terminally GFP tagged MgrB (GFP-MgrB) [18]. Both MgrB
variants with swapped TM regions localized to the cell membrane (S1 Fig) and their
expression levels were comparable to that of wild type (S2 Fig). TM swapping experiments
of two previously studied histidine kinase regulators–SafA and MzrA—indicate that the
transmembrane sequence is dispensable for their function [31, 32]. Indeed, our control
experiments in which we replaced the TM sequences of SafA and MzrA with that of GpA

confirmed these results (S3 Fig). Thus, for SafA and MzrA, the TM region appears to
primarily act as an anchor to direct the regulatory domain of these proteins to the
membrane. We monitored the effect of MgrB and its TM swap variants on PhoQ activity
using a PhoQ/PhoP-regulated transcriptional reporter (P_{mgtA}-*lacZ*) (Fig 2A). Control cells
expressing no MgrB had a high level of β-galactosidase (β-gal) activity, indicating uninhibited
PhoQ, whereas cells expressing WT MgrB had minimal β-gal activity, indicating PhoQ
inhibition by WT MgrB as expected. In contrast to the inhibitory function of the WT MgrB,
both MgrB TM swap variants (MgrB-MalF TM and MgrB-GpA TM) showed high β-gal
activities that were about 80% of the activity of no-MgrB control, indicating that they are
unable to inhibit PhoQ efficiently. These results suggest that the MgrB TM region not only
functions as a membrane anchor but also has additional properties that are critical for MgrB
to inhibit PhoQ.
176
As swapping the TM sequence in MgrB severely affected its activity, we tested whether an
MgrB mutant lacking the periplasmic region inhibits PhoQ. This variant－MgrB-24, in which
N25 is replaced with a stop codon－showed proper membrane localization (S1 Fig), but
could not repress reporter gene expression (Fig 2B), indicating that this truncated variant
composed of the MgrB TM and short N-terminal cytoplasmic regions is not functional. To
assess the minimum length of MgrB needed for efficient repression of PhoQ/PhoP-regulated
reporter gene expression, we tested other C-terminal truncations of MgrB: MgrB-29, MgrB-
34, MgrB-39, MgrB-44, corresponding to replacing the codons for Q30, F35, A40 and I45
with a stop codon, respectively. These mutants displayed similar membrane localization to
that of WT MgrB (S1 Fig). Of these four truncations, MgrB-29 and -34 did not show any
detectable repression, MgrB-39 showed slight repression, and MgrB-44, which lacks only
the last two residues, showed the strongest repression, although last variant was still
compromised in activity compared with WT (Fig 2B). Evidently, amino acid interactions in
both the TM and the periplasmic region of MgrB influence the protein's ability to inhibit
PhoQ.
192
Previous work suggested that the periplasmic region of MgrB is required for PhoQ inhibition
[18]. Since the TM swaps described above change the TM helix length, their loss of activity
could be due to a change in the helical phase at the C-terminus of the TM helix creating
different orientations of the periplasmic region. It is therefore possible that the periplasmic
region of MgrB alone could inhibit PhoQ, while the TM helix only ensures its correct
orientation. Indeed, the chemically synthesized periplasmic region of SafA (SafA[peri]) was
previously shown to be sufficient for PhoQ activation when added exogenously [34].
Therefore, we tested whether the MgrB periplasmic region (MgrB[peri]) could function similarly
to full-length MgrB (Fig 2C) by adding MgrB[peri] to an *E. coli* culture. To enable the MgrB[peri]
peptide to access the periplasm, we used an *imp-4213* strain, which has increased outer
membrane permeability [35]. We monitored PhoQ activity with a GFP reporter fused to the
promoter region of a PhoQ/PhoP regulated gene (pUA66 P_{mgtLA}-*gfp*) in the presence or
absence of chemically synthesized MgrB[peri]. The positive control, chemically synthesized
SafA[peri], enhanced PhoQ activity significantly, consistent with the previous report [34]. In
contrast, MgrB[peri] did not show any inhibitory effect on PhoQ activity in the concentration
range that we tested, though DMSO the solvent for MgrB[peri] peptide showed a slight
inhibition of *gfp* reporter expression. As MgrB[peri] is short and likely capable of sampling
different conformations, our results suggest that, unlike SafA, the MgrB periplasmic region
alone is not sufficient to inhibit PhoQ. Taken together, the above results indicate that both
the MgrB TM and periplasmic regions are required but not sufficient to inhibit PhoQ.
213
214   **MgrB lacking its periplasmic region is capable of physical interaction with PhoQ**
215
Although MgrB-24 (N-terminal region+TM region) does not inhibit PhoQ, it could still play a
role in establishing physical contact with PhoQ. To explore this possibility, we used a

bacterial two-hybrid (BACTH) system based on split adenylyl cyclase [36, 37]. In the assay,
reconstitution of the T18 and T25 fragments restores the activity of adenylate cyclase
(CyaA), which leads to an increase in cAMP levels. This increased cAMP in turn enables
efficient induction of the genes required for maltose catabolism, resulting in pink colonies on
maltose MacConkey plates. To assay interactions of MgrB lacking its periplasmic region with
PhoQ, we prepared N-terminal fusions of MgrB-24 and PhoQ to T18 and T25 fragments,
respectively. As controls, we tested interactions between MgrB–PhoQ, MgrB–*Vc*ToxR
(*Vibrio cholerae* inner membrane protein ToxR) as well as MgrB–T25 fragment alone. Spot
assays of cells expressing T18-MgrB-24 and T25-PhoQ produced a dark pink color after 48
h at 30 °C, whereas cells expressing T18-MgrB-24 and the T25 fragment alone were
colorless (Fig 2D). Spots of cells expressing T18-MgrB-24 and either T25-MgrB or T25-
MgrB-24 were also pink. Control cells expressing T18-MgrB and either T25-PhoQ or T25-
MgrB formed dark pink spots, showing an interaction of MgrB with PhoQ and possibly with
itself, consistent with previous work [18], whereas control cells expressing either T18-MgrB
or T18-MgrB-24 with either T25-*Vc*ToxR or T25 fragment alone remained colorless. Taken
together, these results suggest that the MgrB N-terminus+TM region interacts with PhoQ
and may also interact with itself to facilitate formation of a multimeric complex.

235
236 **A small number of functionally important residues are spread across different regions**
237 **of MgrB**

238
To explore the relative importance of different residues in MgrB, we made alanine
substitutions and analyzed their effects on PhoQ inhibition. Interestingly, a majority of the
substitutions in MgrB did not affect PhoQ/PhoP reporter gene expression, but 11
substitutions spread across different regions showed significant activation of the reporter
gene (Figs 5, S4, and S5), indicating impaired MgrB activity. We confirmed that each of
these 11 MgrB variants was localized to the membrane with expression levels that were
comparable to that of wild-type MgrB (S1 and S2 Fig). Below we discuss these 11 MgrB
variants in more detail.

247
248 *a. Two lysines in the MgrB cytosolic region regulate PhoQ activity.*

249
In the short N-terminal cytosolic region of MgrB (M1 to R5), we made alanine substitutions at
each position except the first methionine. Here, we used the PhoQ/PhoP regulated GFP
reporter (pUA66 P*mgtLA*-*gfp*) to monitor the activities of MgrB variants. Control cells having no
MgrB expression showed a high intensity of GFP fluorescence, while cells expressing wild-
type MgrB had much lower reporter fluorescence (Fig 3A). The K2A and K3A substitutions
displayed the most effect on *gfp* reporter expression, with fluorescence comparable to the
no-MgrB control (Fig 3A). F4A and R5A had about 50% and 80% activity, respectively. K2A
and K3A, however, only retained about 20% activity (Fig 3A, see Materials and Methods for
the definition of normalized activity).

259
To further investigate whether this loss of activity might be caused by deficient protein
interactions, we used *in vivo* acceptor photobleaching FRET [38] to test complex formation
between these two MgrB variants and PhoQ. In this assay, MgrB variants and PhoQ were
tagged with a red fluorescent protein (mCherry) and a green fluorescent protein
(mNeonGreen) respectively and co-expressed in an *E. coli* Δ*phoQ* Δ*mgrB* strain. Complex
formation between tagged proteins typically results in close proximity and consequently
energy transfer between the donor mNeonGreen and the acceptor mCherry, leading to
quenching of the mNeonGreen fluorescence. The latter can be detected as an increase in
the mNeonGreen fluorescence upon photobleaching of mCherry that abolishes FRET (see
Materials and Methods). The apparent FRET efficiency could be defined as the resulting
increase in the donor fluorescence divided by its fluorescence after bleaching [39]. While
apparent FRET efficiencies may vary depending on the optical set-up, for intra-molecular
mNeonGreen-mCherry tandem constructs the reported values are in the range of 15-20%

[40]. We observed about 7.0% FRET efficiency for PhoQ-mNeonGreen/mCherry-MgrB WT,
whereas the negative control, PhoQ-mNeonGreen/mCherry, showed only about 1.5% FRET
efficiency. Interestingly, K2A and K3A mutants showed similar FRET efficiency as MgrB WT
(Fig 3C), suggesting that substitutions at these two positions do not significantly affect the
MgrB/PhoQ complex formation. Therefore, K2 and K3 may be instead directly involved in
regulating PhoQ conformation and promoting a kinase-inactive state of the enzyme.
279
We speculated that MgrB might regulate PhoQ conformation through the positively charged
side chains of K2 and K3. To test this hypothesis, we mutated K2 or K3 to a negatively
charged amino acid. Mutating K2 or K3 individually to aspartate only slightly reduced MgrB
function (S4 Fig A and B). Furthermore, when we substituted these two lysines individually
with 19 other amino acids, many of the mutants retained more than 50% activity (S4 Fig A
and B). Thus, neither K2 nor K3 alone is essential for MgrB function. We then tested the
effect of replacing both K2 and K3 with aspartate. MgrB K2DK3D localized to the cell
membrane similarly to wild-type MgrB (S1 Fig) but with some reduced expression (S2 Fig).
At this expression level, the K2DK3D mutant appeared to be a mild PhoQ activator rather
than an inhibitor and showed a 20% higher reporter activity compared to the no MgrB control
(Fig 3B). Increasing MgrB K2DK3D expression, either by doubling the amount of arabinose
inducer or by using a stronger promoter (*trc* promter), produced a small level of inhibition
that was much less than the level of inhibition of wild-type MgrB (S4 Fig C and D). Double
mutants, in which the lysines were replaced with either of the positively charged amino acids
histidine or arginine (K2HK3H and K2RK3R), on the other hand, retained about 60%
inhibitory activity (Fig 3B). FRET analyses also showed that MgrB K2DK3D had a FRET
efficiency comparable to that of wild type (Fig 3C), indicating that this MgrB variant forms
complexes with PhoQ. Taken together, our results suggest that the two lysines in the MgrB
cytosolic region regulate PhoQ function through their positively charged side chains without
contributing to PhoQ/MgrB complex formation in *E. coli*. We interpret the fact that single
amino acid substitutions of K2 or K3 to uncharged residues has little effect on activity as
indicating that the remaining lysine is sufficient to mediate the required interaction. We also
note that the K3R substitution was inactive (S4 Fig B), which suggests that it is not only the
charge of the lysine sidechains that are important for MgrB activity.
304
305 *b. The TM helix is involved in PhoQ/MgrB complex formation through W20.*
306
Given that the MgrB TM sequence is necessary for its function, we wanted to further identify
specific residues that might contribute to PhoQ inhibition. Polar amino acids such as N and
Q can facilitate TM domain interactions by forming stable hydrogen bonds [41]. Aromatic
residues are known to preferentially occur in the lipid-water interface regions [42, 43] and
could be important for membrane anchoring as well as surface interactions. An MgrB
sequence alignment identified four polar or aromatic residues in the transmembrane region
that are highly conserved: W6, C16, W20 and Q22. Based on these considerations, we
mutated the four highly conserved residues and F24 individually to alanine. The W6A and
C16A substitutions did not affect MgrB function (S5 Fig and Fig 4A); the result for C16A is
consistent with previous observations [20]. The F24A substitutions displayed modest effect
on reporter gene expression (~8-fold higher β-gal activity compared to WT MgrB) and the
change in activity for the Q22A mutant was not statistically significant (Fig 4A). The W20A
substitution, on the other hand, greatly affected MgrB function, with β-gal reporter activity
that was ~30-fold higher than that of WT MgrB (Fig 4A). These results were also reproduced
with the GFP reporter system and untagged MgrB (S5 Fig), except that the Q22A showed
significant activity loss (~70%) in this case. Collectively, the results from both approaches
indicate that W20, F24 and possibly Q22 in the TM region are functionally important.
324
Our bacterial two-hybrid data suggest that the MgrB TM region plays a role in establishing
physical contact with PhoQ. To pinpoint the essential residues in MgrB TM region that are
responsible for MgrB/PhoQ complex formation, we performed FRET analyses on MgrB TM

point mutants (Fig 4B). Compared to wild-type MgrB, which had 7.0% FRET efficiency,
Q22A and F24A showed slightly reduced FRET efficiency, 5.7% and 5.2%, respectively.
W20A, on the other hand, produced a much greater change, with FRET efficiency reduced
to 2.4%, which was comparable to the negative control. These results indicate that W20 in
the TM region is a key residue for MgrB/PhoQ complex formation.
333
334   *c. The periplasmic region has six functionally important residues.*
335
In the periplasmic region, mutants C28A and C39A showed the highest effects (80-90 fold
higher β-gal activity than WT MgrB), consistent with previous data indicating that these
cysteines are important for MgrB activity [20]. In addition, mutants G37A, D31A, F34A and
W47A displayed a significant increase in reporter gene expression – approximately 30-, 8-,
10-, and 10-fold relative to WT MgrB, respectively (Fig 5). These results were also
reproduced with GFP reporter system and untagged MgrB constructs (S5 Fig). The fact that
mutating residues C28, G37 and C39 showed the strongest effect on the reporter activity
suggests that these residues are likely important for structural stability of the MgrB
periplasmic region.
345
346   **A charged residue in the PhoQ TM2 – HAMP junction is involved in MgrB inhibition**
347
Our results suggest that the two N-terminal lysines in MgrB regulate PhoQ activity through
their charged side chains. To further explore which PhoQ residue(s) in PhoQ interact with
these lysines, we screened for PhoQ mutants for which MgrB K2DK3D showed increased
inhibitory activity. Based on the MgrB and PhoQ topologies, we hypothesized that K2 and K3
of MgrB interact with residues in the PhoQ TM2-HAMP region (Fig 6A). We therefore
mutated this region of PhoQ by error-prone PCR and screened for PhoQ mutants that have
reduced activity in the presence of MgrB K2DK3D. About 100 colonies with decreased green
fluorescent reporter protein were selected and the *phoQ* genes were sequenced. Mutants
with missense mutations are highlighted in the supplementary file (PhoQ_RM_screening).
Most of the missense mutations were clustered in/near TM2-HAMP junction and the second
helix of HAMP domain. We focused on the former, as residues in this region are more likely
to be close enough to interact with the lysines in MgrB. Among them, R219 (labeled in Fig
6A), which was the most frequent among identified mutations, appears to be the most
promising residue.
362
We therefore mutated PhoQ R219 to negatively charged amino acids (D and E) as well as
the amino acids with amide side chains (N and Q). Compared to wild-type PhoQ, R219D has
reduced activity (about 70% of WT), while the other three mutants (R219E, R219N and
R219Q) have activities that are either similar to or slightly greater than that of wild type (Fig
6B). Next, we tested the activity of these PhoQ mutants in the presence of MgrB K2DK3D.
Compared to wild-type PhoQ, R219D and R219E are indeed repressed further by MgrB
K2DK3D (Fig 6C), suggesting that reversing the side chain charge of PhoQ R219
compensates the effect of the charge inversion of MgrB N-terminal lysines. The inhibition of
PhoQ R219N and R219Q by MgrB K2DK3D, on the other hand, was comparable to the
inhibition of wild-type PhoQ, indicating that charge but not hydrogen bonding is important for
interactions of R219D and R219E with MgrB K2DK3D. We also found that wild-type MgrB
inhibited all four PhoQ R219 mutants to a similar extent as the inhibition of wild-type PhoQ
(Fig 6D), suggesting the existence of other factors involved in PhoQ inhibition by wild-type
MgrB.
377
378   **Inhibition of *E. coli* PhoQ by MgrB orthologs**
379
MgrB orthologs have previously been identified in several enterobacterial species [18]. To
examine the activities of these MgrB natural variants on *E. coli* PhoQ, we expressed MgrB
from *Klebsiella pneumoniae*, *Yesinia pestis*, *Photorhabdus laumondii*, *Proteus penneri*,

Providencia stuartii, *Serratia sp.MYb239*, *Enterobacter ludwigii* and *Salmonella enterica*
serovar typhimurium in *E. coli* K-12 MG1655. All orthologs except one (MgrB-*E. ludwigii*)
localized to the membrane (S1 Fig F). MgrB-*E. ludwigii*, which is missing the positively
charged residues (K2, K3) in its N-terminus showed defective localization, suggesting the
lack of a signal peptide. We found that most of the orthologs expressed at levels comparable
to that of *E. coli* MgrB, but the ortholog from *E. ludwigii* expressed quite poorly in *E. coli* (~5-
fold lower than *E. coli* MgrB see Fig 7). We then measured reporter gene expression for
PhoP-regulated transcription as a function of activity of each MgrB ortholog (Fig 7).
Unsurprisingly, *E. ludwigii* showed high reporter gene expression, indicating low PhoQ
inhibition. MgrB from *K. pneumoniae, S. sp.MYb239,* and *S. typhimurium* showed small but
significant increases in reporter activity (~4-fold, 2.5-fold, and 2-fold, respectively).
Interestingly, *Y. pestis*, *P. penneri* and *P.stuartii* MgrB produced reporter activities that were
significantly lower than that of *E. coli* MgrB, suggesting that they repress *E. coli* PhoQ more
strongly than does *E. coli* MgrB.
397
398 **MgrB interaction with non-cognate histidine kinases**
399
To test if MgrB is capable of physically interacting with histidine kinases other than PhoQ,
we made constructs of *E. coli* EnvZ, AtoS, CpxA, and PhoR fused to the T25 fragment of
adenylyl cyclase and performed BACTH analysis by spot assays as well as by measuring β-
gal activities of liquid cultures. Intriguingly, cells expressing T18-MgrB and either T25-AtoS, -
EnvZ, or -PhoR showed a high level of activity, although less than the β-gal activity of cells
expressing T18-MgrB and T25-PhoQ (Fig 8). Cells expressing T18-MgrB and T25-CpxA, on
the other hand, displayed a level of β-gal activity only marginally higher than that of the
negative controls. These results suggest that MgrB is capable of physically interacting with
EnvZ, AtoS, and PhoR, at least in the context of the bacterial two-hybrid assay. To test
whether MgrB affects the activity of these histidine kinases, we measured gene expression
from promoters regulated by these histidine kinases, in the presence or absence of plasmid
driven MgrB expression. However, we did not see any effect of MgrB on gene expression
controlled by any of these histidine kinases (data not shown). We also tested several other
histidine kinases: ArcB, BaeS, BasS, CpxA, CusS, EvgS, NtrB, RstB and TorS (S6 Fig).
None of them showed interactions with MgrB that were comparable to that of PhoQ.
415
416 Discussion
417
In this study, we have explored the regions of the small protein MgrB that are important for
its activity as an inhibitor of the PhoQ sensor kinase. Using a combination of biochemical
and biophysical approaches, we identified 11 amino acids in MgrB that, at least in the
context of alanine substitution, are important for the protein's inhibition of PhoQ . These
functionally important residues are spread across different regions of the molecule,
consistent with our observation that TM replacement or periplasmic region truncation leads
to a loss of MgrB function. The fact that the MgrB TM sequence is required for the protein's
activity distinguishes MgrB from at least two other bitopic membrane proteins that regulate
histidine kinases (SafA and MzrA); the TM domains of SafA and MzrA apparently serve
primarily as membrane anchors [31, 32]. In addition, our results indicate that the MgrB TM
region together with the small cytoplasmic region interacts with PhoQ, and that residue W20
in the TM region is essential for MgrB/PhoQ complex formation. Tryptophan is often found in
transmembrane proteins, especially at the ends of TM helixes as a membrane anchor due to
its unique amphipathic side chain [45, 46]. W20 in MgrB does not appear to position near the
computationally predicted membrane/periplasm interface. It is possible that the unique side
chain of W20 interacts with polar and/or aromatic residues in the PhoQ transmembrane
domain, such as the functionally important N202 [47]. The mechanism of how W20 mediates
PhoQ/MgrB complex formation awaits further investigations.
436

We identified two lysines in the MgrB cytosolic region that are important for the protein's
function. Although these residues are not individually essential, replacing both with
negatively charged residues significantly affected MgrB activity. A screen for PhoQ mutants
that increase the inhibitory activity of MgrB K2DK3D identified R219 in PhoQ. When this
residue was mutated to a negatively charged amino acid, the inhibitory activity of MgrB
K2DK3D was partially restored, suggesting the possibility that MgrB inhibitory activity of
PhoQ is in part mediated by charge-charge repulsion between K2K3 in MgrB and R219 in
the PhoQ TM2-HAMP junction. This hypothesis is consistent with the observation that K2
and K3 did not contribute to MgrB/PhoQ complex formation, since the HAMP domain is
conformationally dynamic and transduces signals via conformational changes [44, 48]. It is
also noteworthy that the TM2-HAMP junction has been reported to be essential for signaling
by other membrane proteins such as the chemoreceptor Tar [49]. In fact, some modifications
of the TM2-HAMP junction result in signal inversion in *E. coli* [49]. The partial suppression by
PhoQ R219D indicates the existence of other factors involved in PhoQ/MgrB interaction at
this residue. Mutating arginine to aspartate not only changes the charge but also the length
of the side chain (as well as other properties), which may have an impact on PhoQ/MgrB
interaction. Interestingly, we observed activity inversion (from activation to inhibition) of MgrB
K2DK3D when increasing its expression. Considering its reduced membrane localization
and possible heterogeneous cell population under arabinose induction, it is possible that the
mild activating activity we observed for lower expression level was due to factors other than
the direct effects of the mutant MgrB on PhoQ. Alternatively, different amounts of MgrB
K2DK3D might lead to different energy states or conformations of the PhoQ TM2-HAMP
junction, and thus changes in PhoQ activity. Lastly, we note that some MgrB orthologs have
a longer N-terminal cytosolic region and some do not have positive charged residues at
position 2 and 3. Due to the dynamic nature of the HAMP domain and the TM2-HAMP
junction, it is possible that these MgrB orthologs have different interactions with PhoQ.
463
Previous work indicates that the periplasmic region of MgrB and, in particular, two conserved
cysteines in this region, are critical for PhoQ inhibition [18, 20]. To identify other periplasmic
residues in MgrB that are required for PhoQ inhibition, we analyzed individual alanine
substitutions across the entire periplasmic region. Remarkably, despite the small size of this
region and the level of conservation, substitutions at most positions had little effect on MgrB
activity. Of the 22 residues in the periplasmic region, only six were individually important for
full activity: the two conserved cysteines (C28 and C39), as previously noted [20], as well as
G37, D31, F34 and W47. Furthermore, substitutions at these latter four residues lowered but
did not eliminate MgrB activity. Interestingly, mutations for each of these positions are also
associated with colistin-resistant *Klebsiella* isolates [23, 24, 50-55]. Taken together, our
findings show that the sequence determinants for MgrB activity are not restricted to either
one of the regions – cytosolic, periplasmic or TM, and that functionality is mediated by
collective interactions of key amino acid residues scattered across the polypeptide.
477
We also tested a number of MgrB orthologs derived from other enterobacteria with varying
sequence similarity to *E. coli* MgrB. Most orthologs that we tested efficiently inhibited *E. coli*
PhoQ. The slight decrease in activities of *K. pneumoniae* and *S. typhimurium* MgrB could be
due to the presence of a glycine residue within the TM regions, which may affect the
conformational flexibility of the TM helix and cause a defect in binding and inhibiting PhoQ.
On the other hand, *Y. pestis*, *P. penneri* and *P.stuartii* MgrB appear to be more efficient than
E. coli MgrB in inhibiting *E. coli* PhoQ. It is notable that the tryptophan residue (W20) found
in the TM region of *E. coli* MgrB is replaced by a tyrosine (Y20) in these orthologs. It is
possible that a hydrogen bond involving Y20 or the smaller size of tyrosine relative to
tryptophan affects the interaction of MgrB with PhoQ.
488
Unexpectedly, we also found that MgrB lacking its periplasmic region, as well as full-length
MgrB, interact with several other histidine kinases in addition to PhoQ in a bacterial two-
hybrid assay. However, we did not observe a significant effect of MgrB on gene expression

regulated by these histidine kinases. The TM domains of many dimeric histidine kinases
such as PhoQ and EnvZ are believed to form a four-helix bundle [48]. Our results may
reflect an interaction between the MgrB TM and a conserved conformation of the
transmembrane domain shared by multiple histidine kinases. The cytosolic and periplasmic
regions of MgrB apparently provide additional interactions required for specific regulation of
PhoQ. It will be interesting to determine if there are other small membrane proteins that have
evolved to interact with other histidine kinases using structural features similar to those of
the MgrB TM region. In addition, the results from our study of MgrB, as well as studies of
other small membrane proteins, may provide a platform for reengineering these proteins to
regulate new targets through rational design and directed evolution.
502
503   Materials and Methods
504
Strains, plasmids and cloning
See Supporting information for tables of strains (S1 Table), plasmids (S2 Table), and
primers (S3 Table) used in this study. All strains were derived from *E. coli* K-12 MG1655.
Gene deletions and reporter constructs were transferred between strains using transduction
with P1$_{vir}$. Deletions in the Keio collection [56] that were used for strain construction were
confirmed by PCR using primers flanking the targeted gene. Kanamycin resistance markers
were excised, when required, using the FLP-recombinase-expressing plasmid pCP20
[57]. Strain SAM72 was constructed by first excising the kanamycin resistance cassette from
the *ompC-mCherry* transcriptional reporter strain AFS256 to give SAM71. Kanamycin
resistance from the *mzrA* deletion strain JW3067 was then transduced into SAM71 by P1
transduction. A deletion of *safA* (SAM73) was constructed as described in [28, 57]. SAM73
was then used to construct SAM76 by transducing the *mgtA-lacZ* transcriptional reporter
from TIM199. Strain SAM85, which is a *lacI*^{q} containing version of BTH101 was made by
conjugation of the bacterial two-hybrid host strain BTH101 (F−) with XL1-Blue (F+). This
strain was used instead of BTH101 as it gave reduced background in our MacConkey plate
assays. Strain JingY90 was constructed by introducing *phoQ* deletion to JingY34 using P1
transduction followed by kanamycin resistance cassette removal. Plasmids expressing
gfpA206K fusions to *mgrB* mutants (pSY3-9, pSY72-75, pJC1-7, pTG1-15) were created by
either QuikChange site-directed mutagenesis (Stratagene) or Inverse PCR [58] using pAL38
as template. Plasmids expressing *gfpA206K* fusions of *mzrA* (pSY29)), *safA* (pSY31) and
mgrB orthologs (pSV1-8) were constructed by Gibson assembly [59].The gene fragments for
mzrA and *safA* were PCR amplified from the MG1655 genome, and gene fragments
corresponding to the different *mgrB* orthologs were synthesized commercially (Integrated
DNA Technologies, Inc.).Plasmids pSY32 (MzrA TM swap) and pSY33 (SafA TM swap)
were made by Inverse PCR using pSY29 and pSY31 as templates, respectively. Plasmid
pSY34 expressing T18-MgrB-24 only was created using pAL33 as template by Inverse PCR.
For all the plasmids mentioned above, see S3 Table for specific primer sequences used for
each construction. Plasmids pSY61, pSY64 and pSY68 were constructed as follows: *atoS*,
cpxA, and *phoR* were PCR-amplified using MG1655 genome as the template and the
following primer pairs – atoS-xbaI-U1/atoS-kpnI-L1, cpxA-xbaI-U1/cpxA-kpnI-L1 and phoR-
xbaI-U2/phoR-kpnI-L1, respectively, digested with XbaI/KpnI restriction enzymes and then
cloned into pKT25 at XbaI/KpnI sites. Plasmid pJY151 was constructed by inserting PCR
amplified *mgrB* to SacI/XbaI sites in pBAD33RBS vector. Plasmid pJY569 expressing
mcherry-mgrB was generated by inserting *mgrB* downstream of *mcherry* at BamHI/PstI sites
in pBO2 [60]. Plasmid pJY573 was generated by inserting *mNeonGreen* downstream of
phoQ at SalI/HindIII sites in pBAD33RBS phoQ [14]. Plasmids expressing *mgrB* variants
were constructed using Q5 site-directed mutagenesis kit (NEB). All constructs were
confirmed by DNA sequencing.
543
Media and growth conditions

Liquid cultures were grown at 37 °C with aeration, unless otherwise indicated, in either LB
Miller medium (Fisher Scientific) or in minimal A medium [61] containing 0.2% glucose, 0.1%
casamino acids and 1 mM $MgSO_4$. For routine growth on solid medium, LB or minimal
medium containing bacteriological grade agar (Fisher Scientific) was used. The antibiotics
ampicillin, kanamycin and chloramphenicol were used at concentrations 50–100, 25-50 and
20–34 µg mL^{-1}, respectively. The *lac* and *trc* promoters were induced with isopropyl *β*-D-1-
thiogalactopyranoside (IPTG) at a final concentration of 10 µM in FRET analysis or 1 mM
when indicated. When IPTG was not mentioned in the description of the culture conditions,
the basal transcription from the *trc* promoter was used to drive expression. The araBAD
promoter was induced with 0.008% arabinose. For bacterial two-hybrid screening,
MacConkey/maltose indicator plates were prepared as described [36] using MacConkey
agar base powder (Difco™), 100 µg mL^{-1} ampicillin, 25 µg mL^{-1} kanamycin, 0.5 mM IPTG
and 1% maltose.
558
β-galactosidase reporter gene assay
Cultures of strain AML67 carrying *gfpA206K* fusions of either wild-type *mgrB* (pAL38), *mgrB*
mutants (pSY3-9, pSY72-75, pJC1-7, pTG1-15), *mgrB* orthologs (pSV1-8) or *gfpA206K*-only
(pAL39) plasmids were grown overnight in LB with 50 µg mL^{-1} ampicillin. The next day, they
were diluted 1:1000 into the same medium and grown to OD$_{600}$ ~0.3-0.4 at 37 °C. Cells were
permeabilized with chloroform/SDS and assayed as in [61]. For high-throughput
measurements in a 96-well format, the protocol was modified as described in Thibodeau *et*
al.[62].
567
GFP reporter assay
E. coli imp-4213 strain carrying GFP reporter plasmid (pUA66 P$_{mgtLA}$-*gfp*) was grown
overnight at 37 °C in Luria–Bertani (LB) medium supplemented with 1 or 10 mM $MgSO_4$,
then diluted 1:100 to fresh $MgSO_4$ supplemented LB medium. DMSO, MgrB or SafA C-
terminal peptides (synthesized by Genscript) were added to the culture with indicated
amount. The cultures were grown at 37 °C with vigorous shaking for two hours. The
fluorescence of cells was then monitored with a BD LSRFortessa SORP flow cytometer (BD
Biosciences) and the acquired data were analyzed as described before [14]. To test the
function of MgrB variants, *E. coli* MG1655 *ΔmgrB* strain was transformed with GFP reporter
plasmid (pUA66 P$_{mgtLA}$-*gfp*) [14] and a pBAD33 vector carrying wild-type or mutant *mgrB*
genes. The resulting transformants were grown overnight at 37 °C in LB medium
supplemented with 10 mM $MgSO_4$, then diluted 1:100 to fresh LB medium supplemented
with 1 mM $MgSO_4$ and 0.008% arabinose. The cultures were grown at 37 °C with vigorous
shaking for two hours to reach early log phase (OD=0.4-0.5). The fluorescence of cells was
monitored and the acquired data were analyzed as described before [14]. The normalized
MgrB mutant activity was calculated as following:
Activity^{mut} = (Fluorescence$^{no\_MgrB\_control}$ − Fluorescence$^{MgrB\_mut}$) / (Fluorescence$^{no\_MgrB\_control}$ −
Fluorescence$^{MgrB\_WT}$) x 100%
Where appropriate, 50 µg mL^{-1} kanamycin and 34 µg mL^{-1} chloramphenicol was added to
the growth media.
588
Acceptor photobleaching FRET analysis
The expression of PhoQ-mNeonGreen and mCherry-MgrB was induced in *E. coli*
ΔphoQΔmgrB strain with 0.008% arabinose and 10 µM IPTG, respectively. The cultures
were grown to mid-log phase (OD=0.6) in LB media supplemented with 10 mM $MgSO_4$ and
appropriate antibiotics. Cells were harvested and washed two times with pre-chilled tethering
buffer (20 mM potassium phosphate pH7.0, 1 µM methionine, 10 mM lactic acid and 10 mM
magnesium sulfate). The surface of a glass-bottom 24-well plate (Greiner) was treated with
0.1% poly-L-lysine (Sigma) for 10 minutes at room temperature followed by rinsing with
tethering buffer. Cells were then added to the wells and incubated at room temperature for
10 minutes to allow attachment. Unattached cells were removed by washing two times and
attached cells were overlaid with tethering buffer afterwards.

600
Acceptor photobleaching FRET was performed using a dual layer Nikon Ti-E inverted
fluorescence microscope equipped with a fluorescence lamp (X-cite Exacte, Lumen
Dynamics), Perfect Focus System (PFS) and NIS-Elements AR software (version 4.40
Nikon). Images were acquired through a 40x Plan Apo NA 0.95 objective in the
mNeonGreen (donor) and mCherry (acceptor) channels, excitation power was adjusted both
by controlling the fluorescence lamp output and with ND filters. Specifically, the donor was
excited at 482/18 nm and its emission was detected at 525/50 nm and the acceptor was
excited at 585/29 while its emission was detected at 647/57. Excitation and emission filters
were mounted respectively on excitation and emission filter wheels, while beam splitters
(495LP for mNeonGreen excitation and 605LP for mCherry excitation) were on the lower
dichroic wheel of the microscope. Images were recorded using an iXon 897-X3 EM-CCD
camera (Andor), the acquisition time was set to 1 second for both channels and EM gain
was kept in the range of 100-280. Acceptor (mCherry) photobleaching was induced by a 150
mW DPSS 593 nm laser (ACAL Bfi) that was reflected on the sample by a zt594 DCRB laser
beam splitter (AHF) mounted on the upper dichroic wheel.
616
For each well we acquired two to three sequences of images on isolated sample areas with
the following protocol: (a) first, two images were taken in the acceptor channel; (b) then sixty
images were taken in the donor channel followed by (c) ten seconds acceptor
photobleaching (no image acquisition); afterwards, (d) forty images were taken in the donor
channel; and then (d) two images were taken in the acceptor channel. In order to avoid
donor fluorescence recovery from interrupted illumination, the sample was continuously
illuminated with a 482/18 nm excitation light, even during step (c). FRET efficiency was
calculated as the donor signal increase divided by the total donor signal after acceptor
photobleaching. In order to correct for the donor photobleaching present during steps (b) to
(d), we performed linear fitting (RStudio) of the donor fluorescence signal versus time for
both pre- and post-bleaching curves.
628
Bacterial two-hybrid assays
For the colorimetric spot assay, multiple clones from the transformation plate were
inoculated in 3 mL of LB containing 100 μg mL^{-1} ampicillin and 25 μg mL^{-1} kanamycin.
(Several clones were picked in order to reduce heterogeneity [36]. Cultures were grown
overnight at 30 °C with shaking. The next day, 3 μL of each culture was spotted on
MacConkey/maltose indicator plates and plates were then incubated at 30 °C. For each pair
of plasmids, triplicate experiments were performed. Color change on the indicator plates was
recorded after 48 h of incubation. Alternatively, *β*-gal assays were performed on the
overnight liquid cultures prepared as described for the colorimetric assay.
638
Random mutagenesis of the PhoQ TM2-HAMP region
The GeneMorph II random mutagenesis kit (Stratagene) was used to mutagenize the TM2-
HAMP-encoding region of *phoQ* (in pBAD33 *phoQ*) using primers that specifically target this
region. The library was transformed into an *E. coli ΔphoQΔmgrB* strain carrying a plasmid
GFP reporter (pUA66 *PmgtL-gfp*) and an MgrB K2DK3D expression plasmid (pTrc99a *mgrB*
K2DK3D). The transformants were grown on LB plates containing arabinose to induce PhoQ
expression. *E. coli* colonies with repressed PhoQ have weak green fluorescence. Dim
colonies were selected and their plasmids were sequenced to identify mutations in PhoQ.
Site-directed mutagenesis was performed using Q5 site-directed mutagenesis kit (NEB).
648

649 # Acknowledgements

651 We thank J. Zhu for the BACTH plasmid pKT25-toxR; S. Hoch and S. G. Sierra for technical
652 assistance; members of the Goulian, Sourjik, Zhu and Binns labs for helpful discussions; P.
653 Shah for a critical reading of the manuscript. This work was supported by NIH award

Author contributions

658
659 S.S.Y., J.N.C. V.S., J.Y. and M.G. designed the experiments. S.S.Y., J.Y., G.M., T.G., S.V.
660 and J.N.C. performed the experiments. B.N. contributed new reagents/analytical tools.
661 S.S.Y., T.G., S.V., J.N.C., G.M., V.S., J.Y. and M.G. analyzed the data. S.S.Y., J.Y., V.S.,
662 M.G. wrote the manuscript and all authors edited the manuscript.
663

664   Figures
665
666
667

668
669
670   **Fig 1. The conservation of MgrB sequence**
671   The sequence logo of MgrB generated with bacterial MgrB sequences obtained from KEGG
672   protein BLAST using *E. coli* MgrB as query highlights the degree of amino acid conservation.
673   The MgrB sequences and multiple sequence alignment can be found in the supplementary
674   file (MgrB_alignment).

Fig 2. The MgrB transmembrane domain and periplasmic domain are essential but not sufficient for its function as a PhoQ inhibitor.

A, Schematic representation of transmembrane (TM) domain swap constructs to replace the MgrB TM sequence with TM sequences derived from either the protein MalF or human glycophorin A, GpA. β-galactosidase activities of wild-type MgrB (pAL38), its variants MgrB-MalF TM (pSY3), MgrB-GpA TM (pSY4) and no MgrB control (pAL39) in a $\Delta mgrB$ strain containing a PhoQ/PhoP-regulated transcriptional reporter P_{mgtA}-*lacZ* (AML67) is shown below. Inducer was not added as the leaky expression from the *trc* promoter resulted in sufficient levels for strong inhibition by WT MgrB. Data represent averages and ranges for two independent experiments. MU = Miller Units. **B**, effect of wild-type MgrB and the truncation mutants were measured in a $\Delta mgrB$ strain containing the PhoQ/PhoP-regulated transcriptional reporter P_{mgtA}-*lacZ* (AML67) as in **A**. **C**, the activities of MgrB[Peri] and SafA[Peri] were monitored using PhoQ/PhoP-regulated transcriptional reporter P_{mgtLA}-*gfp*. Peptides or DMSO were added to the *E. coli imp4213* culture with indicated amount. The GFP activities were measured after 2h incubation using flow cytometry. Data points represent averages of three independent experiments and error bars show SDs. **D**, bacterial two-hybrid (BACTH) colorimetric spot assays with cells expressing protein fusions to the T18 and T25 subunits of *B. pertussis* adenylyl cyclase. Representative images showing interactions between MgrB-24 (T18-MgrB-24) and partner proteins PhoQ (T25-PhoQ) and MgrB (T25-MgrB). Interactions between MgrB-PhoQ, MgrB-MgrB, MgrB-T25 only and (MgrB-24)-T25 only were included as positive and negative controls. As an additional control, interactions were tested between MgrB-24 and *Vibrio cholerae* membrane protein ToxR (*Vc*ToxR), and also between MgrB and *Vc*ToxR . Each pair of interactions was assayed in at least three independent trials in a *cyaA*[-] BACTH host strain containing *lacI*[q] (SAM85), on maltose MacConkey indicator plates (results from one representative experiment are shown). The plates were incubated at 30°C for 48 h before images of plates were acquired. The pink color indicates reconstitution of adenylyl cyclase activity. The statistical significances of changes in reporter

735 activity were calculated with t tests by comparing to wild type (in A, B) or to the DMSO
736 negative control (in C) and indicated with asterisks (*** $P \le 0.001$, ** $P \le 0.01$, * $P \le 0.05$, ns
737 = not significant) in A, B and C.
738
739
740

741
742
Fig 3. Two lysines in MgrB cytosolic region regulate PhoQ activity through their charged side chain.
A, the activities of MgrB mutants with single alanine substitution in the cytosolic region were tested using the PhoQ/PhoP-regulated transcriptional reporter P$_{mgtLA}$-*gfp* in a Δ*mgrB* strain. MgrB mutants were constructed in a pBAD33 vector and their expression was induced with 0.008% arabinose. **B**, the activities of MgrB K2K3 double mutants were tested as in **A.** The reporter activity change was significant (*P* = 0.0045) when comparing MgrB K2DK3D to the no_MgrB control. **C**, FRET efficiencies were measured between PhoQ-mNeonGreen and mCherry (-), wild-type MgrB (WT) or MgrB mutants (K2A, K3A, K2DK3D and K2EK3E). MG1655 Δ*phoQ* Δ*mgrB* strain was transformed with pBAD33 *phoQ-mneongreen* and pTrc99a *mcherry-mgrB* variants. Protein expression was induced with 0.008% arabinose and 10 µM IPTG. FRET was measured for each protein pair and the efficiencies were calculated as described in Materials and Methods. All data points represent averages of at least three independent experiments and error bars show SDs. The statistical significances of changes in reporter activity or FRET efficiency were calculated with *t* tests by comparing to wild type or as specified and indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, ns = not significant).

MgrB: MKKFR<u>WVVLVVVVLA</u><u>CLLL</u><u>W</u>A<u>QVF</u>NMMCDQDVQFFSGICAINQFIPW
predicted TM region

Fig 4. W20 in MgrB transmembrane domain is essential for PhoQ/MgrB complex formation.

A, β-galactosidase levels were measured in a $\Delta mgrB$ strain containing the PhoQ/PhoP-regulated transcriptional reporter P$_{mgtA}$-*lacZ* (AML67) and also containing either an empty vector (pAL39), wild-type MgrB (pAL38), or the indicated mutants MgrB W20A (pSY72), Q22A (pSY73) and F24A (pSY74). Inducer (IPTG) was not added as the leaky expression from the *trc* promoter resulted in sufficient levels of protein. Mutants with β-galactosidase activity $\geq$ 5-fold above wild type were selected for further analysis. Data represent averages and ranges for two independent experiments. MU = Miller Units. **B**, FRET efficiencies were measured between PhoQ-mNeonGreen and indicated mCherry-MgrB variants as described in Figure 3C. When compared to wild type, the reduction of FRET efficiencies is significant according to *t* test (*** $P < 0.001$, ** $P < 0.01$). Data points represent averages of at least three independent experiments and error bars show SDs. The statistical significances of changes in reporter activity or FRET efficiency were calculated with *t* tests by comparing to wild type and indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, ns = not significant).

797
798
Fig 5. Effects of site-directed alanine substitutions of MgrB periplasmic residues on
PhoQ activity.
β-galactosidase levels were measured in a $\Delta mgrB$ strain containing the PhoQ/PhoP-
regulated transcriptional reporter P$_{mgtA}$-*lacZ* (AML67) and also containing either an empty
vector (pAL39), wild-type MgrB (pAL38), or the indicated mutants (pJC1 through pJC7 and
pTG1 through pTG15; see Table S2 for details). Inducer (IPTG) was not added as the leaky
expression from the *trc* promoter resulted in sufficient levels of protein. Data for wild-type
MgrB and each mutant was normalized to the no-MgrB control. Mutants with β-galactosidase
activity $\geq$ 5-fold above wild type were selected for further analysis. Averages and standard
deviations for three independent experiments are shown. MU = Miller Units. The statistical
significances of changes in reporter activity were calculated with *t* tests by comparing to wild
type and indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, ns = not significant).

822
823
Figure 6. PhoQ R219 is involved in interactions with MgrB K2 and K3.
Models of TM [14] and cytosolic domains [44] of PhoQ and MgrB are shown in (**A**). The
MgrB TM model was constructed *ab initio* via Rosetta (https://www.rosettacommons.org).
PhoQ R219 residue is labeled and shown with side chain in a zoomed-in view. **B**, the activity
of PhoQ R219 mutants was tested using the PhoQ/PhoP-regulated transcriptional reporter
P$_{mgtLA}$-*gfp* in a Δ*phoQ*Δ*mgrB* strain. PhoQ mutants were expressed from the plasmid
pBAD33 with 0.008% arabinose. The activities of PhoQ R219 mutants were tested in the
presence of MgrB K2DK3D (**C**) or wild type (**D**) using the same GFP reporter as in **B**. MgrB
variants were expressed from a pTrc99a vector with no inducer (IPTG) added, as the leaky
expression from the *trc* promoter resulted in sufficient levels of protein. Averages and
standard deviations for three independent experiments are shown in **B**-**D**. The statistical
significances of changes in reporter activity were calculated with *t* tests by comparing to wild
type and indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, ns = not significant).
837
838
839
840

A

```
Escherichia coli         ---MKKFRWVVLVVVVLACLL-LWAQVFNMMCDQDV-QFFSGICAINQFIPW---
Klebsiella pneumoniae     ---MKKLRWVLLIVIIAGCLL-LWTQMLNVMCDQDV-QFFSGICTINKFIPW---
Yersinia pestis           MLDLNITKLVTTVVIIAACCL-FYLLALDSYCDQGG-TFSTGICAITTIVPW---
Photorhabdus laumondii    ---MDIKKTIAILALLGICLF-LYLLALDSYCDQGE-QFDSGICSVTRYLPF---
Proteus penneri           ---MNAKKIIISLIIALVITFGLYLVALDNFCDRGE-DFQQGLCRFTTLFPSKHH
Providencia stuartii      ---M-KYKLIIGVCIVITCIF-LYLLAMDSLCDQGGENFIFGICQITDLLPF---
Serratia sp. MYb239       ---MLRNKILLIVATVLACLF-FNLLALDSYCDQGG-NFALGICSITRIIPW---
Enterobacter ludwigii     -----------MIIVLVACVV-LWTQTINVMCDQDV-QFFSGVCAINKFIPW---
Salmonella typhimurium    ---MKKFRWVVLGIVVVVCLL-LWAQVFNIMCDQDV-QFFSGICAINKFIPW---
                             . :       ::  **..    *  *:*  ..  .*
```

841
842
Figure 7. Effect of MgrB orthologs on *E. coli* PhoQ activity.
A, Sequence alignment of MgrB orthologs. **B**, β-galactosidase levels of a $\Delta mgrB$ strain
containing the PhoQ/PhoP-regulated transcriptional reporter P_{mgtA}-*lacZ* (AML67) and also
containing either an empty vector (pAL39), wild-type MgrB (pAL38), or a plasmid expressing
the indicated MgrB orthologs. Inducer (IPTG) was not added as the leaky expression from
the *trc* promoter resulted in sufficient levels of protein. Averages and standard deviations for
three independent experiments are shown. MU = Miller Units. The statistical significances of
changes in reporter activity were calculated with *t* tests by comparing to wild type and
indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, ns = not significant). **C**, GFP-
MgrB expression levels for cells expressing MgrB orthologs were imaged by fluorescence
microscopy, and GFP fluorescence levels were quantified (see Methods for details). Data
represent averages and ranges for two independent experiments comprised of 200 cells
each, AU = Arbitrary Units.
856
857
858

Figure 8. MgrB interacts with other histidine kinases in a bacterial two-hybrid assay.
Bacterial two-hybrid (BACTH) assays with cells expressing protein fusions to the T18 and
T25 subunits of *B. pertussis* adenylyl cyclase. β-gal expression from the *lac* promoter was
monitored as a measure of interactions between T18-MgrB and T25 fusion with either PhoQ,
AtoS, CpxA, EnvZ or PhoR. The T25 fragment alone and T25-*Vc*ToxR (*Vibrio cholerae*
ToxR) serve as negative controls. β-gal assays were carried out in a *cyaA*⁻ BACTH host
strain containing *lacI*�q (SAM85). Data represent averages and ranges for two independent
experiments. MU = Miller Units. The statistical significances of changes in reporter activity
were calculated with *t* tests by comparing to the negative control (T18-MgrB/T25 only) and
indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, ns = not significant).

 Supporting information

882
883 **S1 Text. Materials and methods**

884
Fluorescence microscopy
Strains were grown in minimal medium containing with antibiotics as needed and single-cell
measurements were performed essentially as described previously [21]. For MgrB
localization experiments, saturated overnight cultures of AML67 strains carrying *gfpA206K*
fusions of either wild-type *mgrB* (pAL38), *mgrB* mutants (pSY3-9, pSY72-75, pJC1-7, pTG1-
15) , *mgrB* orthologs (pSV1-8) or *gfpA206K*-only (pAL39) were diluted 1:1,000 into fresh
medium containing 100 μg mL^{-1} ampicillin and grown at 37 °C with aeration to an
OD$_{600}$ ~0.2–0.3 (~4-5 h) before visualization. For single-cell fluorescence measurements
from transcriptional reporter strains, cultures were grown as outlined above, then rapidly
cooled in an ice-water slurry and streptomycin was added to a final concentration of
250 μg mL^{-1} to stop protein synthesis. Microscope slides were prepared with 1% agarose
pads and cell fluorescence was measured by imaging and quantified with in-house software
as described previously [21, 63].

898
Membrane protein isolation and Western blot analysis
Cells expressing Flag tagged MgrB variants were harvested and resuspended in lysis buffer
(50mM Tris-Cl pH8.0, 100 mM NaCl and protease inhibitors). The membrane fraction was
collected as described before with modifications [34]. Specifically, cells were disrupted by
homogenization at 4 °C and cell debris were remove by centrifugation at 2,300 x g for 10
minutes. The lysate was centrifuged again at 21,000 x g for 60 minutes at 4 °C to pellet the
membrane fraction. The membrane pellet was washed with 1 ml lysis buffer and dissolved in
SDS sample buffer overnight at 4 °C. The samples were heated at 60 °C for 10 minutes and
the proteins were separated on a 16% tricine gel. The separated proteins were then
transferred to Immobilon-P^{SQ} PVDF membrane (Merckmillipore) in transfer buffer containing
20% methanol and Flag-flexilinker-MgrB was detected with anti-Flag primary antibody
(Sigma) and IRDye 800CW-conjugated secondary antibody (LI-COR). The protein bands
were visualized with the Odyssey CLx imaging system (LI-COR) and quantified with ImageJ.

912
913
914
915 **S1 Table. Strains**
916

Strain	Genotype	Source or Reference
AFS256	MG1655 λ$_{att}$::(P$_{ompC}$-*mCherry FRT-kan-FRT*)	Goulian lab collection
AML67	MG1655 Δ*mgrB*::(*FRT-kan-FRT*) Δ*lacZ* λ$_{att}$::(P$_{mgtA}$-*lacZ cat*)	[18]
BTH101	F-, *cya-99 araD139 galE15 galK16 rpsL1* (Str^r) *hsdR2 mcrA1 mcrB1*	Euromedex
BW25113	*rrnB3 ΔlacZ4787 hsdR514 Δ(araBAD)567 Δ(rhaBAD)568 rph-1*	[64]
JW3067	BW25113 Δ*mzrA*::(*FRT-kan-FRT*)	KEIO collection [56]
MG1655	F$^-$ λ$^-$ ilvG- rfb-50 rph-1	*E. coli* Genetic Stock Center, CGSC# 7740
SAM71	MG1655 λ$_{att}$::(P$_{ompC}$-*mCherry*)	This study
SAM72	MG1655 Δ*mzrA*::(*FRT-kan-FRT*) λ$_{att}$::(P$_{ompC}$-*mCherry*)	This study
SAM73	MG1655 Δ*safA*::(*FRT-kan-FRT*)	This study
SAM76	MG1655 Δ*safA*::(*FRT-kan-FRT*) Δ*lacZ* λ$_{att}$::(P$_{mgtA}$-*lacZ cat*)	This study
SAM85	F+ *lacI^{q}ZΔM15* Tn*10* (Tet^r), *cya-99 araD139 galE15 galK16 rpsL1* (Str^r) *hsdR2 mcrA1 mcrB1*. Strain prepared by conjugation: BTH101 (F-) x XL1-Blue (F+).	This study

TIM199	MG1655 Δ*lacZ* λ_att::(P_*mgtA*-*lacZ cat*)	[18]
XL1-Blue	*recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac* [F´ *proAB lacI*ᑫ*ZΔM15* Tn*10* (Tet^r)]	[65]
Imp-4213	F-, *lptD4213, galK2*(Oc), *λ*⁻, *IN(rrnD-rrnE)1, rpsL200*(strR), *rph-1*	[35]
JingY34	MG1655 Δ*mgrB::FRT*	[14]
JingY90	MG1655 Δ*phoQ::FRT* Δ*mgrB::FRT*	This study

917
918
919 **S2 Table. Plasmids**
920

Plasmid	Relevant genotype	Source or Reference
pAL25	pKT25 P_*lac*-*cyaA*_T25-*mgrB*, Kan^R	[18]
pAL27	pKT25 P_*lac*-*cyaA*_T25-*phoQ*, Kan^R	[18]
pAL30	pBAD33 P_*araBAD*-*phoQ(his)*_6, Cam^R	Goulian lab collection
pAL33	pUT18C P_*lac*-*cyaA*_T18-*mgrB*, Amp^R	[18]
pAL38	pEB52 P_*trc*-*gfpA206K-mgrB*, Amp^R	[18]
pAL39	pEB52 P_*trc*-*gfpA206K*, Amp^R	[18]
pBAD33	*ori*_p15A P_*araBAD*-MCS *araC*, Cam^R	ATCC 87402
pCP20	*λcI857(ts) repA101(ts) oriR101 bla cat λpR-FLP*, Amp^R Cam^R	[66]
pEB52	pTrc99a with the NcoI site removed, Amp^R	Goulian lab collection
pJC1	pEB52 P_*trc*-*gfpA206K-mgrB Q30A*, Amp^R	This study
pJC2	pEB52 P_*trc*-*gfpA206K-mgrB D29A*, Amp^R	This study
pJC3	pEB52 P_*trc*-*gfpA206K-mgrB F34A*, Amp^R	This study
pJC4	pEB52 P_*trc*-*gfpA206K-mgrB G37A*, Amp^R	This study
pJC5	pEB52 P_*trc*-*gfpA206K-mgrB I38A*, Amp^R	This study
pJC6	pEB52 P_*trc*-*gfpA206K-mgrB P46A*, Amp^R	This study
pJC7	pEB52 P_*trc*-*gfpA206K-mgrB W47A*, Amp^R	This study
pKD13	*oriR*_γR6k *bla FRT::kan::FRT*, Kan^R	[57]
pKD46	*repA101(ts) oriR101 bla ParaB-(gam bet exo)*, Amp^R	[57]
pKK14	pKT25 P_*lac*-*cyaA*_T25-*envZ*, Kan^R	Goulian lab collection
pKK15	pKT25 P_*lac*-*cyaA*_T25-*torS*, Kan^R	Goulian lab collection
pKT25	*ori*_p15A P_*lac*-*cyaA*_T25-MCS, Kan^R	Euromedex
pKT25-toxR	pKT25 P_*lac*-*cyaA*_T25-*Vibrio cholerae toxR*, Kan^R	[67]
pSV1	pEB52 P_*trc*-*gfpA206K-Kp_mgrB*, Amp^R	This study
pSV2	pEB52 P_*trc*-*gfpA206K-Yp_mgrB*, Amp^R	This study
pSV3	pEB52 P_*trc*-*gfpA206K-Pl_mgrB*, Amp^R	This study
pSV4	pEB52 P_*trc*-*gfpA206K-Pp_mgrB*, Amp^R	This study
pSV5	pEB52 P_*trc*-*gfpA206K-Ps_mgrB*, Amp^R	This study
pSV6	pEB52 P_*trc*-*gfpA206K-S sp.MYb239_mgrB*, Amp^R	This study
pSV7	pEB52 P_*trc*-*gfpA206K-El_mgrB*, Amp^R	This study
pSV8	pEB52 P_*trc*-*gfpA206K-St_mgrB*, Amp^R	This study
pSY3	pEB52 P_*trc*-*gfpA206K-mgrB-malF TM swap*, Amp^R	This study
pSY4	pEB52 P_*trc*-*gfpA206K-mgrB-gpA TM swap*, Amp^R	This study
pSY5	pEB52 P_*trc*-*gfpA206K-mgrB-24 (N25->stop)* truncation, Amp^R	This study
pSY6	pEB52 P_*trc*-*gfpA206K-mgrB-29 (Q30->stop)* truncation, Amp^R	This study

pSY7	pEB52 P_{trc}-*gfpA206K-mgrB-34 (F35->stop) truncation*, Amp^R	This study
pSY8	pEB52 P_{trc}-*gfpA206K-mgrB-39 (A40->stop) truncation*, Amp^R	This study
pSY9	pEB52 P_{trc}-*gfpA206K-mgrB-44 (I45->stop) truncation*, Amp^R	This study
pSY29	pEB52 P_{trc}-*gfpA206K-mzrA*, Amp^R	This study
pSY31	pEB52 P_{trc}-*gfpA206K-safA*, Amp^R	This study
pSY32	pEB52 P_{trc}-*gfpA206K-mzrA-gpA TM swap*, Amp^R	This study
pSY33	pEB52 P_{trc}-*gfpA206K-safA-gpA TM swap*, Amp^R	This study
pSY34	pUT18C P_{lac}-*cyaA*_{T18}-*mgrB-24* (MgrB truncation mutant lacking periplasmic domain), Amp^R	This study
pSY60	pKT25 P_{lac}-*cyaA*_{T25}-*arcB*, Kan^R	This study
pSY61	pKT25 P_{lac}-*cyaA*_{T25}-*atoS*, Kan^R	This study
pSY62	pKT25 P_{lac}-*cyaA*_{T25}-*baeS*, Kan^R	This study
pSY63	pKT25 P_{lac}-*cyaA*_{T25}-*basS*, Kan^R	This study
pSY64	pKT25 P_{lac}-*cyaA*_{T25}-*cpxA*, Kan^R	This study
pSY65	pKT25 P_{lac}-*cyaA*_{T25}-*cusS*, Kan^R	This study
pSY66	pKT25 P_{lac}-*cyaA*_{T25}-*evgS*, Kan^R	This study
pSY67	pKT25 P_{lac}-*cyaA*_{T25}-*ntrB*, Kan^R	This study
pSY68	pKT25 P_{lac}-*cyaA*_{T25}-*phoR*, Kan^R	This study
pSY69	pKT25 P_{lac}-*cyaA*_{T25}-*rstB*, Kan^R	This study
pSY70	pKT25 P_{lac}-*cyaA*_{T25}-*qseC*, Kan^R	This study
pSY72	pEB52 P_{trc}-*gfpA206K-mgrB W20A*, Amp^R	This study
pSY73	pEB52 P_{trc}-*gfpA206K-mgrB Q22A*, Amp^R	This study
pSY74	pEB52 P_{trc}-*gfpA206K-mgrB F24A*, Amp^R	This study
pSY75	pEB52 P_{trc}-*gfpA206K-mgrB Q22A N25A*, Amp^R	This study
pTG1	pEB52 P_{trc}-*gfpA206K-mgrB N25A*, Amp^R	This study
pTG2	pEB52 P_{trc}-*gfpA206K-mgrB M26A*, Amp^R	This study
pTG3	pEB52 P_{trc}-*gfpA206K-mgrB I45A*, Amp^R	This study
pTG4	pEB52 P_{trc}-*gfpA206K-mgrB C28A*, Amp^R	This study
pTG5	pEB52 P_{trc}-*gfpA206K-mgrB M27A*, Amp^R	This study
pTG6	pEB52 P_{trc}-*gfpA206K-mgrB D31A*, Amp^R	This study
pTG7	pEB52 P_{trc}-*gfpA206K-mgrB V32A*, Amp^R	This study
pTG8	pEB52 P_{trc}-*gfpA206K-mgrB Q33A*, Amp^R	This study
pTG9	pEB52 P_{trc}-*gfpA206K-mgrB F35A*, Amp^R	This study
pTG10	pEB52 P_{trc}-*gfpA206K-mgrB S36A*, Amp^R	This study
pTG11	pEB52 P_{trc}-*gfpA206K-mgrB C39A*, Amp^R	This study
pTG12	pEB52 P_{trc}-*gfpA206K-mgrB I41A*, Amp^R	This study
pTG13	pEB52 P_{trc}-*gfpA206K-mgrB N42A*, Amp^R	This study
pTG14	pEB52 P_{trc}-*gfpA206K-mgrB Q43A*, Amp^R	This study
pTG15	pEB52 P_{trc}-*gfpA206K-mgrB F44A*, Amp^R	This study
pTrc99a	*lacI*^q, P_{trc}-MCS, Amp^R	[68]
pUT18C	pUC19 P_{lac}-*cyaA*_{T18}-MCS, Amp^R	Euromedex
pJY31	pUA66 EC P_{mgtLA}-*gfp*, Kan^R	[14]
	pBAD33RBS, Cm^R	[14]
pJY151	pBAD33RBS *mgrB*, Cm^R	This study
pJY501	pBAD33RBS *mgrB K2A*, Cm^R	This study
pJY502	pBAD33RBS *mgrB K3A*, Cm^R	This study
pJY503	pBAD33RBS *mgrB F4A*, Cm^R	This study
pJY504	pBAD33RBS *mgrB R5A*, Cm^R	This study
pJY505	pBAD33RBS *mgrB W6A*, Cm^R	This study
pJY506	pBAD33RBS *mgrB C16A*, Cm^R	This study
pJY507	pBAD33RBS *mgrB W20A*, Cm^R	This study
pJY508	pBAD33RBS *mgrB Q22A*, Cm^R	This study
pJY509	pBAD33RBS *mgrB F24A*, Cm^R	This study

pJY510	pBAD33RBS *mgrB N25A*, Cm^R	This study
pJY511	pBAD33RBS *mgrB M26A*, Cm^R	This study
pJY512	pBAD33RBS *mgrB M27A*, Cm^R	This study
pJY513	pBAD33RBS *mgrB C28A*, Cm^R	This study
pJY514	pBAD33RBS *mgrB D29A*, Cm^R	This study
pJY515	pBAD33RBS *mgrB D31A*, Cm^R	This study
pJY516	pBAD33RBS *mgrB V32A*, Cm^R	This study
pJY517	pBAD33RBS *mgrB F34A*, Cm^R	This study
pJY518	pBAD33RBS *mgrB F35A*, Cm^R	This study
pJY519	pBAD33RBS *mgrB I38A*, Cm^R	This study
pJY520	pBAD33RBS *mgrB C39A*, Cm^R	This study
pJY521	pBAD33RBS *mgrB I41A*, Cm^R	This study
pJY522	pBAD33RBS *mgrB F44A*, Cm^R	This study
pJY523	pBAD33RBS *mgrB I45A*, Cm^R	This study
pJY524	pBAD33RBS *mgrB W47A*, Cm^R	This study
pJY525	pBAD33RBS *mgrB K2R*, Cm^R	This study
pJY526	pBAD33RBS *mgrB K2H*, Cm^R	This study
pJY527	pBAD33RBS *mgrB K2D*, Cm^R	This study
pJY528	pBAD33RBS *mgrB K2E*, Cm^R	This study
pJY529	pBAD33RBS *mgrB K2S*, Cm^R	This study
pJY530	pBAD33RBS *mgrB K2T*, Cm^R	This study
pJY531	pBAD33RBS *mgrB K2N*, Cm^R	This study
pJY532	pBAD33RBS *mgrB K2Q*, Cm^R	This study
pJY533	pBAD33RBS *mgrB K2C*, Cm^R	This study
pJY534	pBAD33RBS *mgrB K2P*, Cm^R	This study
pJY535	pBAD33RBS *mgrB K2G*, Cm^R	This study
pJY536	pBAD33RBS *mgrB K2V*, Cm^R	This study
pJY537	pBAD33RBS *mgrB K2L*, Cm^R	This study
pJY538	pBAD33RBS *mgrB K2I*, Cm^R	This study
pJY539	pBAD33RBS *mgrB K2M*, Cm^R	This study
pJY540	pBAD33RBS *mgrB K2F*, Cm^R	This study
pJY541	pBAD33RBS *mgrB K2Y*, Cm^R	This study
pJY542	pBAD33RBS *mgrB K2W*, Cm^R	This study
pJY543	pBAD33RBS *mgrB K3R*, Cm^R	This study
pJY544	pBAD33RBS *mgrB K3H*, Cm^R	This study
pJY545	pBAD33RBS *mgrB K3D*, Cm^R	This study
pJY546	pBAD33RBS *mgrB K3E*, Cm^R	This study
pJY547	pBAD33RBS *mgrB K3S*, Cm^R	This study
pJY548	pBAD33RBS *mgrB K3T*, Cm^R	This study
pJY549	pBAD33RBS *mgrB K3N*, Cm^R	This study
pJY550	pBAD33RBS *mgrB K3Q*, Cm^R	This study
pJY551	pBAD33RBS *mgrB K3C*, Cm^R	This study
pJY552	pBAD33RBS *mgrB K3P*, Cm^R	This study
pJY553	pBAD33RBS *mgrB K3G*, Cm^R	This study
pJY554	pBAD33RBS *mgrB K3V*, Cm^R	This study
pJY555	pBAD33RBS *mgrB K3L*, Cm^R	This study
pJY556	pBAD33RBS *mgrB K3I*, Cm^R	This study
pJY557	pBAD33RBS *mgrB K3M*, Cm^R	This study
pJY558	pBAD33RBS *mgrB K3F*, Cm^R	This study
pJY559	pBAD33RBS *mgrB K3Y*, Cm^R	This study
pJY560	pBAD33RBS *mgrB K3W*, Cm^R	This study
pJY561	pBAD33RBS *mgrB K2DK3D*, Cm^R	This study
pJY562	pBAD33RBS *mgrB K2HK3H*, Cm^R	This study

pJY563	pBAD33RBS *mgrB K2RK3R*, Cm^R	This study
pJY564	pBAD33RBS *flag*-flexilinker-*mgrB*, Cm^R	This study
pJY565	pBAD33RBS *flag*-flexilinker-*mgrB K2A*, Cm^R	This study
pJY566	pBAD33RBS *flag*-flexilinker-*mgrB K3A*, Cm^R	This study
pJY567	pBAD33RBS *flag*-flexilinker-*mgrB K2DK3D*, Cm^R	This study
pOB2	pTrc99A *mcherry*, Amp^R	[60]
pJY569	pTrc99A *mcherry-mgrB*, Amp^R	This study
pJY570	pTrc99A *mcherry-mgrB K2A*, Amp^R	This study
pJY571	pTrc99A *mcherry-mgrB K3A*, Amp^R	This study
pJY572	pTrc99A *mcherry-mgrB K2DK3D*, Amp^R	This study
pJY573	pBAD33RBS *phoQ-mneongreen*	This study

921
922 **S3 Table. Primers**
923

Primer	Sequence (5' -> 3')	Use
atoS-xbaI-U1	ACTCTAGAGATGCATTATATGAAGTGGATTTATC	Cloning of *cyaA*$_{T25}$-*atoS*
atoS-kpnI-L1	CATGGTACCCAGTCATACAGTCTGATTTCCCTG	
b1500-red-u1	ACTGATTAACGATTTTTAACGTTATCCGCTAAATAAACATATTTGAAATGATTCCGGGGATCCGTCGACC	Generation of Δ*safA*::(*FRT-kan-FRT*)
b1500-red-l1	ATTTCATATTTATAATTTGCTGTTTGTTTTCAGCCTTGCAAACTATTGATTGTAGGCTGGAGCTGCTTCG	
cpxA-xbaI-U1	ACTCTAGAGATGATAGGCAGCTTAACCG	Cloning of *cyaA*$_{T25}$-*cpxA*
cpxA-kpnI-L1	CATGGTACCCCTGCAAATGCGAAGTTTAACTC	
gfp-Cterm-L1	TTTGTATAGTTCATCCATGCCATGTGTAATCCCAG	Cloning of *gfpA206K-mzrA, -safA*
gfp-mzrA-U1	ACATGGCATGGATGAACTATACAAAATGCAAATACCTCGCATGTCGCTTC	Cloning of *gfpA206K-mzrA*
gfp-safA-U1	ACATGGCATGGATGAACTATACAAAATGCATGCGACCACAGTGAAAAACAAAATC	Cloning of *gfpA206K-safA*
GpA-TM-U1	GTGATTGGCACCATTCTGCTGATTAGCTATAACATGATGTGCGATCAGGATG	Cloning of *gfpA206K-mgrB-gpA TM swap*
GpA-TM-L1	GCCCGCCATCACGCCAAAAATAATCGCGGTAATCCATCGAAACTTTTTCATTTTGTATAG	
MalF-TM-U1	GTGGGTTACCTTGTTGTTTTAATGTACGCAAACATGATGTGCGATCAGGATG	Cloning of *gfpA206K-mgrB-malF TM swap*
MalF-TM-L1	CAGCAGGCCGAGCAGACCTAGCACTGACCACCATCGAAACTTTTTCATTTTGTATAG	
mgrB_stop-pAL38-U1	TGAACTAGAGTCGACCTGCAG	Cloning of *gfpA206K-mgrB-24, mgrB-29, mgrB-34, mgrB-39 and mgrB-44*
mgrB-N25stop-L1	GAATACCTGCGCCCAAAGC	Cloning of *gfpA206K-mgrB-24*, and T18-*mgrB*
mgrB-Q30stop-L1	ATCGCACATCATGTTGAATACCTG	Cloning of *gfpA206K-mgrB-29*

mgrB-F35stop-L1	AAATTGTACATCCTGATCGCACATC	Cloning of *gfpA206K-mgrB-34*
mgrB-A40stop-L1	ACAAATTCCGCTGAAAAATTGTACATCC	Cloning of *gfpA206K-mgrB-39*
mgrB-I45stop-L1	AAACTGGTTAATGGCACAAATTCCGC	Cloning of *gfpA206K-mgrB-44*
mgrB_stop-pAL33-U1	TGAATAAGTAGGGTACCGAGCTC	Cloning of T18-*mgrB*
mgrB W20A-U1	GCGCAGGTATTCAACATGATGTGC	Construction of *gfpA206K-mgrB W20A* in pAL38 by site-directed mutagenesis
mgrB W20A-L1	CGCAAGCAGCAAGCAAGCCAACAC	
mgrB Q22A-U1	GTATTCAACATGATGTGCGATCAGGATG	Construction of *gfpA206K-mgrB Q22A* in pAL38 by site-directed mutagenesis
mgrB Q22A-L1	CGCCGCCCAAAGCAGCAAGCAAGC	
mgrB F24A-U1	AACATGATGTGCGATCAGGATGTAC	Construction of *gfpA206K-mgrB F24A* in pAL38 by site-directed mutagenesis
mgrB F24A-L1	CGCTACCTGCGCCCAAAGCAGCAAG	
mgrB-N25A-U2	ATGATGTGCGATCAGGATGTAC	Cloning of *gfpA206K-mgrB N25A* in pAL38 by site-directed mutagenesis
mgrB-N25A-L2	CGCGAATACCTGCGCCCAAAGC	
mgrB_M26A_U1	GCGCAGGTATTCAACGCGATGTGCGATCAGGATG	Construction of *gfpA206K-mgrB M26A* in pAL38 by site-directed mutagenesis
mgrB_M26A_L1	CATCCTGATCGCACATCGCGTTGAATACCTGCGC	
mgrB_M27A_U1	GCAGGTATTCAACATGGCGTGCGATCAGGATGTAC	Construction of *gfpA206K-mgrB M27A* in pAL38 by site-directed mutagenesis
mgrB_M27A_L1	GTACATCCTGATCGCACGCCATGTTGAATACCTG	
mgrB_C28A_U1	AGGTATTCAACATGATGGCCGATCAGGATGTACAATTTTTC	Construction of *gfpA206K-mgrB C28A* in pAL38 by site-directed mutagenesis
mgrB_C28A_L1	AATTGTACATCCTGATCGGCCATCATGTTGAATACCT	
mgrB_D29A	AGGTATTCAACATGATGTGCGCTCAGGATGTACAATTTTTCAG	Construction of *gfpA206K-mgrB D29A* in pAL38 by site-directed mutagenesis
mgrB_D29A_as	CTGAAAAATTGTACATCCTGAGCGCACATCATGTTGAATACCT	
mgrB_Q30A	AGGTATTCAACATGATGTGCGATGCGGATGTACAATTTTTCAGCG	Construction of *gfpA206K-mgrB Q30A* in pAL38 by site-directed mutagenesis
mgrB_Q30A_as	CGCTGAAAAATTGTACATCCGCATCGCACATCATGTTGAATACCT	

mgrB_D31A_U1	ACATGATGTGCGATCAGGCGGTACAATTT TTCAGCGG	Construction of *gfpA206K-mgrB D31A* in pAL38 by site-directed mutagenesis
mgrB_D31A_L1	CCGCTGAAAAATTGTACCGCCTGATCGCA CATCATGT	
mgrB_V32A_U1	TGATGTGCGATCAGGATGCGCAATTTTTC AGCGGAATTTG	Construction of *gfpA206K-mgrB V32A* in pAL38 by site-directed mutagenesis
mgrB_V32A_L1	ATTCCGCTGAAAAATTGCGCATCCTGATC GCACATC	
mgrB_Q33A_U1	TGTGCGATCAGGATGTAGCATTTTTCAGC GGAATTTG	Construction of *gfpA206K-mgrB Q33A* in pAL38 by site-directed mutagenesis
mgrB_Q33A_L1	CAAATTCCGCTGAAAAATGCTACATCCTG ATCGCAC	
mgrB_F34A	GTGCGATCAGGATGTACAAGCTTTCAGC GGAATTTGTGCC	Construction of *gfpA206K-mgrB F34A* in pAL38 by site-directed mutagenesis
mgrB_F34A_as	GGCACAAATTCCGCTGAAAGCTTGTACAT CCTGATCGCAC	
mgrB_F35A_U1	TCAGGATGTACAATTTGCCAGCGGAATTT GTGCC	Construction of *gfpA206K-mgrB F35A* in pAL38 by site-directed mutagenesis
mgrB_F35A_L1	TGGCACAAATTCCGCTGGCAAATTGTACA TCCTG	
mgrB_S36A_U1	GATGTACAATTTTTCGCCGGAATTTGTGC CATTAAC	Construction of *gfpA206K-mgrB S36A* in pAL38 by site-directed mutagenesis
mgrB_S36A_L1	AATGGCACAAATTCCGGCGAAAAATTGTA CATCC	
mgrB_G37A	GGATGTACAATTTTTCAGCGCAATTTGTG CCATTAACCAGT	Construction of *gfpA206K-mgrB G37A* in pAL38 by site-directed mutagenesis
mgrB_G37A_as	ACTGGTTAATGGCACAAATTGCGCTGAAA AATTGTACATCC	
mgrB_I38A	AGGATGTACAATTTTTCAGCGGAGCTTGT GCCATTAACCAGTTTATCC	Construction of *gfpA206K-mgrB I38A* in pAL38 by site-directed mutagenesis
mgrB_I38A_as	GGATAAACTGGTTAATGGCACAAGCTCCG CTGAAAAATTGTACATCCT	
mgrB_C39A_U1	AATTTTTCAGCGGAATTGCTGCCATTAAC CAGTTTAT	Construction of *gfpA206K-mgrB C39A* in pAL38 by site-directed mutagenesis
mgrB_C39A_L1	ATAAACTGGTTAATGGCAGCAATTCCGCT GAAAAATTG	
mgrB_I41A_U1	TCAGCGGAATTTGTGCCGCTAACCAGTTT ATCCCG	Construction of *gfpA206K-mgrB I41A* in pAL38 by site-directed mutagenesis
mgrB_I41A_L1	ACGGGATAAACTGGTTAGCGGCACAAATT CCGCTG	
mgrB_N42A_U1	GGAATTTGTGCCATTGCCCAGTTTATCCC GTGG	Construction of *gfpA206K-mgrB N42A* in

mgrB_N42A_L1	CCACGGGATAAACTGGGCAATGGCACAA ATTCC	pAL38 by site-directed mutagenesis
mgrB_Q43A_U1	GAATTTGTGCCATTAACGCGTTTATCCCG TGGTGAAC	Construction of *gfpA206K-mgrB Q43A* in pAL38 by site-directed mutagenesis
mgrB_Q43A_L1	GTTCACCACGGGATAAACGCGTTAATGG CACAAATTC	
mgrB_F44A_U1	ATTTGTGCCATTAACCAGGCTATCCCGTG GTGAACTAGAG	Construction of *gfpA206K-mgrB F44A* in pAL38 by site-directed mutagenesis
mgrB_F44A_L1	TCTAGTTCACCACGGGATAGCCTGGTTAA TGGCACAAATTC	
mgrB-I45A-U2	CCGTGGTGAACTAGAGTCGAC	Construction of *gfpA206K-mgrB I45A* in pAL38 by site-directed mutagenesis
mgrB-I45A-L2	CGCAAACTGGTTAATGGCACAAATTCC	
mgrB(pAL38)_P46 A	GCCATTAACCAGTTTATCGCGTGGTGAAC TAGAGTC	Construction of *gfpA206K-mgrB P46A* in pAL38 by site-directed mutagenesis
mgrB(pAL38)_P46 A_as	GACTCTAGTTCACCACGCGATAAACTGGT TAATGGC	
mgrB(pAL38)_W47 A	CATTAACCAGTTTATCCCGGCGTGAACTA GAGTCGACCTG	Construction of *gfpA206K-mgrB W47A* in pAL38 by site-directed mutagenesis
mgrB(pAL38)_W47 A_as	CAGGTCGACTCTAGTTCACGCCGGGATA AACTGGTTAATG	
mzrA-GpA-TM-U1	GTGATTGGCACCATTCTGCTGATTAGCTA TCGCCAGCAAGAGTCTACGCTG	Cloning of *gfpA206K-mzrA-gpA TM swap*
mzrA-GpA-TM-L1	GCCCGCCATCACGCCAAAAATAATCGCG GTAATCTGGCGAAGCGACATGCGAG	
pEB52-U1	ACTAGAGTCGACCTGCAGGCATGCAAG	Cloning of *gfpA206K-mzrA, -safA*
pEB52-L1	GGGTACCGAGCTCGAATTCTCTGTTTC	
pEB52-gfp-U1	AACAGAGAATTCGAGCTCGGTACCC	
pEB52-mzrA-L1	TGCATGCCTGCAGGTCGACTCTAGTTTAT CCGAAGCGATGAGAGTTATCCC	Cloning of *gfpA206K-mzrA*
pEB52-safA-L1	TGCATGCCTGCAGGTCGACTCTAGTTCAG CCTTGCAAACTATTGATAATGAAATGTG	Cloning of *gfpA206K-safA*
phoR-xbaI-U2	ACTCTAGAGATGCTGGAACGGCTGTCGT G	Cloning of *cyaA*$_{T25}$*-phoR*
phoR-kpnI-L1	TGGCTGGTACCGATGACAAAGGCGGATT AATC	
safA-GpA-TM-U1	GTGATTGGCACCATTCTGCTGATTAGCTA TATTGATCAGCTGAGTATTTCAGAAATGG	Cloning of *gfpA206K-safA-gpA TM swap*
safA-GpA-TM-L1	GCCCGCCATCACGCCAAAAATAATCGCG GTAATTTCTTTATAGTTGTCTCTTTGCGTG ATTTTGTTTTTC	
MgrB K2A fw	CGAGCTCATGGCGAAGTTTCGATGGGTC	Construction of *mgrB K2A* by mutagenesis
MgrB K2A rv	AACTCCTCGCTAGCCCAA	
MgrB K3A fw	GCTCATGAAAGCGTTTCGATGGGTCGTTC	Construction of *mgrB K3A* by mutagenesis
MgrB K3A rv	TCGAACTCCTCGCTAGCC	
MgrB F4A fw	CATGAAAAAGGCGCGATGGGTCGTTCTG	Construction of *mgrB*

MgrB F4A rv	AGCTCGAACTCCTCGCTA	*F4A* by mutagenesis
MgrB R5A fw	GAAAAAGTTTGCGTGGGTCGTTCTGGTTGTC	Construction of *mgrB R5A* by mutagenesis
MgrB R5A rv	ATGAGCTCGAACTCCTCG	
MgrB W6A fw	AAAGTTTCGAGCGGTCGTTCTGGTTG	Construction of *mgrB W6A* by mutagenesis
MgrB W6A rv	TTCATGAGCTCGAACTCC	
MgrB W20A fw	CTTGCTGCTTGCGGCGCAGGTATTC	Construction of *mgrB W20A* in pJY151 by mutagenesis
MgrB W20A rv	CAAGCCAACACCACGACA	
MgrB F24A fw	GGCGCAGGTAGCGAACATGATGTGC	Construction of *mgrB F24A* in pJY151 by mutagenesis
MgrB F24A rv	CAAAGCAGCAAGCAAGCC	
MgrB M26A fw	GGTATTCAACGCGATGTGCGATCAGGATGTACAATTTTTC	Construction of *mgrB M26A* in pJY151 by mutagenesis
MgrB M26A rv	TGCGCCCAAAGCAGCAAG	
MgrB M27A fw	ATTCAACATGGCGTGCGATCAGGATGTACAATTTTTCAGC	Construction of *mgrB M27A* in pJY151 by mutagenesis
MgrB M27A rv	ACCTGCGCCCAAAGCAGC	
MgrB C28A fw	CAACATGATGGCCGATCAGGATGTACAATTTTTC	Construction of *mgrB C28A* in pJY151 by mutagenesis
MgrB C28A rv	AATACCTGCGCCCAAAGC	
MgrB D29A fw	CATGATGTGCGCGCAGGATGTACAATTTTTC	Construction of *mgrB D29A* in pJY151 by mutagenesis
MgrB D29A rv	TTGAATACCTGCGCCCAA	
MgrB D31A fw	GTGCGATCAGGCGGTACAATTTTTC	Construction of *mgrB D31A* in pJY151 by mutagenesis
MgrB D31A rv	ATCATGTTGAATACCTGC	
MgrB V32A fw	GATCAGGATGCGCAATTTTTCAGCG	Construction of *mgrB V32A* in pJY151 by mutagenesis
MgrB V32A rv	GCACATCATGTTGAATAC	
MgrBF34A fw	GGATGTACAAGCGTTCAGCGGAATTTG	Construction of *mgrB F34A* in pJY151 by mutagenesis
MgrB F34A rv	TGATCGCACATCATGTTG	
MgrB F35A fw	TGTACAATTTGCGAGCGGAATTTGTG	Construction of *mgrB F35A* in pJY151 by mutagenesis
MgrB F35A rv	TCCTGATCGCACATCATG	
MgrB I38A fw	TTTCAGCGGAGCGTGTGCCATTAAC	Construction of *mgrB I38A* in pJY151 by mutagenesis
MgrB I38A rv	AATTGTACATCCTGATCG	
MgrB C39A fw	CAGCGGAATTGCCGCCATTAACC	Construction of *mgrB C39A* in pJY151 by mutagenesis
MgrB C39A rv	AAAAATTGTACATCCTGATCG	
MgrB I41A fw	AATTTGTGCCGCGAACCAGTTTATCC	Construction of *mgrB I41A* in pJY151 by mutagenesis
MgrB I41A rv	CCGCTGAAAAATTGTACATC	
MgrB F44A fw	CATTAACCAGGCGATCCCGTGGTG	Construction of *mgrB F44A* in pJY151 by mutagenesis
MgrB F44A rv	GCACAAATTCCGCTGAAAAATTG	
MgrB I45A fw	TAACCAGTTTGCGCCGTGGTGATCTAG	Construction of *mgrB I45A* in pJY151 by mutagenesis
MgrB I45A rv	ATGGCACAAATTCCGCTG	
MgrB W47A fw	GTTTATCCCGGCGTGATCTAGAGTC	Construction of *mgrB W47A* in pJY151 by mutagenesis
MgrB W47A rv	TGGTTAATGGCACAAATTC	

MgrB K2R fw	CGAGCTCATGCGTAAGTTTCGATGGGTC	Construction of *mgrB K2X* in pBAD33RBS vector by site-directed mutagenesis
MgrB K2H fw	CGAGCTCATGCATAAGTTTCGATGGGTC	
MgrB K2D fw	CGAGCTCATGGATAAGTTTCGATGGG	
MgrB K2E fw	CGAGCTCATGGAAAAGTTTCGATG	
MgrB K2S fw	CGAGCTCATGAGCAAGTTTCGATGGG	
MgrB K2T fw	CGAGCTCATGACCAAGTTTCGATGGGTCG	
MgrB K2N fw	CGAGCTCATGAACAAGTTTCGATGG	
MgrB K2Q fw	CGAGCTCATGCAGAAGTTTCGATGGG	
MgrB K2C fw	CGAGCTCATGTGCAAGTTTCGATGGGTC	
MgrB K2P fw	CGAGCTCATGCCGAAGTTTCGATGGGTC	
MgrB K2G fw	CGAGCTCATGGGCAAGTTTCGATGGGTC	
MgrB K2V fw	CGAGCTCATGGTGAAGTTTCGATGGGTC	
MgrB K2L fw	CGAGCTCATGCTGAAGTTTCGATGGGTC	
MgrB K2I fw	CGAGCTCATGATTAAGTTTCGATGGGTC	
MgrB K2M fw	CGAGCTCATGATGAAGTTTCGATGGG	
MgrB K2F fw	CGAGCTCATGTTTAAGTTTCGATGGGTCG	
MgrB K2Y fw	CGAGCTCATGTATAAGTTTCGATGGGTC	
MgrB K2W fw	CGAGCTCATGTGGAAGTTTCGATGGGTC	
MgrB K3R fw	GCTCATGAAACGTTTTCGATGGGTCGTTCTGGTTGTC	Construction of *mgrB K3X* in pBAD33RBS vector by site-directed by mutagenesis
MgrB K3H fw	GCTCATGAAACATTTTCGATGGGTCGTTC	
MgrB K3D fw	GCTCATGAAAGATTTTCGATGGGTCGTTC	
MgrB K3E fw	GCTCATGAAAGAATTTCGATGGGTCG	
MgrB K3S fw	GCTCATGAAAAGCTTTCGATGGGTCGTTCTGGTTG	
MgrB K3T fw	GCTCATGAAACCTTTCGATGGGTCGTTCTGGTTGTC	
MgrB K3N fw	GCTCATGAAAAACTTTCGATGGGTCGTTC	
MgrB K3Q fw	GCTCATGAAACAGTTTCGATGGGTC	
MgrB K3C fw	GCTCATGAAATGCTTTCGATGGGTCGTTCTGGTTGTC	
MgrB K3P fw	GCTCATGAAACCGTTTCGATGGGTCGTTCTGG	
MgrB K3G fw	GCTCATGAAAGGCTTTCGATGGGTCGTTCTGGTTG	
MgrB K3V fw	GCTCATGAAAGTGTTTCGATGGGTCGTTC	
MgrB K3L fw	GCTCATGAAACTGTTTCGATGGGTCGTTCTGG	
MgrB K3I fw	GCTCATGAAAATTTTTCGATGGGTCGTTCTGGTTG	
MgrB K3M fw	GCTCATGAAAATGTTTCGATGGGTC	
MgrB K3F fw	GCTCATGAAATTTTTTCGATGGGTCGTTCTGGTTGTC	
MgrB K3Y fw	GCTCATGAAATATTTTCGATGGGTCGTTCTG	
MgrB K3W fw	GCTCATGAAATGGTTTCGATGGGTCGTTCTG	
MgrB K2DK3D fw	CGAGCTCATGGATGATTTTCGATGGGTCGTTC	Construction of *mgrB K2DK3D* by mutagenesis

MgrB K2RK3R fw	CGAGCTCATGCGTCGTTTTCGATGGGTCGTTCTGG	Construction of *mgrB K2RK3R* by mutagenesis
MgrB K2HK3H fw	CGAGCTCATGCATCATTTTCGATGGGTCGTTCTG	Construction of *mgrB K2HK3H* by mutagenesis
MgrB K2K3 rv	AACTCCTCGCTAGCCCAA	Construction of *mgrB K2XK3X* by mutagenesis
PhoQ TM2 fw	GAG CTA AAA AGT TCC TAT ATG GTC GAGCTAAAAAGTTCCTATATGGTC	Construction of PhoQ TM2-HAMP region random mutagenesis library
PhoQ HAMP rv	GGTGAGCGTCGTACGGTATTT	
PhoQ R219Dfw	GTGGAGTTTAgatCCCATCGAAGCCCTGGCAAAAGAAGTCCG	Construction of PhoQ R219X mutants by site-directed mutagenesis
PhoQ R219rv	CAGGCGGCGACCCACAGC	
PhoQ R219Efw	GTGGAGTTTAgaaCCCATCGAAGCCCTGGCAAAAGAAGTCCG	
PhoQ R219Nfw	GTGGAGTTTAaacCCCATCGAAGCCCTGGCAAAAGAAG	
PhoQ R219Qfw	GTGGAGTTTAcagCCCATCGAAGCCCTGGCAAAAG	

Supplementary Figures

S1 Fig. Visualization of cellular localization of MgrB and its variants by microscopy. **A**,
Representative fluorescence microscopy images of cells expressing wild-type MgrB, GFP-
only, and MgrB variants where the transmembrane region is swapped with transmembrane
domain from either human GpA or MalF, respectively. **B**, Representative green fluorescence
images of cells expressing C-terminal truncations of MgrB – MgrB-24 (TM only construct, no
periplasmic domain), MgrB-29, MgrB-34, MgrB-39 and MgrB-44. **C**, Representative
fluorescence microscopy images of cells expressing mCherry-only, mCherry tagged wild-
type MgrB and MgrB variants with indicated mutations. **D**, Representative green
fluorescence images of cells expressing TM region single Ala mutants of MgrB – W20A,
Q22A and F24A. **E**, Representative green fluorescence images of cells expressing
periplasmic single Ala mutants of MgrB – C28A, D31A, F34A, G37A, C39A and W47A. *E.*
coli MG1655 with *mgrB* deletion is the strain background in C and AML67 is the strain
background in A, B, D and E. **F**, Representative green fluorescence images of cells
expressing MgrB orthologs (EC- *E. coli*, ST- *Salmonella enterica serovar typhimurium*, EL-
Enterobacter ludwigii, SM- *Serratia sp.MYb239*, PS- *Providencia stuartii*, PP- *Proteus*
penneri, PL- *Photorhabdus laumondii*, YP- *Yesinia pestis* and KP- *Klebsiella pneumoniae*).
Inducer was not added as the leaky expression from the *trc* promoter resulted in sufficient
levels of protein except for constructs in panel C, where 10 µM IPTG was added.
947
948

949
950
S2 Fig. GFP-MgrB expression levels for MgrB variants relative to wild type.
GFP-MgrB expression levels for cells expressing (**A**) MgrB TM hybrids, (**B**) MgrB truncation
mutants and (**D**) the site-directed single alanine mutants were imaged by fluorescence
microscopy and GFP fluorescence levels were quantified (see Methods for details). Data
represent averages and ranges for two independent experiments comprised of at least 50
cells each, AU = Arbitrary Units. **C**, The expression level of the FLAG tagged wild-type and
MgrB mutants were monitored by western blot. A representative western blot is below. The
band intensities were quantified using ImageJ and normalized to wild-type MgrB after taking
account of total protein amount in each sample. Data represent averages of three
independent experiments and error bars show SDs.
961
962

963
964
S3 Fig. Transmembrane sequences of histidine kinase regulators, SafA and MzrA, are
dispensable for their function.
A, Activities of wild-type SafA and its variant where the transmembrane region is swapped
with transmembrane domain from human glycophorin A (GpA) measured as a function of β-
galactosidase levels in a $\Delta safA$ strain containing PhoQ/PhoP-regulated transcriptional
reporter P$_{mgtA}$-*lacZ* (SAM76), MU = Miller Units. **B**, Activities of wild-type MzrA and its variant
where the transmembrane region is swapped with transmembrane domain from human GpA
measured as a function of red fluorescence levels in a $\Delta mzrA$ strain containing EnvZ/OmpR-
regulated transcriptional reporter P$_{ompC}$-*mCherry* (SAM72), AU = Arbitrary Units. Inducer
(IPTG) was not added as the leaky expression from the *trc* promoter resulted in sufficient
levels of protein. Data represent averages and ranges for two independent experiments.
976
977

A

B

C   **D**

S4 Fig. **The activities of MgrB variants with mutations in the cytosolic region.**
The normalized activities of MgrB mutants with single alanine substitutions at K2 (**A**) and K3
(**B**) were tested using the PhoQ/PhoP-regulated transcriptional reporter P*mgtLA*-*gfp* in a
Δ*mgrB* strain (see Materials and methods for the definition of normalized activity). MgrB
mutants were constructed in a pBAD33 vector and their expression was induced with
0.008% arabinose. The activities of MgrB K2DK3D were tested as in **A** but with double the
amount of arabinose inducer (0.016%) (**C**) or with a different plasmid expression construct
(**D**). Specifically, in **D**, MgrB variants were expressed from the basal activity of the *trc*
promoter (i.e. without inducer) in the plasmid pTrc99a. Mutants with normalized MgrB
activity less than 0.4 were selected for further analysis. Data points represent averages of at
least three independent experiments and error bars show SDs. The statistical significances
of changes in reporter activity or normalized MgrB activity were calculated with *t* tests by
comparing to wild type and indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, ns
= not significant).

997
998
S5 Fig. **The normalized activities of MgrB variants with alanine substitutions at**
indicated positions.
The activities of MgrB mutants with single alanine substitution were tested using
PhoQ/PhoP-regulated transcriptional reporter P*mgtLA*-*gfp* in a Δ*mgrB* strain. MgrB mutants
were constructed in a pBAD33 vector and their expression was induced with 0.008%
arabinose. The normalized activity was calculated and plotted (see Materials and methods
for the definition of normalized activity). Mutants with normalized MgrB activity less than 0.4
(as indicated by the red dashed line) were selected for further analysis. All data points
represent averages of at least three independent experiments and error bars show SDs. The
statistical significances of changes in normalized MgrB activity were calculated with *t* tests
by comparing to wild type and indicated with asterisks (*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$,
ns = not significant).
1011

1012

1013

S6 Fig. **MgrB interacts with other histidine kinases in a bacterial two-hybrid assay.**

Bacterial two-hybrid (BACTH) assays with cells expressing protein fusions to the T18 and

T25 subunits of *B. pertussis* adenylyl cyclase. *β*-gal expression from the *lac* promoter was

monitored as a measure of interactions between T18-MgrB and T25 fusion with either ToxR,

ArcB, BaeS, CpxA, CusS, EvgS, NtrB, RstB and TorS. The statistical significances of

changes in reporter activity were calculated with *t* tests by comparing to the negative control

(T18-MgrB/T25 only) and indicated with asterisks (*** $P ≤ 0.001$, ** $P ≤ 0.01$, * $P ≤ 0.05$, ns =

not significant).

1022

1023

1024

1025

1026

1027

1028

1029

1030 References

1031

1.   Duval M, Cossart P. Small bacterial and phagic proteins: an updated view on a

rapidly moving field. Curr Opin Microbiol. 2017;39:81-8. doi: 10.1016/j.mib.2017.09.010.

PubMed PMID: 29111488.

2.   Storz G, Wolf YI, Ramamurthi KS. Small proteins can no longer be ignored. Annu

Rev Biochem. 2014;83:753-77. doi: 10.1146/annurev-biochem-070611-102400. PubMed

PMID: 24606146; PubMed Central PMCID: PMCPMC4166647.

3.   VanOrsdel CE, Kelly JP, Burke BN, Lein CD, Oufiero CE, Sanchez JF, et al.

Identifying New Small Proteins in Escherichia coli. Proteomics. 2018;18(10):e1700064. doi:

10.1002/pmic.201700064. PubMed PMID: 29645342; PubMed Central PMCID:

PMCPMC6001520.

4.   Weaver J, Mohammad F, Buskirk AR, Storz G. Identifying Small Proteins by

Ribosome Profiling with Stalled Initiation Complexes. MBio. 2019;10(2). doi:

10.1128/mBio.02819-18. PubMed PMID: 30837344; PubMed Central PMCID:

PMCPMC6401488.

5.   Nakahigashi K, Takai Y, Kimura M, Abe N, Nakayashiki T, Shiwa Y, et al.

Comprehensive identification of translation start sites by tetracycline-inhibited ribosome

profiling. DNA Res. 2016;23(3):193-201. doi: 10.1093/dnares/dsw008. PubMed PMID:
27013550; PubMed Central PMCID: PMCPMC4909307.
6.   Carol Smith JGC, Jing Wang, Keith M. Derbyshire, Todd A. Gray, Joseph T. Wade.
Pervasive translation in Mycobacterium tuberculosis. bioRxiv. 2019. doi:
https://doi.org/10.1101/665208.
7.   Sberro H, Fremin BJ, Zlitni S, Edfors F, Greenfield N, Snyder MP, et al. Large-Scale
Analyses of Human Microbiomes Reveal Thousands of Small, Novel Genes. Cell.
2019;178(5):1245-59 e14. doi: 10.1016/j.cell.2019.07.016. PubMed PMID: 31402174.
8.   Hemm MR, Paul BJ, Schneider TD, Storz G, Rudd KE. Small membrane proteins
found by comparative genomics and ribosome binding site models. Mol Microbiol.
2008;70(6):1487-501. doi: 10.1111/j.1365-2958.2008.06495.x. PubMed PMID: 19121005;
PubMed Central PMCID: PMCPMC2614699.
9.   Stock AM, Robinson VL, Goudreau PN. Two-Component Signal Transduction. Annu
Rev Biochem. 2000;69(1):183-215. doi:
papers2://publication/doi/10.1146/annurev.biochem.69.1.183.
10.   Groisman EA. The Pleiotropic Two-Component Regulatory System PhoP-PhoQ. J
Bacteriol. 2001;183(6):1835-42. doi: papers2://publication/doi/10.1128/JB.183.6.1835-
1842.2001.
11.   Garcia Vescovi E, Soncini FC, Groisman EA. Mg^{2+} as an extracellular signal:
environmental regulation of *Salmonella* virulence. Cell. 1996;84(1):165-74. PubMed PMID:
8548821.
12.   Prost LR, Daley ME, Le Sage V, Bader MW, Le Moual H, Klevit RE, et al.
Activation of the bacterial sensor kinase PhoQ by acidic pH. Mol Cell. 2007;26(2):165-74.
doi: 10.1016/j.molcel.2007.03.008. PubMed PMID: 17466620.
13.   Bader MW, Sanowar S, Daley ME, Schneider AR, Cho U, Xu W, et al. Recognition
of antimicrobial peptides by a bacterial sensor kinase. Cell. 2005;122(3):461-72. doi:
10.1016/j.cell.2005.05.030. PubMed PMID: 16096064.
14.   Yuan J, Jin F, Glatter T, Sourjik V. Osmosensing by the bacterial PhoQ/PhoP two-
component system. Proc Natl Acad Sci U S A. 2017;114(50):E10792-E8. doi:
10.1073/pnas.1717272114. PubMed PMID: 29183977; PubMed Central PMCID:
PMCPMC5740661.
15.   Goulian M. Two-component signaling circuit structure and properties. Curr Opin
Microbiol. 2010;13(2):184-9. doi: papers2://publication/doi/10.1016/j.mib.2010.01.009.
16.   Wang H, Yin X, Wu Orr M, Dambach M, Curtis R, Storz G. Increasing intracellular
magnesium levels with the 31-amino acid MgtS protein. Proc Natl Acad Sci U S A.
2017;114(22):5689-94. doi: 10.1073/pnas.1703415114. PubMed PMID: 28512220; PubMed
Central PMCID: PMCPMC5465934.
17.   Alix E, Blanc-Potard AB. Peptide-assisted degradation of the Salmonella MgtC
virulence factor. EMBO J. 2008;27(3):546-57. doi: 10.1038/sj.emboj.7601983. PubMed
PMID: 18200043; PubMed Central PMCID: PMCPMC2241655.
18.   Lippa AM, Goulian M. Feedback inhibition in the PhoQ/PhoP signaling system by a
membrane peptide. PLoS Genet. 2009;5(12):e1000788. doi: 10.1371/journal.pgen.1000788.
PubMed PMID: 20041203; PubMed Central PMCID: PMCPMC2789325.
19.   Salazar ME, Podgornaia AI, Laub MT. The small membrane protein MgrB regulates
PhoQ bifunctionality to control PhoP target gene expression dynamics. Mol Microbiol. 2016.
doi: 10.1111/mmi.13471. PubMed PMID: 27447896.
20.   Lippa AM, Goulian M. Perturbation of the oxidizing environment of the periplasm
stimulates the PhoQ/PhoP system in Escherichia coli. J Bacteriol. 2012;194(6):1457-63. doi:
10.1128/JB.06055-11. PubMed PMID: 22267510; PubMed Central PMCID:
PMCPMC3294871.

21.   Yadavalli SS, Carey JN, Leibman RS, Chen AI, Stern AM, Roggiani M, et al.
Antimicrobial peptides trigger a division block in Escherichia coli through stimulation of a
signalling system. Nat Commun. 2016;7:12340. doi: 10.1038/ncomms12340. PubMed
PMID: 27471053; PubMed Central PMCID: PMCPMC4974570.
22.   Kidd TJ, Mills G, Sa-Pessoa J, Dumigan A, Frank CG, Insua JL, et al. A Klebsiella
pneumoniae antibiotic resistance mechanism that subdues host defences and promotes
virulence. EMBO Mol Med. 2017;9(4):430-47. doi: 10.15252/emmm.201607336. PubMed
PMID: 28202493; PubMed Central PMCID: PMCPMC5376759.
23.   Cannatelli A, Giani T, D'Andrea MM, Di Pilato V, Arena F, Conte V, et al. MgrB
inactivation is a common mechanism of colistin resistance in KPC-producing Klebsiella
pneumoniae of clinical origin. Antimicrob Agents Chemother. 2014;58(10):5696-703. doi:
papers2://publication/doi/10.1128/AAC.03110-14.
24.   Olaitan AO, Morand S, Rolain JM. Mechanisms of polymyxin resistance: acquired
and intrinsic resistance in bacteria. Front Microbiol. 2014;5:643. Epub 2014/12/17. doi:
10.3389/fmicb.2014.00643. PubMed PMID: 25505462; PubMed Central PMCID:
PMCPMC4244539.
25.   Poirel L, Jayol A, Bontron S, Villegas MV, Ozdamar M, Turkoglu S, et al. The mgrB
gene as a key target for acquired resistance to colistin in Klebsiella pneumoniae. J
Antimicrob Chemother. 2015;70(1):75-80. doi: 10.1093/jac/dku323. PubMed PMID:
25190723.
26.   Eguchi Y, Itou J, Yamane M, Demizu R, Yamato F, Okada A, et al. B1500, a small
membrane protein, connects the two-component systems EvgS/EvgA and PhoQ/PhoP in
Escherichia coli. Proc Natl Acad Sci USA. 2007;104(47):18712-7. doi:
papers2://publication/doi/10.1073/pnas.0705768104.
27.   Eguchi Y, Ishii E, Hata K, Utsumi R. Regulation of acid resistance by connectors of
two-component signal transduction systems in Escherichia coli. J Bacteriol.
2011;193(5):1222-8. doi: 10.1128/JB.01124-10. PubMed PMID: 21193607; PubMed Central
PMCID: PMCPMC3067605.
28.   Roggiani M, Yadavalli SS, Goulian M. Natural variation of a sensor kinase
controlling a conserved stress response pathway in Escherichia coli. PLoS Genet.
2017;13(11):e1007101. Epub 2017/11/16. doi: 10.1371/journal.pgen.1007101. PubMed
PMID: 29140975; PubMed Central PMCID: PMCPMC5706723.
29.   Burton NA, Johnson MD, Antczak P, Robinson A, Lund PA. Novel aspects of the
acid response network of E. coli K-12 are revealed by a study of transcriptional dynamics.
Journal of Molecular Biology. 2010. doi: papers2://publication/uuid/EE91AF7A-5760-40B4-
B549-BE44E24F43E7.
30.   Gerken H, Charlson ES, Cicirelli EM, Kenney LJ, Misra R. MzrA: a novel modulator
of the EnvZ/OmpR two-component regulon. Mol Microbiol. 2009;72(6):1408-22. doi:
papers2://publication/doi/10.1111/j.1365-2958.2009.06728.x.
31.   Ishii E, Eguchi Y, Utsumi R. Mechanism of activation of PhoQ/PhoP two-component
signal transduction by SafA, an auxiliary protein of PhoQ histidine kinase in Escherichia coli.
Biosci Biotechnol Biochem. 2013;77(4):814-9. doi:
papers2://publication/doi/10.1271/bbb.120970.
32.   Gerken H, Misra R. MzrA-EnvZ interactions in the periplasm influence the
EnvZ/OmpR two-component regulon. J Bacteriol. 2010. doi:
papers2://publication/uuid/0796C281-CF57-426D-AB5B-52D4AAA89C30.
33.   Leeds JA, Beckwith J. Lambda repressor N-terminal DNA-binding domain as an
assay for protein transmembrane segment interactions in vivo. J Mol Biol. 1998;280(5):799-
810. Epub 1998/07/22. doi: 10.1006/jmbi.1998.1893. PubMed PMID: 9671551.

34.   Eguchi Y, Ishii E, Yamane M, Utsumi R. The connector SafA interacts with the
multi-sensing domain of PhoQ in Escherichia coli. Mol Microbiol. 2012;85(2):299-313. doi:
10.1111/j.1365-2958.2012.08114.x. PubMed PMID: 22651704.
35.   Sampson BA, Misra R, Benson SA. Identification and characterization of a new gene
of Escherichia coli K-12 involved in outer membrane permeability. Genetics.
1989;122(3):491-501. PubMed PMID: 2547691; PubMed Central PMCID:
PMCPMC1203724.
36.   Battesti A, Bouveret E. The bacterial two-hybrid system based on adenylate cyclase
reconstitution in Escherichia coli. Methods. 2012;58(4):325-34. doi:
papers2://publication/doi/10.1016/j.ymeth.2012.07.018.
37.   Karimova G, Pidoux J, Ullmann A, Ladant D. A bacterial two-hybrid system based on
a reconstituted signal transduction pathway. Proc Natl Acad Sci U S A. 1998;95(10):5752-6.
Epub 1998/05/20. PubMed PMID: 9576956; PubMed Central PMCID: PMCPMC20451.
38.   Kentner D, Sourjik V. Dynamic map of protein interactions in the Escherichia coli
chemotaxis pathway. Mol Syst Biol. 2009;5:238. doi: 10.1038/msb.2008.77. PubMed PMID:
19156130; PubMed Central PMCID: PMCPMC2644175.
39.   Sourjik V, Vaknin A, Shimizu TS, Berg HC. In vivo measurement by FRET of
pathway activity in bacterial chemotaxis. Methods Enzymol. 2007;423:365-91. doi:
10.1016/S0076-6879(07)23017-4. PubMed PMID: 17609141.
40.   Meiresonne NY, van der Ploeg R, Hink MA, den Blaauwen T. Activity-Related
Conformational Changes in d,d-Carboxypeptidases Revealed by In Vivo Periplasmic Forster
Resonance Energy Transfer Assay in Escherichia coli. MBio. 2017;8(5). doi:
10.1128/mBio.01089-17. PubMed PMID: 28900026; PubMed Central PMCID:
PMCPMC5596342.
41.   Fink A, Sal-Man N, Gerber D, Shai Y. Transmembrane domains interactions within
the membrane milieu: principles, advances and challenges. Biochim Biophys Acta.
2012;1818(4):974-83. Epub 2011/12/14. doi: 10.1016/j.bbamem.2011.11.029. PubMed
PMID: 22155642.
42.   von Heijne G. Membrane-protein topology. Nature Reviews Molecular Cell Biology.
2006;7:909. doi: 10.1038/nrm2063
https://www.nature.com/articles/nrm2063 - supplementary-information.
43.   Ulmschneider MB, Sansom MS. Amino acid distributions in integral membrane
protein structures. Biochim Biophys Acta. 2001;1512(1):1-14. Epub 2001/05/04. PubMed
PMID: 11334619.
44.   Matamouros S, Hager KR, Miller SI. HAMP Domain Rotation and Tilting
Movements Associated with Signal Transduction in the PhoQ Sensor Kinase. MBio.
2015;6(3):e00616-15. doi: 10.1128/mBio.00616-15. PubMed PMID: 26015499; PubMed
Central PMCID: PMCPMC4447245.
45.   Yau WM, Wimley WC, Gawrisch K, White SH. The preference of tryptophan for
membrane interfaces. Biochemistry. 1998;37(42):14713-8. doi: 10.1021/bi980809c. PubMed
PMID: 9778346.
46.   de Jesus AJ, Allen TW. The role of tryptophan side chains in membrane protein
anchoring and hydrophobic mismatch. Biochim Biophys Acta. 2013;1828(2):864-76. doi:
10.1016/j.bbamem.2012.09.009. PubMed PMID: 22989724.
47.   Goldberg SD, Clinthorne GD, Goulian M, DeGrado WF. Transmembrane polar
interactions are required for signaling in the Escherichia coli sensor kinase PhoQ. Proc Natl
Acad Sci U S A. 2010;107(18):8141-6. doi: 10.1073/pnas.1003166107. PubMed PMID:
20404199; PubMed Central PMCID: PMCPMC2889538.
48.   Bhate MP, Molnar KS, Goulian M, DeGrado WF. Signal transduction in histidine
kinases: insights from new structures. Structure. 2015;23(6):981-94. Epub 2015/05/20. doi:

10.1016/j.str.2015.04.002. PubMed PMID: 25982528; PubMed Central PMCID:
PMCPMC4456306.
49.   Bi S, Jin F, Sourjik V. Inverted signaling by bacterial chemotaxis receptors. Nat
Commun. 2018;9(1):2927. doi: 10.1038/s41467-018-05335-w. PubMed PMID: 30050034;
PubMed Central PMCID: PMCPMC6062612.
50.   Olaitan AO, Diene SM, Kempf M, Berrazeg M, Bakour S, Gupta SK, et al.
Worldwide emergence of colistin resistance in Klebsiella pneumoniae from healthy humans
and patients in Lao PDR, Thailand, Israel, Nigeria and France owing to inactivation of the
PhoP/PhoQ regulator mgrB: an epidemiological and molecular study. Int J Antimicrob
Agents. 2014;44(6):500-7. Epub 2014/09/30. doi: 10.1016/j.ijantimicag.2014.07.020.
PubMed PMID: 25264127.
51.   Cheng YH, Lin TL, Pan YJ, Wang YP, Lin YT, Wang JT. Colistin resistance
mechanisms in Klebsiella pneumoniae strains from Taiwan. Antimicrob Agents Chemother.
2015;59(5):2909-13. Epub 2015/02/19. doi: 10.1128/AAC.04763-14. PubMed PMID:
25691646; PubMed Central PMCID: PMCPMC4394772.
52.   Bakthavatchalam YD, Pragasam AK, Biswas I, Veeraraghavan B. Polymyxin
susceptibility testing, interpretative breakpoints and resistance mechanisms: An update. J
Glob Antimicrob Resist. 2018;12:124-36. Epub 2017/10/01. doi: 10.1016/j.jgar.2017.09.011.
PubMed PMID: 28962863.
53.   Nordmann P, Jayol A, Poirel L. A Universal Culture Medium for Screening
Polymyxin-Resistant Gram-Negative Isolates. J Clin Microbiol. 2016;54(5):1395-9. Epub
2016/03/18. doi: 10.1128/JCM.00446-16. PubMed PMID: 26984971; PubMed Central
PMCID: PMCPMC4844728.
54.   Bonura C, Giuffre M, Aleo A, Fasciana T, Di Bernardo F, Stampone T, et al. An
Update of the Evolving Epidemic of blaKPC Carrying Klebsiella pneumoniae in Sicily, Italy,
2014: Emergence of Multiple Non-ST258 Clones. PLoS One. 2015;10(7):e0132936. Epub
2015/07/16. doi: 10.1371/journal.pone.0132936. PubMed PMID: 26177547; PubMed Central
PMCID: PMCPMC4503429.
55.   Haeili M, Javani A, Moradi J, Jafari Z, Feizabadi MM, Babaei E. MgrB Alterations
Mediate Colistin Resistance in Klebsiella pneumoniae Isolates from Iran. Front Microbiol.
2017;8:2470. Epub 2018/01/13. doi: 10.3389/fmicb.2017.02470. PubMed PMID: 29326662;
PubMed Central PMCID: PMCPMC5741654.
56.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of
Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst
Biol. 2006;2:2006 0008. doi: 10.1038/msb4100050. PubMed PMID: 16738554; PubMed
Central PMCID: PMC1681482.
57.   Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci USA. 2000;97(12):6640-5.
doi: papers2://publication/doi/10.1073/pnas.120163297.
58.   Ochman H, Ajioka JW, Garza D, Hartl DL. Inverse polymerase chain reaction.
Biotechnology (N Y). 1990;8(8):759-60. Epub 1990/08/01. PubMed PMID: 1366903.
59.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods.
2009;6(5):343-5. Epub 2009/04/14. doi: 10.1038/nmeth.1318. PubMed PMID: 19363495.
60.   Besharova O, Suchanek VM, Hartmann R, Drescher K, Sourjik V. Diversification of
Gene Expression during Formation of Static Submerged Biofilms by Escherichia coli. Front
Microbiol. 2016;7:1568. doi: 10.3389/fmicb.2016.01568. PubMed PMID: 27761132;
PubMed Central PMCID: PMCPMC5050211.
61.   Miller JH. A Short Course in Bacterial Genetics: CSHL Press; 1992.

62.   Thibodeau SA, Fang R, Joung JK. High-throughput beta-galactosidase assay for
bacterial cell-based reporter systems. Biotechniques. 2004;36(3):410-5. doi:
10.2144/04363BM07. PubMed PMID: 15038156.
63.   Miyashiro T, Goulian M. Stimulus-dependent differential regulation in the
Escherichia coli PhoQ PhoP system. Proc Natl Acad Sci USA. 2007;104(41):16305-10. doi:
papers2://publication/doi/10.1073/pnas.0700025104.
64.   Haldimann A, Wanner BL. Conditional-replication, integration, excision, and
retrieval plasmid-host systems for gene structure-function studies of bacteria. J Bacteriol.
2001;183(21):6384-93. Epub 2001/10/10. doi: 10.1128/JB.183.21.6384-6393.2001. PubMed
PMID: 11591683; PubMed Central PMCID: PMCPMC100134.
65.   Bullock WO, Fernandez JM, Short JM. XL1-Blue: a high efficiency plasmid
transforming recA *Escherichia coli* strain with beta-galactosidase selection. BioTechniques.
1987;5:376-9.
66.   Cherepanov PP, Wackernagel W. Gene disruption in Escherichia coli: Tc R and Km
R cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant.
Gene. 1995. doi: papers2://publication/uuid/31A60367-CF87-4BF6-B690-93462755E5ED.
67.   Hay AJ, Yang M, Xia X, Liu Z, Hammons J, Fenical W, et al. Calcium Enhances Bile
Salt-Dependent Virulence Activation in Vibrio cholerae. Infect Immun. 2017;85(1). Epub
2016/11/17. doi: 10.1128/IAI.00707-16. PubMed PMID: 27849180; PubMed Central
PMCID: PMCPMC5203667.
68.   Amann E, Ochs B, Abel KJ. Tightly regulated tac promoter vectors useful for the
expression of unfused and fused proteins in Escherichia coli. Gene. 1988;69(2):301-15.
PubMed PMID: 3069586.
1269

F sequencing failed
NS nonsense mutation
FS frameshift
S silent mutation
IS insertion
D deletion

Label Number	Sample Name	Primer Name	sequencing results
90073451	RMphoQ001	pBADfw	F
90073452	RMphoQ002	pBADfw	NS
90073453	RMphoQ003	pBADfw	F
90073454	RMphoQ004	pBADfw	NS
90073455	RMphoQ005	pBADfw	F
90073456	RMphoQ006	pBADfw	NS
90073457	RMphoQ007	pBADfw	R219G
90073458	RMphoQ008	pBADfw	F
90073459	RMphoQ009	pBADfw	F
90073460	RMphoQ010	pBADfw	R262H
90073461	RMphoQ011	pBADfw	F
90073462	RMphoQ012	pBADfw	NS
90073463	RMphoQ013	pBADfw	F
90073464	RMphoQ014	pBADfw	NS
90073465	RMphoQ015	pBADfw	F
90073466	RMphoQ016	pBADfw	F
90073467	RMphoQ017	pBADfw	F
90073468	RMphoQ018	pBADfw	F
90073469	RMphoQ019	pBADfw	F
90073470	RMphoQ020	pBADfw	F

Label Number	Sample Name	Primer Name	sequencing results
90073499	RMphoQ2_001	pBADfw	NS
90073500	RMphoQ2_002	pBADfw	I196F, S217N
90073501	RMphoQ2_003	pBADfw	IS
90073502	RMphoQ2_004	pBADfw	R262S
90073503	RMphoQ2_005	pBADfw	L250M
90073504	RMphoQ2_006	pBADfw	WT
90073505	RMphoQ2_007	pBADfw	F
90073506	RMphoQ2_008	pBADfw	WT
90073507	RMphoQ2_009	pBADfw	WT
90073508	RMphoQ2_010	pBADfw	F
90073509	RMphoQ2_011	pBADfw	WT
90073510	RMphoQ2_012	pBADfw	F
90073511	RMphoQ2_013	pBADfw	L199H, E237V, S249R
90073512	RMphoQ2_014	pBADfw	W211C, L247Q
90073513	RMphoQ2_015	pBADfw	W215L
90073514	RMphoQ2_016	pBADfw	D
90073515	RMphoQ2_017	pBADfw	R262L
90073516	RMphoQ2_018	pBADfw	F
90073517	RMphoQ2_019	pBADfw	FS
90073518	RMphoQ2_020	pBADfw	FS

ID	Name	Vector	Variant
90073471	RMphoQ021	pBADfw	F
90073472	RMphoQ022	pBADfw	F
90073473	RMphoQ023	pBADfw	R219G
90073474	RMphoQ024	pBADfw	F
90073475	RMphoQ025	pBADfw	K259R
90073476	RMphoQ026	pBADfw	N235Z
90073477	RMphoQ027	pBADfw	WT
90073478	RMphoQ028	pBADfw	WT
90073479	RMphoQ029	pBADfw	WT
90073480	RMphoQ030	pBADfw	WT
90073481	RMphoQ031	pBADfw	V198A, R219C
90073482	RMphoQ032	pBADfw	WT
90073483	RMphoQ033	pBADfw	WT
90073484	RMphoQ034	pBADfw	S
90073485	RMphoQ035	pBADfw	WT
90073486	RMphoQ036	pBADfw	WT
90073487	RMphoQ037	pBADfw	WT
90073488	RMphoQ038	pBADfw	V198E, E230L
90073489	RMphoQ039	pBADfw	WT
90073490	RMphoQ040	pBADfw	WT
90073491	RMphoQ041	pBADfw	WT
90073492	RMphoQ042	pBADfw	V198A, R219C
90073519	RMphoQ2_021	pBADfw	F
90073520	RMphoQ2_022	pBADfw	F
90073521	RMphoQ2_023	pBADfw	A213T
90073522	RMphoQ2_024	pBADfw	D
90073523	RMphoQ2_025	pBADfw	WT
90073524	RMphoQ2_026	pBADfw	F
90073525	RMphoQ2_027	pBADfw	F
90073526	RMphoQ2_028	pBADfw	FS
90073527	RMphoQ2_029	pBADfw	F
90073528	RMphoQ2_030	pBADfw	WT
90073529	RMphoQ2_031	pBADfw	FS
90073530	RMphoQ2_032	pBADfw	NS
90073531	RMphoQ2_033	pBADfw	WT
90073532	RMphoQ2_034	pBADfw	WT
90073533	RMphoQ2_035	pBADfw	F
90073534	RMphoQ2_036	pBADfw	L247Q mixed with WT
90073535	RMphoQ2_037	pBADfw	WT
90073536	RMphoQ2_038	pBADfw	WT
90073537	RMphoQ2_039	pBADfw	WT
90073538	RMphoQ2_040	pBADfw	WT
90073539	RMphoQ2_041	pBADfw	WT
90073540	RMphoQ2_042	pBADfw	FS

Label Numbe	Sample Name	Primer Name	sequencing results
90073546	RMphoQ3_001	pBADfw	F
90073547	RMphoQ3_002	pBADfw	I221T
90073548	RMphoQ3_003	pBADfw	NS
90073549	RMphoQ3_004	pBADfw	R262C
90073550	RMphoQ3_005	pBADfw	WT
90073551	RMphoQ3_006	pBADfw	V251I
90073552	RMphoQ3_007	pBADfw	FS
90073553	RMphoQ3_008	pBADfw	S217C
90073554	RMphoQ3_009	pBADfw	WT
90073555	RMphoQ3_010	pBADfw	A223T
90073556	RMphoQ3_011	pBADfw	D274N
90073557	RMphoQ3_012	pBADfw	FS
90073558	RMphoQ3_013	pBADfw	FS
90073559	RMphoQ3_014	pBADfw	WT
90073560	RMphoQ3_015	pBADfw	FS
90073561	RMphoQ3_016	pBADfw	W211R
90073562	RMphoQ3_017	pBADfw	L210M
90073563	RMphoQ3_018	pBADfw	WT
90073564	RMphoQ3_019	pBADfw	FS
90073565	RMphoQ3_020	pBADfw	D274N
90073566	RMphoQ3_021	pBADfw	S217C
90073567	RMphoQ3_022	pBADfw	V206N
90073568	RMphoQ3_023	pBADfw	T270R
90073569	RMphoQ3_024	pBADfw	L203Q
90073570	RMphoQ3_025	pBADfw	FS
90073571	RMphoQ3_026	pBADfw	F
90073572	RMphoQ3_027	pBADfw	L209P

Label Numbe	Sample Name	Primer Name	sequencing results
90073573	RMphoQ3_028	pBADfw	WT
90073574	RMphoQ3_029	pBADfw	WT
90073575	RMphoQ3_030	pBADfw	NS
90073576	RMphoQ3_031	pBADfw	R256L
90073577	RMphoQ3_032	pBADfw	W216R, E230D
90073578	RMphoQ3_033	pBADfw	IS
90073579	RMphoQ3_034	pBADfw	WT
90073580	RMphoQ3_035	pBADfw	F
90073581	RMphoQ3_036	pBADfw	F
90073582	RMphoQ3_037	pBADfw	IS
90073583	RMphoQ3_038	pBADfw	A201S

Label Numbe	Sample Name	Primer Name	sequencing results
90073594	RMphoQ4_001	pBADfw	FS
90073595	RMphoQ4_002	pBADfw	A214T
90073596	RMphoQ4_003	pBADfw	FS
90073597	RMphoQ4_004	pBADfw	FS
90073598	RMphoQ4_005	pBADfw	L254R
90073599	RMphoQ4_006	pBADfw	FS
90073600	RMphoQ4_007	pBADfw	F195L
90073601	RMphoQ4_008	pBADfw	FS
90073602	RMphoQ4_009	pBADfw	IS
90073603	RMphoQ4_010	pBADfw	WT
90073604	RMphoQ4_011	pBADfw	FS
90073605	RMphoQ4_012	pBADfw	IS
90073606	RMphoQ4_013	pBADfw	F195L