Single cell variations in expression of codominant alleles A and B on RBC of AB blood group individuals.

Anjali Bajpai[1,3], Badanapuram Sridevi[1], Vidya Jonnalagadda[2], Pathma Mutthukotti[1], Rachel A Jesudasan[1,4]*

[1]Center for Cellular and Molecular Biology, Habsiguda, Hyderabad –500 007, T.S. India.

[2]Bhavan's Vivekanada College, Sainikpuri, Secunderabad – 500094. T.S.

Anjali Bajpai anjalii@iitk.ac.in

Vidya Jonnalagadda vsjonn@gmail.com

Badanapuram Sridevi ravisri7@yahoo.com

Rachel A Jesudasan rachellike@gmail.com

Pathma Muthukotti pathmamuthukotti@gmail.com

Present address

[3]Dept Biological Sciences and Bioengineering, Indian Institute of Technology, Kanpur, India
[4]Department of Genetics, Osmania University, Hyderabad – 500007, Telangana. India.

*Corresponding Author

rachellike@gmail.com

Telephone   +91-9440407163

Running Title: Mosaicism of A and B antigens in AB RBC

Key words: ABO locus, codominant alleles, mosaicism, differential expression, immunofluorescence.

Abstract

One of the key questions in biology is whether all the cells of a "cell type" have more or less the same phenotype, especially with relation to non-imprinted autosomal loci. This could be answered by single-cell studies in heterozygous organisms. We studied the ABO locus to assess the relative levels of expression of the co-dominant alleles I^A and I^B in 2512 RBC from 24 individuals with AB blood group using Cy3- and FITC- tagged antibodies. Contrary to the general belief that all RBC in AB individuals express both antigens in comparable amounts, they segregated into 4 groups: showing normal distribution for both antigens, either antigen, and neither antigen; the deviation from normal distribution could not be correlated to maternal/paternal origin. Surprisingly, very few people showed any correlation between the amounts of these two antigens on RBC. In fact, the ratio of antigen A to B in the entire set of samples spanned over 6 orders of magnitude. This variability in amount of the antigens A and/or B, combined with a lack of correlation between the amounts of these two antigens resulted in unique staining patterns for RBC, generating widespread mosaicism in the RBC population of AB blood group individuals.

Introduction

Alleles I^A and I^B that are responsible for the production of antigens A and B[1-4] respectively, of the ABO blood group are cited as classic examples of co-dominant expression of Mendelian traits[5]. The A and B antigens are produced by the differential glycosylation of a common substrate, H, present on the surface of all human cells. Allele I^A encodes α1-3 N-acetylgalactosaminyl transferase which adds an N-acetyl galactosamine residue to H disaccharides present on glycolipids or glycoproteins; this modified form is

termed antigen A. Allele I^B encodes α1-3 galactosyl transferase which adds a galactose residue at the same position of the H antigen; this modified form is termed antigen B. Null alleles at this locus are termed i, and the unmodified substance H is termed O antigen. Alleles I^A and I^B are both dominant over allele i, but are co-dominant with respect to each other. Recently, natural and engineered enzymes with dual specificity (cis-AB alleles) have been reported that can modify H antigen to produce both A and B antigens. However, an individual molecule of substance H can be modified to produce either A antigen or B antigen, but not both.

Cells of individuals with genotype $I^A I^B$ express both, A antigen and B antigen, on their surfaces. It is generally believed that these two antigens are present in more or less equivalent amounts, as judged by agglutination reactions. Certain types of I^A and I^B alleles are known to be 'weak', either due to lower level of transcription and/or poor catalytic activity of the encoded polypeptide product[6,7]. For example, I^A and I^B alleles containing missense mutations showed variation in expression when transfected into HeLa cells in terms of mean antigen amount as well as number of antigen expressing cells[8]. Nevertheless, one would expect that all the cells within an $I^A I^B$ individual would be fairly homogenous in terms of the stoichiometric ratios of the A and B antigens.

Here we report the quantification by immunofluorescence, of A and B antigens on 2512 peripheral red blood cells (RBC) from 24 individuals of AB blood type. We observed that a majority of the subjects (19 of the 24) in this cohort did not have RBC that stained equally for antigens A and B; in most cells, each RBC stained substantially more for either antigen A or antigen B. Quantification of both antigens on each cell revealed a wide variation in the amount of A and/or B antigen in most samples. Of the 24 individuals tested, only 8 showed appreciable or strong positive correlation (Carl Pearson correlation coefficient r >

0.5). The effect of these two levels of variation in quantity of surface antigens is that most individuals of AB blood group in this study group were phenotypic mosaics for these proteins. These observations challenge the presumption of stoichiometric expression of these two alleles within the RBC population and may be indicative of epigenetic mechanisms that can render these two alleles functionally non-equivalent within a cell.

Materials and Methods

Ethics Statement

Blood samples were collected with informed consent of the subjects, after obtaining clearance from the ethical committee, Centre for Cellular and Molecular Biology (CCMB), Hyderabad, India. The subjects were healthy donors from the institute. All the experimental procedures followed were as per strict guidelines and regulations issued by ethical committee, CCMB.

Immunofluorescent detection of antigens A and B on RBC: 100 μl of peripheral blood was fixed on poly-lysine coated cover slips using 3.7% formaldehyde. The cells were blocked with 2% BSA and incubated serially in (a) mouse monoclonal Anti-A antibody (1:10 dilution; Tulip Diagnostics (P) Ltd. India) for 90 min., (b) Cy3-tagged anti-mouse antibody (1:500 dilution; Amersham Pharmacia) for 60 min, (c) mouse monoclonal Anti-B antibody (1:10 dilution; Tulip Diagnostics (P) Ltd. India) for 90 min, and (d) FITC tagged anti-mouse antibody (1:200 dilution; Vector Labs). After each incubation, the slides were washed in 1x PBS (Phosphate Buffered Saline) for 15 min. After the final wash, slides were mounted using a glycerol-based anti-fade reagent, Vectashield (Vector Labs). Control reactions carried out to rule out non-specific binding were incubation of (a) anti-A antibody to RBC of O and B blood group types, (b) anti-B antibody to RBC of O and A blood group types, and (b) anti-

mouse secondary antibody to RBC of all four blood groups (O, A, B, and AB) in the absence of any primary antibody. All procedures were done at room temperature.

Image capture: Slides were scanned at 40x (dry) and 100x (oil) magnifications using the dual scanning, multi-track mode of a confocal microscope (Zeiss, LSM510 META). The numerical aperture used was 0.95 for 40x and 1.4 for 100x. Images containing 5 Z-stacks were captured for quantification (100x magnification) on Axioplan-2 using Axiovision version-4 camera (Carl Zeiss). The intensity of fluorescence of FITC and Cy3 was quantified from the projected images using the LSM-FCS Version 3.2SP software (Carl Zeiss) and corrected for background fluorescence. Typically, around 100 cells were analyzed from each sample slide.

Statistical analysis: Data were analyzed using MS Excel and R Statistical Software. Outliers in box plots were identified using default parameters of R. The Shapiro test was used to determine the probability that a given set of fluorescence values conformed to a normal distribution. For each subject, the Karl Pearson correlation coefficient was calculated using basic functions in Excel.

Data availability statement

The authors affirm that all data necessary for confirming the conclusions of the article are present within the article, figures, and tables.

Results and discussion

Unequal staining for A and B antigens on RBC from AB blood group individuals

The RBC from 24 (15 males and 9 females) AB blood group individuals were fixed on slides and incubated with anti-A and anti-B antibodies followed by fluorescently tagged

secondary antibodies; Cy3 (red fluorescence) for antigen A, and FITC (green fluorescence) for antigen B. Contrary to expectations of finding equivalent amounts of antigen A and antigen B on each RBC, we observed that RBC from AB individuals differed with respect to the fluorescent staining pattern on their surface on co-localization of the A and B antigens (Fig. 1).

The samples appeared to fall into four distinct categories. In group (i), most of the RBC were stained for antigen B but differentially stained for antigen A (Fig. 2, A-F). In group (ii), almost all RBC were stained for antigen A but only some cells showed staining for antigen B; in other words, staining for antigen B was differential, i.e., quantitatively different from cell to cell (Fig. 2, G-L). In group (iii), RBC appeared to be stained differentially for both antigens (Fig. 2, M-R). Finally, in group (iv), all the RBC stained equally for both antigens (Fig. 2, S-X).

Distribution of A and B antigens on RBC from AB blood group individuals

To determine the extent of expression of the antigens on individual cells, we used densitometry to quantify the fluorescence for both dyes. These RBCs were studied with respect to the number of cells showing differences and the quantitative variation between the two antigens from single cells. The data was analyzed to check for statistically significant differences, if any. The fluorescence measurements of the A and B antigens were checked for normal distribution, using the Quantile-Quantile (Q-Q) plots, which plot the distribution of the observed values against an expected normal distribution. The Q-Q plots show that the fluorescence did not show a normal distribution for most of the samples (Fig. 3).

These data were also plotted as a box plot to show the distribution of fluorescence for antigen A (red bars) and antigen B (green bars) (Fig. 4). Using the Shapiro Wilk test to assess

for normality of the distribution, we found that the samples could again be assigned to 4 classes: (i), where only antigen B shows a normal distribution (subject # 19); (ii), where only antigen A shows a normal distribution (subjects #2, 3, 5, 7, 13, 17, 20 and 25); (iii), where neither antigen shows a normal distribution (subjects # 9, 11, 15, 18, 1, 24, 6, 12, 16, 23 and 26) and (iv), where both antigens show a normal distribution (subjects # 4, 8, 14, and 21).

Another interesting observation was that most samples showed no correlation between the amounts of antigens A and B on individual cells. As noted in Figure 4, only 2 samples (subjects # 4 and 19) showed a value above 0.9 for Karl Pearson correlation coefficient (r), and 3 other samples (subjects # 8, 6, and 26) had r>0.6.

Range and ratio of expression of A and B antigens on RBC from AB blood group individuals

Considering that these two antigens are expressed from alleles on homologous chromosomes, this lack of correlation shows that the expression of these antigens is not under stringent control. The extent of wide variation in the relative amount of antigens A and B on individual RBC is evident from a box plot of the ratio of fluorescence from antigen A to that from antigen B (Fig. 5). If these two alleles were expressed at comparable amounts on all cells, we would expect to see a narrow range of ratios (the ratio would be 1 if the fluorescence due to the two dyes – Cy3 and FITC – had molar equivalence). Yet we see the ratios spanning 5 orders of magnitude, from 1/100 to over 1000. This is consistent with essentially independent levels of expression of these antigens and/or rates of loss during the lifetime of the RBC.

Our observations are consistent with a previous report by Balea and colleagues[9] where flow cytometry of RBC from AB individuals revealed distinct sub-population of cells with

equal concentrations of antigens A and B, or with maximum concentration of antigen A and minimum concentration of antigen B, or with maximum concentration of antigen B and minimum concentration of antigen A. The use of scanning fluorescence microscopy in our study has revealed more layers of heterogeneity within population of RBC in individuals of AB blood group. In many individuals, the level of heterogeneity is so high that it appears that no two RBC are alike with respect to antigens A and B on their surface. This degree of mosaicism appears to be unparalleled in human tissues.

In this cohort of 24 subjects, 12 showed normal distribution for antigen A while only 4 showed normal distribution for antigen B. This was an unexpected finding: differences have been reported in the promoter strength as well as the level of transcripts from various I^A and I^B alleles, but one would expect that the products of strongly or weakly expressed alleles would show a normal distribution nevertheless, even if the absolute amount of antigen A was not the same as that of antigen B. There is no synthesis of new molecules of these antigens in circulating RBC, given that these cells are devoid of ER where the galactosyl transferases encoded by I^A and I^B alleles function. Scant expression of the A or B antigen on circulating RBC is consistent with the corresponding enzyme being fully active only in few cells; the reasons for a possible sporadic activity, however, are not clear.

Is the ABO locus imprinted?

To probe if the non-normal distribution of antigens from the I^A and/or I^B alleles in certain individuals was potentially due to epigenetic mechanisms dependent on the parent of origin, we attempted to correlate the alleles to paternal or maternal allele origin. The blood group of the parents of some of the subjects in this test group was known. Comparison of the blood group of the parents with the test individuals showed that there was no relationship between the allele showing differential expression and the parent of origin (Table 1).

Serial number	Subject number	Category	Mother	Father
1	13	1	B	A
2	10	2	A	B
3	3	2	AB	AB
4	5	2	AB	AB
5	12	2	AB	(not known)
6	1	3	AB	AB
7	21	4	A	B
8	6	4	A	B
9	11	4	A	B
10	24	4	AB	B

Table 1: Blood groups of selected subjects and their parents

The alleles of the ABO blood group genes localizing to chromosome 9 in humans are classic examples of co-dominant gene expression[10]. Earlier studies in the lab reported a difference in the occurrence of sister chromatid exchanges on the chromosomes of *Muntiacus muntjak*[11]. More recent studies show that indeed there are a number of genes, which show allelic variation in gene expression within an individual, elucidating the fact that this could be a more common phenomenon than envisaged[12,13]. Allele-specific differences in gene expression have been reported and currently this concept is more widely accepted[14,15]. Epigenetic factors such as single nucleotide polymorphisms in regulatory elements, parent of origin imprinting, differential methylation etc. is believed to effect such differences in gene expression[16]. Variation in transcription has been observed within cells of a population. Yet the variability at the cytoplasmic level is tightly regulated by the nucleus[17]. In the RBC of the AB blood group, we observe a variation in the protein. The long term follow up of few

subjects over a span of 6-7 months shows that the expression pattern of these antigens appears to be fixed for a given individual (data not shown).

The mechanisms responsible for this heterogeneity are not clear. Different mechanisms are known to result in mosaicism in a tissue: position effect variegation due to variable heterochromatin formation, epigenetic modifications such as methylation, hyper-mutation, cellular volume and number of mitochondria within a cell[17]. Since the RBC lack DNA, it is not possible to determine if the sporadic or non-coordinated transcription of the alleles is due to any of these mechanisms or a novel pathway. A and B antigens are expressed on all cells[18,19]; it is possible that characterization of clones from single cells taken from some other somatic tissues may reveal modifications in the genotype that can be correlated to the phenotype, but such analyses will be technologically challenging.

It is also not clear whether this heterogeneity has any physiological significance. A variety of functions have been proposed for the ABO antigens; certain blood types have been associated with increased risk for certain medical conditions, or infections (Reviewed in[20]). The revelation of extensive heterogeneity in the quantitative profiles of these antigens suggests that the use of quantitative immunofluorescence instead of the traditional method of agglutination reactions to determine the blood group of individuals may allow more thorough correlation of these risk factors with specific antigen profiles.

The differential and combinatorial expression from autosomal homologues, can give rise to cell-specific expression, not only giving each cell a unique signature, but accounting as well for individual specific gene expression. This may be one of the mechanisms by which nature can bring about phenotypic variation within a single genotype. It would be interesting to investigate if the differential response to drugs by individuals can be understood better by studying combinatorial expression of sets of genes.

Conflict of Interest Statement

Authors state that they have no conflict of interest.

Author Contributions Statement

Conceived and designed the experiments: RAJ. Performed the experiments: AB, BS, PM.

Analyzed the data: RAJ, AB, VJ. Wrote the paper: RAJ, VJ.

Funding

This project was supported by intramural funding from CSIR to RAJ,

Acknowledgements

We thank Dr. R. Sarkar for help with statistics, Ms. R. Nandini for technical help with microscopy and CSIR for financial assistance to RAJ and AB.

References

Ginsburg, V. Enzymatic basis for blood groups in man. *Adv Enzymol Relat Areas Mol Biol* **36**, 131-149 (1972).

Race, C., Ziderman, D. & Watkins, W. M. An alpha-d-galactosyltransferase associated with the blood-group B character. *Biochem J* **107**, 733-735 (1968).

Tuppy, H. & Staudenbauer, W. L. Microsomal incorporation of N-acetyl-D-galactosamine into blood group substance. *Nature* **210**, 316-317 (1966).

Watkins, W. M. & Morgan, W. T. Possible genetical pathways for the biosynthesis of blood group mucopolysaccharides. Vox Sang 1959:4:97-119. *Vox Sang* **64**, 245 (1993).

Von Dungern, E. & Hirschfeld, L. Ueber Vererbung gruppenspezifischer Strukturen des Blutes. *Molecular and General Genetics MGG* **5**, 196-197 (1911).

Yamamoto, F. *et al.* Molecular genetic analysis of the ABO blood group system: 1. Weak subgroups: A3 and B3 alleles. *Vox Sang* **64**, 116-119 (1993).

Chester, M. A. & Olsson, M. L. The ABO blood group gene: a locus of considerable genetic diversity. *Transfus Med Rev* **15**, 177-200, doi:10.1053/tmrv.2001.24591 (2001).

Seltsam, A. & Blasczyk, R. Missense mutations outside the catalytic domain of the ABO glycosyltransferase can cause weak blood group A and B phenotypes. *Transfusion* **45**, 1663-1669 (2005).

Balea, M. *et al.* Pattern of antigens A and B distribution on the surface of AB group red blood cells--an argument in support of the concept of "genetic oscillation". *Rom J Intern Med* **30**, 269-279 (1992).

Bennett, E. P., Steffensen, R., Clausen, H., Weghuis, D. O. & Vankessel, A. G. Genomic cloning of the human histo-blood group ABO locus. *Biochemical and biophysical research communications* **206**, 318-325 (1995).

Rachel, A., Sharma, T. & Menon, V. Differences in sister-chromatid exchange frequency between homologous chromosomes in Muntiacus muntjak. *Mutation Research Letters* **283**, 193-198 (1992).

Lo, H. S. *et al.* Allelic variation in gene expression is common in the human genome. *Genome research* **13**, 1855-1862 (2003).

Pant, P. K. *et al.* Analysis of allelic differential expression in human white blood cells. *Genome research* **16**, 331-339 (2006).

Eckersley-Maslin, M. A. & Spector, D. L. Random monoallelic expression: regulating gene expression one allele at a time. *Trends in Genetics* **30**, 237-244 (2014).

Borel, C. *et al.* Biased allelic expression in human primary fibroblast single cells. *The American Journal of Human Genetics* **96**, 70-80 (2015).

Crowley, J. J. *et al.* Analyses of allele-specific gene expression in highly divergent mouse crosses identifies pervasive allelic imbalance. *Nature genetics* (2015).

Battich, N., Stoeger, T. & Pelkmans, L. Control of transcript variability in single mammalian cells. *Cell* **163**, 1596-1610 (2015).

Keitschewski, I. & Schwarzmann, L. Die gruppenspezifische Differenzierung der menschlichen Organe. *Journal of Molecular Medicine* **6**, 2081-2086 (1927).

Witebsky, E. & Okabe, K. Über den Nachweis von Gruppenmerkmalen in den Organen des Menschen. *Zeitschr. f. Immunitf. u. exp. Therap*, 359 (1927).

Greenwell, P. Blood group antigens: molecules seeking a function? *Glycoconj J* **14**, 159-173 (1997).

Figure legends

Figure 1: Mosaicism for antigens A and B on RBC from different subjects revealed by immunofluorescent staining: Panels **a-h** represent RBC from eight different individuals. RBC were fixed on cover slips and stained with antibodies to stain antigen A with Cy3-tag (red fluorescence) and antigen B with FITC-tag (green fluorescence). Overlay of red and green fluorescence appears as yellow fluorescence. RBC from eight subjects are shown at 100X magnification.

Figure 2: Immunofluorescent staining for antigens A and B on RBC from individuals of AB blood group: RBC were fixed on cover slips and stained with antibodies to stain antigen A with Cy3-tag (red fluorescence) and antigen B with FITC-tag (green fluorescence). Left panel shows co-localization of antigens A and B (yellow fluorescence); middle panel shows staining for antigen A; and right panel shows staining for antigen B. Cells from representative samples from each category are shown at 100X and 40 X magnifications. A-F: cells from

group (i) - stained predominantly for antigen B and differential staining for antigen A; G-L: cells from group (ii) - stained predominantly for antigen A and differential staining for antigen B; M-R: cells from group (iii) - differential staining for both antigens; S-X: cells from group (iv) - equal and uniform staining for both antigens.

Figure 3. Figure shows the Q-Q plots depicting the distribution of A and B antigens from four representative individuals. The Panels **a** and **b** show the distribution of antigen A and B respectively of an AB individual who shows a differential expression of A antigen. The antigen A shows a distribution away from the normal (**a**), and antigen B (**b**) shows a normal distribution. Panels **c, d** show the distribution of an individual with differential expression of antigen B, which shows a skewed distribution (**c**) and antigen A shows a normal distribution (**d**). The panels (**e**) and (**f**) display skewed distribution of both A and B antigens in an individual who shows differential expression of both the antigens. Panels (**g**) and (**h**) show normal distribution of both the antigens A and B in an individual with equal expression of both the antigens. The Q-Q plot plots the expected normal distribution on the Y axis against the observed values on the X axis.

Figure 4: Comparison of distribution of fluorescence (arbitrary units) corresponding to antigens A and B on RBC: Box plot showing distribution for fluorescence intensity for antigens A (red boxes) and B (green boxes) stained with Cy3- and FITC- tagged antibodies. The value of Karl Pearson coefficient of correlation (r) and number of RBC quantified (n) is shown for each subject. Outliers were identified using the boxplot() function of R. The box marks the limits of the upper and lower quartiles, and the median is indicated as a line within the box. Whiskers extend to 1.5*IQR (interquartile range). Filled circles indicate outliers.

Figure 5: Fluorescence ratios for all individuals. Box plots and outliers were marked as for Figure 4.

Figure 1

Figure 2

| | MERGED | ANTI A | ANTI B |

Figure 3

Figure 4

Figure 5