Illumina-based sequencing framework for accurate detection and mapping of influenza virus defective interfering particle-associated RNAs

Fadi G. Alnaji[1], Jessica R. Holmes[2,3], Gloria Rendon[2,3], J. Cristobal Vera[1,2], Chris Fields[2,3], Brigitte E. Martin[1], and Christopher B. Brooke*[1,2]

[1]Department of Microbiology, University of Illinois at Urbana-Champaign
[2]Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign
[3]High Performance Biological Computing at the Roy J. Carver Biotechnology Center, University of Illinois at Urbana-Champaign
* Corresponding author (cbrooke@illinois.edu)

Abstract

The mechanisms and consequences of defective interfering particle (DIP) formation during influenza virus infection remain poorly understood. The development of next generation sequencing (NGS) technologies has made it possible to identify large numbers of DIP-associated sequences, providing a powerful tool to better understand their biological relevance. However, NGS approaches pose numerous technical challenges including the precise identification and mapping of deletion junctions in the presence of frequent mutation and base-calling errors, and the potential for numerous experimental and computational artifacts. Here we detail an Illumina-based sequencing framework and bioinformatics pipeline capable of generating highly accurate and reproducible profiles of DIP-associated junction sequences. We use a combination of simulated and experimental control datasets to optimize pipeline performance and demonstrate the absence of significant artifacts. Finally, we use this optimized pipeline to generate a high-resolution profile of DIP-associated junctions produced during influenza virus infection and demonstrate how this data can provide insight into mechanisms of DIP formation. This work highlights the specific challenges associated with NGS-based detection of DIP-associated sequences, and details the computational and experimental controls required for such studies.

Importance

Influenza virus defective interfering particles (DIPs) that harbor internal deletions within their genomes occur naturally during infection in humans and cell culture. They have been hypothesized to influence the pathogenicity of the virus; however, their specific function remains elusive. The accurate detection of DIP-associated deletion junctions is crucial for understanding DIP biology but is complicated by an array of technical issue that can bias or confound results. Here we demonstrate a combined experimental and computational framework for detecting DIP-associated deletion junctions using next generation sequencing (NGS). We detail how to validate pipeline performance and provide the bioinformatics pipeline for groups interested in using it. Using this optimized pipeline, we

44 detect hundreds of distinct deletion junctions generated during IAV infection, and use
45 these data to test a long-standing hypothesis concerning the molecular details of DIP
46 formation.
47
INTRODUCTION
Influenza A virus (IAV) DIPs were first described over 60 years ago, and are classically
defined by their ability to interfere with the production of wild-type virus(1, 2). This ability
has been linked to the ability of DI RNAs to both outcompete wild-type (WT) genomic
RNAs for resources and packaging into virions, as well as to more potently stimulate the
induction of anti-viral immunity through cytosolic RNA sensors (3–6). DIPs have also been
implicated in influencing the outcome of influenza virus infection in humans(7). The
specific mechanisms and broader functional consequences of DIP formation during IAV
infection remain poorly understood.
57
IAV DIPs are characterized by the presence of large internal deletions in one or more
genome segments that disrupt essential open reading frames while retaining the
sequences required for replication and packaging(5). As such, the mapping of DIP-
associated deletions has helped to define the minimum sequences required for genome
replication and packaging (8, 9). These deletions are believed to result from a poorly
defined process by which the viral RNA-dependent RNA polymerase (RdRp) ceases RNA
polymerization at one site of the viral RNA template (donor site), only to resume at another
site downstream (acceptor site), resulting in a failure to copy an internal stretch of the WT
template (10). Until recently, the ability to characterize these DIP-associated deletion
junction sites (breakpoints) has been limited based on the need to clone and Sanger
sequence individual DIP-associated RNAs. As a result, the number of individual DIP-
associated RNA sequences that have been analyzed has been relatively small, hindering
efforts to define the factors that govern DIP deletion formation.
71
The advent of next generation sequencing (NGS) has increased the number of individual
recombinant sequences that can be identified within a given sample by orders of
magnitude. However, the identification and analysis of DIP-associated RNAs by NGS
poses new challenges, including the successful alignment of junction-containing (or
junction-spanning) reads to the viral reference sequence, the precise definition and
localization of DIP-associated deletion breakpoints, and the differentiation of true DIP
deletion sequences from the artifactual recombinants that can form during reverse
transcription, PCR, and/or sequencing. Without careful optimization and validation, these
issues can easily compromise efforts to define the genetic profile of DIP populations.
81
Here, we describe the development and validation of an Illumina-based sequencing
framework for the identification and analysis of influenza virus DIP-associated deletion
junctions. The bioinformatics pipeline combines the Bowtie 2 alignment algorithm with the
ViReMa (Virus Recombination Mapper) algorithm developed by Andrew Routh and a
collection of additional scripts for data processing and analysis (11, 12). We used
simulated NGS datasets and a panel of experimental control samples to optimize and

quantify the sensitivity, precision, and reproducibility of our pipeline. Subsequently, we
used the optimized pipeline to fine-tune the experimental protocol from sample
preparation to RNA sequencing to better detect and map DIP-associated deletions
generated during experimental IAV infection. This work highlights the computational and
experimental controls needed for Illumina-based NGS studies of viral recombination, and
provides an optimized, user-friendly sequencing and bioinformatics pipeline for the
identification and analysis of IAV DIP-associated sequences. Higher resolution analysis
of these deletion sequences can shed light on both the specific molecular mechanisms
of DIP formation, as well as how DIPs may affect the overall behavior of viral populations.
97
RESULTS
Overview of the pipeline
The sequencing framework we describe here encompasses sample preparation,
sequencing, and data analysis (**Fig 1A**). In brief, we generate 8-segment, full-length
amplicons from viral samples and sequence these using the Illumina MiSeq sequencing
platform. Datasets are quality-filtered and aligned to the viral reference genome using
Bowtie 2 in a conservative manner that disallows soft clipping. Thus, reads containing
deletion junctions fail to align, and are fed into the ViReMa algorithm to detect DIP-
associated deletion junctions. Finally, the identified junctions are mapped to the viral
genome and output as a matrix containing the segment name, junction start and end sites,
and NGS read support that can easily be analyzed using additional software tools. Below,
we outline the approaches we have taken to optimize and validate the various steps in
the process.
111

Fig 1. Overview of sequencing/bioinformatics framework. A combination of experimental and computational approaches was extensively optimized in a stepwise manner to establish a pipeline for detecting and analyzing DIP-associated junctions within IAV **(A)** Flowchart outlining the pipeline steps. **(B)** Simple depiction of the possible types of NGS reads in relation to a deletion junction within a sample. Arrows represent individual NGS reads, and the black circle denotes the location of a deletion junction.

112
113

Optimization of analysis pipeline using simulated data

All bioinformatic pipelines have the potential to introduce artifacts and biases during data analysis. Therefore, we first aimed to optimize the sensitivity and precision of our bioinformatics pipeline using simulated NGS datasets where we absolutely know the identity and frequency of all DIP-associated deletion sequences present. IAV DIP-associated deletions can be found in nearly all (if not all) genome segments at a wide range of frequencies (13, 14). To mimic this natural variation, we used MetaSim to generate a panel of Illumina MiSeq-based NGS simulated datasets that contain DIP-associated deletions in all genome segments at varying frequencies and locations (see **Table 1, Fig S1**). We used a simple Perl script to randomly generate deletion junctions within the terminal ~600nts of A/California/07/09 (Cal07), since these regions have been shown to be hotspots for DIP-associated deletions(9, 13, 15). We also generated a negative control dataset that lacks deletions to quantify the occurrence of false positives generated by the pipeline. Critically, we introduced a nucleotide substitution frequency of ~1% into these datasets, based on the published Illumina MiSeq empirical error model(16, 17). Each dataset comprised ~1million 2x250nts paired-end reads, mirroring the read depth that we expect per sample on a typical sequencing run.

Dataset name	Total paired-end reads	Total junction count	Total junction NGS reads	Total WT NGS individual reads	Junction count							
					PB2	PB1	PA	HA	NP	NA	M	NS
Cal07-400	~1 million (2X250). Total 2 million individual reads	400	1838898	116548	50	50	50	50	50	50	50	50
Cal07-200		200	1774920	225080	25	25	25	25	25	25	25	25
Cal07		0	0	2000000	0	0	0	0	0	0	0	0

Table 1. Description of the simulated datasets used in this study.

Optimization of alignment

We first optimized the filtering of reads that contain deletion junctions (**Fig 1B,** R1), from those that don't include junctions (**Fig 1B,** R2,R3). To do this, we aligned all reads to the WT reference genome using Bowtie 2. Reads that successfully align should not contain deletion junctions and are saved for further analysis, while reads that fail to align are fed through the ViReMa algorithm. The performance of this alignment step is highly dependent upon the mismatch penalty scores that are used during alignment. If mismatch penalties are too stringent, reads with random mutations or base calling errors will fail to align and be sent to ViReMa, increasing both the chances of false positives and the total computational time per sample; too lenient, and true junction-spanning reads will successfully align and be excluded from downstream analysis.

We used a junction-rich simulated dataset (Cal07-400) to test the effects of varying the alignment penalty score on the output of ViReMa (**Fig 2A**). We observed that a penalty score of 0.3 minimized the number of unaligned reads (and thus potential for false positives) without diminishing the number junction-spanning reads detected. This value was used for all subsequent analysis.

Optimization of ViReMa operation

We next optimized the sensitivity and precision with which the pipeline detects deletion junctions. The ability of ViReMa to accurately map true junction-containing reads is affected by three factors. The first is the method the algorithm uses to identify breakpoints. ViReMa extracts and aligns a seed sequence of 20-30nts (the default value of 25 was used in this study) from the beginning of each read and begins aligning the downstream nucleotides. If at any point the downstream alignment fails (as would be the case for a deletion breakpoint), ViReMa generates a new seed sequence starting from that location for realignment. Thus, breakpoints cannot be detected if they occur within the terminal 25 nts of a read.

Fig 2. Optimization of bioinformatics pipeline using simulated data. (A) Quantification of the effects of varying the bowtie 2 penalty score on the number of junction spanning reads detected by ViReMa in the Cal07-400 simulated dataset (black line; dashed line represents the actual number of junction-spanning reads present in the dataset). The percentages of reads that aligned to reference genome (purple) and failed to align (green) are also shown for each penalty score. **(B)** The effects of ViReMa --X and --N parameters on the percentage of junction spanning reads present in the simulated dataset that were successfully detected. Dashed line shows the maximum theoretical sensitivity (~81.5%), based on the ViReMa seed length of 25nts. **(C)** The effects of ViReMa --X and --N parameters on the number of accurately (green) and inaccurately mapped (purple) deletion junctions reported by the pipeline using the Cal07-200 simulated dataset. The maximum possible number of accurate junctions and the minimum number of inaccurate junctions (resulting from junctions adjacent to direct repeat sequences) are shown for comparison **(D)** Effects of varying the minimum read support cutoff (RSC) on junction detection. Analysis performed on the Cal07-200 simulated dataset using N1X8 ViReMa values.

The second factor is the presence of short direct repeats adjacent to the junction site. These repeats result in a situation where multiple potential breakpoints can give rise to the same final sequence, making precise definition of the true breakpoints impossible (**Fig S2 and S3A**). ViReMa deals with these 'fuzzy' regions through the parameter 'Defuzz', which can be set to report the junction either to the 5' end, 3' end, or the middle of the ambiguous region. For consistency's sake, we pushed all fuzzy junctions towards the 3' of the ambiguous region. The effects of direct repeats on breakpoint mapping are impossible to avoid and vary somewhat between IAV genome segments. Importantly, while this effect reduces the precision of breakpoint mapping, it does not affect the ability of the pipeline to determine the actual sequences of DIP-associated RNAs.

The third factor is the potential for base calling errors or mutations to result in erroneous junction mapping (**Fig S3B**). Even though reported junctions in this category are derived from real junctions, they can be viewed as false positives in that they are reported as

distinct junctions that do not actually exist in the viral population. Altogether, these three
factors set a ceiling on the maximum number of deletion junctions that can be accurately
detected and mapped. Using our simulated datasets, we knew how many deletion
junctions actually existed, exactly where they were located, and whether or not they were
adjacent to direct repeats (see Materials and Method) that could result in incorrect
mapping. This allowed us to systematically optimize the sensitivity and precision of the
software pipeline.
183
We tested how varying the ViReMa operating parameters affected both junction-spanning
read detection and actual junction reporting. We used the Cal07-200 dataset to challenge
ViReMa across a range of --N parameter (number of mismatches allowed) and --X
parameter (mismatch distance from the putative junction location) values. We first asked
how varying the --N and --X parameters influenced the total number of junction-spanning
reads detected (**Fig 2B**). We found that using N=0 (--X is irrelevant at this condition)
significantly decreased the number of junction-spanning reads detected compared with
non-zero --N and --X values. We next asked how increasing the --N and --X values
affected the number of accurately and inaccurately mapped junctions reported (**Fig 2C**).
We observed a clear correlation between the --X parameter and junction-mapping
precision, as increasing the --X value decreased the number of inaccurately mapped
junctions. Overall, we found that using N=1 and X=8 reduced inaccurate junction mapping
to the minimum amount possible, given the occurrence of direct repeats adjacent to
23.5% (47 of 200) of junctions in the dataset.
198
We next asked whether setting a minimum read support cutoff (RSC) to report a junction
affected the numbers of both accurate and inaccurate junctions that the pipeline identified.
Requiring that a given junction be represented within a minimum number of reads can
decrease the number of erroneously mapped junctions arising from base calling errors
but could also result in some true junctions being lost due to insufficient read coverage.
We aligned our simulated Cal07-200 dataset with Bowtie 2 and used the resulting
unaligned reads to challenge ViReMa using different RSC values (**Fig 2D**). We found that
the number of true junctions reported by the pipeline was very close to the theoretical
maximum, with minimal drop-off across the range of RSCs tested. In contrast, we
observed that the number of inaccurately reported junctions was highly sensitive to the
RSC value used. An RSC of >30 was needed to lower the number of inaccurately reported
junctions to the minimal limit (determined by the number of 'fuzzy' junctions with adjacent
direct repeats in the dataset).
212
213 Altogether, these data highlight the importance of optimizing RSC values and the ViReMa
214 --N and --X parameters for maximizing the sensitivity of junction detection while
215 minimizing the number of false positives. We set our default values at RSC>30, --N=1,
216 and --X=8 for subsequent analysis.
217
Validation of sequencing pipeline
After optimizing the bioinformatics component of our pipeline using simulated datasets,

we examined the ability of the pipeline to detect DIP-associated deletions within complex
viral populations from experimental samples. Our overall strategy was based on the
universal, eight segment RT-PCR approach pioneered by Zhou et al. (18). Critically, there
are a number of steps within the library preparation and sequencing steps that have the
potential to introduce artifacts that can compromise junction detection and analysis. In
particular, we were concerned about the potential for recombination during reverse
transcription, PCR, and/or sequencing to generate junctions that will be called by the
pipeline (19, 20). To address this, we prepared several control sample libraries,
sequenced them on the MiSeq, and ran the results through our optimized pipeline.
229
To quantify false positive generation during the PCR and/or sequencing steps, we
constructed libraries without using actual viral RNA or reverse transcriptase. To do this,
we generated an equimolar ratio mixture of full length PCR amplicons from each of the
eight IAV genome segments, using reverse genetics plasmids encoding the gene
segments from A/Puerto Rico/8/1934 (PR8) as templates. These amplicons were gel
purified to ensure correct, full-length size, and then used as template for the universal
amplification PCR and subsequent library preparation. Our analysis pipeline detected no
breakpoints in this control, indicating that none of the steps in our pipeline from PCR
onwards were significant sources of false positive signals.
239
We next sequenced a recombinant Cal07 stock that was grown under low MOI conditions
to minimize the frequency of DIPs (21). We performed two independent RNA extractions
and reverse transcription reactions on this stock to serve as technical replicates (named
Par1 and Par2). ViReMa detected 6 and 7 DIP-associated deletion junctions from Par1
and Par2, respectively, with junction-spanning reads representing ~0.1-0.2% of the total
reads **(Fig 3A)**. The majority of these reads were derived from a single shared deletion
junction in HA (indicated by the following nomenclature: 615_1132_HA). 4 other DIP
junctions were shared between replicates, each with low NGS read depth (ranging
between 19 and 94). Two unshared junctions in Par1 and one in Par2 were actually
reported in both replicates but failed to reach the level of detection in one replicate.
250
The significant overlap in the specific junctions that were reported from the two replicates
suggested that these junctions were produced by the viral polymerase (and were thus
bona fide DIP-associated sequences) rather than by the reverse transcriptase. However,
the generation of the same junction in independent RT reactions could also indicate the
existence of strong hotspots for RT recombination. To more directly address the potential
contribution of RT-derived recombinants, we performed two independent experiments.
First, we compared the junctions detected in HA segment libraries generated from Par 1
using two different RT enzymes, Invitrogen Superscript III and Agilent AccuScript.
Second, we performed *in vitro* transcription of a plasmid-derived Cal07 HA segment using
T7 RNA polymerase, which then was used as a template for RT-PCR to produce the
amplicon library for sequencing. The IAV polymerase was not involved in this control; thus
any deletions detected will have been generated by T7 polymerase or the RT enzyme.
263

The junctions reported from libraries generated by the two RT enzymes had significant
overlap and were both dominated by 615_1132_HA (**Fig 3B**). In contrast, we detected
none of the Par1-derived junctions in the library generated from T7-transcribed HA (**Fig**
3B). Although the read depth coverage was comparable to Par1, 615_1132_HA was
completely absent, and the three junctions that were detected had minimal read support
and were not seen in virus-derived libraries. Altogether, these results suggest that the
formation of deletion junctions during the reverse transcription reaction is rare, and that
the Par1-derived junctions we observed are most likely derived from true DIPs present
within our viral stock, despite the stock having been prepared at low MOI. This highlights
the difficulty in producing a completely DIP-free virus preparation.
274

Fig 3. DIP-associated deletion junctions present in virus working stock. We performed two
independent RNA extractions, RT reactions, PCR amplifications, and library preparations from a
single recombinant Cal07 working stock grown at low MOI (Par1 and Par2). **(A)** Comparison of
deletion junctions detected in Par1 and Par2 samples. Purple blocks represent instances where
junction was detected but read support was below RSC **(B)** Comparison of HA segment junctions
detected in two libraries generated from independent RT reactions using two different RT
enzymes, along with a library generated from in vitro T7-transcribed viral RNA. White blocks
denote no detection

275
276

Generation of DIP-enriched populations through high MOI passage
277
To test the ability of the pipeline to detect real DIP-associated RNAs, we enriched for
DIPs through serial undiluted passage of Cal07 in MDCK cells. We confirmed the
presence of DIPs by amplifying full-length genomic cDNA at each passage and examining
the size distribution of PCR products by gel electrophoresis **(Fig 4A)**, as previously
described (21). The gradual disappearance of the polymerase segments, which form the
majority of DIPs, and the appearance of a smear below the shortest IAV segment (NS
~0.9kb) were consistent with the accumulation of DIPs over passage. Based on these
results, we picked P1, P3, and P6 as representative samples for sequencing.

We further confirmed the presence of DIPs by plotting the read coverage of the aligned
reads from passages 1, 3, and 6 (**Fig 4B**). These coverage plots clearly reveal the
characteristic pattern of DIP-rich populations, with much lower depth of read alignment in
the middle portion of the segment compared with the termini. As expected, the number of
DIP-associated deletion junctions detected by the pipeline also increased across
passages, reaching the highest level at passage 6 (**Fig 4C**). To confirm that these
junction-containing sequences were derived from virion rather than cellular RNA, we
measured the number of reads that aligned to the host (canine) genome in our samples.
We found very few reads derived from the canine genome in all the passages, compared
with about 40% of the reads from RNA extracted from infected cells (**Fig S4**).

Fig 4. Generation of DIP-rich populations through high MOI passage. Cal07 was serially passaged 6 times (P1-P6) in MDCK cells at sustained high MOI. **(A)** PCR products from the indicated Cal07 populations following 8-segment whole genome amplification, visualized on a 1% agarose gel. The accumulation of deletion junctions is reflected by the disappearance of the polymerase segments (~2.3kb) and the appearance of a smear below the NS segment (~0.9kb) ranging from ~0.3-0.8kb. **(B)** Coverage depth of aligned reads from the indicated passages for PB2 and NS genome segments. The coverage was normalized to the read coverage of the parental sample (Par1). **(C)** Fold increase in the number of deletion junctions (left y axis) and total read support for those junctions (right y axis) over the parental sample (Par1).

298
299

Reproducibility of pipeline performance

Reproducibility of pipeline performance
Multiple steps in the combined experimental/computational pipeline could introduce
stochasticity into the pipeline performance, thus diminishing overall consistency and
reproducibility of output. To examine the reproducibility of our pipeline's performance, we
sequenced two separate extractions of a single P6 population (Hereafter known as L1-
P6-Rep1 and L1-P6-Rep2, where L refers to lineage) and compared the pipeline outputs
between the two replicates (**Fig S5**). We found that the normalized read support values
of individual junctions were highly correlated between the two replicate samples, whether
the replicates were sequenced on the same MiSeq flowcell (Spearman R = 0.92) or
separate ones (Spearman R = 0.91). Thus, the combined steps from RNA extraction to
sequence analysis introduce minimal noise into the pipeline output, and pipeline
performance is highly reproducible between experiments.

Optimization of minimum read support cutoffs

Optimization of minimum read support cutoffs

Our experiments using simulated datasets revealed the importance of setting minimal
RSCs for maximizing the accuracy of pipeline performance, and suggested that the
optimal RSC may differ between datasets. We next attempted to optimize RSC values for
our experimental dataset where we did not actually know the precise location and number
of junctions present in the population (as we did with our simulated datasets). To quantify
precision in junction detection for our experimental dataset, we assumed that base calling
errors and mutations that result in inaccurate junction reporting would be stochastic and
thus read support for these inaccurate junctions would be highly variable between
technical replicates. In contrast, read support for real junctions should be consistent
between replicates.
324
We assessed the effects of varying the RSC on the degree of correlation between
junctions identified in L1-P6-Rep1 and L1-P6-Rep2. We varied the RSC values from 1 to
50 for each individual genome segment, and examined the effect on the number of
reported junctions (**Fig 5**). We observed a similar pattern to that observed for our
simulated data, where raising the RSC to 10 or higher resulted in a large drop-off in the
number of reported junctions. We next determined the RSC value that yielded the highest
degree of correlation between the two replicates. We identified distinct optimal RSC cutoff
values for each segment: 20, 20, 30, 30, and 15 for PB2, PB1, PA, HA, and NA,
respectively. The average of these values was used as an RSC for the remaining
segments where no enough junctions were detected to perform the correlation test (see
below).
336
337 We do not expect these values to be universal, as they likely are influenced by a number
338 of factors that will vary between individual sequencing runs. Also, for different
339 applications, it may be beneficial to lower the RSC to improve detection sensitivity at the
340 cost of precision. Thus, we suggest running two technical replicates with each NGS run
341 to be used as reference to establish optimal per-segment RSC values for that run.
342

Fig 5. Determination of optimal read support cutoffs for experimental data. Plots showing the numbers of deletion junctions reported in the indicated genome segments for two technical library preparation and sequencing replicates generated from a single DIP-rich viral population (L1-P6-Rep1 and L2-P6-Rep2; left y axis). Black dots represent the results of Spearman correlation tests between the replicates at each RSC condition (right y axis). Blue dots indicate the point with the highest degree of correlation and minimum decrease of junction count for each genome segment.

343
344

Analysis of DIP-associated deletion profiles
We next examined the overall diversity of DIP-associated deletion junctions within the P6
populations from the two independent lineages (L1-P6-Rep1 and L2-P6), and found
dozens of distinct deletion junctions scattered across the viral genome in both lineages
(**Fig 6A**). Junctions were not evenly distributed across the genome segments, as few to
no junctions were detected in the NP, M, or NS segments. Within each segment, the read
support for individual junctions varied significantly (**Fig 6B**). When we compared the
deletion junction repertoires between the two passage lineages, we observed that a
significant fraction of the detected junctions was shared between the two, and that these
shared junctions exhibited a high degree of correlation in terms of read support (**Fig 6C**).
These data suggest that specific DIP-associated deletions may be consistently formed by
Cal07. While there was substantial diversity in terms of the number of distinct deletion
junctions present, when we plotted the locations of those these junctions within the
genome segments, we observed that they were largely confined within clear hotspots
towards the termini of the segments with few exceptions (**Fig 6D**).
359
360

Fig 6. DIP-associated junctions analysis. (A) The total number of detected junctions in the individual genome segments for the two independent passage 6 populations. **(B)** Stacked column charts showing the proportional abundance of each deletion junction per segment between lineage 1 (including both technical replicates) and lineage 2 at passage 6. Each color bar represents a unique junction within each segment, whose height reflects the relative NGS read support, normalized to the total number of NGS junction-spanning reads for the indicated segment. **(C)** Comparison between L1-P6-REP1 and L2-P6 in relation to the number of shared junctions (upper panel) and the correlation between their NGS reads (lower panel) **(D)** Parallel coordinates diagrams showing the deletion junctions in P6-Rep1 mapped to their actual respective locations on each segment. Each individual junction is represented by a black line that connects the donor and acceptor sites of the breakpoint.

361
362

Effect of varying template input on pipeline performance
We next asked whether the amount of cDNA template that goes into the library
preparation affects the sensitivity and stochasticity of junction detection by the pipeline.
We serially diluted both the amount of viral RNA template used in the RT reaction and the
amount of cDNA template used in the PCR and compared pipeline outputs from the DIP-
rich L1-P6-Rep1 population. We first tested the correlation of detected DIP-associated
junctions between a limited number of dilutions ranging from 1:3 to 1:15. We observed
that the correlation of read support values between specific junctions across dilutions was
more consistent when cDNA was diluted, rather than RNA, suggesting that RNA dilution
may increase the stochasticity of downstream PCR amplification (**Fig S6A**).

Based on this, we performed whole genome PCR using a dilution series of L1-P6-Rep1-
derived cDNA (spanning roughly $4*10^8$ to $4*10^6$ NP genome equivalents per PCR) as
template (**Fig 7A**). We observed that there is an optimal amount of input cDNA template
for maximizing junction detection. Diluting the input cDNA 1:120 (corresponding to ~$4*10^6$
NP genome equivalents) increased the number of detected junctions over 4-fold
compared with undiluted input. Although the number of DIP-associated junctions was
increased, the distribution of junctions across segments and their mapped locations were
consistent with our earlier results (**Fig S6B and Fig 6**).

Further dilution of input template beyond 1:120 resulted in a decrease in sensitivity.
Importantly, dilution across the range tested did not result in a failure to detect any of the
junctions reported in the undiluted sample. We also observed that the correlation of read
support values between specific junctions across dilutions tracked closely with the
sensitivity (**Fig 7B**). Altogether, these observations indicate that optimization of the cDNA

template input amount can significantly improve the sensitivity of DIP-associated junction detection.

Fig 7. Effects of viral template input on the detection of DIP-associated junctions. We serially diluted cDNA generated from the L1-P6-Rep1 sample, and compared sequencing results between libraries generated with these dilutions as templates. **(A)** For each dilution, the total numbers of detected junctions (purple) are shown, along with the number of specific junctions detected that were also detected in the undiluted sample (grey). The copy number of viral cDNA molecules included in downstream PCR and library preparation for each dilution was determined by RT-qPCR (green; right y axis). **(B)** Read support values for all deletion junctions common across the diluted and undiluted samples were normalized to the total number of deletion junction-spanning reads for each sample and used to perform a Spearman correlation between all pairs of samples using R cor function.

Lack of association between direct repeats and junction formation

Direct repeat sequences (detailed in **Fig S2 and S3A**) are common across the IAV genome and have previously been hypothesized to contribute to DIP-associated deletion formation by promoting viral polymerase slippage (10, 15). We leveraged the large number of DIP-associated deletion junctions that we identified in this study to test this hypothesis. We asked whether the deletion junctions in the DIP-enriched sample L1-P6-Rep1 were found more frequently adjacent to direct repeats than would be expected if the junctions were located randomly in the viral genome. We compared the frequency of deletion junctions associated with direct repeats between the L1-P6-Rep1 and L2-P6 populations (where all deletions are formed by the viral RdRp) and the Cal07-200 simulated dataset, where all deletions are randomly localized (**Fig 8**). The frequency of direct repeats of varying lengths at junction sites in the real viral populations was not significantly different than that seen in the simulated data, indicating that direct repeat sequences are not enriched at DIP-associated junctions and arguing against a significant role for direct repeats in DIP formation.

	Chi square P value
Simulated vs P6-Rep1	0.65
Simulated vs Bio-Rep	0.79
P6-Rep1 vs Bio-Rep	0.95

Fig 8. Direct repeat sequences are not over-represented at DIP-associated deletion junctions. The percentages of deletion junctions within the polymerase segments that occurred at unique sites or at sites with direct nucleotide repeats with lengths 2-4nts was compared between L1-P6-Rep1, L2-P6, and the Cal07-200 simulated dataset. The number of junctions was plotted and compared by Chi Square. The table shows the Chi Square P values between every possible pair of the samples.

409
410
DISCUSSION
Sensitive and accurate detection of DIP-associated sequences within viral populations is
critical for defining how DIPs form and function during IAV infection. Here we outline a
pipeline to detect DIP-associated junctions within viral populations using Illumina-based
short read sequencing, and validate its performance using a combination of simulated
and experimental control datasets.
417
Our primary goal was to develop and optimize a reasonably simple and straightforward
sequencing framework that accounts for the potential artifacts that can potentially
confound NGS-based DIP detection efforts. We chose the Illumina sequencing platform
because it is widely available, easy to use, conducive to sample multiplexing, and
because it has a relatively low rate of base-calling errors. One concern that we had initially
was that recombination during reverse transcription, PCR, or sequencing might make
identification of bona fide DIP-associated junctions a challenge. Two recently developed
technologies, CirSeq and ClickSeq, largely eliminate this issue, but also significantly
increase the amount of labor involved in library preparation (22, 23). We observed that
the occurrence of non-viral recombination that occurs during our library preparation and
sequencing procedures was vanishingly small, and can effectively be ignored. Thus, while
both Cirseq and ClickSeq are enormously useful in certain circumstances, our data

indicates that such methods are not required to generate highly accurate and sensitive
profiles of IAV DIPs.
432
A significant shortcoming of the method we detail here is that the measured read support
for individual deletion junctions does not necessarily reflect the actual frequencies of
these deletions within the viral population. This is due to both biasing of PCR amplification
towards shorter products, as well as the uneven distribution of read coverage across the
viral genome. For situations where the accurate measurement of individual DIP genotype
frequencies is critical, we recommend pairing a cDNA barcoding method such as primer
ID (24, 25) with a platform capable of long-read sequencing, such as PacBio or Oxford
Nanopore (26). Alternatively, direct sequencing of viral RNA using the Oxford Nanopore
platform may also prove to be useful for accurate measurement of junction frequencies
(27).
443
When we used our pipeline to examine DIP-enriched viral populations generated through
serial high-MOI passage, we detected dozens of distinct DIP-associated deletion
junctions, revealing a high degree of diversity within the DIP population. Although the
majority of these DIP-associated junctions were derived from the polymerase segments
as expected, we also detected a substantial proportion of deletions within the HA and NA
segments, but not the NP, M, and NS segments. The non-random distribution of junctions
across the genome segments mirrors what has been reported elsewhere, and highlights
how little we know about the specific molecular mechanisms that regulate DIP formation.
452
453 We hope that the approach detailed here, and the associated bioinformatics pipeline
454 prove useful to other groups interested in defective interfering particle biology. Our
455 approach is optimized for influenza virus sequences; however, the approaches and
456 controls detailed here can easily be adapted to other RNA virus systems.
457
MATERIALS AND METHODS
Viruses and Cells
Madin-Darby canine kidney cells (MDCK; obtained from Dr. Jonathan Yewdell) and
human embroyonic kidney 293 cells (293T; obtained from Dr. Joanna Shisler) were grown
in Minimal Essential Medium (MEM) + GlutaMAX (Gibco), supplemented with 8.3% fetal
bovine serum (Seradigm), at 37°C and 5% CO2. Recombinant A/California/07/09 (Cal07)
virus was rescued via the standard 8-plasmid reverse genetics approach. Briefly, 60-90%
confluent 293T cells were transfected with 500ng of the following plasmids (pDZ::PB2,
pDZ::PB1, pDZ::PA, pDZ::HA, pDZ::NP, pDZ::NA, pDZ::M, pDZ::NS) using JetPRIME
(Polyplus) according to the manufacturer's instructions. Cal07 reverse genetics plasmids
were originally obtained as A/California/04/2009-encoding plasmids from Dr. Jonathan
Yewdell. We introduced A660G and A335G substitutions into the HA and NP plasmids,
respectively, to convert them to match the amino acid sequence of A/California/07/2009
HA and NP (NCBI accession# CY121680, CY121683). A seed stock was prepared by
amplifying a plaque isolate from the rescue supernatants. Virus working stocks were
generated by infecting MDCK cells with seed stock at an MOI of 0.001 TCID50/cell and

collecting and clarifying supernatants at 48 hpi.

Generation of DIP stocks through high MOI passage
476   Confluent MDCK cells in 96-well plates were infected with IAV Cal07 at an MOI of 5
477   TCID50/cell. Supernatants (200μl total per well) were harvested at 24 hpi (passage 1)
478   and pooled. 100μl/well of this pooled supernatant was used to infect a 96 well plate of
479   fresh MDCK cells to generate the next passage. This process was repeated 6 times to
480   produce passages 1-6 in two independent lineages (1 96-well plate per lineage).

IAV Genome amplification
Viral RNA was extracted from 140μl of cell culture supernatant using the QIAamp viral
RNA kit (Qiagen) and eluted in 60μl distilled H2O (dH2O). For cDNA reactions, 3μL of
RNA was mixed with 1μL (2μM) MBTUni-12 primer (5'-
ACGCGTGATCAGCRAAAGCAGG-3') + 1μL (10μM) dNTPs + 8μL dH2O. The mixture
was incubated for 5 minutes at 65°C and then placed on ice for 2 min. Subsequently, the
mixture was removed from ice and the following was added: 1μL SuperScript III RT
(Invitrogen), 4μL of 5X First-Strand Buffer (Comes with SSIII kit), 1μL of DTT, 1μL RNase-
in (Invitrogen). The reaction was incubated at 45°C for 50 min, followed by a 15 min
incubation at 70°C for inactivation. 5μL of cDNA product was mixed with the following for
PCR amplification: 2.5μL (10μM) MBTUni-12_4R primer (5'-
ACGCGTGATCAGCRAAAGCAGG-3'), 2.5μL (10μM) MBTUni-13 primer (5'-
ACGCGTGATCAGTAGAAACAAGG-3'), 0.5μL Phusion polymerase (NEB), 10μL - 5x HF
buffer, 1μL (10mM dNTPs mix), and 28.5μL dH2O. The PCR reaction conditions used:
98°C (30 s) followed by 25 cycles of 98°C (10 s), 57°C (30 s) and 72°C (1:30 min), a
terminal extension of 72°C (5 min), and a final 10°C hold. PCR products were purified
using the PureLink PCR purification kit (Invitrogen) with the <300nt cutoff option and
eluted in 30μL dH2O. There was no difference in deletion junction detection when we
purified the PCR products with the lower cutoff option (data not shown).

NGS library preparation
We started with ~20ng of the PCR products in a volume of 50μl. The Covaris M220
sonicator (Covaris) was used to fragment the DNA. Three different conditions were used
to generate different average fragment lengths of 300, 500, 700 base pairs (bp): (I) 300
bp = Peak Power 50, Duty Factor 20 and Cycles/Burst 200 for 2:40 min, (II) 500 bp =
Peak Power 50, Duty Factor 10 and Cycles/Burst 200 for 1:30 min, and (III) ~600 bp =
fragment length Peak Power 50, Duty Factor 10 and Cycles/Burst 200 for 1 min. In our
hands, the fragmentation length did not have any effect on our sequencing results (data
not shown). For the sake of consistency, we used the 300 bp fragmentation length. To
confirm the PCR products, we visualized the amplicons on a Fragment Analyzer (AATI)
with the DNF-486 high sensitivity NGS kit before and after fragmentation. Next, we used
KAPA Hyper Prep kit (Roche) to construct the libraries according to the manual. To
eliminate the possibility of index hopping (or index switching), we used the TruSeq Unique
Dual Indexes (UDI) from Illumina. The Adapter ligation step was carried out with 5μl of
Truseq UDIs diluted 1:10 with 10nM Tris. For maximum efficiency we increased the
ligation time to 30mins. We then performed 3 cycles of PCR with the Kapa library

amplification primers diluted 1:5 in water followed by a cleanup step with 40μl of AxyPrep
Mag PCR beads (Thermofisher). We then mixed the libraries at an equimolar ratio and
carried out a qPCR to accurately quantitate the library pool and maximize the number of
clusters in the sequencing flowcell. A size selection step was not needed. Finally, the
pooled libraries were sequenced with paired-ends 2x250nt reads on an Illumina MiSeq
using V2 chemistry. The fastq files were generated and demultiplexed with the bcl2fastq
v2.20 Conversion Software (Illumina).
524
Simulated Datasets
All the simulated datasets used in this study were generated by MetaSim (v0.9.1) (28),
a genomic and metagenomics simulator. Several reference library sequences composed
of WT reference sequences of IAV Cal07 or PR8 (see Table S1 for NCBI accession
numbers), mixed with a defined DIP sequence population - generated randomly within
the first and last 600 nts of all the segments - were used in Metasim for data simulation.
The configurations were fixed across all datasets to maintain the preferable conditions.
The reference sequences were fragmented into 350 nts fragments length with a standard
deviation of +/- 50 and were simulated into ~1 million 2x250nts paired-end reads per
sample, with a total mutation rate of ~1% based on the published Illumina empirical error
model, and corresponds to substitutions as the indel error rate is negligible within Illumina
MiSeq. One dataset was simulated with no DIP sequences as a control sample for any
computational artifacts. Metasim generated two FASTA files of 1 million reads per file per
sample (~2 million single-end reads = 1 million paired-end reads), which subsequently
were used for the optimization process.
Sequencing analysis of DIP-associated junctions
The raw sequencing reads were quality-filtered by Trimmomatic (v0.36) (Parameters:
`ILLUMINACLIP:TruSeq3-PE-2.fa:2:15:10` `SLIDINGWINDOW:3:20`
`LEADING:28 TRAILING:28`) (29) and any reads shorter than 75nts were removed from
the datasets. The paired reads were concatenated into one file and treated as single-end
when aligned end-to-end to the WT reference sequences using Bowtie2 (v2.3.1)
(Parameters: --score-min `L,0,-0.3`). Subsequently, the algorithm ViReMa (v0.10) was
used to analyze the remaining un-aligned reads (putative junction-spanning reads)
(Parameters `-DeDup --MicroInDel_Length 20 --Defuzz 3 --N 1 --X 8`).
Next, the DIP-associated deletion junctions and their read support were extracted from
ViReMa output files and sorted per segment, using an in-house Perl scrip, for data
analysis and visualization. To detect any MDCK genome leakage, the datasets were
aligned against the dog genome (assembly CanFam3.1). All scripts are available at
https://github.com/BROOKELAB/Influenza-virus-DI-identification-pipeline.
554
Quantification of sensitivity and precision
To calculate the actual number of junction-spanning reads in Fig 2A, reads that derived
from DIP-associated sequences were counted by their FASTA headers, which contain
the source of each read, produced by MetaSim. To calculate the maximum theoretical
sensitivity of ViReMa (Fig 2B) based on seed length of 25nts and two allowed mutations
(--N = 2), the number of mutations was subtracted from the seed length, which on its turn

was multiplied by 2 to account for both termini ((25-2)*2=46). Subsequently, this number was subtracted from the possible cutting site of a 250nts read and divided by the total number of cutting sites and multiplied by 100 ((249-46)/249*100)=81.5%). To calculate the number of accurately and inaccurately mapped junctions in Fig 2C,D, the seed sequences of the Cal07-200 dataset were used against ViReMa with --N set to 0, and the remaining parameters were kept the same. These sequences were generated initially to establish the seed for MetaSim to simulate sequencing, therefore their lengths are varied between ~350-1800. The long ones were trimmed to <1000nts, so ViReMa would take them as reads (the maximum default read length that ViReMa could take is ~1024nts) and, critically, the junction locations were maintained. The junctions that occurred within the first or last 25nts were removed (4 junction sequences). Finally, the junctions that accurately mapped were counted, which found to be 149 versus 47 inaccurately mapped junctions.

Correlation analysis
For the correlation tests, the NGS read support count for each DIP-associated junction was normalized to the total detected junction-spanning reads of every sample. Next, the correlation was calculated based on Spearman rank correlation using either R (cor function) or an online tool at:
http://www.biostathandbook.com/spearman.html

ACKNOWLEDGMENTS
We are grateful to other members of the lab for helpful comments and critical reading of the manuscript, as well as to Dr. Alvaro Hernandez and the staff at the high-through sequencing and genotyping unit within the Roy J. Carver Biotechnology Center for excellent advice and technical assistance. This work was generously funded by the Defense Advanced Research Projects Agency under contract DARPA-16-35-INTERCEPT-FP-018.

REFERENCES
1. Von Magnus P. 1954. Incomplete forms of influenza virus. Adv Virus Res 2:59–79.

2. von MAGNUS P. 1951. Propagation of the PR8 strain of influenza A virus in chick embryos. II. The formation of incomplete virus following inoculation of large doses of seed virus. Acta Pathol Microbiol Scand 28:278–293.

3. Rezelj VV, Levi LI, Vignuzzi M. 2018. The defective component of viral populations. Curr Opin Virol 33:74–80.

4. Baum A, Sachidanandam R, García-Sastre A. 2010. Preference of RIG-I for short viral RNA molecules in infected cells revealed by next-generation sequencing. Proc Natl Acad Sci U S A 107:16303–16308.

5.   Nayak DP, Chambers TM, Akkina RK. 1985. Defective-interfering (DI) RNAs of
influenza viruses: origin, structure, expression, and interference. Curr Top Microbiol
Immunol 114:103–151.

6.   Brooke CB. 2017. Population Diversity and Collective Interactions during Influenza
Virus Infection. J Virol 91.

7.   Vasilijevic J, Zamarreño N, Oliveros JC, Rodriguez-Frandsen A, Gómez G,
Rodriguez G, Pérez-Ruiz M, Rey S, Barba I, Pozo F, Casas I, Nieto A, Falcón A.
2017. Reduced accumulation of defective viral genomes contributes to severe
outcome in influenza virus infected patients. PLOS Pathog 13:e1006650.

8.   Sherry L, Punovuori K, Wallace LE, Prangley E, DeFries S, Jackson D. 2016.
Identification of cis-acting packaging signals in the coding regions of the influenza B
virus HA gene segment. J Gen Virol 97:306–315.

9.   Hutchinson EC, von Kirchbach JC, Gog JR, Digard P. 2010. Genome packaging in
influenza A virus. J Gen Virol 91:313–328.

10.   Jennings PA, Finch JT, Winter G, Robertson JS. 1983. Does the higher order
structure of the influenza virus ribonucleoprotein guide sequence rearrangements in
influenza viral RNA? Cell 34:619–627.

11.   Routh A, Johnson JE. 2014. Discovery of functional genomic motifs in viruses
with ViReMa-a Virus Recombination Mapper-for analysis of next-generation
sequencing data. Nucleic Acids Res 42:e11.

12.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods 9:357–359.

13.   Jennings PA, Finch JT, Winter G, Robertson JS. 1983. Does the higher order
structure of the influenza virus ribonucleoprotein guide sequence rearrangements in
influenza viral RNA? Cell 34:619–627.

14.   Janda JM, Davis AR, Nayak DP, De BK. 1979. Diversity and generation of
defective interfering influenza virus particles. Virology 95:48–58.

15.   Saira K, Lin X, DePasse JV, Halpin R, Twaddle A, Stockwell T, Angus B, Cozzi-
Lepri A, Delfino M, Dugan V, Dwyer DE, Freiberg M, Horban A, Losso M, Lynfield R,
Wentworth DN, Holmes EC, Davey R, Wentworth DE, Ghedin E, INSIGHT FLU002
Study Group, INSIGHT FLU003 Study Group. 2013. Sequence analysis of in vivo
defective interfering-like RNA of influenza A H1N1 pandemic virus. J Virol 87:8064–
8074.

16.   Loman NJ, Misra RV, Dallman TJ, Constantinidou C, Gharbia SE, Wain J, Pallen
MJ. 2012. Performance comparison of benchtop high-throughput sequencing
platforms. Nat Biotechnol 30:434–439.

17.   Schirmer M, Ijaz UZ, D'Amore R, Hall N, Sloan WT, Quince C. 2015. Insight into
biases and sequencing errors for amplicon sequencing with the Illumina MiSeq
platform. Nucleic Acids Res 43:e37.

18.   Zhou B, Donnelly ME, Scholes DT, St George K, Hatta M, Kawaoka Y,
Wentworth DE. 2009. Single-reaction genomic amplification accelerates sequencing
and vaccine production for classical and Swine origin human influenza a viruses. J
Virol 83:10309–10313.

19.   Görzer I, Guelly C, Trajanoski S, Puchhammer-Stöckl E. 2010. The impact of
PCR-generated recombination on diversity estimation of mixed viral populations by
deep sequencing. J Virol Methods 169:248–252.

20.   Lahr DJG, Katz LA. 2009. Reducing the impact of PCR-mediated recombination
in molecular evolution and environmental studies using a new-generation high-
fidelity DNA polymerase. BioTechniques 47:857–866.

21.   Xue J, Chambers BS, Hensley SE, López CB. 2016. Propagation and
Characterization of Influenza Virus Stocks That Lack High Levels of Defective Viral
Genomes and Hemagglutinin Mutations. Front Microbiol 7:326.

22.   Routh A, Head SR, Ordoukhanian P, Johnson JE. 2015. ClickSeq:
Fragmentation-Free Next-Generation Sequencing via Click Ligation of Adaptors to
Stochastically Terminated 3'-Azido cDNAs. J Mol Biol 427:2610–2616.

23.   Acevedo A, Andino R. 2014. Library preparation for highly accurate population
sequencing of RNA viruses. Nat Protoc 9:1760–1769.

24.   Kosik I, Ince WL, Gentles LE, Oler AJ, Kosikova M, Angel M, Magadán JG, Xie
H, Brooke CB, Yewdell JW. 2018. Influenza A virus hemagglutinin glycosylation
compensates for antibody escape fitness costs. PLOS Pathog 14:e1006796.

25.   Jabara CB, Jones CD, Roach J, Anderson JA, Swanstrom R. 2011. Accurate
sampling and deep sequencing of the HIV-1 protease gene using a Primer ID. Proc
Natl Acad Sci U S A 108:20166–20171.

26.   Jaworski E, Routh A. 2017. Parallel ClickSeq and Nanopore sequencing
elucidates the rapid evolution of defective-interfering RNAs in Flock House virus.
PLOS Pathog 13:e1006365.

27.   Garalde DR, Snell EA, Jachimowicz D, Sipos B, Lloyd JH, Bruce M, Pantic N,
Admassu T, James P, Warland A, Jordan M, Ciccone J, Serra S, Keenan J, Martin

668  S, McNeill L, Wallace EJ, Jayasinghe L, Wright C, Blasco J, Young S, Brocklebank
669  D, Juul S, Clarke J, Heron AJ, Turner DJ. 2018. Highly parallel direct RNA
670  sequencing on an array of nanopores. Nat Methods 15:201–206.

671 28. Richter DC, Ott F, Auch AF, Schmid R, Huson DH. 2008. MetaSim—A
672  Sequencing Simulator for Genomics and Metagenomics. PLOS ONE 3:e3373.

673 29. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for
674  Illumina sequence data. Bioinforma Oxf Engl 30:2114–2120.

675 30. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis
676  G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The
677  Sequence Alignment/Map format and SAMtools. Bioinforma Oxf Engl 25:2078–
678  2079.

679 31. Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing
680  genomic features. Bioinforma Oxf Engl 26:841–842.

681