1

A simple microbiome in the European common cuttlefish, *Sepia officinalis*

Holly L. Lutz[a,b,c,†,*], S. Tabita Ramírez-Puebla[d,†], Lisa Abbo[d], Amber Durand[d], Cathleen

Schlundt[d], Neil Gottel[a,b], Alexandra K. Sjaarda[e], Roger T. Hanlon[d], Jack A. Gilbert[a,b,c] , Jessica

L. Mark Welch[d,*]

[a]Department of Pediatrics, University of California San Diego, La Jolla, California, USA.

[b]Scripps Institution for Oceanography, UCSD, La Jolla, California, USA

[c]Integrative Research Center, Field Museum of Natural History, Chicago, Illinois, USA

[d]Marine Biological Laboratory, Woods Hole, MA 02543

[e]Biological Sciences Division, University of Chicago, Chicago, IL 60637

[†] co-first authors, contributed equally to this work

*to whom correspondence should be addressed. Email address: hlutz@fieldmuseum.org or

jmarkwelch@mbl.edu

ABSTRACT

The European common cuttlefish, *Sepia officinali*s, is used extensively in biological and

biomedical research yet its microbiome remains poorly characterized. We analyzed the

microbiota of the digestive tract, gills, and skin in mariculture-raised *S. officinalis* using a

combination of 16S rRNA amplicon sequencing, qPCR and fluorescence spectral

imaging. Sequencing revealed a highly simplified microbiota consisting largely of two

single bacterial amplicon sequence variants (ASVs) of Vibrionaceae and Piscirickettsiaceae. The

esophagus was dominated by a single ASV of the genus *Vibrio*. Imaging revealed bacteria in the

family Vibrionaceae distributed in a discrete layer that lines the esophagus. This *Vibrio* was also

the primary ASV found in the microbiota of the stomach, cecum, and intestine, but occurred at

lower abundance as determined by qPCR and was found only scattered in the lumen rather than

in a discrete layer via imaging analysis. Treatment of animals with the commonly-used antibiotic

enrofloxacin led to a nearly 80% reduction of the dominant *Vibrio* ASV in the esophagus but did

not significantly alter the relative abundance of bacteria overall between treated versus control

animals. Data from the gills was dominated by a single ASV in the family Piscirickettsiaceae,

which imaging visualized as small clusters of cells. We conclude that bacteria belonging to the

Gammaproteobacteria are the major symbionts of the cuttlefish *Sepia officinalis* cultured from

eggs in captivity, and that the esophagus and gills are major colonization sites.

IMPORTANCE

Microbes can play critical roles in the physiology of their animal hosts, as evidenced in

cephalopods by the role of *Vibrio (Aliivibrio) fischeri* in the light organ of the bobtail squid and

the role of Alpha- and Gammaproteobacteria in the reproductive system and egg defense in a

variety of cephalopods. We sampled the cuttlefish microbiome throughout the digestive tract,

gills, and skin and found dense colonization of an unexpected site, the esophagus, by a microbe

of the genus *Vibrio*, as well as colonization of gills by Piscirickettsiaceae. This finding expands

the range of organisms and body sites known to be associated with *Vibrio* and is of potential

significance for understanding host-symbiont associations as well as for understanding and

maintaining the health of cephalopods in mariculture.

KEYWORDS:

microbiome, fluorescence *in situ* hybridization, Cephalopoda, Vibrionaceae, Piscirickettsiaceae,

enrofloxacin

1. INTRODUCTION

Symbiotic associations between invertebrates and bacteria are common. Among cephalopods, the

most intensely studied association is the colonization of the light organ of the bobtail squid

Euprymna scolopes by the bioluminescent bacterium *Vibrio (Aliivibrio) fischeri* in a highly

specific symbiosis (1). A more diverse but still characteristic set of bacteria colonize the

accessory nidamental gland from which they are secreted into the egg jelly coat and likely

protect the eggs from fungal and bacterial attack (2). The accessory nidamental gland and egg

cases of the squid *Doryteuthis* (*Loligo) pealeii* and the Chilean octopus (*Octopus mimus*) have

also been reported to contain Alphaproteobacteria and Gammaproteobacteria (3, 4). These

associations indicate that bacteria can play a key role in the physiology of cephalopods.

*Sepia officinali*s, the European common cuttlefish (hereafter cuttlefish), is used

extensively in biological and biomedical research (5-7) and is a model organism for the study of

rapid adaptive camouflage (8-11). Cuttlefish are also widely represented among zoos and

aquaria, and play an important role in educating the public about cephalopod biology and life

history (12). Little is known about the association of bacterial symbionts with cuttlefish, and

whether such associations may play a role in the health or behavior of these animals.

Understanding the importance, or lack thereof, of the cuttlefish microbiome will not only shed

light on the basic biology of this model organism but will also have important implications for

future husbandry practices and research design.

Using a combination of 16S rRNA amplicon sequencing, fluorescence *in situ*

hybridization (FISH), and quantitative PCR (qPCR), we characterized the gastrointestinal tract

(GI), gill, skin, and fecal microbiota of the common cuttlefish in wild-bred, captive-raised

animals (5) housed at the Marine Biological Laboratory (Woods Hole, MA). We observed a

highly simplified microbiome dominated by Vibrionaceae in the gastrointestinal tract and

Piscirickettsiaceae in the gills. We treated a subset of cuttlefish with antibiotic enrofloxacin,

commonly used among aquaria veterinarians, and found both ASVs to remain dominant in

esophagus and gill microbiota, suggesting they are resilient to this antibiotic. The simplicity of

this system makes it a promising model for further exploration of the factors driving host-

symbiont associations in marine invertebrates.

2. RESULTS

2. 1 Two taxa dominate the *S. officinalis* microbiome.

We sampled 27 healthy adult cuttlefish (*Sepia officinalis*) from the mariculture laboratory

at Marine Biological Laboratory (Woods Hole, MA). The study comprised two time periods. The

first (20-21 June 2017) was a pilot survey in which three individuals were sampled. The second

(25 September -10 October 2017) was an experiment involving 24 individuals, of which 16 were

exposed to repeated doses of the antibiotic enrofloxacin and 8 individuals served as untreated

controls. 16S rRNA amplicon sequencing of the GI tract, gills, and skin of all 27 animals

revealed a highly simplified microbiota dominated by bacterial amplicon sequence variants

(ASVs) in the Vibrionaceae and Piscirickettsiaceae families, regardless of treatment with

enrofloxacin.

In particular, results showed a consistent and highly simplified microbiota in the

esophagus (Figure 1; Table 1). A single ASV in the genus *Vibrio* (referred to as ASV1 in

subsequent figures and tables) made up the majority of the 16S rRNA sequence data from the

esophagus of the three pilot investigation individuals (mean $92\% \pm 10\%$) and of 24 individuals

sampled four months later (control group mean $100\% \pm 1\%$; treatment group mean $94\% \pm 10\%$).

Thus, this ASV represents a dominant constituent of the esophagus microbiota stably over two

time periods in the study and after exposure to antibiotic treatment with enrofloxacin. Another

ASV of the related genus *Photobacterium* (Vibrionaceae) (referred to as ASV2 in Table 1) was

present in the esophagus community in the pilot investigation animals (mean $7\% \pm 10\%$).

Combined, the two Vibrionaceae ASVs (ASV1 and ASV2) in the three pilot animals constituted

>99% of the esophagus community.

The major *Vibrio* ASV1 was also a major constituent of downstream sites in the GI tract,

although present at lower abundance measured both as relative abundance in 16S sequencing

data and by quantitative PCR (qPCR) (Figure 1; Table 1). qPCR revealed a high abundance of

Vibrio cell copies in the esophagus (average 520.4 ± 410 copies/ng of total DNA including host

DNA) relative to more distal portions of the GI tract that included stomach, cecum, and intestine

(combined average 25.6 ± 77.8 copies/ng) ($p < 0.005$, $\chi^2 = 16.5$, df =3 Kruskal-Wallis) (Fig. 1B).

Comparison of qPCR measures of ASVs in the genus *Vibrio* between the esophagus of treatment

and control animals revealed a striking and significant decrease of nearly 80% in the quantity (p

< 0.02, df = 10.8, Welch two sample t-test). We did not observe a significant difference in *Vibrio*

ASV1 quantity between other organs of the digestive tract with antibiotic treatment (Fig. 1B).

Relative abundance of *Vibrio* ASV1 in the esophagus and stomach also did not differ

significantly between antibiotic and control groups (Welch two sample t-test, $p > 0.10$), but did

differ significantly among the cecum (Welch two sample t-test, $p = 0.00235$) and intestine

(Welch two sample t-test, $p = 1.99e{-}05$) of treatment versus control groups. Analysis of GI tracts

(esophagus, stomach, cecum, and intestine samples combined) between treated versus control

animals from the second period of study revealed significant differences in weighted UniFrac β-

diversity between the two groups (PERMANOVA Pr(>F) = 0.006, F = 5.63, df = 1), despite the

Vibrio ASV1 remaining dominant in most organs. These differences in measured relative

abundance and β-diversity may result from stochastic variation in low-abundance sequences, as

the non-*Vibrio* portion of the 16S rRNA sequence data from the GI tract consisted of an

assortment of taxa that varied between individuals or between the two time points of the study

and thus suggested transient organisms rather than stable microbial colonization.

Samples from gills were dominated by a single highly abundant ASV in the family

Piscirickettsiaceae (referred to as ASV3 in subsequent figures and tables), which made up an

average relative abundance of $96.9\% \pm 2.5\%$ in the gills. In samples from other body sites this

ASV3 was detected only sporadically and at low relative abundance (mean 0.2%, range 0 to

5.8%) (Fig. 1A). An additional Vibrio ASV, ASV4, was a major constituent of the microbiota of

the shrimp used as food for the cuttlefish and was also detectable in some samples from stomach,

cecum, and intestine (Figure 1A). Skin samples did not exhibit much similarity to GI tract or

gills with respect to microbiome composition, with most common ASVs found in other

anatomical sites comprising $< 20\%$ (mean $17.2\% \pm 2.9\%$) relative abundance of the

microbiome (Table S1).

In addition to surveying internal organs, we collected fecal samples from the 24 animals

from the second time period of our study. These samples were collected daily for each

individual throughout the course of the antibiotic treatment experiment (see methods).

Comparison of weighted UniFrac dissimilarity of fecal samples from experimental animals

revealed significant differences in beta-diversity between cuttlefish fecal samples, seawater, and

shrimp (PERMANOVA Pr(>F) = 0.001, F = 5.26, df = 2) upon which the animals were fed.

These results, paired with the differences we observed in compositional relative abundance

between organs, seawater, and shrimp provide additional support for our finding that bacterial

communities associated with cuttlefish differ from those found in their seawater environment and

food source (Fig. 2).

2.2 Imaging shows spatial structure of the microbiota in cuttlefish esophagus and scattered

distribution elsewhere in the gastrointestinal tract.

Fluorescence *in situ* hybridization (FISH) revealed a striking organization of bacteria

distributed in a layer lining the interior of the convoluted esophagus of cuttlefish (Fig. 3A-C).

Hybridization with the near-universal Eub338 probe showed bacteria in high density in a layer

~20-40 μm thick at the border between host tissue and lumen. Staining with fluorophore-

conjugated wheat germ agglutinin revealed a mucus layer that covered the epithelium and

generally enclosed the bacteria (Fig. 3). To verify the identity of these bacteria we employed a

nested probe set including Eub338 as well as probes for Alphaproteobacteria,

Gammaproteobacteria, and probes we designed specifically for Vibrionaceae (Vib1749 and

Vib2300, Table 2). Bacterial cells imaged in the esophagus showed signal from all probes

expected to hybridize with Vibrionaceae, suggesting that the bacteria observed in this organ are a

near-monoculture of this taxon (Fig. 4B-E). A probe targeted to Alphaproteobacteria was

included in the FISH as a negative control and, as expected, did not hybridize with the cells (Fig.

4F). As an additional control to detect non-specific binding of probes, we performed an

independent FISH with a set of probes labeled with the same fluorophores as the experimental

probe set but conjugated to oligonucleotides not expected to hybridize with the cuttlefish

microbiota (Table 2). No signal from this non-target probe set was detected (Fig. 4G-H)

supporting the interpretation that the signal observed in the esophagus results from a specific

interaction of the Vibrionaceae-targeted oligonucleotides with the visualized bacteria.

In other parts of the digestive tract we observed a sparser distribution of bacteria without

obvious spatial organization. Bacteria in the intestine were present not in a layer but scattered

throughout the lumen and mixed with the luminal contents (Fig. 5). Similarly, in the cecum, we

observed bacteria in low abundance in the lumen (Fig. 6). FISH was also applied to the stomach,

posterior salivary gland (poison gland) and duct of the salivary gland, but no bacteria were

detected (not shown).

Fluorescence *in situ* hybridization to cross-sections of the gills revealed clusters of

bacteria at or near the surface of the tissue (Fig. 7). These bacteria hybridized with the Eub338

near-universal probe and a probe for Gammaproteobacteria (Fig. 7B, C, Table 2) but not

Alphaproteobacteria (not shown), consistent with the identification of the clusters of gill bacteria

as members of the gammaproteobacterial family Piscirickettsiaceae.

3. DISCUSSION

We sampled the cuttlefish microbiome of the digestive tract, gills, and skin and found

dense colonization of an unexpected site, the esophagus, by a bacterium of the genus *Vibrio*.

Both imaging and 16S rRNA gene sequencing showed a near-monoculture of *Vibrionaceae* in

the esophagus, with imaging showing dense colonization of the interior lining of the esophagus

with a single morphotype that hybridized to probes targeting *Vibrionaceae*. In the remainder of

the GI tract, both imaging and 16S rRNA sequencing indicated a less consistent microbiota.

Sequencing also showed lower relative abundance of the dominant *Vibrio* ASV, and qPCR

confirmed a significantly lower total abundance of *Vibrio* cell copies in the distal GI tract.

Imaging revealed sparse and sporadic colonization in the intestine and cecum, with scattered

cells in the lumen and no clear colonization of the epithelium. In light of these results, we

conclude that the GI tract of laboratory cultured *Sepia officinalis* has a highly simplified

microbiome dominated by the genus *Vibrio*.

Diverse associations with *Vibrio* and the Vibrionaceae are known from cephalopods.

Among the most extensively investigated is the mutualistic association of the bioluminescent

Vibrio (Aliivibrio) fischeri with the light organ of the bobtail squid *Euprymna scolopes* (1, 13).

Other well-known symbioses include the colonization of the cephalopod accessory nidamental

gland with Alpha- and Gammaproteobacteria, which enables the host to secrete a layer of

bacteria into the protective coating of the egg capsule (3, 14-18). Thus, colonization by

Gammaproteobacteria and specifically by Vibrionaceae is common in cephalopods, yet

colonization of the GI tract, and particularly the esophagus, was unexpected.

Bacteria from genus *Vibrio* and the related Vibrionaceae genus *Photobacterium* are

frequent colonizers of the GI tracts of marine fishes (19, 20) and are prominent in the

gastrointestinal microbiota of *Octopus vulgaris* paralarvae (21). Vibrionaceae have been reported

to produce chitinases, proteases, amylase, and lipase (20), suggesting the possibility that

colonization of the digestive tract by the Vibrionaceae serves to aid in host digestion (20). If the

Vibrio and *Photobacterium* ASVs serve this function, their localization in high density in the

esophagus, near the beginning of the digestive tract, may serve to seed the distal gut;

colonization of the lining of the esophagus may provide a reservoir that permits the microbes to

avoid washout from the gut by continually re-populating the lumen of downstream gut chambers.

An alternative explanation is that the colonization of the esophagus, and the rest of the

gut, is pathogenic or opportunistic. Various *Vibrio* species are known pathogens of cephalopods,

causing skin lesions and sometimes mortality in squids and octopuses (22-24). The genus *Vibrio*

includes representative species that are pathogenic to corals (*V. coralliilyticus*), fish (*V.*

salmonicida), diverse marine organisms (*V. harveyi*) and humans (*V. alginolyticus*, *V. cholerae*,

V. parahaemolyticus, and *V. vulnificus)* (25, 26). Likewise, the genus *Photobacterium* contains

pathogenic as well as commensal representatives (27). A previous study of the microbiota of

Octopus vulgaris paralarvae found that recently hatched paralarvae had a high-diversity

microbiome that changed, in captivity, to a lower-diversity microbiome with abundant

Vibrionaceae (21). Whether the Vibrionaceae are an integral part of cuttlefish physiology in

nature or whether they represent opportunistic colonists of these laboratory-reared organisms is a

question for future research.

Our sequence data from gills were dominated by a single ASV classified as

Piscirickettsiaceae that was in low abundance at other body sites. The Piscirickettsiaceae are a

family within the Gammaproteobacteria (28) that includes the salmon pathogen *P. salmonis*.

Rickettsia-like organisms have been described from the gills of clams and oysters (29, 30) as

well as associated with the copepod *Calanus finmarchicus* (31). In recent years

Piscirickettsiaceae have been identified in high-throughput sequencing datasets from seawater

and sediment as a taxon that may be involved in biodegradation of oil and other compounds (32-

38). Whether taxa in this family colonize the gills of cuttlefish and other organisms as symbionts

or as opportunistic pathogens is a subject for future investigation.

Studies of wild *S. officinalis* microbiota will be informative for understanding natural

host-symbiont associations under natural conditions, as compared to the mariculture-reared

animals in the present study. *S. officinalis* in the eastern Atlantic and Mediterranean are known to

prey on small mysids (crustaceans) in their first few weeks post-hatching; then as juveniles and

adults they prey mainly on marine fishes and crabs. The sparseness and simplicity of the gut

microbiota observed in our study may have been in part a result of the monodiet of grass shrimp

(*Palaemonetes*) we employed. It remains to be seen whether differences in diet and natural

variation in environmental conditions influence the association of microbial symbionts with *S.*

officinalis in the wild.

Because cuttlefish behavior is well-studied and there exist standardized methods for

documenting multiple behaviors (8), we hypothesized that these animals may provide a unique

opportunity to study microbes and the gut-brain axis – the effect of gut microbiota on behavior

(39) – in an invertebrate system. Therefore, in parallel with our study of the microbiome of

various organ systems, we conducted extensive preliminary experiments to study the effect of

antibiotic treatment on the behavior of *S. officinalis*. These results were largely negative, perhaps

due to the highly simplified microbiota we observed and its resilience to the antibiotic employed.

These results may prove helpful to the cephalopod husbandry community, as they suggest that

application of the commonly used antibiotic enrofloxacin is compatible with maintenance of

normal behavioral and microbiome characteristics of this species

4. MATERIALS AND METHODS

4. 1 Sampling and antibiotic treatment

Our study included 27 cuttlefish that were bred in the wild and moved to captivity prior

to hatching. Animals were held in water tables connected to a single open-filtration system fed

by filtered seawater. Animals were euthanized via immersion into a 10% dilution of ethanol in

seawater and were then dissected under sterile conditions within a biosafety cabinet using

autoclaved tools to obtain samples for microbial analyses. The gastrointestinal tract was

dissected into four components: esophagus, stomach, cecum, and intestine. Gill tissue and skin

from the mantle was sampled as well (~ 0.5g per sample). All tissues were stored in separate

sterile cryogenic tubes and flash-frozen in liquid nitrogen.

Following a pilot study of three individual cuttlefish, we included 24 cuttlefish (16 test, 8

control) in an experiment designed to test the effect of the antibiotic treatment on the

composition of the cuttlefish microbiome. The experimental design consisted of administering

antibiotic to animals in the treatment group (n = 16) via injection into the food source (grass

shrimp, *Palaemonetes* sp.), which was then fed to the animals. Prior to feeding, shrimp were

injected with enrofloxacin (Baytril®; 22.7 mg/mL, Bayer HealthCare LLC, Shawnee Mission,

KS, USA) using a 0.5 cc, U-100 insulin syringe with an attached 28 g x 1/2" needle (Covidien

LLC, Mansfield, MA, USA). The antibiotic dosage was 10 mg/kg rounded up to the nearest

hundredth mL. The antibiotic was injected into the coelomic cavity of the shrimp which were

then immediately fed to the cuttlefish once daily for 14 days. We maintained 8 animals as

controls, none of which received antibiotic treatment. Experimental animals were held in two

separate water tables and control animals a third, all of which were connected to the same open-

filtration system fed by filtered seawater. Within each water table, animals were isolated into

individual holding pens via plastic panels. The experimental period lasted for 14 days (1 – 15

Oct 2017), during which fecal samples were collected from each individual daily; fecal samples

were collected via pipetting free-floating material from the tank of each individual animal into a

sterile pipette.

To assess the extent to which cuttlefish microbial symbionts were shared with their

environment and food sources, 1L water samples were taken from each of three water tables in

which animals were held on the day of euthanasia and filtered using a 0.22 micron Sterivex filter

for DNA extraction; grass shrimp used as the food source throughout the duration of the

experiment were collected in 1.8ml sterile cryotubes on the same day that water was sampled,

and were frozen at -20˚C until prepared for DNA extraction.

4.2 DNA extraction, sequencing, qPCR, and16S rRNA gene statistical analyses

DNA extractions were performed on cuttlefish tissue biopsies, water, and whole shrimp

using the MoBio PowerSoil 96 Well Soil DNA Isolation Kit (Catalog No. 12955-4, MoBio,

Carlsbad, CA, USA). We used the standard 515f and 806r primers (49-51) to amplify the V4

region of the 16S rRNA gene, using mitochondrial blockers to reduce amplification of host

mitochondrial DNA. Sequencing was performed using paired-end 150 base reads on an

Illumina HiSeq sequencing platform. Following standard demultiplexing and quality filtering

using the Quantitative Insights Into Microbial Ecology pipeline (QIIME2) (52) and vsearch8.1

(53), ASVs were identified using the Deblur method (19) and taxonomy was assigned using the

Greengenes Database (May 2013 release; http://greengenes.lbl.gov). Libraries containing fewer

than 1000 reads were removed from further analyses. All 16S rRNA sequence data and metadata

will be made available via the QIITA platform prior to publication. Statistical analyses and

figure production were produced with the aid of several R packages including vegan (40), dplyr

(41), ggplot2 (42) and Adobe Illustrator® CC 2019. Code for analyses and figures is available at

www.github.com/hollylutz/CuttlefishMP. We performed qPCR analyses targeting the genus

Vibrio to quantify the abundance of bacterial cells throughout the GI tract. Conditions for qPCR

were borrowed from Thompson et al. (43), using the primers 567F and 680R and performing two

replicates of qPCR analysis that were combined to produce the final results reported in this

study.

4.3 Sample collection, fixation and sectioning for imaging

Samples from esophagus, stomach, intestine and cecum of 9 cuttlefish (1 from the pilot

and 8 from the second period experiment) were dissected and divided in order to include the

same individuals in both microscopy and sequencing analyses. Immediately after dividing,

samples for imaging were fixed with 2% paraformaldehyde in 10 mM Tris pH 7.5 for 12 h at 4

ºC, washed in PBS, dehydrated through an ethanol series from 30, 50, 70, 80 and 100%, and

stored at -20 ºC. Samples were dehydrated with acetone for 1 h, infiltrated with Technovit 8100

glycol methacrylate (EMSdiasum.com) infiltration solution 3 times 1 hour each followed by a

final infiltration overnight under vacuum, transferred to Technovit 8100 embedding solution and

solidified for 12 h at 4 ºC. Blocks were sectioned to 5 um thickness and applied to Ultrastick

slides (Thermo Scientific). Sections were stored at room temperature until FISH was performed.

4.4 Fluorescence *in situ* hybridization (FISH)

Hybridization solution [900 mM NaCl, 20 mM Tris, pH 7.5, 0.01% SDS, 20% (vol/vol)

formamide, each probe at a final concentration of 2 μM] was applied to sections and incubated at

46 ºC for 2 h in a chamber humidified with 20% (vol/vol) formamide. Slides were washed in

wash buffer (215 mM NaCl, 20 mM Tris, pH 7.5, 5mM EDTA) at 48 ºC for 15 min. Samples

were incubated with wheat germ agglutinin (20 ug ml^{-1}) conjugated with Alexa Fluor 488 and

DAPI (1 ug ml^{-1}) at room temperature for 30 minutes after FISH hybridization to label mucus

and host nuclei, respectively. Slides were dipped in ice-cold deionized water, air dried, mounted

in ProLong Gold antifade reagent (Invitrogen) with a #1.5 coverslip, and cured overnight in the

dark at room temperature before imaging. Probes used in this study are listed in Table 2.

4.5 Image acquisition and linear unmixing

Spectral images were acquired using a Carl Zeiss LSM 780 confocal microscope with a

Plan-Apochromat 40X, 1.4 N.A. objective. Images were captured using simultaneous excitation

with 405, 488, 561, and 633 nm laser lines. Linear unmixing was performed using the Zeiss ZEN

Black software (Carl Zeiss) using reference spectra acquired from cultured cells hybridized with

Eub338 probe labeled with the appropriate fluorophore and imaged as above. Unmixed images

were assembled and false-colored using FIJI software (Schindelin *et al.*, 2012).

4.6 Data availability

All 16S rRNA sequences and sample metadata will be made available via the QIITA platform

prior to publication.

FIGURE LEGENDS

Figure 1. A single *Vibrio* taxon dominates the esophagus, and a single Piscirickettsiaceae

taxon dominates the gills of the European common cuttlefish in captivity. (A) Relative

abundance of top bacterial taxa found among cuttlefish organs. Shrimp for feeding and seawater

from holding tanks are also shown. ASVs are labeled according to the finest level of taxonomic

resolution provided by the Greengenes database; bacteria not included in the top 8 taxa are

pooled as "other". Bars correspond to individual 16S rRNA sequence libraries from the pilot

investigation animals (labeled "A"), experimental animals in the control category (labeled "B"),

and experimental animals in the antibiotic treatment category (labeled "C"); only libraries with

>1000 read depth are shown. (B) Quantity of *Vibrio* cells per nanogram of DNA measured using

Vibrio-specific primers (567F and 680R (43)). Asterisk indicates significant difference between

organs ($p < 0.05$, Welch two sample t-test). (C) Anatomical depiction of the gastrointestinal tract

of *S. officinalis*, illustration modified from Plate XI of Tompsett (44).

Figure 2. Principal Coordinates Analysis (PCoA) of weighted UniFrac β-diveristy

comparing shrimp, tank water, and (A) fecal samples of treatment cuttlefish and (B) fecal

samples of control cuttlefish. Fecal samples from control animals (B) show a more tightly

clustered pattern than do fecal samples from animals treated with enrofloxacin (A).

Figure 3. Spatial organization of bacteria in the esophagus of the European common

cuttlefish, *S. officinalis*. The image shown is a cross-section of esophagus that was embedded in

methacrylate, sectioned, and subjected to fluorescence *in situ* hybridization (FISH). (A) Bacteria

(magenta) lining the interior of the esophagus in association with the mucus layer (wheat germ

agglutinin staining, green). (B) and (C) are enlarged images of the regions marked with

arrowheads in (A) where bacteria extend past the edge of the mucus layer. Blue: Host nuclei.

Scale bar =100 μm (A); 20 μm (B) and (C).

Figure 4. Fluorescence *in situ* hybridization identifies bacteria in the esophagus of *S.*
***officinalis* as Vibrionaceae**. A methacrylate-embedded section was hybridized with a nested
probe set containing probes for most bacteria, Gammaproteobacteria, Alphaproteobacteria, and
Vibrionaceae. (A) near-universal probe showing a similar bacterial distribution as in Figure 3.
(B, C, D) Enlarged images of the region marked by the dashed square in (A) showing
hybridization with near-universal, Vibrionaceae, and Gammaproteobacteria probes, respectively.
(E) Merged image of B, C, and D showing an exact match of the signal from those three probes.
(F) Alphaproteobacteria probe showing no hybridization. (G) An independent hybridization with
the non-target probe set as a control; no signal is observed. (H) enlarged image of the dashed
square in (G). Scale bar=20 μm (A, G); 5 μm (B-F, H).

Figure 5. Fluorescence *in situ* hybridization in intestine of *S. officinalis*. A methacrylate-
embedded section was hybridized with the near-universal probe and stained with fluorophore-
labeled wheat germ agglutinin to visualize mucus. (A) Bacteria (magenta) are sparsely
distributed through the lumen. (B) Enlarged image of the dashed square in (A). (C) An
independent FISH control with a non-target probe (Hhaem1007); no signal was detected. Scale
bar= 20 μm (A, C); 5 μm (B).

Figure 6. Fluorescence *in situ* hybridization in cecum of *S. officinalis*. (A) Bacteria are
observed in low abundance in the lumen of cecum. (B) Enlarged image of dashed square in (A).
Scale bar=20 μm (A); 5 μm (B).

Figure 7. Fluorescence *in situ* hybridization in gills of *S. officinalis*. Bacteria are observed in small clusters at or near the surface of the gill. (A) Overview image. (B, C): enlarged images of the dashed square in (A) showing hybridization with near-universal and Gammaproteobacteria probes, respectively. (D) Merged image of (B) and (C) showing colocalization of the signal from those two probes. Scale bar=20 μm (A); 5 μm (B-D).

ACKNOWLEDGEMENTS

We thank Alan Kuzirian, Louie Kerr, Wyatt Arnold, Madeline Kim, Elle Hill, Tyler He, Tifani Anton, and Eric Edsinger. HLL was supported by NSF 1611948. TR and JMW were supported by NSF 1650141. RTH thanks the Sholley Foundation for partial support. Amber Durand was supported by the Woods Hole Partnership Education Program. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

1.   **McFall-Ngai M.** 2014. Divining the essence of symbiosis: insights from the squid-vibrio model. PLoS Biol **12:**e1001783.
2.   **Kerwin AH, Nyholm SV.** 2017. Symbiotic bacteria associated with a bobtail squid reproductive system are detectable in the environment, and stable in the host and developing eggs. Environ Microbiol **19:**1463-1475.

3.   **Barbieri E, Bruce J. Paster, Deborah Hughes, Ludek Zurek, Duane P. Moser, Andreas**
Teske, and Mitchell L. Sogin 2001. Phylogenetic characterization of epibiotic bacteria in
the accessory nidamental gland and egg capsules of the squid Loligo pealei
(Cephalopoda: Loliginidae). Environmental Microbiology **3:**151-167.

4.   **Iehata S, Valenzuela F, Riquelme C.** 2016. Evaluation of relationship between Chilean
octopus (Octopus mimusGould, 1852) egg health condition and the egg bacterial
community. Aquaculture Research **47:**649-659.

5.   **Panetta D, Solomon M, Buresch K, Hanlon RT.** Small-scale rearing of cuttlefish (Sepia
officinalis) for research purposes. Marine and Freshwater Behaviour and Physiology
50:115-124.

6.   **Messenger JB.** 2001. Cehpalopod chromatophores: neurobiology and natural history.
Biological Review **76:**473-528.

7.   **Darmaillacq AS, Dickel L, Mather J.** 2014. Cephalopod cognition. Cambridge University
Press, Cambridge, UK.

8.   **Hanlon RT, Messenger JB.** 1988. Adaptive coloration in young cuttlefish (*Sepia officinalis*
L): the morhpology and development of body patterns and their relation to behaviour.
Phil Trans R Soc B **320:**437-487.

9.   **Buresch KC, Ulmer KM, Akkaynak D, Allen JJ, Mäthger LM, Nakamura M, Hanlon RT.**
2015. Cuttlefish adjust body pattern intensity with respect to substrate intensity to aid
camouflage, but do not camouflage in extremely low light. Journal of Experimental
Marine Biology and Ecology **462:**121-126.

10.   **Yu C, Li Y, Zhang X, Huang X, Malyarchuk V, Wang S, Shi Y, Gao L, Su Y, Zhang Y, Xu H,**
Hanlon RT, Huang Y, Rogers JA. 2014. Adaptive optoelectronic camouflage systems with
designs inspired by cephalopod skins. Proc Natl Acad Sci U S A **111:**12998-13003.

11.   **Chiao CC, Chubb C, Hanlon RT.** 2015. A review of visual perception mechanisms that
regulate rapid adaptive camouflage in cuttlefish. Journal of Comparative Physiology A:
Neuroethology, Sensory, Neural, and Behavioral Physiology **201:**933-945.

12.   **Tonkins BM, Tyers AM, Cooke GM.** 2015. Cuttlefish in captivity: An investigation into
housing and husbandry for improving welfare. Applied Animal Behaviour Science
168:77-83.

13.   **Koch EJ, Miyashiro T, McFall-Ngai MJ, Ruby EG.** 2014. Features governing symbiont
persistence in the squid-vibrio association. Molecular Ecology **23:**1624-1634.

14.   **Bloodgood RA.** 1977. The squid accessory nidamental gland: ultrastructure and
association with bacteria. . Tissue & Cell **9:**197-208.

15.   **Pichon D, Isabelle Domart-Coulon, and Renata Boucher-Rodoni** 2017. Cephalopod
bacterial associations: characterization and isolation of the symbiotic complex in the
Accessory Nidamental Glands. . Boll Malacol **43:**96-102.

16.   **Collins AJ, LaBarre BA, Won BS, Shah MV, Heng S, Choudhury MH, Haydar SA, Santiago**
J, Nyholm SV. 2012. Diversity and partitioning of bacterial populations within the
accessory nidamental gland of the squid Euprymna scolopes. Appl Environ Microbiol
78:4200-4208.

17.   **Gromek SM, Andrea M. Suria, Matthew S. Fullmer, Jillian L. Garcia, Johann Peter**
Gogarten, Spencer V. Nyholm, and Marcy J. Balunas. 2016. Leisingera sp. JC12, a

478   Bacterial Isolate from Hawaiian Bobtail Squid Eggs, Produces Indigoidine and
479   Differentially Inhibits Vibrios. Frontiers in Microbiology **7:**1342.

480   18.   **Lum-Kong A, Hastings TS.** 1992. The accessory nidamental glands of Loligo forbesi
481   (Cephalopoda: Loliginidae): characterization of symbiotic bacteria and preliminary
482   experiments to investigate factors controlling sexual maturation. . J Zool, Lond **228:**395-
483   403.

484   19.   **Tarnecki AM, Burgos FA, Ray CL, Arias CR.** 2017. Fish intestinal microbiome: diversity
485   and symbiosis unravelled by metagenomics. J Appl Microbiol doi:10.1111/jam.13415.

486   20.   **Egerton S, Sarah Culloty, Jason Whooley, Catherine Stanton, and R. Paul Ross** 2018.
487   The Gut Microbiota of marine fish. Microbiology **9:**837.

488   21.   **Roura A, Doyle SR, Nande M, Strugnell JM.** 2017. You Are What You Eat: A Genomic
489   Analysis of the Gut Microbiome of Captive and Wild Octopus vulgaris Paralarvae and
490   Their Zooplankton Prey. Front Physiol **8:**362.

491   22.   **Hanlon RT, Forsythe JW.** 1990. 1. Diseases of Mollusca: Cephalopoda. 1.1. Diseases
492   caused by microorganisms and 1.3 Structural abnormalities and neoplasia., p 23-46 and
493   203-204. *In* Kinne O (ed), Diseases of Marine Animals, vol III. Biologische Anstalt
494   Helgoland, Hamburg.

495   23.   **Forsythe JW, Hanlon RT, Lee PG.** 1990. A formulary for treating cephalopod mollusc
496   diseases. San Diego: Academic Press.

497   24.   **Ford LA, Alexander SK, Cooper KM, Hanlon RT.** 1986. Bacterial populations of normal
498   and ulcerated mantle tissue of the squid, Lolliguncula brevis. Journal of Invertebrate
499   Pathology **48:**13-26.

500   25.   **Takemura AF, Chien DM, Polz MF.** 2014. Associations and dynamics of Vibrionaceae in
501   the environment, from the genus to the population level. Front Microbiol **5:**38.

502   26.   **Pruzzo C, Huq A, Colwell RR, Donelli G.** 2005. Pathogenic Vibrio Species in the Marine
503   and Estuarine Environment, p 217-252. *In* Belkin S, Colwell RR (ed), Oceans and Health:
504   Pathogens in the Marine Environment doi:10.1007/0-387-23709-7_9. Springer US,
505   Boston, MA.

506   27.   **Labella AM, Arahal DR, Castro D, Lemos ML, Borrego JJ.** 2017. Revisiting the genus
507   Photobacterium: taxonomy, ecology and pathogenesis. Int Microbiol **20:**1-10.

508   28.   **Mauel MJ, S.J. Giovannoni, and J.L. Fryer** 1999. Phylogenetic analysis of Piscirickettsia
509   salmonis by 16S, internal transcribed spacer (ITS) and 23S ribosomal DNA sequencing. .
510   Diseases of Aquatic Organisms **35:**115-123.

511   29.   **Azevedo C, Villalba A.** Extracellular Giant Rickettsiae Associated with Bacteria in the Gill
512   of Crassostrea gigas (Mollusca, Bivalvia). Journal of Invertebrate Pathology **58:**75-81.

513   30.   **Wen C-M, Guang-Hsiung Kou, and Shiu-Nan Chen** 1994. Rickettsiaceae-like
514   Microorganisms in the Gill and Digestive Gland of the Hard Clam, Meretrix lusoria
515   Röding. Journal of Invertebrate Pathology **64:**138-142.

516   31.   **Moisander P, Andrew D. Sexton, and Meaghan C. Daley.** 2014. Stable Associations
517   Masked by Temporal Variability in the Marine Copepod Microbiome. . PLoS ONE
518   **10:**e0138967.

519   32.   **Lu X, Sun S, Hollibaugh JT, Mou X.** 2015. Identification of polyamine-responsive
520   bacterioplankton taxa in South Atlantic Bight. Environ Microbiol Rep **7:**831-838.

33. **Wang K, Zhang D, Xiong J, Chen X, Zheng J, Hu C, Yang Y, Zhu J.** 2015. Response of
bacterioplankton communities to cadmium exposure in coastal water microcosms with
high temporal variability. Appl Environ Microbiol **81:**231-240.
34. **Hamdan LJ, Salerno JL, Reed A, Joye SB, Damour M.** 2018. The impact of the Deepwater
Horizon blowout on historic shipwreck-associated sediment microbiomes in the
northern Gulf of Mexico. Sci Rep **8:**9057.
35. **Kamalanathan M, Xu C, Schwehr K, Bretherton L, Beaver M, Doyle SM, Genzer J,**
Hillhouse J, Sylvan JB, Santschi P, Quigg A. 2018. Extracellular Enzyme Activity Profile in
a Chemically Enhanced Water Accommodated Fraction of Surrogate Oil: Toward
Understanding Microbial Activities After the Deepwater Horizon Oil Spill. Front
Microbiol **9:**798.
36. **Lofthus S, Netzer R, Lewin AS, Heggeset TMB, Haugen T, Brakstad OG.** 2018.
Biodegradation of n-alkanes on oil-seawater interfaces at different temperatures and
microbial communities associated with the degradation. Biodegradation **29:**141-157.
37. **Ribicic D, McFarlin KM, Netzer R, Brakstad OG, Winkler A, Throne-Holst M, Storseth**
TR. 2018. Oil type and temperature dependent biodegradation dynamics - Combining
chemical and microbial community data through multivariate analysis. BMC Microbiol
18:83.
38. **Wang B, Liu H, Tang H, Hu X.** 2018. Microbial ecological associations in the surface
sediments of Bohai Strait. Journal of Oceanology and Limnology **36:**795-804.
39. **Carabotti M, Scirocco A, Maselli MA, Severi C.** 2015. The gut-brain axis: interactions
between enteric microbiota,
central and enteric nervous systems. Annals of Gastroenterology **28:**203-209.
40. **Oksanen J., Blanchet F. G., Friendly M., Kindt R., Legendre P., McGlinn D., Minchin P.**
R., O'Hara R. B., Simpson G. L., Solymos P., Henry M., Stevens H., Szoecs E., H. W.
2018. vegan: Community Ecology Package, vR package version 2.5-2. https://CRAN.R-
project.org/package=vegan.
41. **Wickham H, François R, Henry L, Müller K.** 2018. dplyr: A Grammar of Data
Manipulation, vR package version 0.7.6. https://CRAN.R-project.org/package=dplyr.
42. **Wickham H.** 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New
York, NY.
43. **Thompson J, Randa M, Marcelino L, Tomita-Mitchell A, Lim E, Polz M.** 2004. Diversity
and dynamics of a north Atlantic coastal *Vibrio* community. Applied Environmental
Microbiology **70:**4103-4110.
44. **Tompsett DH.** 1939. L. M. B. C. Memoirs on Typical British Marine Plants & Animals. The
University Press of Liverpool.
557
558 42. **Amann RI, Ludwig, W., Schleiffer, K.** 1995. Phylogenetic identification and in situ
559 detection of individual microbial cells without cultivation. Microbiological Reviews **59:**143-169.
560
561 43. **Neef A.** 1997. Anwendung der in situ-Einzelzell- Identifizierung von Bakterien zur
562   Populations analyse in komplexen mikrobiellen BiozönosenUniversity of Munich.
563

564   44.   **Manz W, Amann R, Ludwig W, Wagner M, Schleifer KH.** 1992. Phylogenetic
565   oligodeoxynucleotide probes for the major subclasses of Proteobacteria: problems and
566   solutions. Systematic and Applied Microbiology **15:**593-600.
567

568   45.   **Chalmers NI, Palmer RJ, Jr., Cisar JO, Kolenbrander PE.** 2008. Characterization of a
569   Streptococcus sp.-Veillonella sp. community micromanipulated from dental plaque. J
570   Bacteriol **190:**8145-8154.
571

572   46.   **Valm A. M. MW, J. L., Rieken, C. W., Hasegawa, Y., Sogin, M. L., Oldenbourg, R.,**
573   **Dewhirst, F. E., Borisy, G. G. .** 2011. Systems-level analysis of microbial community
574   organization through combinatorial labeling and spectral imaging.
575

A

Control Group Fecal vs Shrimp, Water

Legend:
- Control
- Shrimp
- Water

Y-axis: [10.2%] (0.3, 0.2, 0.1, 0.0, -0.1)
X-axis: [57.8%] (0.00, 0.25, 0.50)

B

Treatment Group Fecal vs Shrimp, Water

Legend:
- Treatment
- Shrimp
- Water

Y-axis: [9.2%] (0.1, 0.0, -0.1)
X-axis: [52%] (-0.1, 0.0, 0.1, 0.2)

Figure 3. Spatial organization of bacteria in the esophagus of *S. officinalis*. The image shown is a cross-section of esophagus that was embedded in methacrylate, sectioned, and subjected to fluorescence *in situ* hybridization. (A) Bacteria (magenta) lining the interior of the esophagus in association with the mucus layer (wheat germ agglutinin staining, green). (B) and (C) are enlarged images of the regions marked with arrowheads in (A) where bacteria extend past the edge of the mucus layer. Blue: Host nuclei. Scale bar =100 μm in (A) and 20 μm in (B) and (C).

Bacteria (Eub338-Rhodamine Red-X)　　Mucus (Wheat Germ Agglutinin-Alexa Fluor 488)　　Host nuclei (DAPI)

Figure 4. Fluorescence *in situ* hybridization identifies bacteria in the esophagus of *S. officinalis* as Vibrionaceae. A methacrylate-embedded section was hybridized with a nested probe set containing probes for most bacteria, Gammaproteobacteria, Alphaproteobacteria, and Vibrionaceae. (A) Near-universal probe showing a similar bacterial distribution as in figure 2. (B - D) Enlarged images of the region marked by the dashed square in (A), showing hybridization with near-universal, Vibrionaceae, and Gammaproteobacteria probes, respectively. (E) Merged image of (B, C, and D) showing an exact match of the signal from those three probes. (F) Alphaproteobacteria probe showing no hybridization. (G) An independent hybridization with the non-target probe set as a control; no signal is observed. (H) enlarged image of the region marked by the dashed square in (G). Scale bar=20 μm (A, G); 5 μm

Specific probe set

Figure 5. Fluorescence *in situ* hybridization in intestine of *S. officinalis.* A methacrylate-embedded section was hybridized with the near-universal probe and stained with fluorophore-labeled wheat germ agglutinin to visualize mucus (green). (A) Bacteria (magenta) are sparsely distributed through the lumen. (B) enlarged image of the region marked by the dashed square in (A). (C) An independent FISH control with a non-target probe (Hhaem1007); no signal was detected. Scale bar= 20 μm (A, C); 5 μm (B).

Bacteria probe   Non-target probe

Figure 6. Fluorescence *in situ* hybridization in cecum of *S. officinalis*. A methacrylate-embedded section was hybridized with the near-universal probe and stained with fluorophore-labeled wheat germ agglutinin to visualize mucus (green). (A) Bacteria are observed in low abundance in the lumen of cecum. (B) Enlarged image of region marked by the dashed square in (A). Scale bar=20 μm (A); 5 μm (B).

🟪	Bacteria (Eub338-Rhodamine Red-X)	🟦	Host nuclei (DAPI)
🟩	Mucus (Wheat Germ Agglutinin Alexa Fluor-488)	⬜	Autofluorescence

Figure 7. Fluorescence *in situ* hybridization in gills of *S. officinalis*. A methacrylate-embedded section was hybridized with the near-universal probe and stained with fluorophore-labeled wheat germ agglutinin to visualize mucus (green). Bacteria are observed in small clusters. (A) Overview image. (B, C): enlarged images of the region marked by the dashed square in (A) showing hybridization with near-universal, and Gammaproteobacteria probes, respectively. (D) Merged image of (B), and (C) showing an exact match of the signal from those two probes. Display in (B, C, and D) was adjusted for clarity. Scale bar=20 μm (A); 5 μm (B-D).

Table 1. Relative abundance of three most abundant ASVs found among two sampling periods of the European common cuttlefish. Averages correspond to relative abundance of individual ASVs from each anatomical site and period. Period 1 consisted of three individuals, Period 2 consisted of eight individuals.

Anatomical Site	ASV1 Vibrionaceae, *Vibrio* sp.			ASV2 Vibrionaceae, *Photobacterium* sp.			ASV3 Piscirickettsiaceae, *unknown*		
	Pilot	Control	Antibiotic	Pilot	Control	Antibiotic	Pilot	Control	Treatment
Esophagus	0.92 ± 0.10	1.00 ± 0.01	0.94 ± 0.10	0.07 ± 0.10	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
Stomach	0.39 ± 0.34	0.43 ± 0.37	0.19 ± 0.31	0.25 + 0.42	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
Cecum	0.24 ± 0.14	0.44 ± 0.23	0.06 ± 0.09	0.28+0.35	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
Intestine	0.22 ± 0.42	0.57 ± 0.16	0.06 ± 0.06	0.50 ± 0.57	0.01 ± 0.01	0 ± 0	0 ± 0	0 ± 0	0 ± 0
Gills	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.01 ± 0.01	0 ± 0	0.79 ± 0.11	0.97 ± 0.03	0.82 ± 0.19
Skin	0.02 ± 0.03	0.08 ± 0.11	0.01 ± 0.02	0.03 ± 0.02	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.01 ± 0.02

Table 2. FISH probes used in this study

	Probe name	Fluorophore(s)	Target organism	Sequence 5' – 3'	Target position	Reference
Experimental probe set	Eub338-I	Alexa 555 or Rhodamine Red-X	Most Bacteria	GCTGCCTCCCGTAGGAGT	16S, 338-355	Amann et al. 1990
	Vib1749	Texas Red-X	Vibrionaceae family	AGCCACCTGGTATCTGCGACT	23S, 1749-1769	Schlundt *et al*., in prep
	Vib2300	Texas Red-X	Vibrionaceae family	TAACCTCACGATGTCCAACCGTG	23S, 2299-2321	Schlundt *et al*., in prep
	Alf968	Atto 620 or Dy490	Alphaproteobacteria	GGTAAGGTTCTGCGCGTT	16S, 968-985	Neef, A., 1997
	Gam42a	Atto 647N or Cy5	Gammaproteobacteria	GCCTTCCCACATCGTTT	23S, 1027-1043	Manz et al. 1992
Non-target control probe set	Vei488	Alexa 555	*Veillonella*	CCGTGGCTTTCTATTCCG	16S, 488-505	Chalmers et al. 2008
	PorTan34	Texas Red-X	*Porphyromonas* and *Tannerella*	GTTAAGCCTATCGCTAGC	16S, 34-51	Mark Welch *et al*., in prep.
	Fus714	Atto 620	*Fusobacterium*	GGCTTCCCCATCGGCATT	16S, 714-731	Valm et al. 2011
	Lept568	Atto 647N	*Leptotrichia*	GCCTAGATGCCCTTTATG	16S, 568-585	Valm et al. 2011
Non-target probe	Hhaem1007	Rhodamine Red-X	*Haemophilus haemolyticus*	AGGCACTCCCATATCTCTACAG	16S, 1007-1028	Mark Welch *et al*., in prep.