The detection of novel and resident marine non-indigenous species using environmental

DNA metabarcoding of seawater and sediment

Mr. Luke E. Holman[1]* (ORCID: 0000-0002-8139-3760)

Dr. Mark de Bruyn[2,3]

Prof. Simon Creer[2]

Prof. Gary Carvalho[2]

Dr. Julie Robidart[4]

Dr. Marc Rius[1,5] (ORCID: 0000-0002-2195-6605)

[1] Ocean and Earth Science, National Oceanography Centre Southampton, University of

Southampton, United Kingdom

[2] Molecular Ecology and Fisheries Genetics Laboratory, School of Natural Sciences, Bangor

University, United Kingdom

[3] The University of Sydney, School of Life and Environmental Sciences, Australia

[4] Ocean Technology and Engineering Group, National Oceanography Centre Southampton,

United Kingdom

[5] Centre for Ecological Genomics and Wildlife Conservation, University of Johannesburg, South

Africa

*Corresponding Author

Email: l.e.holman@soton.ac.uk

Abstract

1. Biodiversity monitoring in aquatic ecosystems is challenging as it requires taxonomic

expertise and is difficult to automate. One expedient approach is the use of environmental DNA

(eDNA) surveys to infer species incidence from nucleic acids found in environmental samples.

An advantage of eDNA surveys is that they allow rapid detection of non-indigenous species

(NIS), lessening the time from introduction to discovery and increasing the likelihood of

successful control and / or eradication. Despite this, relatively few studies have compared eDNA

and traditional surveys for the study of NIS, or examined the differences between

metabarcoding of different environmental sample types.

2. We evaluated the ability of eDNA to detect a broad range of native and NIS in urban coastal

environments, and compared the results with previously published traditional surveys. We

collected water and sediment samples from the same sites and then performed eDNA

metabarcoding of 18S rRNA and COI genes.

3. We found very different patterns of biodiversity in water and sediment samples; with sediment

containing more than two times the number of operational taxonomic units (OTUs) than

seawater in some cases. The findings showed the presence of as much variation in assemblage

diversity among environmental sample types as amongst geographically-segregated sampling

sites. Additionally, species detection within phyla was not consistent in water or sediment

samples, indicating that at a broad scale sample type does not perfectly predict taxa detected.

4. We found almost perfect agreement in species detection from eDNA and traditional surveys.

Additionally, eDNA metabarcoding detected three previously undocumented species

introductions. Finally, our work provided a novel high-resolution biodiversity dataset for urban

marine environments.

5. *Synthesis and applications* Our study showed that the type of eDNA sample dramatically

affects the detected biodiversity and that eDNA metabarcoding is accurate for the detection of

notorious NIS. Natural resource managers, ecological practitioners and researchers should

consider the benefits of integrating molecular tools such as eDNA into applied ecology.

Keywords

18S, COI; eDNA; sediment; metabarcoding; non-indigenous; non-native.

Introduction

Anthropogenic activities are causing a global decrease in biodiversity (Sala and Knowlton 2006; Butchart et al. 2010) that negatively affects ecosystem services and function (Worm et al. 2006). Such impacts create an urgent need for tools that rapidly and accurately monitor species diversity. Traditional biodiversity surveys have been criticised for their lack of standardisation and taxonomic resolution (Oliver & Beattie 1993; Fitzpatrick *et al.* 2009). One approach that has the potential to overcome limitations of traditional surveys is the use of nucleic acids found in environmental samples, such as water or sediment, to infer presence or absence of living organisms in the local environment. Such genetic material, known as environmental DNA (hereafter eDNA), is a poly-disperse mixture of tissue, cells, subcellular fragments and extracellular DNA lost to the environment by organisms (Turner *et al.* 2014; Sassoubre *et al.* 2016). Studies using eDNA focus on detection from water samples using targeted (single species) methods such as qPCR (Dougherty *et al.* 2016; Simpson *et al.* 2017; Wood *et al.* 2017; Kim *et al.* 2018), or community (multi-species) methods such as metabarcoding (Borrell *et al.* 2017; Grey *et al.* 2018; Lacoursière-Roussel *et al.* 2018). Samples can be collected with minimal training and once the methodology is optimised, surveys are highly amenable to automation (McQuillan & Robidart 2017). Thus, eDNA surveys are highly informative (but see specific considerations to ensure validity, Goldberg *et al.* 2016) and can complement traditional methods (Deiner *et al.* 2017). Recent work has identified a vast range of viable protocols for the collection, extraction and detection of target nucleic acids from different environmental samples (Deiner et al., 2018; Spens *et al.* 2016; Sellers *et al.* 2018). Despite the finding that sediment has been shown to harbour 8-1800 times more eDNA compared to water samples in freshwater ecosystems (Turner, Uy & Everhart 2015) relatively few eDNA studies incorporate multiple environmental sample types (Shaw *et al.* 2016). Furthermore, no work to date has compared

species richness estimates derived from metazoan target eDNA metabarcoding between

seawater and marine sediment samples.

Non-indigenous species (NIS) are those that have been transported from their native range

through human action into a novel geographic location. The impacts of NIS are well documented

and they can pose a severe threat to natural ecological systems, agriculture, biodiversity and

human health (Bax *et al.* 2003; Lovell, Stone & Fernandez 2006; Ricciardi *et al.* 2013; Mazza *et*

al. 2014). Most marine NIS have spread globally via vectors such as transoceanic shipping,

aquaculture, the construction of canals interconnecting large water bodies and capture aquarium

industry (Molnar *et al.* 2008; Nunes *et al.* 2014). At fine (10s of km) geographical scales, other

vectors such as recreational boating significantly enhance the spread and subsequent impact of

NIS (Clarke Murray *et al*. 2011). Along coastal areas, NIS studies have highlighted the

importance of monitoring marinas and harbours (Ashton *et al.* 2006), as these are hotspots of

NIS. In these habitats, sessile marine NIS often outcompete native species and dominate

artificial hard substrata (Glasby *et al.* 2007; Dafforn *et al*. 2009). Marinas and harbours have

distinct ecological and physico-chemical conditions compared to the surrounding natural

environment (Rivero *et al.* 2013; Foster *et al.* 2016). Consequently, there is a need for specific

sampling and surveying protocols to study both native and NIS in these ecologically distinct

environments.

The detection of NIS has been performed traditionally by surveys standardised by time or by

reaching a species discovery asymptote (Ashton *et al.* 2006; Campbell, Gould & Hewitt 2007;

Bishop *et al.* 2015). Moreover, traditional NIS surveys require expert taxonomic skills that are

inherently time consuming and are not typically amenable to automation (Darling *et al.* 2017). In

addition, the results of such surveys reflect only species that are being targeted at the time, with

no ability to retrospectively separate erroneously grouped species in light of new discoveries.

This is particularly important considering that between 9,000-35,000 marine species (2.7% of

the total number of estimated marine species) are cryptic (i.e., morphologically similar but

genetically distinct) (Appeltans *et al.* 2012). Indeed, many common global sessile marine

invaders are morphologically indistinguishable and contain cryptic lineages as revealed by

genetic studies, highlighting the need for the use of genetic tools to accurately assess species

invasions (Pérez-Portela *et al.* 2013; Rius *et al.* 2017). The use of eDNA as monitoring tool of

NIS has great potential, mainly because it tackles limitations of the existing biodiversity

monitoring tools.

Here we tested the efficacy of eDNA metabarcoding for exploring the biodiversity of several

distinct marinas. We first documented the differences in both alpha and beta diversity of different

eDNA sample types (seawater and sediment). We then compared the eDNA metabarcoding

results with traditional methods and identified a number of key NIS that are both introduced in

the study region and / or elsewhere. Subsequently, we identified a number of previously

unrecorded NIS in the study region. Finally we discuss the strengths and weaknesses of eDNA

metabarcoding for detecting marine NIS.

Methods

Study sites

Four marinas were selected from around the United Kingdom to represent variation in modelled

invasion potential (Pearce, Peeler & Stebbing 2012), known NIS present (Bishop *et al.* 2015)

and surrounding benthic habitat type (Calewaert *et al.* 2016). Importantly, all chosen marinas

have been surveyed previously for NIS and so there is a good understanding of the expected

NIS in these areas (Wood, Bishop & Yunnie 2015a; b; Wood *et al.* 2016). Marina access was

contingent on anonymity and so marina names and exact locations are not presented, Fig.1a

shows approximate locations. Marina *TQ* is an open marina subject to tides and varying salinity

located in Southampton Water on the north coast of the English Channel. Marina *PQ* is a loch

marina open during high tide to the Bristol Channel and the Celtic Sea. Marina *TB* is located at

the mouth of the River Blackwater open to the North Sea. Marina *HH* is located on the Isle of

Anglesey and is open to the Celtic Sea.

Environmental DNA sampling

Surveys were conducted during May 2017. A total of 24 sampling points were randomly selected

within each site. At each point 50ml of water was collected from 10cm below the surface using a

sterile 60ml luer lock syringe and filtered through a 0.22mm polyethersulfone Sterivex filter

(Merck Millipore, Massachusetts USA). After collecting seawater from eight locations (400ml

total volume) the filter was changed, resulting in a total of three filters per site. To test the effect

of different sample preservation methods sampling was performed in duplicate. One set of three

filters had ~1.5ml sterile Longmire's solution (100mM Tris,10mM EDTA, 10mM NaCl, 0.5%

SDS) applied in the inlet valve (Renshaw et al. 2015). The second set of three filters were kept

on ice for no longer than eight hours before being frozen at -20°C. During the surveys, a

sediment sample was collected at the start and then after every 3rd water sample, for a total of 9

per site. We used a UWITEC Corer (UWITEC, Mondsee, Austria) to collect a sediment core

(600mm tall x 60mm diameter). Using a sterile disposable spatula, a subsample of 10-20g of

sediment was taken from the top 2cm of the core, taking care to avoid sampling the sides of the

core. The subsample was stored in a sterile plastic zip container and kept on ice for no longer

than eight hours before begin frozen at -80°C. Due to equipment malfunction no sediment

sample could be taken for Site HH. Disposable gloves were changed after collection of each

sample. All reused equipment was washed thoroughly and soaked in 10% bleach between sites,

before rinsing in DNAse-free sterile water.

eDNA extraction

DNA extractions were performed in a PCR–free cleanroom, separate from main laboratory

facilities. No high copy templates, cultures or amplicons were permitted in this sterile laboratory.

DNA extractions from water samples followed Spens et al. (2016) using the SX$_{CAPSULE}$ method.

Briefly, preservative solution was removed from the outlet and filters were dried at room

temperature for two hours, 720µl Qiagen buffer ATL (Qiagen, Hilden, Germany) and 80µl

Proteinase K was added to the filter and all samples were digested overnight at 56ºC. After

digestion, samples were processed using the Qiagen DNeasy Blood and Tissue Kit as per

manufacturer instructions. The final elution was 200µl PCR grade water.

Sediment extractions were performed using the Qiagen DNeasy Powermax Soil Kit using the

manufacturer recommended protocol. For each site the nine samples were mixed to form three

pooled samples; for each extraction, 10g of pooled sample was processed. A total of ten

samples were processed, three each per site with a single no blank, as per manufacturer's

instructions.

Inhibition testing

To ensure extracted DNA was free of PCR inhibitors a Primer Design Real-Time PCR Internal

Control Kit (PrimerDesign, Southampton, UK) was used. qPCR reactions were performed

following the manufacturer's protocol. Inhibition due to co-purified compounds from DNA

extraction protocols would produce an increase in cycle threshold number in comparison to no

template controls. All samples were successfully processed and no samples showed indication

of PCR inhibition.

Primer selection & library preparation

Two sets of primers were chosen for metabarcoding the environmental samples: a 313bp

section of the standard DNA barcoding region of the cytochrome c oxidase subunit I gene (COI)

using primers described in Leray et al. (2013); and a variable length target of the hypervariable

v4 region of the nuclear small subunit ribosomal DNA (18S) using primers from Zhan *et al.*

(2013). Sequencing libraries were prepared using a 2-step PCR approach as detailed in Bista *et*

al. (2017), this method amplifies the target region in PCR 1 annealing universal adapters onto

which sample specific indices and sequencing primers are annealed in PCR 2. In contrast to

Bista *et al*. (2017) we used unique dual-matched indexes for PCR 2 to avoid index crosstalk

associated with combinatorial indexing (MacConaill et al. 2018). PCR 1 was prepared in a PCR–

free room, separate from main laboratory facilities. PCR conditions and reaction volumes are

detailed in Supplementary Information 1. Blank filters, DNA extraction kits and positive controls

where collected, extracted and sequenced as the experimental treatments (detailed in

Supplementary Information 1). Samples were pooled at an equimolar ratio and sequenced using

the Illumina MiSeq instrument (Illumina, San Diego, USA) with a V3 2 x 300bp kit.

Bioinformatic analyses

Samples were demultiplexed using the Illumina MiSeq control software (v 2.6.2.1). The

demultiplexed data was analysed using a custom pipeline written in the R programming

language (R Core Team 2018) hosted at https://github.com/leholman/metabarTOAD, the steps

are as follows. Forward and reverse paired end reads were merged using the -fastq_mergepairs

option of USEARCH v.10.0.240 (Edgar 2013) with maximum difference of 15, percent identity of

80% and quality filter set at maximum expected errors of 1. Both the forward and reverse primer

sequences were matched using Cutadapt v.1.16 (Martin 2011) and only sequences containing

both primer regions were retained. Sequences were discarded if they were outside of a defined

length boundary (303-323bp for COI, 375-450bp for 18S) using Cutadapt. Sequences were then

pooled, singletons were discarded and sequences were quality filtered with a maximum

expected error of 1 using the -fastq_filter option of vsearch v.2.4.3 (Rognes *et al.* 2016).

Sequences were then denoised and chimeras filtered using the unoise3 algorithm implemented

in USEARCH. The resultant OTUs were curated using the LULU package v.0.1.0 in R (Frøslev

et al. 2017). An OTU by sample table was produced by mapping the merged and trimmed reads

against the curated OTUs, reporting a single best hit within 97% of the query sequence. The

OTU x sample table was filtered in R as follows. To minimise the chance of spurious OTUs

being included in the final dataset any record with less than 3 raw reads were changed to zero

and any OTU that did not appear in more than one sample was removed from the analysis.

OTUs found in negative controls were removed from the analysis.

Taxonomic assignment

Assigning correct taxonomy to an unknown set of marine sequences can be challenging as

large databases require vast computational resources for query matching; many databases

contain errors and the taxonomy of some marine groups is uncertain. With such limitations in

mind, we assigned taxonomy using a BLAST v.2.6.0+ search (Camacho *et al.* 2009) returning

the single best hit from databases within 97% of the query. The MIDORI database

(UNIQUE_20180221) (Machida *et al.* 2017) was used for the COI data and the SILVA database

(SSU r132) (Quast *et al.* 2013) was used for the 18S rRNA data. The match taxa tool from the

World Register of Marine Species (WoRMS) (WoRMS Editorial Board 2018) was used to filter

the data for marine species and check the classification. Remaining annotations were checked

against the World Register of Introduced Marine Species (WRIMS) (Ahyong *et al.* 2018) to

determine non-indigenous status.

Statistical analyses

All statistical analyses were conducted in R v3.5.0. The Vegan R package v.2.5.2 (Oksanen *et*

al. 2011) was used to rarefy samples to the minimum sample read depth for each amplicon. The

number of OTUs per site/condition was calculated as the number of OTUs with a non-zero

number of normalized reads after summing the reads across all three site level replicates. To

test if there was a significant difference between the number of OTUs generated by sediment

and water eDNA, individual non-summed replicate sample data was used to build a linear

regression model following the formula *number_of_OTUs~sedimentorwater*site* implemented in

R using the function lm(). Non-metric multidimensional scaling ordination plots were generated

from Bray-Curtis dissimilarity values derived using Vegan. A Permutation Analysis of Variance

(PERMANOVA) (Balakrishnan et al. 2014) was performed using the Bray Curtis dissimilarity

following the model *dissimlairty_matrix~sedimentorwater*site* implemented in R using the

function *adonis* from the *vegan* package. OTUs with taxonomic assignment were separated into

those found in sediment, water or both media and the OTUs were then collapsed at the Phylum

level to explore taxonomic patterns of detection in water or sediment. Phyla with less than eight

OTUs were combined. To test for non-random counts of species detection between water and

sediment within taxa an exact binomial test was performed between counts of species detected

in water and sediment. Half the number of counts for species detected in both water and

sediment were added to water and sediment, with non-integer values conservatively rounded up

to the nearest whole number. A Bonferroni correction for multiple comparisons was applied

across the p values from the exact binomial tests. Records from manual surveys previously

conducted for non-native invertebrates at the sample sites (Wood, Bishop & Yunnie 2015a; b;

Wood *et al.* 2016) were compared with the detected species from metabarcoding data.

Results

Raw sequencing results & OTU generation

Sequencing produced a total of 17.8 million paired end reads, with 15.2 million sequences

remaining after paired end read merging and quality filtering. The average number of sequences

per sample after filtering (excluding control samples) was 200,185 ± 64,019 (s.d). No template

control samples contained an average of 811 ± 3,402 (s.d) sequences. One control sample

contained ~15,000 sequences that mapped to an operational taxonomic unit (OTU) that had

100% identity match to a sequence of a terrestrial fungi (Genbank: FJ804151.1), excluding this

OTU gives an average of 51 ± 94 (s.d) sequences per no-template control sample. Denoising

produced 8,069 OTUs for COI and 2,433 for 18S with 6,435 and 1,679 remaining respectively

after LULU curation. Taxonomic annotation identified 395 OTUs from the 18S rRNA dataset

against the SILVA database and 219 OTUs from the COI dataset against the MIDORI database.

Taxonomic data from WoRMS could be retrieved for 204 of the annotated COI OTUs and 138 of

the 18S OTUs.

Biodiversity detection

The effect of different water eDNA sample preservation techniques differed between the target

amplicons. The 18S rRNA amplicon produced significantly more OTUs in samples preserved by

freezing compared to Longmire's preservation method, while the COI amplicon showed no

significant difference between preservation treatments (see Supplementary Information 2 for

details). As a conservative approach all subsequent analyses used sample data from frozen

samples. The minimum number of reads per sample was 137,624 for the COI dataset and

117,915 for the 18S dataset and so samples were rarefied to this number of reads. More OTUs

in total were detected in the sediment samples compared to the water samples across all sites

and both markers as shown in Figure 1b,d. In all cases, both water and sediment samples

detected unique OTUs but the mean proportion of unique OTUs detected in water was lower

(49.2%) in comparison to sediment (73.8%). A 2-way ANOVA testing the effect of eDNA type of

environmental sample on number of OTUs generated indicated a significant effect (p<0.001) of

sample type for both 18S rRNA and COI (See Supplementary Information 3 for full model

output). Ordination plots of Bray-Curtis dissimilarity (Fig. 1c,e) showed that OTUs in eDNA found

in sediment and water differ in community structure as much as among sites in ordination space.

Additionally, the PERMANOVA model indicated highly significant differences (p<0.001) between

sites and eDNA medium in both the 18S rRNA and COI datasets. Furthermore, eDNA detection

medium in the PERMANOVA model explained 23.2% and 32.5% of the variation in the 18S and

COI data respectively, while the site explained 34.2% and 30.5% in the COI and 18S rRNA data

(See Supplementary Information 4 for full model output). At phylum level (Figure 2), taxonomy

does not perfectly predict medium of detection, however a binomial goodness of fit test shows

non-random detection proportions in the Nematoda (Bonferroni corrected p=0.005), with eDNA

detections mostly in sediment.

Detection of non-indigenous species

In total 16 NIS to the study region and 30 species documented as NIS in other countries were

detected across the four sites (see Supplementary Table 1 for full list). Out of the detected NIS

seven were present in the list of 21 NIS previously detected in manual non-native invertebrate

surveys at the sites. As shown in Fig. 3 the results of the eDNA surveys closely matched those

of the manual survey results. Only a single detection differed from the manual surveys, an eDNA

false-negative detection of *Bugula neritina* at Site HH. Remapping of the cleaned reads to the

Bugula neritina COI region (Genbank Accession: KY235450.1) indicated that 5 reads from a

single replicate corresponded with *Bugula neritina.* These reads were lost during data filtering

and so did not feature in the final dataset. A detection of note was 199 reads from the sediment

of Site TQ mapping to an OTU corresponding with *Arcuatula senhousia* (Asian Date Mussel), a

previously undocumented NIS for the UK. Targeted visual surveys on tidal mudflats within two

kilometres of Marina TQ confirmed the presence of this species in proximity to the sampling site.

Furthermore, COI sequences generated from these physical samples (Genbank Accession:

MH924820 and MH924821) matched to known *A. senhousia* sequences confirming the eDNA

detection (see Supplementary Information 5 for details of DNA barcoding). Additionally, the

nematode *Cephalothrix simula* and the oligochaete *Paranais frici* were also detected using

eDNA at site TQ. Both are new species introductions to the United Kingdom, previously

undocumented in academic literature.

Discussion

We demonstrated that very different community composition can be detected in seawater and

sediment samples, affirming the importance of integrating multiple sample types for obtaining a

full assessment of community composition. We also found that eDNA metabarcoding shows

excellent concordance with traditional methods for the detection of NIS. Finally, we

demonstrated that eDNA metabarcoding can detect novel species introductions. This study

emphasises the need to have a thorough understanding of the effects of environmental sample

types both at the level of whole community and for specific species of concern. eDNA

metabarcoding continues to be used in both regional and global biodiversity surveys.

The majority of research using eDNA to detect aquatic macrofauna is based on the collection of

water samples, while sediment samples have received comparatively less attention. We found

dramatic differences in species richness in sediment and water samples, observing a

consistently greater number of OTUs detected in sediment compared to water. Shaw *et al.*

(2016) found that sediment 12S rRNA metabarcoding detected fewer fish compared to water in

a freshwater lotic environment. Our results indicated a similar trend when considering only fish

species; with more fish being detected in seawater samples compared to sediment samples (5

in water, 2 in sediment and water), but the opposite when considering all OTUs. More broadly,

taxonomy at the level of phylum did not predict if a species will be detected in water, sediment or

both sample types (except the Nematoda, whose members are predominantly benthic

inhabitants). It is likely, as seen in the case of fish above, that at lower taxonomic levels the

species-specific ecology of eDNA (sensu Barnes & Turner 2015) will result in convergent eDNA

occupancy in different sample types. However, further work is needed to clarify how eDNA

partitions into adjacent environmental samples across the tree of life. Our study showed that at

the level of phyla detection was not significantly different between sediment and water for most

taxa. Similarly, we showed that for most NIS both water or sediment samples served as an

excellent media for detection.

Current eDNA metabarcoding research has identified large variation in the detected diversity

across small spatial scales in both sediment (Nascimento *et al.* 2018) and seawater (O'Donnell

et al. 2017). Additionally fractionation of environmental samples (i.e. sorting samples by particle

size class) can produce significant differences in the metabarcoding results between fractions

(Wangensteen *et al.* 2018a; b) indicating that reliable variation can be discovered at a scale of

site. Here we found similar patterns, with site and eDNA sample type containing approximately

equivalent OTU biodiversity. Future research should explore how eDNA extracted using different

sample types and extraction methods affects the detection of biodiversity, especially as eDNA

metabarcoding moves from an experimental technique to a routine monitoring tool (Pawlowski *et*

al. 2018; Aylagas *et al.* 2018).

We found that eDNA metabarcoding of water samples accurately detects many NIS species, as

seen in previous work (Borrell *et al.* 2017; Grey *et al.* 2018; Lacoursière-Roussel *et al.* 2018).

Additionally, we identified the presence of NIS in sediment samples for the first time. In

comparing these data to those collected using traditional methods we found almost perfect

agreement in NIS presence. The single false-negative was found to be a result of bioinformatic

384 parameters, identifying that choices made during sequence processing can have an effect on

385 the detectability of species in eDNA samples. Indeed, this has previously been shown in

386 metabarcoding of bulk samples (Scott *et al.* 2018) and work is urgently needed to determine

387 optimal bioinformatic parameters, the effects of primer binding sites and the role of DNA

388 barcodes in reference databases for the detection of NIS in eDNA samples. It is therefore

389 important to combine eDNA metabarcoding with traditional surveys where possible, as both

390 methods provide reciprocal validation data (e.g. important NIS may be missed using molecular

391 techniques or eDNA metabarcoding detects rare species that often missed by traditional

392 surveys). We identified several NIS currently unrecorded in the UK and confirmed the detection

393 with targeted local surveys to confirm the presence of *A. senhousia*. The case of *A. senhousia* is

394 particularly relevant when evaluating eDNA metabarcoding for NIS detection as the species is

395 spreading globally (Bachelet *et al.* 2009), has the potential to dramatically alter benthic

396 biodiversity (Crooks 2001; Mistri 2003) and may have long remained undetected. Future

397 research should evaluate the sensitivity of eDNA metabarcoding for the detection of novel

398 species introductions, which may allow an unprecedented level of accuracy during early stages

399 of invasion. The incorporation of autonomous sampling (McQuillan & Robidart 2017) and eDNA

400 biobanking (Jarman, Berry & Bunce 2018) could also help the rapid detection of NIS by

401 improving DNA reference databases for specific geographical regions that have high biosecurity

402 risk, providing an invaluable resource for biodiversity managers and researchers alike.

Author Contributions

L.E.H. and M.R. designed the experiment, L.E.H. collected samples, generated and analysed

the data, designed all figures and wrote the first draft of the paper. All authors contributed

critically to the drafts and gave final approval for publication.

Acknowledgements

We are grateful to the Bishop Lab (Marine Biological Association) and affiliates for excellent NIS

survey data and for sharing information on marinas. We acknowledge staff in the Environmental

Sequencing Facility at the National Oceanography Centre Southampton for advice and

assistance during library preparation. We acknowledge the Department of Geography &

Environment from the University of Southampton for access to coring equipment and lab space.

LH was supported by the Natural Environmental Research Council (grant number

NE/L002531/1).

Data accessibility

Raw Illumina sequencing data is available from the European Nucleotide Archive under

accession number (pending).

Associated metadata, script and intermediate files can be found on GitHub with the following

DOI: 10.5281/zenodo.1453959.

References

Ahyong, S., Costello, M.J., Galil, B.S., Gollasch, S., Hutchings, P., Katsanevakis, … Zhan, A.

(2018) http://www.marinespecies.org/introduced

Anderson, M.J. (2014) Permutational Multivariate Analysis of Variance (PERMANOVA).

Wiley Stats Ref: Statistics Reference Online (eds N. Balakrishnan, T. Colton, B. Everitt,

W. Piegorsch, F. Ruggeri, & J.L. Teugels), pp. 1–15. John Wiley & Sons, Ltd,

Chichester, UK.

Appeltans, W., Ahyong, S.T., Anderson, G., Angel, M.V., Artois, T., Bailly, N., … Costello,

M.J. (2012) The magnitude of global marine species diversity. *Current Biology*, **22**,

2189–2202.

Ashton, G., Boos, K., Shucksmith, R. & Cook, E. (2006) Rapid assessment of the distribution

of marine non-native species in marinas in Scotland. *Aquatic Invasions*, **1**, 209–213.

Aylagas, E., Borja, Á., Muxika, I. & Rodríguez-Ezpeleta, N. (2018) Adapting metabarcoding-

based benthic biomonitoring into routine marine ecological status assessment

networks. *Ecological indicators*, **95**, 194–202.

Bachelet, G., Blanchet, H., Cottet, M., Dang, C., de Montaudouin, X., de Moura Queirós, A.,

Gouillieux, B. & Lavesque, N. (2009) A round-the-world tour almost completed: first

records of the invasive mussel *Musculista senhousia* in the north-east Atlantic

(southern Bay of Biscay). *Marine Biodiversity Records*, **2**, 2002–2005.

Barnes, M.A. & Turner, C.R. (2015) The ecology of environmental DNA and implications for

conservation genetics. *Conservation Genetics* , **17**, 1–17.

Bax, N., Williamson, A., Aguero, M., Gonzalez, E. & Geeves, W. (2003) Marine invasive alien

species: a threat to global biodiversity. *Marine Policy*, **27**, 313–323.

Bishop, J., Wood, C., Yunnie, A. & Griffiths, C. (2015) Unheralded arrivals: non-native sessile

invertebrates in marinas on the English coast. *Aquatic Invasions*, **10**, 249–264.

Bista, I., Carvalho, G.R., Walsh, K., Seymour, M., Hajibabaei, M., Lallias, D., Christmas, M. &

Creer, S. (2017) Annual time-series analysis of aqueous eDNA reveals ecologically

relevant dynamics of lake ecosystem biodiversity. *Nature Communications*, **8**, 14087.

Borrell, Y.J., Miralles, L., Do Huu, H., Mohammed-Geba, K. & Garcia-Vazquez, E. (2017)

DNA in a bottle—Rapid metabarcoding survey for early alerts of invasive species in

ports. *PloS One*, **12**, e0183347.

Butchart, S.H.M., Walpole, M., Collen, B., van Strien, A., Scharlemann, J.P.W., Almond,

R.E.A., … Watson, R. (2010) Global biodiversity: indicators of recent declines. *Science*,

328, 1164–1168.

Calewaert, J.-B., Weaver, P., Gunn, V., Gorringe, P. & Novellino, A. (2016) The European

Marine Data and Observation Network (EMODnet): Your Gateway to European Marine

and Coastal Data. *Quantitative Monitoring of the Underwater Environment: Results of*

the International Marine Science and Technology Event MOQESM´14 in Brest, France

(eds B. Zerr, L. Jaulin, V. Creuze, N. Debese, I. Quidu, B. Clement, & A. Billon-Coat),

pp. 31–46. Springer International Publishing, Cham.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K. & Madden,

T.L. (2009) BLAST+: architecture and applications. *BMC Bioinformatics*, **10**, 421.

Campbell, M.L., Gould, B. & Hewitt, C.L. (2007) Survey evaluations to assess marine

bioinvasions. *Marine Pollution Bulletin*, **55**, 360–378.

Clarke Murray, C., Pakhomov, E.A. & Therriault, T.W. (2011) Recreational boating: a large

unregulated vector transporting marine invasive species: Transport of NIS by

recreational boats. *Diversity and Distributions*, **17**, 1161–1172.

Crooks, J.A. (2001) Assessing Invader Roles Within Changing Ecosystems: Historical and

Experimental Perspectives on an Exotic Mussel in an Urbanized Lagoon. *Biological*

Invasions, **3**, 23–36.

Dafforn, K.A., Johnston, E.L. & Glasby, T.M. (2009) Shallow moving structures promote

marine invader dominance. *Biofouling*, **25**, 277–287.

Darling, J.A., Galil, B.S., Carvalho, G.R., Rius, M., Viard, F. & Piraino, S. (2017)

Recommendations for developing and applying genetic tools to assess and manage

biological invasions in marine ecosystems. *Marine Policy*, **85**, 56–64.

Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., …

Bernatchez, L. (2017) Environmental DNA metabarcoding: Transforming how we

survey animal and plant communities. *Molecular Ecology*, **26**, 5872–5895.

Deiner, K., Lopez, J., Bourne, S., Holman, L.E.**,** Seymour, M., Gray, E.K., … Lodge, D.M. *In*

press. Optimizing the detection of marine taxonomic richness using environmental DNA

metabarcoding: the effects of filter material, pore size, and extraction method.

Metabarcoding & Metagenomics

Dougherty, M.M., Larson, E.R., Renshaw, M.A., Gantz, C.A., Egan, S.P., Erickson, D.M. &

Lodge, D.M. (2016) Environmental DNA (eDNA) detects the invasive rusty crayfish

Orconectes rusticus at low abundances. *The Journal of applied ecology*, **53**, 722–732.

Edgar, R.C. (2013) UPARSE: highly accurate OTU sequences from microbial amplicon

reads. *Nature Methods*, **10**, 996–998.

Fitzpatrick, M.C., Preisser, E.L., Ellison, A.M. & Elkinton, J.S. (2009) Observer bias and the

detection of low-density populations. *Ecological applications: a publication of the*

Ecological Society of America, **19**, 1673–1679.

Foster, V., Giesler, R.J., Wilson, A.M.W., Nall, C.R. & Cook, E.J. (2016) Identifying the

physical features of marina infrastructure associated with the presence of non-native

species in the UK. *Marine biology*, **163**, 173.

Frøslev, T.G., Kjøller, R., Bruun, H.H., Ejrnæs, R., Brunbjerg, A.K., Pietroni, C. & Hansen,

A.J. (2017) Algorithm for post-clustering curation of DNA amplicon data yields reliable

biodiversity estimates. *Nature Communications*, **8**, 1188.

Glasby, T.M., Connell, S.D., Holloway, M.G. & Hewitt, C.L. (2007) Nonindigenous biota on

artificial structures: could habitat creation facilitate biological invasions? *Marine Biology*,

151, 887–895.

Goldberg, C.S., Turner, C.R., Deiner, K., Klymus, K.E., Thomsen, P.F., Murphy, M.A., …

Taberlet, P. (2016) Critical considerations for the application of environmental DNA

methods to detect aquatic species ed M. Gilbert. *Methods in Ecology and Evolution*, **7**,

1299–1307.

Grey, E.K., Bernatchez, L., Cassey, P., Deiner, K., Deveney, M., Howland, K.L., …Lodge,

D.M. (2018) Effects of sampling effort on biodiversity patterns estimated from

environmental DNA metabarcoding surveys. *Scientific Reports*, **8**, 8843.

Jarman, S.N., Berry, O. & Bunce, M. (2018) The value of environmental DNA biobanking for

long-term biomonitoring. *Nature Ecology & Evolution*.

Kim, P., Kim, D., Yoon, T.J. & Shin, S. (2018) Early detection of marine invasive species,

Bugula neritina (Bryozoa: Cheilostomatida), using species-specific primers and

environmental DNA analysis in Korea. *Marine Environmental Research*, **139**, 1–10.

Lacoursière-Roussel, A., Howland, K., Normandeau, E., Grey, E.K., Archambault, P., Deiner,

K., Lodge, D.M., Hernandez, C., Leduc, N. & Bernatchez, L. (2018) eDNA

metabarcoding as a new surveillance approach for coastal Arctic biodiversity. *Ecology*

and Evolution, **26**, 146.

Leray, M., Yang, J.Y., Meyer, C.P., Mills, S.C., Agudelo, N., Ranwez, V., Boehm, J.T. &

Machida, R.J. (2013) A new versatile primer set targeting a short fragment of the

mitochondrial COI region for metabarcoding metazoan diversity: application for

characterizing coral reef fish gut contents. *Frontiers in Zoology*, **10**, 34.

Lovell, S.J., Stone, S.F. & Fernandez, L. (2006) The Economic Impacts of Aquatic Invasive

Species: A Review of the Literature. *Agricultural and Resource Economics Review*, **35**,

195–208.

MacConaill, L.E., Burns, R.T., Nag, A., Coleman, H.A., Slevin, M.K., Giorda, … Thorner, A.R.

(2018) Unique, dual-indexed sequencing adapters with UMIs effectively eliminate index

cross-talk and significantly improve sensitivity of massively parallel sequencing. *BMC*

Genomics, **19**, 30.

Machida, R.J., Leray, M., Ho, S.-L. & Knowlton, N. (2017) Metazoan mitochondrial gene

sequence reference datasets for taxonomic assignment of environmental samples.

Scientific Data, **4**, 170027.

Martin, M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal*, **17**, 10–12.

Mazza, G., Tricarico, E., Genovesi, P. & Gherardi, F. (2014) Biological invaders are threats to

human health: an overview. *Ethology Ecology & Evolution*, **26**, 112–129.

McQuillan, J.S. & Robidart, J.C. (2017) Molecular-biological sensing in aquatic environments:

recent developments and emerging capabilities. *Current Opinion in Biotechnology*, **45**,

43–50.

Mistri, M. (2003) The non-indigenous mussel *Musculista senhousia* in an Adriatic lagoon:

effects on benthic community over a ten year period. *Journal of the Marine Biological*

Association of the United Kingdom, **83**, 1277–1278.

Molnar, J.L., Gamboa, R.L., Revenga, C. & Spalding, M.D. (2008) Assessing the global threat

of invasive species to marine biodiversity. *Frontiers in Ecology and the Environment*, **6**,

485–492.

Nascimento, F.J.A., Lallias, D., Bik, H.M. & Creer, S. (2018) Sample size effects on the

assessment of eukaryotic diversity and community structure in aquatic sediments using

high-throughput sequencing. *Scientific Reports*, **8**, 11737.

Nunes, A.L., Katsanevakis, S., Zenetos, A. & Cardoso, A.C. (2014) Gateways to alien

invasions in the European seas. *Aquatic Invasions*, **9**, 133–144.

O'Donnell, J.L., Kelly, R.P., Shelton, A.O., Samhouri, J.F., Lowell, N.C. & Williams, G.D.

(2017) Spatial distribution of environmental DNA in a nearshore marine habitat. *PeerJ*,

5, e3044.

Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R.B., … Wagner,

H. (2011) vegan: Community ecology package. *R package version*, 117–118.

Oliver, I. & Beattie, A.J. (1993) A Possible Method for the Rapid Assessment of Biodiversity.

Conservation Biology, **7**, 562–568.

Pawlowski, J., Kelly-Quinn, M., Altermatt, F., Apothéloz-Perret-Gentil, L., Beja, P., Boggero,

A.,… Kahlert, M. (2018) The future of biotic indices in the ecogenomic era: Integrating

(e)DNA metabarcoding in biological assessment of aquatic ecosystems. *The Science of*

the Total Environment, **637-638**, 1295–1310.

Pearce, F., Peeler, E. & Stebbing, P. (2012) *Modelling the Risk of the Introduction and*

Spread of Non-Indigenous Species in the UK and Ireland; Project Report for E5405W.

CEFAS.

Pérez-Portela, R., Arranz, V., Rius, M. & Turon, M. (2013) Cryptic speciation or global

spread? The case of a cosmopolitan marine invertebrate with limited dispersal

capabilities. *Scientific Reports*, **3**, 3197.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. &

Glöckner, F.O. (2013) The SILVA ribosomal RNA gene database project: improved

data processing and web-based tools. *Nucleic Acids Research*, **41**, D590–6.

R Core Team. (2018) *R: A Language and Environment for Statistical Computing*. ISBN 3-

900051-07-0.

Renshaw, M.A., Olds, B.P., Jerde, C.L., Mcveigh, M.M. & Lodge, D.M. (2015) The room

temperature preservation of filtered environmental DNA samples and assimilation into a

phenol-chloroform-isoamyl alcohol DNA extraction. *Molecular Ecology Resources*, **15**,

168–176.

Ricciardi, A., Hoopes, M.F., Marchetti, M.P. & Lockwood, J.L. (2013) Progress toward

understanding the ecological impacts of nonnative species. *Ecological Monographs*, **83**,

263–282.

Rius, M., Teske, P.R., Maríquez, P.H., Suárez-Jiménez, R., McQuaid, C.D. & Castilla, J.

(2017) Ecological dominance along rocky shores, with a focus on intertidal ascidians.

Oceanography and Marine Biology. pp. 55–86.

Rivero, N.K., Dafforn, K.A., Coleman, M.A. & Johnston, E.L. (2013) Environmental and

ecological changes associated with a marina. *Biofouling*, **29**, 803–815.

Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. (2016) VSEARCH: a versatile open

source tool for metagenomics. *PeerJ*, **4**, e2584.

Sala, E. & Knowlton, N. (2006) Global Marine Biodiversity Trends. *Annual Review of*

Environment and Resources, **31**, 93–122.

Sassoubre, L.M., Yamahara, K.M., Gardner, L.D., Block, B.A. & Boehm, A.B. (2016)

Quantification of Environmental DNA (eDNA) Shedding and Decay Rates for Three

Marine Fish. *Environmental Science & Technology*, **50**, 10456–10464.

Scott, R., Zhan, A., Brown, E.A., Chain, F.J.J., Cristescu, M.E., Gras, R. & MacIsaac, H.J.

(2018) Optimization and performance testing of a sequence processing pipeline applied

to detection of nonindigenous species. *Evolutionary Applications*, **11**, 891–905.

Sellers, G.S., Di Muri, C., Gómez, A. & Hänfling, B. (2018) Mu-DNA: a modular universal

DNA extraction method adaptable for a wide range of sample types. *Metabarcoding*

and Metagenomics, **2**, e24556.

Shaw, J.L.A., Clarke, L.J., Wedderburn, S.D., Barnes, T.C., Weyrich, L.S. & Cooper, A.

(2016) Comparison of environmental DNA metabarcoding and conventional fish survey

methods in a river system. *Biological Conservation*, **197**, 131–138.

Simpson, T.J.S., Dias, P.J., Snow, M., Muñoz, J. & Berry, T. (2017) Real-time PCR detection

of *Didemnum perlucidum* (Monniot, 1983) and *Didemnum vexillum* (Kott, 2002) in an

applied routine marine biosecurity context. *Molecular Ecology Resources*, **17**, 443–453.

Spens, J., Evans, A.R., Halfmaerten, D., Knudsen, S.W., Sengupta, M.E., Mak, S.S.T.,

Sigsgaard, E.E. & Hellström, M. (2016) Comparison of capture and storage methods for

aqueous macrobial eDNA using an optimized extraction protocol: advantage of

enclosed filter. *Methods in Ecology and Evolution*, **8**, 635-645.

Turner, C.R., Barnes, M.A., Xu, C.C.Y., Jones, S.E., Jerde, C.L. & Lodge, D.M. (2014)

Particle size distribution and optimal capture of aqueous macrobial eDNA. *Methods in*

Ecology and Evolution, **5**, 676–684.

Turner, C.R., Uy, K.L. & Everhart, R.C. (2015) Fish environmental DNA is more concentrated

in aquatic sediments than surface water. *Biological Conservation*, **183**, 93–102.

Wangensteen, O.S., Cebrian, E., Palacín, C. & Turon, X. (2018a) Under the canopy:

Community-wide effects of invasive algae in Marine Protected Areas revealed by

metabarcoding. *Marine Pollution Bulletin*, **127**, 54–66.

Wangensteen, O.S., Palacín, C., Guardiola, M. & Turon, X. (2018b) DNA metabarcoding of

littoral hard-bottom communities: high diversity and database gaps revealed by two

molecular markers. *PeerJ*, **6**, e4705.

Wood, C.A., Bishop, J.D.D., Rennocks, L. & Crundwell, R. (2016) *RAS 2015 Non-Native*

Species Rapid Assessment Surveys in English Marinas (E Anglia & W Coast). Report

to the Bromley Trust.

Wood, C.A., Bishop, J.D.D. & Yunnie, A.L.E. (2015a) *RAS 2014 Non-Native Species Rapid*

Assessment Surveys in English Marinas. Report to the Bromley Trust.

Wood, C.A., Bishop, J.D.D. & Yunnie, A.L.E. (2015b) *Comprehensive Reassessment of NNS*

in Welsh Marinas. Report to Natural Resources Wales.

Wood, S.A., Zaiko, A., Richter, I., Inglis, G.J. & Pochon, X. (2017) Development of a real-time

polymerase chain reaction assay for the detection of the invasive Mediterranean

fanworm, Sabella spallanzanii, in environmental samples. *Environmental science and*

pollution research international, **24**, 17373–17382.

Worm, B., Barbier, E.B., Beaumont, N., Duffy, J.E., Folke, C., Halpern, … Watson, R. (2006)

Impacts of biodiversity loss on ocean ecosystem services. *Science*, **314**, 787–790.

WoRMS Editorial Board. (2018) World Register of Marine Species,

http://www.marinespecies.org

Zhan, A., Hulák, M., Sylvester, F., Huang, X., Adebayo, A.A., Abbott, C.L., … Macisaac, H.J.

(2013) High sensitivity of 454 pyrosequencing for detection of rare species in aquatic

communities. *Methods in Ecology and Evolution,* **4**, 558–565.

Fig. 1. *a) Map of United Kingdom indicating the geographic position of the sampled sites, a legend is*

provided below indicating the four sites (PQ, TB, TQ and HH) and colours for water and sediment eDNA

samples for each site. Barplots detailing number of OTUs detected across sampling sites and eDNA

sample type for COI b) and 18S rRNA d) metabarcoding of UK marinas, the break in bars indicates the

number of shared OTUs between sediment and water eDNA samples. Non-metric multidimensional

scaling ordination plots based on Bray-Curtis dissimilarities between: c) COI and e) 18S rRNA

metabarcoding of marina sediment and water eDNA samples.

Fig. 2. *Horizontal stacked barchart detailing proportion of OTUs detected in eDNA from sediment, water*

or both across the top 10 phyla for pooled 18S rRNA and COI metabarcoding of UK marinas.

669   **Fig. 3.** *Presence absence diagram for seven non-indigenous species across four sampling sites. For each*

670   *species-location the left semi-circle indicates the detection using eDNA metabarcoding surveys of 18S*

671   *rRNA and COI fragments, and the right semi-circle indicates the detection from traditional manual*

672   *surveys. Blue indicates a positive detection for that species-location and red indicates no detection.*

675