Cellular organization of visual information processing channels in the mouse
visual cortex

Xu Han[1,2,*], Ben Vermaercke[1,3], Vincent Bonin[1,2,4,5,6,*]

[1] Neuro-Electronics Research Flanders, Kapeldreef 75, 3001 Leuven, Belgium.

[2] KU Leuven, Department of Biology & Leuven Brain Institute, 3000 Leuven, Belgium.

[3] VIB-KU Leuven Center for Brain & Disease Research, 3000 Leuven, Belgium.

[4] VIB, 3001 Leuven, Belgium.

[5] imec, 3001 Leuven, Belgium

[6] Lead contact Vincent.Bonin@nerf.be

[*] For correspondence: xu.han@nerf.be (X.H.), Vincent.Bonin@nerf.be (V.B)

Abstract

Visual processing and behavior depend on specialized neural representations and information
channels that encode distinct visual information and enable distinct computations. Our
understanding of the neural substrate, however, remain severely limited by sparse recordings and
the restricted range of visual areas and visual stimuli considered. We characterized in the mouse
the multidimensional spatiotemporal tuning properties of > 30,000 layer 2/3 pyramidal neurons
across seven areas of the cortex. The dataset reveals population specialized for processing of
oriented and non-oriented contrast, spatiotemporal frequency, and motion speed. Areal analysis
reveals profound functional diversity and specificity as well as highly specific representations of
visual processing channels in distinct visual areas. Clustering analysis shows a branching of
visual representations along the posterior to anterior axis, and between lateral and dorsal areas.
Overall, this dataset provides a cellular-resolution atlas for understanding organizing principles
underlying sensory representations across the cortex.

Summary (for reviewers)

Visual representations and visual channels are the cornerstones of mammalian visual processing
and critical for a range of life sustaining behaviors. However, the lack of data sets spanning

multiple visual areas preclude unambiguous identification of visual processing streams and the
sparse, singular recording data sets obtained thus far are insufficient to reveal the functional
diversity of visual areas and to study visual information channels. We characterized the tunings
of over 30,000 cortical excitatory neurons from 7 visual areas to a broad array of stimuli and
studied their responses in terms of their ability to encode orientation, spatiotemporal contrast and
visual motion speed. We found all mouse visual cortical areas convey diverse information but
show distinct biases in terms of numbers of neurons tuned to particular spatiotemporal features.
Neurons in visual areas differ in their spatiotemporal tuning but also in their relative response to
oriented and unoriented contrast. We uncovered a population that preferentially responds to
unoriented contrast and shows only weak responses to oriented stimuli. This population is
strongly overrepresented in certain areas (V1, LM and LI) and underrepresented in others (AL,
RL, AM, and PM). Spatiotemporal tunings are broadly distributed in all visual areas indicating that
all areas have access to broad spatiotemporal information. However, individual areas show
specific biases. While V1 is heavily biased in favor of low spatial and temporal frequencies, area
LM responds more strongly to mid-range frequencies. Areas PM and LI are biased in favor of
slowly-varying high-resolution signals. By comparison, anterior areas AL, RL and AM are heavily
biased in favor of fast-varying, low to mid spatial frequency signals. Critically, theses biases
express themselves in vastly different number of cells tuned to particular features, suggesting
differential sampling of visual processing channels across areas. Comparing across areas, we
found divergent visual representations between anterior and posterior areas, and between lateral
and dorsal areas, suggesting the segregated organization of cortical streams for distinct
information processing.

Introduction

The visual cortex is one of the most elaborate sensory processing network and processing in its
multiple areas are important for a range of perceptual, cognitive and motor abilities (Dipoppa et
al., 2018; Marques et al., 2018; Minderer et al., 2019; Steinmetz et al., 2019). How distinct visual
areas integrate and carry information from distinct visual channels is fundamental to their function
and constrains the computations they can perform. However, their functional organization and
composition particularly at the cellular level is largely unknown.

Studies of the mouse cortex have the potential to shed significant new light on visual information
processing streams and their contributions to perception and behavior. The mouse visual cortex
contains over ten higher visual areas (HVAs), which receive direct input from primary visual cortex

(V1) and form parallel representations of the visual field (Garrett et al., 2014; Wang and
Burkhalter, 2007; Zhuang et al., 2017). HVAs display rich functional response properties and
connectivity allowing for a wide range of associations and behaviors (Gămănuţ et al., 2018;
Juavinett and Callaway, 2015; La Chioma et al., 2019; Marshel et al., 2011; Minderer et al., 2019;
Olcese et al., 2013; Roth et al., 2012; Smith et al., 2017; Tohmi et al., 2014). HVAs show
differences in visual tuning properties including orientation selectivity, spatiotemporal frequency
and motion speed preference (Andermann et al., 2011; Marshel et al., 2011; Murakami et al.,
2017). Areas PM and AL, for example, show marked preferences for slow and fast varying stimuli
and neurons projecting to these areas show functional differences that are biased towards the
preferences of target areas (Glickfeld et al., 2013; Han et al., 2018; Kim et al., 2020, 2018). The
specificity of visual representations and long-range connections are suggestive of specialized
subnetworks involving distinct visual processing channels.

In primates there is anatomical and functional evidence of visual processing channels, namely
ventral and dorsal streams, that are specialized in processing distinct visual features at different
levels of complexity along the visual hierarchy (Felleman and Van Essen, 1991; Nassi and
Callaway, 2009). Whether such distinct visual processing channels exist in the mouse cortex and
the extent to which distinct cortical areas partake in these processing channels is yet to be
determined. Existing datasets, whether in rodents or in primates although compelling, vastly
undersample the cellular functional diversity of visual cortical areas. To study diversity in visual
areas and reveal visual information processing channels, it is essential to directly compare
response patterns across visual areas and to employ a broad enough stimulus set and large
neuronal sampling to allow characterization of a wide range of tuning properties of visual neurons.
Such sampling would allow classification of visual neurons into functional cell types and
identification of shared cell classes across visual areas to uncover distinct processing channels
(Baden et al., 2016; Piscopo et al., 2013; Román Rosón et al., 2019). The vast majority of studies
in mouse visual cortical areas used gratings stimuli, which only robustly activate a fraction of the
neurons. A recent large-scale study (de Vries et al., 2020) recorded responses to broader sets of
stimuli and found different visual areas and cell types show differential stimulus selectivity, but did
not address visual channels.

To characterize the functional diversity of mouse visual areas and probe their functional specificity
and involvement distinct visual information processing channels, we characterized the
multidimensional tuning properties of > 30,000 layer 2/3 pyramidal neurons in 7 mouse visual
areas. We found that all mouse visual cortical areas encode diverse albeit distinct visual

information. The distinct encoding and functional diversity are seen in overall bias of visual tuning
and the sampling of visual features. Individual visual areas encode distinct visual motion speeds
and also differentially partake in distinct visual information processing channels. Altogether, this
study shows profound diversity and specificity of representations in mouse visual cortical areas
and provides a cellular-resolution blueprint for understanding organizing principles underlying
visual cortical processing.

Results

Characterizing layer 2/3 neurons with cellular imaging and visual noise stimuli

We used large-scale chronic cellular imaging (Figure 1A-C) to probe the visual tuning properties
of layer 2/3 cortical excitatory neurons in 7 mouse visual areas (V1; LM: lateromedial; AL:
anterolateral; RL: rostrolateral; AM: anteromedial; PM: posteromedial; LI: laterointermediate). We
focused on tuning for orientation and spatiotemporal features that are characteristic of visual
information processing channels in the mammalian visual system. In the first part of this paper,
we examine, for each of the studied visual areas, both the biases in neuronal tuning and the
neural representation of the visual feature space. In the second part, we examine the implications
of cortical neurons' visual tuning for encoding of visual motion speed and for the organization of
visual information processing channels in the mouse cortex.

To characterize neuronal tunings for visual features in mouse visual cortical areas, we used large-
field stochastic visual stimuli designed to activate diverse orientation and spatiotemporal
frequency channels. The stimuli were made of spectrally-filtered random noise with filters of
specific center spatial and temporal frequencies, center orientation, and orientation bandwidth
(Figure supplement 1D; Materials and Methods), resulting in dynamic textures that varied in
orientation, spatiotemporal frequency and spatial elongation (anisotropy). We catalogued in
separate imaging sessions responses to isotropic (non-oriented) textures and anisotropic
(oriented) textures of different spatiotemporal frequencies (Figure 1D). To characterize
functionally-diverse populations and probe visual processing channels, we tested 30 different
combinations of spatial and temporal frequencies (spatial frequencies: 0.02, 0.04, 0.08, 0.16, 0.32
cpd; temporal frequencies: 0.5, 1, 2, 4, 8, 16 Hz; data sets 1 and 2). The power spectrum of the
isotropic textures (Figure 1D, left) is evenly distributed across spatial orientations and therefore
well suited to probe neurons tuned to non-oriented features. The anisotropic stimuli (Figure 1D,
right) had power restricted to a narrow band of orientations (5 deg, full width at half maximum,
FWHM) and are well suited to probe the properties of orientation tuned neurons. To cover a full

range of orientations, the stimulus center orientation was varied slowly (45 deg/s) over stimulation
epochs. In a separate experiment, we recorded responses to stimuli of different elongation and
spatial frequencies (data set 3). The visual stimuli were presented on a visual display positioned
in the right visual field (azimuth 0~100 deg; elevation -30~50 deg) in 4-second epochs interleaved
with 4-second epochs of gray screen of the same mean luminance. The stimuli were presented
four times in randomized order with stimuli in repeated trials obtained using distinct pseudo-
random generator seeds. Data sets were registered to common reference to match somatic ROIs
across experiments.

To target cortical excitatory neurons, we employed transgenic mice (Thy1-GCaMP6s GP4.12,
Dana et al., 2014) that broadly express the calcium indicator GCaMP6s in a large subset of
pyramidal neurons (Figure 1C). As anesthesia profoundly alters visual response properties (Lien
and Scanziani, 2013; Suzuki and Larkum, 2020), recordings were made in awake animals. Mice
were head-fixed comfortably on a treadmill, yielding recordings in quiet wakefulness allowing for
occasional locomotor behavior. Locomotion epochs were too infrequent to be examined. All data
were included irrespective of the animal's locomotion state. The cellular imaging was targeted to
the retinotopic locations in the mouse cortex that corresponded to the visual display center. To
identify the borders of visual areas at high spatial resolution, we performed widefield and cellular
calcium imaging of responses to retinotopic stimuli (Figure supplement 1A-C). V1 and surrounding
higher visual areas were assigned using the reversals in the retinotopic representations (Wang
and Burkhalter, 2007; Zhuang et al., 2017). Borders between areas were further refined by
manually inspecting cellular responses to retinotopic stimuli in 2-photon images. 2-photon imaging
was focused 100–300 μm below the pia to target cortical pyramidal neurons within layer 2/3.

Figure 1. Spectral noise stimuli elicited rich responses in multiple visual cortical areas. (A) Cartoon
showing V1 and six higher visual areas investigated in this study. (B) Experimental setup for 2-photon
calcium imaging on awake mice. Mice were placed on a treadmill while cellular responses to spectral noise
stimuli were recorded using 2-photon calcium imaging. (C) Example field of view of the V1 L2/3 pyramidal
neurons of Thy-GCaMP6s (GP4.12) mice. Scale bar: 100um. (D) Example cells showing distinct responses
to ISO and ANISO stimuli (isotropic vs anisotropic noise stimuli; left and right icons). ISO and ANISO stimuli
have varying spatiotemporal frequencies, and non-oriented and oriented textures, respectively. See
methods and figure supplement 1 for a detailed description. Gray traces show the median stimulus-
triggered responses across trials. The median absolute deviations across trials are shown as light gray

shadows. Cell 1 and 2 selectively responded to ISO and ANISO stimuli respectively, while cell 3 showed
similar responses to both stimuli. Blue bars indicate 4-sec stimulus epochs. ΔF/F: ratio of fluorescent
changes. (E) Model fitting of spatiotemporal frequency tunings. Neurons' responses to the preferred stimuli
are used for fitting. Yellow: ISO response; green: ANISO responses. The calcium responses during the
stimulus epoch are averaged to generate the raw response maps (upper left), which are then fitted to two-
dimensional Gaussian models (bottom left). The size of the dot indicates the normalized response
amplitude. A high-resolution fit is generated by oversampling (right panel). Contours indicate the region of
frequency eliciting responses greater than 50% of the maximal amplitude. (F) Distribution of peak
responses of reliable cells and unselected cells. Distributions are smoothed with a Gaussian kernel
(bandwidth=0.05). Color bars indicate median values. Gray scale bars indicate 2.5% of cells of each
distribution. (G) Distribution of ANISO preference index of each area. Gaussian kernel bandwidth=0.05.
Gray scale bars indicate 2% of cells for each distribution. Pairwise statistical comparisons were shown in
the right panel (KS test with Bonferroni correction). See Table 1 for sample sizes.

The following figure supplements are available for figure 1:

Figure supplement 1: Retinotopic mapping and spectral noise stimuli.

Figure supplement 2: Population statistics of response to ISO and ANISO stimuli.

Specialized populations for processing of oriented and non-oriented contrast

Out of 78,930 identified layer 2/3 neurons (n = 10 mice), 58% showed a stimulus-evoke response

to at least one of the visual stimuli tested (46169/78930 cells, > 3sd of the baseline ΔF/F, see

Materials and Methods, data sets 1–3). To characterize the neurons' visual tunings, we identified

neurons showing strong, reliable responses to the visual stimuli (figure supplement 2A-B). We

correlated the raw ΔF/F activity time courses in visual stimulation epochs across repeated trials

and selected the cells showing pronounced trial-to-trial correlations (75th percentile of correlation

coefficients > 0.3, see Methods). This criterion served to identify reliably responding cells and

exclude weak responders from which clear visual tunings could not be extracted (Figure 1F).

Across visual areas, about a third of the probed neurons showed visual responses that fulfilled

this criterion (31053/78930 cells, data sets 1–3) yielding samples of 1800 to 9400 functionally-

characterized neurons for each visual area (Table 1).

The responses were highly tuned to specific ranges of spatiotemporal frequencies, indicative of

visual processing channels (Figure 1D, traces). To characterize these tunings, we fitted 2-

dimensional Gaussian functions (Figure 1E) to individual neurons' trial-averaged amplitudes of

stimulus-evoked responses and extracted response descriptors including peak response

magnitude, spatial and temporal bandwidths, and preferred and cutoff spatiotemporal

frequencies. These function fits captured the magnitudes of responses as well their dependence

on spatiotemporal frequency (Figure 1E, left; Figure supplement 2C). For neurons that responded

to both isotropic and anisotropic stimuli, spatiotemporal tuning estimates were highly correlated

(Figure 1E, bottom; Figure supplement 2D), therefore, we used the fits from the data set with

highest peak responses (Figure 1E, thick lines). In total we good fits were obtained for a total of
24986 neurons across areas (normalized residual of fits < 0.1, Table 1).

In addition to spatiotemporal tuning, the neurons also show a variety of responses for isotropic
and anisotropic stimuli (Figure 1D). While some cells show similarly strong responses to both
types of stimuli (Figure 1D, bottom), many have clear preferences, responding strongly to one
stimulus type and weakly to the other (Figure 1D, middle vs. top). A fraction of cells shows a clear
preference for anisotropic stimuli (Figure 1D, middle). A preference for anisotropic stimuli is
indicative of pronounced sensitivity to oriented edges or orientation tuning, whereas a lack of this
preference is indicative of a lack of orientation tuning. Surprisingly, a separate population of visual
cortical neurons shows a strong preference for isotropic stimuli, which may reflect tuning for a
non-oriented feature. To quantify the neurons' preference for isotropic and anisotropic stimuli, we
extracted peak response amplitudes from model fits, and computed an anisotropy preference
index (API), which we defined as the ratio of the difference in peak responses to the stimuli over
their sum (Figure 1G). The index has a value of -1 for perfectly isotropic preferring neurons, 1 for
perfectly anisotropic preferring neurons and 0 for neurons without preference.

The neurons show highly diverse preferences for isotropic and anisotropic stimuli (Figure 1G --
Figure supplement 2E). Across the population, 32% of the cells showed a clear preference for
isotropic stimuli (7928/24986 cells; API < -1/3), whereas 20% showed a clear preference for
anisotropic stimuli (5045/24986 cells; API > 1/3). The remaining 48% showed only weak
preferences (12013/24986 cells). Importantly, anisotropic and isotropic preferring cells are
observed across all visual areas, however, there were specific biases (Figure 1G). Areas V1, LM
and LI have a pronounced bias for isotropic stimuli. In contrast, areas AL, RL, AM and PM show
less prevalence of isotropic-preferring cells and an enrichment of cells without preference.

Thus, layer 2/3 visual cortical populations show pronounced preferences for isotropic and
anisotropic stimuli. This may reflect specific visual channels specialized for oriented and non-
oriented contrast.

Specializations for oriented and non-oriented contrast in dorsal and lateral areas

To examine these visual processing channels in further detail, we characterized, in a separate
dataset (22546 cells showing reliable responses, 10 mice), the responses to isotropic stimuli and
3 different levels of anisotropy (orientation bandwidth: infinite, 40, 10, 5 deg FWHM; Figure 2A).
We sought to determine whether responses to stimulus elongation form a continuum or a

dichotomy. We sampled a range of spatial frequencies (0.04, 0.08, 0.16, 0.32 cpd, 2 Hz), and

examined the amplitudes of responses as a function of elongation (Figure 2B).

Figure 2. Distinct preferences for oriented and non-oriented stimuli in dorsal and lateral areas. (A)
Example noise stimuli showing increasing stimulus anisotropies and texture elongation corresponding to
increasingly narrow orientation bandwidths (B) Example cells showing distinct responses to combinations
of spatial frequencies and stimulus anisotropies (upper panels). These cells correspond to the example
cells in Figure 1D, showing matching response preferences for ISO and ANISO stimuli. Tuning curves for
the stimulus anisotropy are measured at the preferred spatial frequencies (red curves) and are shown in
the lower panels. (C) Population tuning curves of randomly selected 500 cells from each area, sorted by
peak-normalized response amplitude. (D) Average tuning curves for stimulus anisotropy across population
of each area. Mean ± sem. Inset: scheme of cortical areas. See Table 1 for sample sizes.

We found across the population a dichotomy of responses to spatial elongation (Figure 2C).

Anisotropic preferring cells did not respond to isotropic stimuli (Figure 2B, middle, 2C) and show

nearly invariant responses with elongation. Isotropic preferring neurons (Figure 2B, left and right,

2C) instead show a steep fall-off of the response amplitude with elongation. We observed in the

population little evidence of tuning to specific length scales. Comparing population response

patterns across visual areas (Figure 2C), we found that areas V1, LM, and LI show the largest

fractions of cells tuned to isotropic stimuli, whereas areas PM, AL, RL and AM show an enrichment

of orientation tuned neurons. The striking differences in the tuning of these areas was also seen
in the average tuning curves (Figure 2D).

Taken together the results indicate that visual cortex contains populations specialized for
processing of oriented and non-oriented contrast and that these populations are non-uniformly
distributed across visual areas.

252 Diversity and specificity of spatiotemporal visual channels in visual areas

We next examined the diversity and specificity of spatiotemporal channels in mouse visual areas.
Previous studies reported biases in spatiotemporal tunings across visual areas, particularly in
areas AL and PM which show biases in favor of fast and slow stimuli respectively (Andermann et
al., 2011; Marshel et al., 2011; Murakami et al., 2017). These biases could reflect specific
sampling of specific spatiotemporal channels. To examine in detail the diversity and specificity of
spatiotemporal tunings in visual areas, lower and higher, we examined population responses to
spatiotemporal stimuli, including the numbers of neurons activated as a function of spatiotemporal
frequency, and the distributions of spatiotemporal tuning types (lowpass, bandpass, or high-pass),
preferred frequencies and tuning bandwidths.

We observed a profound diversity of spatiotemporal channels in mouse visual areas. Each area
shows highly diverse tunings, with each neuron individually responding to distinct ranges of
spatiotemporal frequencies, but as a population, covering broad swaths of the spatiotemporal
spectrum (Figure 3A). This diversity in neuronal tunings is particularly evident in the broad
distributions of peak and cutoff spatiotemporal frequencies (Figures 3D; Figure 4A,C; Figure
supplement 4A,B), but is also seen in the spatiotemporal tuning types (Figures 3C) and tuning
bandwidths (Figure supplement 4C). Critically, the diversity was observed across all of the visual
areas, not only seen in lower visual areas V1 and LM, but also in more specialized areas (e.g. AL
and PM).

Although each area is comprised of neurons with diverse spatiotemporal preferences, the biased
distributions of these neurons among visual areas give rise to striking functional specificity such
that each area displays a unique set of activation patterns in response to spatiotemporal stimuli
(Figure 3).

First, V1 and LM, show profound specializations for processing low and mid-range spatio-
temporal frequencies, respectively. V1 and LM are differentially activated by low and mid-range
frequencies (Figure 3B, **V1** vs **LM**) and differ strikingly in the numbers of spatiotemporally lowpass

and bandpass cells (Figure 3C, **V1** vs **LM**) and in their distributions of peak spatiotemporal

frequencies (Figure 3D, **V1** vs **LM**). Second, areas PM and LI which are located on opposite sides

of V1 share a pronounced preference for low temporal (0.5–4Hz), high spatial frequencies (0.08–

0.32 cpd), which links them to processing of high spatial resolution visual signals. This is seen in

the population activation pattern, and the distributions of passband properties and peak

frequencies (Figure 3B-D, **PM** and **LI**).

Figure 3. Diversity and specificity of spatiotemporal tunings in visual areas. (A) Superimposed
surfaces showing diverse spatiotemporal tunings of 20 randomly selected cells. Inset: scheme of cortical
areas. (B) Proportion of neurons responding to certain spatiotemporal frequencies. (C) Proportions of
spatiotemporal passband types. H: highpass. B: bandpass. L: lowpass. (D) Scatter plots showing the
distribution of preferred frequencies of the populations across areas. The inner box delineates the range of
frequency explored in this study, dots at the boarders are scattered outward to visualize the density. See
Table 1 for sample sizes.

By comparison, areas AL, RL and AM, which are located anterior to V1, share a striking

preference for low resolution, fast varying visual signals. These areas show weak activation at

low temporal (1 Hz and below) and high spatial frequencies (0.08 cpd and above) (Figure 3B, **AL**,

RL and **AM**) and broadly distributed peak frequencies with a mode around 0.08 cpd and 4 Hz

(Figure 3D, **AL**, **RL** and **AM**). They also show a preponderance of temporally bandpass and high-

pass neurons and a sparsity of spatially high-pass cells (Figure 3C, **AL**, **RL** and **AM**). Moreover,

anterior visual areas have their own distinguishing features (Figure 3B-D, **AL**, **RL** and **AM**). Area

RL shows an excess of spatially-lowpass and temporally-highpass cells and the strongest

activation to stimuli with high temporal frequencies and low spatial frequencies. Area AM, in

comparison, shows tunings shifted towards mid-range spatial frequencies and weaker responses

to very high temporal frequencies. Area AL comprises a higher diversity of tunings without clear

biases seen in areas RL and AM.

These differences between spatial and temporal properties of distinct visual areas of the mouse

cortex are robust and surprising. Each area shows a highly distinct distribution of peak spatial and

temporal frequencies (Figure 4A,C). Although the preferences measured with ISO and ANISO

stimuli are on a par with a study of response properties measured with gratings in V1, AL and PM

in awake animals (Figure 4B,D, red vs. black)(Andermann et al., 2011), they differ strikingly from

properties measured in anesthetized mice with discrepancies that can exceed one octave (Figure

4B,D, red vs. gray) (Marshel et al., 2011).

Figure 4: Neurons encoded higher spatial acuity and faster stimuli in wakefulness than in anesthesia. (A) Distributions of preferred spatial frequencies across areas. The width of each column is normalized to the maxima of the distribution. Distributions are smoothed with a Gaussian kernel (bandwidth=0.1). The median values are shown as colored bars. Gray scale bars indicate 2.5% of cells of each distribution. (B) Comparison of geometric means of neurons' preferred spatial frequencies acquired in this study (red bars) and previous studies in anesthetized (gray bars) and awake mice (black bars). Statistical differences between the current datasets and the 'anesthesia' dataset are shown at the bottom row. KS tests with Bonferroni correction. P<0.001, ***; P<0.01, **; P<0.05, *; p ≥ 0.05, n.s. (C-D) Temporal frequency analysis, same as (A-B). See Table 1 for sample sizes.

The following figure supplement is available for figure 4:

Figure supplement 3: Spatiotemporal tuning parameters.

These results indicate that spatiotemporal tuning properties of layer 2/3 cortical pyramidal neurons in each mouse visual area show high level of diversity, suggesting distributed processing with a wide range of visual information shared across areas. At the same time unique distributions of tuning parameters in each area creates functional specialization, allowing differential involvement of each area in spatiotemporal information processing. The specialization of HVAs suggests two separate branches of spatiotemporal representations where anterior areas (AL, RL,

AM) are involved in processing of fast-moving large objects while lateral areas (LI and PM) are
specialized in encoding fine details of still or slowly moving scenes.

Higher visual areas encode complementary motion speed representations

The distinct spatiotemporal tunings observed in different visual areas suggest that they might
encode distinct visual motion speeds. A previous study (Andermann et al., 2011) reported an
abundance of speed-tuned cells in area PM, which is biased towards low temporal and high
spatial frequencies (Figure 3B). To identify similar biases in our dataset that spans multiple visual
areas, we computed a spatiotemporal tuning index ξ extracted from the spatiotemporal tuning fits
to the responses to isotropic noise stimuli (Materials and methods). This parameter describes the
dependency between neurons' spatial and temporal tunings, which expresses as the slant of
spatiotemporal response profiles (Figure 5A). Neurons with $\xi \geq 0.5$ are speed tuned because they
show consistent speed tuning curves across broad ranges of spatial frequencies (Figure
supplement 4A,B). We examined the distributions of speed tuning indices across visual areas and
examined responses of speed-tuned cells to stimuli of different speeds.

We found a striking specialization for visual motion speed in mouse visual areas (Figure 5A). Area
V1 shows highly diverse tunings but only a small proportion of speed-tuned cells (25%, 835/3335
cells). Relative to V1, anterior visual areas AL, RL, and AM show a clear enrichment of speed-
tuned cells (42~48%), whereas areas LM, PM and LI show a less profound enrichment (35~37%).
Differences across visual areas are highly significant (Figure 5A, right).

Moreover, speed-tuned neurons encode distinct ranges of speed in different areas (Figure 5B, t-
tests). V1 and LM are broadly tuned for the intermediate speeds (peak at 12.5 ~ 25 deg/s). Areas
AL and AM primarily respond to fast-moving stimuli (peak at ~100 deg/s); RL neurons selectively
respond to extremely fast stimuli (peak at 200~400 deg/s). In contrast, areas PM and LI mainly
respond to slow motion (peak at 6.25 deg/s). Therefore, individual higher visual areas are
specialized for specific regimes of motion speed and they collectively establish a broad
representation of speed that might be critical for a wide range of visual behaviors.

Figure 5. Complementary representation of visual motion speed in higher visual areas. (A) Distribution of speed tuning index ξ of each area. Colored bars show median values. Neurons above the threshold (ξ ≥0.5) are deemed speed-tuned, otherwise untuned. The right panel shows statistical differences of the distributions across areas (KS tests with Bonferroni correction). (B) Population average speed tuning curves of each areas. Each line is the average of peak-normalized speed tuning curves of speed-tuned cells within each area. Error bars: standard error of the mean. (C) Enhanced speed discriminability in higher visual areas. Discrimination tasks were performed between cross-speed pairs of isotropic stimuli, which have 4-fold difference in speed. The discriminability is estimated as the number of cells required to reach 90% classification accuracy (red lines). Higher visual areas (except PM) show regional increases in the discrimination ability comparing to V1.

The following figure supplement is available for figure 5:

Figure supplement 4: Speed tuning analysis.

Enhanced speed discrimination capacity in higher visual areas

To assess whether the visual motion tuning properties endow visual areas with distinct encoding abilities, we compared how well the stimulus categories (spatiotemporal frequencies or speed) can be decoded from the activities of neuronal populations. We trained SVM (support vector machines) classifiers to discriminate stimulus pairs using the responses to ISO stimuli from random cell samples. All recording trials were randomly divided into two halves that were respectively used for training and testing the classifiers (Methods). The discrimination accuracy for each stimulus pair was measured with different population sizes, from which the population size required to reach 90% accuracy was interpolated and used as a proxy of discrimination power of each visual area in discriminating stimulus pairs. To measure the discriminability of speed, we tested how well the classifiers discriminate cross-speed stimulus pairs, which are neighboring stimulus pairs lying orthogonally to the iso-speed lines. The cross-speed pairs have small differences in the spatial (1 octave) and temporal frequency (1 octave), and a modest (4-fold) difference in speed (Figure 5C). Switching the combination of the spatial and temporal frequencies of cross-speed pairs yield iso-speed pairs (Figure supplement 4C), which also have small differences in the spatiotemporal frequency but identical visual motion speed. The difference between the discriminability of cross-speed and iso-speed pairs isolate speed discriminability, which is intermingled with the discriminability of spatiotemporal frequencies.

Higher visual areas showed better speed discrimination power than V1 in distinct frequency domains (Figure 5C). V1 showed relatively uniform ability to discriminate across cross-speed pairs, with a slight increase at lower spatial frequencies. Area LM showed better discrimination at slow to intermediated speed ranges (6.25 – 200 deg/s). Area AL clearly separated slow stimuli (12.5-25 deg/s) from intermediate speed (50-100 deg/s). Areas AM and RL were specialized for speed discrimination at intermediate frequencies. Area LI showed an elevated discrimination power for slow speed pairs (3.1-100 deg/s). Area PM showed a similar yet less pronounced pattern comparing to LI. Moreover, the frequency-specific enhancement of speed discrimination performances in higher visual areas was not merely a consequence of individual areas' frequency preferences, because the discriminability of cross-speed pairs remarkably outperformed the discriminability of iso-speed stimulus pairs in the same range of frequency space (Figure 5C; compare to Figure supplement 4C;). Altogether, these results highlight the area-specific enhancement of speed discriminability in higher visual areas, suggesting differential roles of higher visual areas in the visual motion processing.

Asymmetric sampling of visual information channels in mouse visual areas

We next examined the representations of visual processing channels in mouse visual areas. In the previous sections we characterized independently how specific aspects of visual scenes are represented in individual visual areas. To identify visual processing channels and examine multidimensional tunings, we employed a non-supervised clustering approach (spectral clustering, see Method and Materials) to group neurons according to their responses to isotropic and anisotropic stimuli of different spatiotemporal frequencies. The clustering approach also allows to determine whether different visual feature subspaces are captured by distinct cell populations in the visual cortex, and how these functional cell types are represented across visual areas.

We found neuronal responses in visual areas can be divided into broad functional cell classes. To cluster responses, we repeatedly sampled randomly 2000 response vectors to give each area equal weight. Each response vector contained the amplitudes of responses to ISO and ANISO stimuli of all combinations of spatial and temporal frequencies (Figure 6A). To determine the number of clusters needed to capture the diversity of responses across visual areas, we computed the total within-cluster variability as a function of number of clusters. The resulting Scree plot showed an inflection point at 12 clusters (Figure supplement 5A), indicating that the population response patterns to visual stimuli in visual areas can be described by approximately 12 functional cell groups, representing a unique combination of visual features, such as distinct spatiotemporal tunings and differential responses to isotropic and anisotropic stimuli (Figure 6A–B; Figure supplement 6).

We observed differential preferences for isotropic and anisotropic stimuli across identified cell groups (Figure supplement 6B), indicating distinct tuning for spatial elongation. To examine this, we further computed each group's average tuning for stimulus anisotropy from the responses to elongated textures, which were not used for the clustering. The resulting tuning curves showed remarkable consistency with individual groups' preferences for ISO versus ANISO stimuli (Figure supplement 6E), confirming independently that the dichotomy of selectivity for non-oriented versus oriented contrast is a distinguishing feature between functional groups.

Figure 6: Asymmetric distribution of visual channels across cortical areas. (A) Matrices showing individual areas are characterized by distinct composition of functional cell types. Each matrix comprises the response profiles of 2000 randomly selected neurons. Each row represents a single cell's peak-normalized responses to ISO and ANISO stimuli. Cells are clustered into 12 functional groups based on the similarity of response profiles using a spectral clustering approach (Materials and methods). The black lines delineate neighboring groups and connect the same groups across matrices. The stimulus parameters are annotated below the V1 matrix. (B) Heat map showing the cosine distances between the response profiles of the whole dataset. Black boxes delineate the 12 clusters. (C) Dot plot presenting the proportion of functional groups within and across areas. The groups are reordered to demonstrate the similarities and differences between areas. (D) The dissimilarity between areas are visualized as a Hierarchical tree based on the Euclidean distances estimated from the population composition across areas.

The following figure supplements are available for figure 6:

Figure supplement 5: Clustering.

Figure supplement 6: Summary of functional groups.

Although each visual area encompasses cells belonging to each visual information channel, the

proportion of cells belonging to each channel varied widely across visual areas (Figure 6C), such

that each area was characterized by a distinct proportion of cells showing different isotropic and

anisotropic preferences and spatiotemporal properties (Figure 6A). To quantify the similarities of

visual representations in visual areas, we calculated the Euclidian distance of population

compositions between pairs of areas, from which we computed a hierarchical tree of visual cortical

areas (Figure 6D). This comparison revealed V1 to be most similar to area LM, showing relative

uniform compositions of functional cell types. Areas PM and LI showed unique enrichment of

some groups and sparsity of some other groups, positioning them in a different branch from V1

and LM. These patterns were reversed in anterior areas, showing a vast segregation from

posterior areas (Figure 6C,D). These results indicate that mouse visual areas differentially sample

from distinct spatiotemporal frequencies channels. These channels have simple stereotyped

responses properties that capture a large fraction of the diversity seen across visual areas.

Discussion

This study provides a detailed functional characterization of the visual features driving visual

neuron responses and a comprehensive comparison of the cellular basis of representations

across seven areas of the mouse visual cortex. Our findings reveal highly distributed visual

information channels, representing distinct spatiotemporal and spatial orientation features in the

visual cortex. Biased distribution of these processing channels across areas suggests differential

role of individual areas in specialized visual computations. Considering the accumulating

evidence of higher-order computations in the mouse higher visual cortex (Akrami et al., 2018;

Burgess et al., 2016; Meijer et al., 2019; Morcos and Harvey, 2017), it is essential to understand

how area-specific representations of visual features arise along the visual hierarchy, and how

information from multiple modalities are integrated in the higher order cortex for complex

computations, such as visual motion and shape processing. The results and implications of this

study, provide a necessary basis for future studies investigating circuitry mechanisms for visual

perception and behaviors.

Comparison of spatiotemporal selectivity with previous studies

The preferred spatiotemporal frequencies observed in the current study are remarkably higher

than in an earlier study in anesthetized mice (Marshel et al., 2011), but comparable to what has

been reported in awake mice (Andermann et al., 2011). Moreover, our findings are not limited to

a specific mouse line (Thy1-GCaMP6s mice), recordings we performed in another GCaMP6

reporter mouse line (TRE-GCaMP6s line G6s2) (Wekselblatt et al., 2016), which labels a broader
population of pyramidal neurons, yielded similar results (data not shown). The lower ranges of
preferred temporal frequencies observed in anesthetized mice (Marshel et al., 2011; Roth et al.,
2012; Tohmi et al., 2014; Van Den Bergh et al., 2010) might result from suppressed
thalamocortical synaptic transmission (Reinhold et al., 2015). Another factor that should be taken
into consideration is the differential neuronal populations sampled in these two studies. In
Marshel's study (Marshel et al., 2011), excitatory and inhibitory neurons were ubiquitously labeled
with the synthetic calcium indicator (Oregon Green Bapta-1 AM); whereas here we sampled from
a subset of excitatory neurons in Thy1-GCaMP6s transgenic mice (Dana et al., 2014). As V1
interneurons prefer lower spatial frequencies than layer 2/3 excitatory neurons (Niell and Stryker,
2008), the population preference would shift towards lower frequencies with the inclusion of
interneurons. However, it is unclear whether such a difference between excitatory and inhibitory
neurons also exist in higher visual areas, and consequently leads to the lower preferred spatial
frequencies as a population. Further comparisons of neuronal response properties in wakefulness
and anesthesia (Adesnik, 2017; Durand et al., 2016; Greenberg et al., 2008; Haider et al., 2013)
can explain the influence of different brain states on the neural representation and transformation
among visual cortical areas. Nevertheless, studying neuronal physiology in the awake brain will
be of great value for understanding neural computations for perception and behavior.

Selectivity for stimulus anisotropy
We also found cortical neurons to be highly selective for stimulus anisotropy, suggesting diverse
spatial integration properties. Many neurons show linear spatial summation (or 'length
summation', (Schumer and Movshon, 1984)), such that their responses increase as the oriented
textures elongate. Meanwhile, many neurons prefer short stimuli, reminiscent of 'end-stopping'
cells, which preferentially respond to stimuli of certain lengths (Hubel and Wiesel, 1965; Gilbert,
1977). This length tuning property may be attributed to 'surround suppression', where one
receptive field is inhibited by the stimulation at the surround. Diverse length tuning curves may
result from differential ratios of the excitation on classical receptive fields and the inhibitory effect
of the receptive field surround (Adesnik, 2017; Adesnik et al., 2012; Vaiceliunaite et al., 2013).

We found diverse spatial integration properties distributed across the cortical areas with area-
specific biases, which might participate in distinct neural computations. V1 and LM show relatively
strong biases for non-oriented stimuli, suggesting strong surround suppression that is also
observed in the primate (El-Shamayleh et al., 2013; Hubel and Livingstone, 1987; Shushruth et
al., 2009), the cat (DeAngelis et al., 1994) and the mouse (Adesnik, 2017; Adesnik et al., 2012;

Murgas et al., 2020; Nienborg et al., 2013; Vaiceliunaite et al., 2013; Van Den Bergh et al., 2010).

Surround suppression is involved in a number of neural computations, including shape

processing. In the primate, 'end-stopping' behavior increases along the ventral processing stream

and is suggested to benefit the coding for curvatures (Ponce et al., 2017). Recent studies also

revealed the lateral areas in the rat also show less sharpness of the orientation tuning and

insensitivity to the phase of the drifting gratings, which might contribute to the invariant

representations of visual shapes of different scales and shapes in these lateral areas (Matteucci

et al., 2019; Tafazoli et al., 2017; Vermaercke et al., 2014). These observations agree to the

strong preferences for non-oriented stimuli in the lateral area LI observed in this study. Future

studies characterizing the role of nonlinear spatial integration properties in building invariant

representations of visual shapes will benefit the understanding of computational mechanisms of

object processing.

Functional organization of mouse higher visual areas

Our findings revealed a clear asymmetry in representation of the spatiotemporal features in the

mouse visual cortex which may indicate a functional organization that is adapted to encode

ethologically relevant visual information. We uncovered a graded representation of fast to slow

motion along the anterior-posterior axis in the mouse higher visual cortex. Anterior areas,

especially areas RL and A, mainly represent the lower visual field (Zhuang et al., 2017). The

preferences for large-scale, fast motion of these areas might benefit the encoding for fast optic

flows near the ground during animal's exploratory behaviors. Supporting this theory, a recent

study found area RL is specialized for binocular disparities corresponding to nearby visual stimuli

(La Chioma et al., 2019) and might contain a visuo-tactile map of near space (Olcese et al., 2013).

In contrast, area PM mainly represents the visual periphery, which is dominated by strong but

relatively slow visual flows during animals' navigation. In an additional set of experiment (data not

shown), using isotropic noise stimuli with global motions, we found PM neurons, with the similar

preference for slow speed, respond more robustly to coherent, global motion than to incoherent,

local motion, suggesting the specialization for encoding slow optic flows in the visual periphery

during navigation. Area PM might relay information about optic flows via the dense, direct efferent

projections to the retrosplenial cortex (Wang et al., 2012), which preferentially responds to slow

motion (Murakami et al., 2015) and is involved in spatial navigation (Mao et al., 2017). Moreover,

area-specific encodings of behavioral variables and the distinct projections of dorsal areas

suggest their differential involvements in visuomotor behaviors (Minderer et al., 2019; Wang et

al., 2012; Wang and Burkhalter, 2013). Altogether, these data suggest the dorsal higher visual

areas of mouse are specialized for ethological encoding of visual motion, reminiscent of the dorsal
streams in the primate visual system (Van Essen and Maunsell, 1983).

The ventral processing stream has been found a conserved organization across many species,
including primates, cats and rats (Matteucci et al., 2019; Tafazoli et al., 2017; Van Essen and
Maunsell, 1983; Vermaercke et al., 2014). Here we found functional evidence suggesting it also
exists in the mouse lateral cortex. Lateral area LI shows high spatial acuity and 'end-stopping'
properties. These properties are suitable for encoding spatial details and curvatures, which are
critical for object recognition (Lu et al., 2018; Ponce et al., 2017). However, in the mouse visual
system the presence of higher-order functions, such as transformation-tolerant representation of
visual objects (Tafazoli et al., 2017; Vermaercke et al., 2014), has not yet been tested. Other
lateral areas, including postrhinal area (POR) and posterior area, were hinted to have similar
preferences for low temporal and high spatial frequencies (Murakami et al., 2017), consistent with
the anterior-posterior cortical gradient for fast-to-slow motion information. Recent studies suggest
POR as a hub region, associating visual objects with behaviorally relevant variables, such as
reward values and object location (Burgess et al., 2016; Furtak et al., 2012). How do these
properties coordinate with other visual properties, such as selectivity for moving objects (Beltramo
and Scanziani, 2019; Bennett et al., 2019) and for slow stimuli (Murakami et al., 2017), and the
behavioral implications of such a functional specialization remain to be determined. Strong
anatomical connections of these lateral areas to temporal and parahippocampal regions (Wang
et al., 2012), supports the idea that mouse lateral visual cortex might be the involved in shape
processing, although species-specific differences might also exist.

Based on the feedforward and feedback characteristics of the intracortical connectivity, (Felleman
and Van Essen, 1991), area LM is generally placed at the first stage of higher-order visual
processing (D'Souza et al., 2016; Harris et al., 2019). It receives strong feedforward inputs from
V1 (Glickfeld et al., 2013; Wang and Burkhalter, 2007) and in return provides strong feedback
projections, which might facilitate predictive coding of sensory information (Keller et al., 2020;
Nguyen et al., 2018). Area LM has been proposed to be the gateway for the ventral stream (AL
as the gateway for the dorsal stream), given its relatively denser projections to the ventral areas
(Wang et al., 2012, 2011) and population responses similar to the ventral areas (wide field imaging
in Murakami et al., 2017; Smith et al., 2017). However our data show that LM contains highly
diverse neurons that exhibit both dorsal and ventral properties, resembling the primate V2 (Van
Den Bergh et al., 2010). In addition, LM neurons send strong projections to both dorsal and ventral
areas (Wang et al., 2012, 2011), routing target-specific information (Glickfeld et al., 2013). These

functional and anatomical characteristics argue for LM's role as the divided gateway of dorsal and
ventral streams.

The functional specialization of visual cortical areas must depend on specific intracortical
connectivity. Previous anatomical studies revealed a nearly 'all-to-all' diagram of intracortical
connections in the primate and rodent visual cortex (Gămănuţ et al., 2018; Wang et al., 2012),
suggesting intensive crosstalk between functionally specialized areas. The information
transmitted through the connections, however, remain unclear. Individual projection neurons
either broadcast the same information across multiple areas, or selectively target a single area
(Han et al., 2018). The connectivity between higher visual areas can be functionally target-
specific, in the similar manner as the projections from lower to higher cortical areas (Glickfeld et
al., 2013; Kim et al., 2018). Therefore, neurons sharing similar response properties might
interconnect via long-range projections and form cortical subnetworks that spread the common
information. Here we uncovered the widespread distribution of shared response types across the
visual cortex, suggesting multiple distributed cortical subnetworks. The similarity of response type
compositions between areas might be predictive of the strength of anatomical connectivity
between visual cortical areas (Gămănuţ et al., 2018; Wang et al., 2012). Future studies combining
in vivo cellular physiology along with circuit mapping and manipulation should provide important
insights to the organization rules of connectivity and information flow between visual cortical
areas.

Materials and Methods

Animals and Surgery

599
All experiments were approved by the Animal Ethics Committee of KU Leuven. Standard
craniotomy surgeries were performed to gain optical access to the visual cortex (Goldey and
Andermann, 2014). C57BL/6J-Tg (Thy1-GCaMP6s)GP4.12Dkim/J mice (Dana et al. 2014; 5 male
and 5 female) between 2 and 3-month-old were anesthetized with isoflurane (2.5%–3% induction,
1%–1.25% surgery). A custom-made titanium frame was mounted to the skull, and a craniotomy
was made to expose the left visual cortex. The 5mm-wide craniotomy was covered by a set of
cover glasses. Buprenex and Cefazolin were administered postoperatively when the animal
recovered from anesthesia after surgery (2 mg/kg and 5 mg/kg respectively; every 12 hours for 2
days).

Behavior monitoring

All mice were habituated at least 3 days before data acquisition. During imaging experiments,
mice were head restrained on a treadmill, passively viewing the display directed to the right eye.
We recorded animals' right eye movement using a macro lens (Zoom 7000, Navitar) mounted to
a GigE camera (Prosilica GC660, Allied Vision Technologies), we tracked the running speed and
monitored the animals' general behavior (MAKO G-030b, Allied Vision Technologies). Mice
showed no consistent behavioral or gaze responses to the visual stimulation.

Visual Stimulation

Visual stimuli were generated in Matlab (The Mathworks) and delivered with customized software
written in python. Stimuli were spherically corrected and displayed on a gamma-corrected LCD
display (22'', Samsung 2233RZ). The screen was oriented parallel to the right eye of the animal
with 18 cm in distance (covering 80 deg in elevation by 105 deg in azimuth).

Spectral noise stimuli (Figure1D, 2A -- Figure supplement 1D) were created by applying a set of
parametrized filters to random pink-noise fields. Narrow bandpass filters (bandwidth: 1 octave)
for spatial and temporal frequencies determine the scale and flickering frequencies of the textures.
The orientation bandwidth determines the length/width ratio of spatial patterns, which we refer to
as stimulus anisotropy. Isotropic noise stimuli, which are spectrally band-passed noise with no
angular filter, resemble clouds of moving dots (Figure supplement 1D upper row; ISO stimuli).
Narrowing the orientation bandwidth leads to elongation of textures, yielding higher stimulus
anisotropy (Figure supplement 1D). While isotropic textures contain no global motion, oriented
textures were designed to slowly rotate (45 deg per second, clockwise) to activate neurons tuned
to different orientations.

This study uses three stimulus sets generated by combining the above stimulus dimensions. For
the spatiotemporal frequency assay, we used ISO and ANISO stimulus sets, which contained
non-oriented and highly elongated textures (orientation bandwidth: 5 deg) respectively (Figure
1D). Both stimulus sets comprised 30 combinations of spatial frequencies (0.02, 0.04, 0.08, 0.16,
0.32 cpd) and temporal frequencies (0.5, 1, 2, 4,8,16 Hz). For the stimulus anisotropy assay, the
stimulus set contains the combinations of 4 spatial frequencies (0.04, 0.08, 0.16, 0.32 cpd) and 4
orientation bandwidths (infinite, 40, 10, 5 deg FWHM), with a fixed temporal frequency at 2 Hz
(Figure 2A,B). Each stimulus condition was presented for 4 second, intertwined with 4-second
equiluminant gray screen. In each of the four pseudorandomized trials, a different seed was used
to generate a unique set of random pink noise, with varying phases yet constant frequency spectra
across trials.

Two-photon Calcium Imaging

A customized two-photon microscopy (Neurolabware) was used. The calcium indicator GCaMP6s was excited by a Ti:Sapphire excitation laser (MaiTai DeepSee, Spectra-Physics) operating at 920 nm. The green fluorescence of GCaMP6s was collected with a photomultiplier tube (PMT, Hamamatsu) through a bandpass filter (510/84 nm, Semrock). Images (720x512 pixel per frame) were collect at 31 Hz with a 16x objective (Nikon). Volume imaging was achieved using a focus tunable lens (EL-10-30-TC, Optotune; staircase mode). We recorded activity from neurons typically between 100 to 300 μm deep below pia.

Widefield Calcium Imaging

Widefield fluorescent images were acquired with a 2x objective (NA = 0.055, Mitutoyo, Edmund Optics). Blue LED light (479nm, ThorLabs) was used for illumination passing an excitation filter (469/35 nm Thorlabs) and a dichroic filter centered at 498 nm. The green fluorescence was collected with an EMCCD camera (EM-C^2, QImaging) via a bandpass filter (525/39 nm filter, Thorlabs). The image acquisition was controlled with customized software written in python.

Retinotopic mapping

To identify visual cortical areas *in vivo*, we performed retinotopic mapping with 1-photon wide field imaging using dynamic noise stimuli partially visible through a circling round aperture ("circling patch") or a translating rectangular aperture ("traveling bar"). The circling patch stimuli is a small patch of contrast (20 deg in diameter) circling along an elliptic trajectory on the display (azimuth: -40 to 40 deg; elevation: -30 to 30 deg; 20 sec per cycle with 20 repetitions; Figure supplement 1 A). The traveling bar stimuli comprised a narrow bar (13 deg wide) sweeping across the screen in 4 cardinal directions (10 sec per sweep with 20 repetitions). An isotropic noise background (0.08 cpd, 2 Hz) was embedded in the patch or bars. The phase maps were generated by calculating the phase/position of the stimuli evoking maximal neural responses for each image pixel. Each area has a complete representation of the elliptic trajectory of the circling patch, resulting pinwheel-like retinotopic maps (Figure supplement 1B). The traveling bar stimuli generated azimuth and altitude phase maps, which were used to generate sign maps with the methods described in the previous study (Figure supplement 1C) (Juavinett et al., 2017). Areas are identified according to the common delineation (Wang and Burkhalter, 2007; Zhuang et al., 2017). The boarders between areas were further refined with 2-photon imaging of cellular responses to retinotopic stimuli to assign neurons into different areas.

Data Analysis

674 All subsequent data analysis was performed in MATLAB (The Mathworks, Natick, MA).

Selection for visual cells

For the cellular imaging, raw images were reconstructed and corrected for brain motion artefacts
using custom MATLAB routines (Bonin et al., 2011). Regions of interest (ROI) were selected at
somatic regions showing significant calcium activities during visual stimulation. Cellular
fluorescence time courses were generated by averaging all pixels in a ROI, followed by
subtracting the neuropil signals in the surround shell. Responses were defined as the averaged
$\Delta F/F$ during the stimulus epoch, where ΔF is the change in the fluorescent signal and F is the
baseline fluorescence (1s before stimulation). Stimulus-evoked responses were identified when
median cross-trial response curves showed response magnitude higher than 3x standard
deviation of the baseline activity for over one second. A cell was classified as responsive if it
shows stimulus-evoked responses to at least one stimulus condition. Response reliability r was
measured as the 75 percentiles of the correlation coefficients between stimulus-evoked
fluorescence traces across trials. Neurons above the threshold of 0.3 were deemed reliable.

Tuning curves

689 The anisotropy preference index (API) is expressed as $API = \frac{R(aniso)-R(iso)}{R(aniso)+R(iso)}$, where R(aniso) and
690 R(iso) are individual neurons' peak response amplitudes to ANISO and ISO stimuli respectively.

691 For the spatial and temporal frequency assay, responses were fit to two-dimensional elliptical
692 Gaussian models (Priebe et al. 2003; Andermann et al. 2011):

$$R(sf,tf) = A \exp\left(\frac{-(\log_2 sf - \log_2 sf_0)^2}{2(\sigma_{sf})^2}\right) \exp\left(\frac{-(\log_2 tf - \log_2 tf_p(sf))^2}{2(\sigma_{tf})^2}\right),$$

where $\log_2 tf_p(sf) = \xi(\log_2 sf - \log_2 sf_0) + \log_2 tf_0$, and A is the peak response amplitude, sf_0 and tf_0 are the
preferred spatial and temporal frequencies, and σ_{sf} and σ_{tf} are the spatial and temporal frequency
tuning widths. The dependence of temporal frequency tunings on the spatial frequency is captured
by a power-law exponent ξ.

Neurons are categorized as lowpass, bandpass and highpass based on whether the responses
to the minimal and maximal tested frequencies are greater than half maximal responses. Lowpass
cells tend to respond to frequencies lower than the tested spectrum; while highpass cells are likely
to respond to higher frequencies. Bandpass neurons respond mainly within the tested spectrum.

Estimates of cutoffs for the spatial and temporal frequency were measured at cross-sections at
$R(sf, tf_0)$ and $R(sf_0, tf)$, respectively. We estimated low cutoffs from bandpass and highpass cells,
high cutoffs from bandpass and lowpass cells, and bandwidths from bandpass cells.

Population coding analysis

To examine the information encoded in individual visual cortical areas, we decoded stimulus
categories from neuronal population activities using linear classifiers (Hung et al., 2005;
Vermaercke et al., 2014). Support vector machines were trained and tested in pair-wise
classification for all possible pairs (30 stimulus conditions result in 435 unique pairs). To compare
the decoding performance with the result of cell-by-cell speed tuning analysis, we focused on the
neurons responded to ISO stimuli. Visual responses across four trials were split into training and
testing groups (half-half) and the performance was measured as the proportion of correct
classification decisions to the testing groups (standard cross validation). The decoding analysis
were repeated for 100 times with random sampling. To test the scaling of decoding performance
as a function of population size, we measured the decoding performances as a function of
numbers of sampled neurons (logarithmic increase from 1 to 1000; without replacement) per area.
To compare the discriminability for specific stimulus pairs, we measured the number of neurons
required to reach the classification accuracy at 90% by interpolate the growth curves of
performances as a function of population size. To assess the differences in the overall and feature
specific decoding performance between visual areas, we compared the decoding performances
between iso-speed stimulus pairs and between cross-speed pairs (Figure 6C – Figure supplement
4C). Iso-speed pairs were neighboring textures lying along the iso-speed lines. Cross-speed pairs
are neighboring textures lying orthogonally to iso-speed lines and have a 4-fold difference in
speed.

Clustering

To assess whether cortical neurons form functional cell types, we used a spectral clustering
approach. We randomly selected 2000 neurons with replacement from individual areas to avoid
bias. Each neuron is expressed as a vector of responses to ISO and ANISO stimuli of all
combinations of spatial and temporal frequencies (response amplitude normalized to range from
0 to 1). The response matrix of all selected neurons was analyzed using a spectral clustering
algorithm (Ingo, 2020). To determine whether there are discrete functional types, we plotted the
overall variances (within-cluster sum-of-square variances) as a function of number of clusters.
The resulting Scree plot showed an inflection point at 12 clusters (Figure supplement 5A),

suggesting 12 broad functional cell types. We measured the similarity between neurons within individual clusters and across clusters, based on the cosine distance between response vectors, and found high similarities within clusters and high differences between clusters. These clusters showed distinct response properties and differential distributions across areas (Figure supplement 6). To quantify the similarity/differences between visual areas in terms of the proportions of functional cell types, we calculated the average Euclidean distances between areas and expressed them in a dendrogram (Figure 6D). We ran the clustering algorithm, with a constrain of 12 clusters, with random sampled pools for 100 times. The resulted dendrograms were highly consistent (Figure supplement 5C).

Acknowledgements

We are grateful to members of the Bonin and Farrow laboratories for feedback, discussions and comments on the manuscript, Joao Couto for establishing the imaging setup and support with image processing, Karl Farrow and Mark Andermann for comments on an earlier version of the manuscript, and Asli Ayaz for editing and valuable suggestions for the manuscript writing. This work was supported by Neuro-Electronics Research Flanders (VB), Research Foundation Flanders (FWO) fellowship 12E4314N (BV), grant G0D0516N (VB), KU Leuven Research Council grant C14/16/048 (VB) and a personal fellowship award from Chinese Scholarship Council (XH).

Author Contributions

Conceptualization & Methodology, X.H., B.V. and V.B.; Investigation, X.H. and B.V.; Data Analysis, X.H., B.V, and V.B. ; Writing – Review & Editing, X.H., B.V. and V.B.; Funding Acquisition, V.B.; Resources, V.B.; Supervision, V.B.

Competing interests

The authors declare no competing interests.

References

Adesnik H. 2017. Synaptic Mechanisms of Feature Coding in the Visual Cortex of Awake Mice. *Neuron* **95**:1147-1159.e4. doi:10.1016/j.neuron.2017.08.014

Adesnik H, Bruns W, Taniguchi H, Huang ZJ, Scanziani M. 2012. A neural circuit for spatial summation in visual cortex. *Nature* **490**:226–31. doi:10.1038/nature11526

Akrami A, Kopec CD, Diamond ME, Brody CD. 2018. Posterior parietal cortex represents sensory history and mediates its effects on behaviour. *Nature* **554**:368–372. doi:10.1038/nature25510

Andermann ML, Kerlin AM, Roumis DK, Glickfeld LL, Reid RC. 2011. Functional specialization of mouse higher visual cortical areas. *Neuron* **72**:1025–39. doi:10.1016/j.neuron.2011.11.013

Baden T, Berens P, Franke K, Roman-Roson M, Bethge M, Euler, Baden, Tom Berens, Philipp Franke, Katrin Roman-Roson, Miroslav Bethge ME, Baden T, Berens P, Franke K, Roman-Roson M, Bethge M, Euler. 2016. The functional diversity of mouse retinal ganglion cells. *Nature* **529**:1–21. doi:10.1038/nature16468

Beltramo R, Scanziani M. 2019. A collicular visual cortex: Neocortical space for an ancient midbrain visual structure. *Science (80-)* **363**:64–69. doi:10.1126/science.aau7052

Bennett C, Gale SD, Garrett ME, Newton ML, Callaway EM, Murphy GJ, Olsen SR. 2019. Higher-Order Thalamic Circuits Channel Parallel Streams of Visual Information in Mice. *Neuron* **102**:477-492.e5. doi:10.1016/j.neuron.2019.02.010

Bonin V, Histed MH, Yurgenson S, Reid RC. 2011. Local diversity and fine-scale organization of receptive fields in mouse visual cortex. *J Neurosci* **31**:18506–21. doi:10.1523/JNEUROSCI.2974-11.2011

Burgess CR, Ramesh RN, Sugden AU, Levandowski KM, Minnig MA, Fenselau H, Lowell BB, Andermann ML. 2016. Hunger-Dependent Enhancement of Food Cue Responses in Mouse Postrhinal Cortex and Lateral Amygdala. *Neuron* **91**:1154–1169. doi:10.1016/j.neuron.2016.07.032

D'Souza RD, Meier AM, Bista P, Wang Q, Burkhalter A. 2016. Recruitment of inhibition and excitation across mouse visual cortex depends on the hierarchy of interconnecting areas. *Elife* **5**:e19332. doi:10.7554/eLife.19332

Dana H, Chen T-W, Hu A, Shields BC, Guo C, Looger LL, Kim DS, Svoboda K. 2014. Thy1-GCaMP6 Transgenic Mice for Neuronal Population Imaging In Vivo. *PLoS One* **9**:e108697. doi:10.1371/journal.pone.0108697

de Vries SEJ, Lecoq JA, Buice MA, Groblewski PA, Ocker GK, Oliver M, Feng D, Cain N, Ledochowitsch P, Millman D, Roll K, Garrett M, Keenan T, Kuan L, Mihalas S, Olsen S, Thompson C, Wakeman W, Waters J, Williams D, Barber C, Berbesque N, Blanchard B,

Bowles N, Caldejon SD, Casal L, Cho A, Cross S, Dang C, Dolbeare T, Edwards M, Galbraith
J, Gaudreault N, Gilbert TL, Griffin F, Hargrave P, Howard R, Huang L, Jewell S, Keller N,
Knoblich U, Larkin JD, Larsen R, Lau C, Lee E, Lee F, Leon A, Li L, Long F, Luviano J, Mace
K, Nguyen T, Perkins J, Robertson M, Seid S, Shea-Brown E, Shi J, Sjoquist N,
Slaughterbeck C, Sullivan D, Valenza R, White C, Williford A, Witten DM, Zhuang J, Zeng
H, Farrell C, Ng L, Bernard A, Phillips JW, Reid RC, Koch C. 2020. A large-scale
standardized physiological survey reveals functional organization of the mouse visual cortex.
Nat Neurosci **23**:138–151. doi:10.1038/s41593-019-0550-9

DeAngelis GC, Freeman RD, Ohzawa I. 1994. Length and width tuning of neurons in the cat's
primary visual cortex. *J Neurophysiol* **71**:347–374. doi:8158236

Dipoppa M, Ranson A, Krumin M, Pachitariu M, Carandini M, Harris KD. 2018. Vision and
Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. *Neuron*
98:602-615.e8. doi:10.1016/j.neuron.2018.03.037

Durand S, Iyer R, Mizuseki K, de Vries S, Mihalas S, Reid RC. 2016. A Comparison of Visual
Response Properties in the Lateral Geniculate Nucleus and Primary Visual Cortex of Awake
and Anesthetized Mice. *J Neurosci* **36**:12144–12156. doi:10.1523/JNEUROSCI.1741-
16.2016

El-Shamayleh Y, Kumbhani RD, Dhruv NT, Movshon JA. 2013. Visual response properties of V1
neurons projecting to V2 in macaque. *J Neurosci* **33**:16594–605.
doi:10.1523/JNEUROSCI.2753-13.2013

Felleman DJ, Van Essen DC. 1991. Distributed hierarchical processing in the primate cerebral
cortex. *Cereb Cortex* **1**:1–47.

Furtak SC, Ahmed OJ, Burwell RD. 2012. Single neuron activity and theta modulation in postrhinal
cortex during visual object discrimination. *Neuron* **76**:976–88.
doi:10.1016/j.neuron.2012.10.039

Gămănuţ R, Kennedy H, Toroczkai Z, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Burkhalter
A. 2018. The Mouse Cortical Connectome, Characterized by an Ultra-Dense Cortical Graph,
Maintains Specificity by Distinct Connectivity Profiles. *Neuron* **97**:698-715.e10.
doi:10.1016/j.neuron.2017.12.037

Garrett ME, Nauhaus I, Marshel JH, Callaway EM. 2014. Topography and Areal Organization of
Mouse Visual Cortex. *J Neurosci* **34**:12587–12600. doi:10.1523/JNEUROSCI.1124-14.2014

Gilbert CD. 1977. Laminar differences in receptive field properties of cells in cat primary visual
cortex. *J Physiol* **268**:391–421.

Glickfeld LL, Andermann ML, Bonin V, Reid RC. 2013. Cortico-cortical projections in mouse visual
cortex are functionally target specific. *Nat Neurosci* **16**:219–26. doi:10.1038/nn.3300

Goldey GJ, Andermann ML. 2014. Simultaneous Two-Photon Calcium Imaging of Entire Cortical
Columns 23–32.

Greenberg DS, Houweling AR, Kerr JNDD. 2008. Population imaging of ongoing neuronal activity
in the visual cortex of awake rats. *Nat Neurosci* **11**:749–751. doi:10.1038/nn.2140

Haider B, Häusser M, Carandini M. 2013. Inhibition dominates sensory responses in the awake
cortex. *Nature* **493**:97–102. doi:10.1038/nature11665

Han Y, Kebschull JM, Campbell RAA, Cowan D, Imhof F, Zador AM, Mrsic-Flogel TD. 2018. The
logic of single-cell projections from visual cortex. *Nature* **556**:51–56.
doi:10.1038/nature26159

Harris JA, Mihalas S, Hirokawa KE, Whitesell JD, Choi H, Bernard A, Bohn P, Caldejon S, Casal
L, Cho A, Feiner A, Feng D, Gaudreault N, Gerfen CR, Graddis N, Groblewski PA, Henry
AM, Ho A, Howard R, Knox JE, Kuan L, Kuang X, Lecoq J, Lesnar P, Li Y, Luviano J,
McConoughey S, Mortrud MT, Naeemi M, Ng L, Oh SW, Ouellette B, Shen E, Sorensen SA,
Wakeman W, Wang Q, Wang Y, Williford A, Phillips JW, Jones AR, Koch C, Zeng H. 2019.
Hierarchical organization of cortical and thalamic connectivity. *Nature* **575**:195–202.
doi:10.1038/s41586-019-1716-z

Hubel DH, Livingstone MS. 1987. Segregation of form, color, and stereopsis in primate area 18.
J Neurosci **7**:3378–3415.

Hung CP, Kreiman G, Poggio T, DiCarlo JJ. 2005. Fast readout of object identity from macaque
inferior temporal cortex. *Science (80-)* **310**:863–866. doi:10.1126/science.1117593

Ingo. 2020. Fast and efficient spectral clustering. *MATLAB Cent File Exch*.

Juavinett AL, Callaway EM. 2015. Pattern and Component Motion Responses in Mouse Visual
Cortical Areas. *Curr Biol* **25**:1–6. doi:10.1016/j.cub.2015.05.028

Juavinett AL, Nauhaus I, Garrett ME, Zhuang J, Callaway EM. 2017. Automated identification of
mouse visual areas with intrinsic signal imaging. *Nat Protoc* **12**:32–43.

doi:10.1038/nprot.2016.158

Keller AJ, Roth MM, Scanziani M. 2020. Feedback generates a second receptive field in neurons
of the visual cortex. *Nature* 1–5. doi:10.1038/s41586-020-2319-4

Kim EJ, Zhang Z, Huang L, Ito-Cole T, Jacobs MW, Juavinett AL, Senturk G, Hu M, Ku M, Ecker
JR, Callaway EM. 2020. Extraction of Distinct Neuronal Cell Types from within a Genetically
Continuous Population. *Neuron* 1–9. doi:10.1016/j.neuron.2020.04.018

Kim MH, Znamenskiy P, Iacaruso MF, Mrsic-Flogel TD. 2018. Segregated Subnetworks of
Intracortical Projection Neurons in Primary Visual Cortex. *Neuron* **100**:1313-1321.e6.
doi:10.1016/j.neuron.2018.10.023

La Chioma A, Bonhoeffer T, Hübener M. 2019. Area-Specific Mapping of Binocular Disparity
across Mouse Visual Cortex. *Curr Biol* **29**:2954-2960.e5. doi:10.1016/j.cub.2019.07.037

Lien AD, Scanziani M. 2013. Tuned thalamic excitation is amplified by visual cortical circuits. *Nat
Neurosci* **16**:1315–23. doi:10.1038/nn.3488

Lu Y, Yin J, Chen Z, Gong H, Liu Y, Qian L, Li X, Liu R, Andolina IM, Wang W. 2018. Revealing
Detail along the Visual Hierarchy: Neural Clustering Preserves Acuity from V1 to V4. *Neuron*
417–428. doi:10.1016/j.neuron.2018.03.009

Mao D, Kandler S, McNaughton BL, Bonin V. 2017. Sparse orthogonal population representation
of spatial context in the retrosplenial cortex. *Nat Commun* **8**. doi:10.1038/s41467-017-00180-
9

Marques T, Summers MT, Fioreze G, Fridman M, Dias RF, Feller MB, Petreanu E. 2018. A Role
for Mouse Primary Visual Cortex in Motion Perception. *Curr Biol* **28**:1703-1713.e6.
doi:10.1016/j.cub.2018.04.012

Marshel JH, Garrett ME, Nauhaus I, Callaway EM. 2011. Functional specialization of seven
mouse visual cortical areas. *Neuron* **72**:1040–1054. doi:10.1016/j.neuron.2011.12.004

Matteucci G, Marotti RB, Riggi M, Rosselli FB, Zoccolan D. 2019. Nonlinear processing of shape
information in rat lateral extrastriate cortex. *J Neurosci* **39**:1649–1670.
doi:10.1523/JNEUROSCI.1938-18.2018

Meijer GT, Mertens PEC, Pennartz CMA, Olcese U, Lansink CS. 2019. The circuit architecture of
cortical multisensory processing: Distinct functions jointly operating within a common

anatomical network. *Prog Neurobiol* **174**:1–15. doi:10.1016/j.pneurobio.2019.01.004

Minderer M, Brown KD, Harvey CD. 2019. The Spatial Structure of Neural Encoding in Mouse
Posterior   Cortex   during   Navigation.   *Neuron*   **102**:232-248.e11.
doi:10.1016/j.neuron.2019.01.029

Morcos AS, Harvey CD. 2017. Evidence Accumulation in Cortex **19**:1672–1681.
doi:10.1038/nn.4403.History-dependent

Murakami T, Matsui T, Ohki K. 2017. Functional Segregation and Development of Mouse Higher
Visual Areas. *J Neurosci* **37**:9424–9437. doi:10.1523/jneurosci.0731-17.2017

Murakami T, Yoshida T, Matsui T, Ohki K. 2015. Wide-field Ca2+ imaging reveals visually evoked
activity in the retrosplenial area. *Front Mol Neurosci* **08**:1–12. doi:10.3389/fnmol.2015.00020

Murgas KA, Wilson AM, Michael V, Glickfeld LL. 2020. Unique spatial integration in mouse
primary   visual   cortex   and   higher   visual   areas.   *J   Neurosci*   1997–19.
doi:10.1523/JNEUROSCI.1997-19.2020

Nassi JJ, Callaway EM. 2009. Parallel processing strategies of the primate visual system. *Nat*
Rev Neurosci **10**:360–72. doi:10.1038/nrn2619

Nguyen J, Marques T, Fioreze G, Petreanu L. 2018. The functional organization of cortical
feedback inputs to primary visual cortex. *Nat Neurosci*. doi:10.1038/s41593-018-0135-z

Niell CM, Stryker MP. 2008. Highly selective receptive fields in mouse visual cortex. *J Neurosci*
28:7520–36. doi:10.1523/JNEUROSCI.0623-08.2008

Nienborg H, Hasenstaub A, Nauhaus I, Taniguchi H, Huang ZJ, Callaway EM. 2013. Contrast
dependence and differential contributions from somatostatin- and parvalbumin-expressing
neurons   to   spatial   integration   in   mouse   V1.   *J   Neurosci*   **33**:11145–54.
doi:10.1523/JNEUROSCI.5320-12.2013

Olcese U, Iurilli G, Medini P. 2013. Cellular and synaptic architecture of multisensory integration
in the mouse neocortex. *Neuron* **79**:579–93. doi:10.1016/j.neuron.2013.06.010

Piscopo DM, El-Danaf RN, Huberman AD, Niell CM. 2013. Diverse Visual Features Encoded in
Mouse   Lateral   Geniculate   Nucleus.   *J   Neurosci*   **33**:4642–4656.
doi:10.1523/JNEUROSCI.5187-12.2013

Ponce CR, Hartmann TS, Livingstone MS, Ponce XCR, Hartmann XTS, Livingstone XMS. 2017.

End-Stopping Predicts Curvature Tuning along the Ventral Stream, The Journal of Neuroscience. doi:10.1523/JNEUROSCI.2507-16.2017

Reinhold K, Lien AD, Scanziani M. 2015. Distinct recurrent versus afferent dynamics in cortical visual processing. *Nat Neurosci* 1–15. doi:10.1038/nn.4153

Román Rosón M, Bauer Y, Kotkat AH, Berens P, Euler T, Busse L. 2019. Mouse dLGN Receives Functional Input from a Diverse Population of Retinal Ganglion Cells with Limited Convergence. *Neuron* **102**:462-476.e8. doi:10.1016/j.neuron.2019.01.040

Roth MM, Helmchen F, Kampa BM. 2012. Distinct functional properties of primary and posteromedial visual area of mouse neocortex. *J Neurosci* **32**:9716–26. doi:10.1523/JNEUROSCI.0110-12.2012

Schumer RA, Movshon MSL a. JA. 1984. Length summation in simple cells of cat striate cortex. *Vis Res* **24**:565–571.

Shushruth S, Ichida JM, Levitt JB, Angelucci A. 2009. Comparison of Spatial Summation Properties of Neurons in Macaque V1 and V2. *J Neurophysiol* **102**:2069–2083. doi:10.1152/jn.00512.2009

Smith IT, Townsend LB, Huh R, Zhu H, Smith SL. 2017. Stream-dependent development of higher visual cortical areas. *Nat Neurosci* **20**:200–208. doi:10.1038/nn.4469

Steinmetz NA, Zatka-Haas P, Carandini M, Harris KD. 2019. Distributed coding of choice, action and engagement across the mouse brain. *Nature* 1–8. doi:10.1038/s41586-019-1787-x

Suzuki M, Larkum ME. 2020. General Anesthesia Decouples Cortical Pyramidal Neurons. *Cell* **180**:666-676.e13. doi:10.1016/j.cell.2020.01.024

Tafazoli S, Safaai H, De Franceschi G, Rosselli FB, Vanzella W, Riggi M, Buffolo F, Panzeri S, Zoccolan D. 2017. Emergence of transformation-tolerant representations of visual objects in rat lateral extrastriate cortex. *Elife* **6**:1–39. doi:10.7554/eLife.22794

Tohmi M, Meguro R, Tsukano H, Hishida R, Shibuki K. 2014. The Extrageniculate Visual Pathway Generates Distinct Response Properties in the Higher Visual Areas of Mice. *Curr Biol* 1–11. doi:10.1016/j.cub.2014.01.061

Vaiceliunaite A, Erisken S, Franzen F, Katzner S, Busse L. 2013. Spatial integration in mouse primary visual cortex. *J Neurophysiol* **110**:964–72. doi:10.1152/jn.00138.2013

Van Den Bergh G, Zhang B, Arckens L, Chino YM. 2010. Receptive-field properties of V1 and V2
neurons in mice and macaque monkeys. *J Comp Neurol* **518**:2051–70.
doi:10.1002/cne.22321

Van Essen DC, Maunsell JHR. 1983. Hierarchical organization and functional streams in the
visual cortex. *Trends Neurosci* **6**:370–375. doi:10.1016/0166-2236(83)90167-4

Vermaercke B, Gerich FJ, Ytebrouck E, Arckens L, Op de Beeck HP, Van den Bergh G. 2014.
Functional specialization in rat occipital and temporal visual cortex. *J Neurophysiol*
jn.00737.2013. doi:10.1152/jn.00737.2013

Wang Q, Burkhalter A. 2013. Stream-Related Preferences of Inputs to the Superior Colliculus
from Areas of Dorsal and Ventral Streams of Mouse Visual Cortex. *J Neurosci* **33**:1696–
1705. doi:10.1523/JNEUROSCI.3067-12.2013

Wang Q, Burkhalter A. 2007. Area Map of Mouse Visual Cortex. *Jounal Comp Neurol* **505**:337–
351. doi:10.1002/cne

Wang Q, Gao E, Burkhalter A. 2011. Gateways of ventral and dorsal streams in mouse visual
cortex. *Sci Signal* **31**:1905–1918. doi:10.1523/JNEUROSCI.3488-10.2011

Wang Q, Sporns O, Burkhalter A. 2012. Network analysis of corticocortical connections reveals
ventral and dorsal processing streams in mouse visual cortex. *J Neurosci* **32**:4386–4399.
doi:10.1523/JNEUROSCI.6063-11.2012

Wekselblatt JB, Flister ED, Piscopo DM, Niell CM. 2016. Large-scale imaging of cortical dynamics
during sensory perception and behavior. *J Neurophysiol* jn.01056.2015.
doi:10.1152/jn.01056.2015

Zhuang J, Ng L, Williams D, Valley M, Li Y, Garrett M, Waters J. 2017. An extended retinotopic
map of mouse cortex. *Elife* **6**:1–29. doi:10.7554/eLife.18372

Area	# Mice	# Cells	Responsive (% all cells) Dataset 1-3	Reliable responses (% all cells)					SFTF good fits		
				Dataset 1 (ISO)	Dataset 2 (ANISO)	Dataset 3 (Elongation)	Dataset 1&2	Dataset 1-3	Dataset 1	Dataset 2	Dataset 1&2
V1	7	15658	12515(80)	5736(37)	5476(35)	7309(47)	7498(48)	9447(60)	5421	5031	7126
LM	8	22795	13516(59)	6069(27)	5924(26)	6939(30)	8395(37)	9472(42)	5661	5222	7895
AL	6	9694	4936(51)	1821(19)	2308(24)	2023(21)	2849(29)	3158(33)	1732	2111	2705
RL	6	6792	4261(63)	1548(23)	1812(27)	1985(29)	2360(35)	2800(41)	1471	1693	2252
AM	6	7399	3625(49)	1197(16)	1353(18)	1574(21)	1755(24)	2105(28)	1178	1312	1724
PM	6	9275	3623(39)	940(10)	1100(12)	1200(13)	1505(16)	1802(19)	927	1079	1482
LI	7	7317	3693(50)	1422(19)	1113(15)	1516(21)	1905(26)	2269(31)	1320	1027	1802
Total	10	78930	46169(58)	18733(24)	19086(24)	22546(29)	26267(33)	31053(39)	17710	17475	24986

Table 1. Population for data analysis. Table summarizing the number of mice and cells imagined for each area, responsive cells (response amplitude > 3sd of baseline ΔF/F0 for over 1 sec for at least one stimulus condition), cells showing reliable responses (75th percentile of cross-trial correlation coefficient > 0.3) to individual and combinations of datasets, cells with good fits of spatiotemporal responses.