Hidden 'risk' in polygenic scores: clinical use today could exacerbate health

disparities

Alicia R. Martin[1,2,3], Masahiro Kanai[1,2,3,4,5], Yoichiro Kamatani[5,6], Yukinori Okada[5,7,8],

Benjamin M. Neale[1,2,3], Mark J. Daly[1,2,3,9]

[1] Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston,

MA 02114, USA

[2] Program in Medical and Population Genetics, Broad Institute of Harvard and MIT,

Cambridge, MA 02142, USA

[3] Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT,

Cambridge, MA 02142, USA

[4] Department of Biomedical Informatics, Harvard Medical School, Boston, MA 02115,

USA

[5] Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences,

Yokohama 230-0045, Japan

[6] Kyoto-McGill International Collaborative School in Genomic Medicine, Graduate

School of Medicine, Kyoto University, Kyoto 606-8507, Japan

[7] Department of Statistical Genetics, Osaka University Graduate School of Medicine,

Suita 565-0871, Japan

[8] Laboratory of Statistical Immunology, Immunology Frontier Research Center (WPI-

IFReC), Osaka University, Suita 565-0871, Japan

[9] Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki,

Finland

Abstract

Polygenic risk scores have the potential to improve health outcomes for a variety of

complex diseases and are poised for clinical translation, driven by the low cost of

genotyping (<$50 per person), the ability to predict genetic risk of many diseases with a

single test, and the dramatically increasing scale and power of genetic studies that aid

prediction accuracy. However, the major *ethical* and *scientific* challenge surrounding

clinical implementation is the observation that they are currently of far greater predictive

value in individuals of recent European ancestry than others. The better performance of

such risk scores in European populations is an inescapable consequence of the heavily

biased makeup of genome-wide association studies, with an estimated 79% of

participants in all existing genetic studies being of European descent. Empirically,

polygenic risk scores perform far better in European populations, with prediction

accuracy reduced by approximately 2- to 5-fold in East Asian and African American

populations, respectively. This highlights that—unlike specific clinical biomarkers and

prescription drugs, which may individually work better in some populations but do not

ubiquitously perform far better in European populations–clinical uses of prediction today

would systematically afford greater improvement to European populations. Early

diversifying efforts, however, show promise in levelling this vast imbalance, even when

non-European sample sizes are considerably smaller than the best-powered studies to

date. Polygenic risk scores provide a new opportunity to improve health outcomes for

many diseases in all populations, but to realize this full potential equitably, we must

prioritize greater inclusivity of diverse study participants in genetic studies and open

access to resulting summary statistics to ensure that health disparities are not increased

for those already most underserved.

Keywords: health disparities, genetic risk prediction, polygenic risk scores, diversity,

population genetics, statistical genetics

Polygenic risk scores (PRS), which predict traits using genetic data, are of burgeoning

interest to the clinical community as researchers demonstrate their growing power to

improve clinical care, genetic studies of a wide range of phenotypes increase in size

and power, and genotyping costs plummet to less than US$50. Many earlier criticisms

of limited prediction power are now recognized to have been chiefly an issue of small

sample size, which is no longer the case for many outcomes [1]. For example, integrated

models of PRS together with other lifestyle and clinical factors have enabled clinicians

to more accurately quantify the risk of heart attack for patients; consequently, they have

more effectively targeted the reduction of LDL cholesterol and by extension heart attack

by prescribing statins to patients at the greatest overall risk of cardiovascular disease [2-6].

While we share enthusiasm about the potential of PRS to improve health outcomes

through their eventual routine implementation as clinical biomarkers, we consider the

consistent observation that they are currently of far greater predictive value in

individuals of recent European descent than in others to be the major *ethical* and

scientific challenge surrounding clinical translation and, at present, the most critical

limitation to genetics in precision medicine. The scientific basis of this imbalance has

been demonstrated in population genetics simulations, theoretically, and empirically

across many traits and diseases [7-18].

All studies to date using well-powered genome-wide association studies (GWAS) to

assess the predictive value of PRS in European and non-European descent populations

have made a consistent observation: PRS predict individual risk far better in Europeans

than non-Europeans. In complex traits including height, body mass index (BMI),

educational attainment, schizophrenia, and major depression, existing PRS computed

with the largest available GWAS results predict outcomes far more accurately in new

samples of European-descent than they do in non-Europeans, with the clearest study

examples in East Asians and African Americans [11,12,14-20]. Rather than chance or

biology, this is a predictable consequence of the fact that the genetic discovery efforts to

date heavily overrepresent European populations globally. The correlation between true

and genetically predicted phenotypes decays with genetic divergence from the makeup

of the discovery GWAS, meaning that the accuracy of polygenic scores in different

populations is highly dependent on the study population representation in the largest

existing 'training' GWAS. Here, we document study biases that underrepresent non-

European populations in current GWAS, and explain the fundamental concepts

contributing to reduced phenotypic variance explained with increasing genetic

divergence from populations included in GWAS.

Predictable basis of disparities in polygenic risk score accuracy

Predictable basis of disparities in polygenic risk score accuracy

The lack of generalizability of genetic studies across global populations arises from the

overwhelming abundance of European descent studies—according to the GWAS

catalog [21-24], ~79% of all GWAS participants are of European descent despite making

up only 16% of the global population (**Figure 1**). More concerningly, the fraction of non-

European individuals in GWAS has stagnated or declined since late 2014 (**Figure 1**),

suggesting that we are not on a trajectory to correct this imbalance. These numbers

provide a composite metric of study availability, accessibility, and use—i.e., cohorts that

have been included in numerous studies are represented multiple times, which may

disproportionately include cohorts of European descent. The relative sample

compositions of GWAS result in highly predictable disparities in prediction accuracy;

statistical and population genetics theory predicts that genetic risk prediction accuracy

will decay with increasing genetic divergence between the original GWAS sample and

target of prediction, a function of population history [9,10]. This pattern can be attributed to

several statistical observations which we detail below: 1) GWAS favor the discovery of

genetic variants that are common in the study population; 2) linkage disequilibrium (LD)

differentiates marginal effect size estimates for highly polygenic traits across

populations, even when causal variants are the same; and 3) demographic and

environmental differences may drive differential forces of natural selection that in turn

drive differences in causal genetic architecture. Of note, the first two of these degrade

prediction performance across populations substantially even when there exist no

biological, environmental, or diagnostic differences.

Common discoveries and low-hanging fruit

First, the power to discover an association in a genetic study depends on the effect size

and frequency of the variant [25]. This power dependence means that the most significant

associations tend to be more common in the populations in which they are discovered

than in other populations [9,26]. For example, GWAS catalog variants are on average

more common in European populations compared to East Asian and African

populations (**Figure 2B**), an observation not representative of genomic variants at large.

Understudied populations offer low-hanging fruit for genetic discovery because variants

that are common in these groups but rare or absent in European populations could not

be discovered even with very large European sample sizes. Some examples include

SLC16A11 and *HNF1A* associations with type II diabetes in Latino populations, *APOL1*

associations with end-stage kidney disease, and associations with prostate cancer in

African descent populations [27-30]. If we assume that causal genetic variants have an

equal effect across all populations—an assumption with some empirical support that

offers the best case scenario for transferability [31-35]—Eurocentric GWAS biases mean

that variants that are common in European populations are preferentially discovered

and associated with risk, and thus account for a larger fraction of the variance in

polygenic risk [9]. Furthermore, imputation reference panels share the same biases as in

GWAS, and imputing sites that are common in European populations but rarer in other

populations is challenging when the catalog of non-European haplotypes is substantially

smaller. These issues are insurmountable through statistical methods alone, but rather

motivate substantial investments in more diverse populations to produce similar-sized

GWAS of biomedical phenotypes as well as sequenced reference panels in other

populations.

Linkage disequilibrium

Second, the correlation structure of the human genome, i.e. LD, varies across

populations due to demographic history (**Figure 2A,C-E**). These LD differences in turn

drive differences in effect size estimates (i.e. predictors) from GWAS across

populations, even when causal effects are the same. (Mathematically, the marginal

GWAS estimate $\hat{\beta}_j = \sum_{k=1}^{m} r_{j,k} \beta_k + \epsilon_j$, where $\hat{\beta}_j$ are effect size estimates at SNP *j*, $r_{j,k}$ is

pairwise SNP LD between SNPs *j* and *k*, β_k is the causal SNP effect at nearby SNP *k*,

and ϵ is residual error from bias or noise). While differences in effect size estimates due

to LD differences may typically be small for most regions of the genome, PRS sum

across these effects, also aggregating these population differences. Statistical methods

that account for LD differences across populations may help improve risk prediction

accuracy within each population. While empirical studies suggest that causal effect

sizes tend to be shared [31,32], it may not be feasible to fine-map most variants to a single

locus to solve issues of low generalizability, even with very large GWAS (i.e., millions).

This is because complex traits are highly polygenic, meaning most of our prediction

power comes from small effects that do not meet genome-wide significance and/or

cannot be fine-mapped, even in the best-powered GWAS to date [36].

History, selection, the environment, and complex interactions

Lastly, other environmental, demographic, and cohort considerations may further

worsen prediction accuracy differences across populations in less predictable ways.

GWAS ancestry study biases and LD differences across populations are extremely

challenging to address, but these issues actually make many favorable assumptions

that all causal loci have the same impact and are under equivalent selective pressure in

all populations. In contrast, other effects on polygenic adaptation or risk scores such as

natural selection can impact populations differently based on their unique histories.

Additionally, residual uncorrected population stratification may impact risk prediction

accuracy across populations, but the magnitude of its effect is currently unclear. These

effects are particularly challenging to disentangle, as has clearly been demonstrated for

height, where evidence of polygenic adaptation is under question [37,38]. Comparisons of

geographically stratified phenotypes like height across populations with highly divergent

genetic backgrounds and mean environmental differences, such as differences in

resource abundance during development across continents, are especially prone to

uninterpretable results [39]. Related to stratification, most polygenic scoring methods do

not explicitly address recent admixture and none consider recently admixed individuals'

unique local mosaic of ancestry—further methods development in this space is needed.

Furthermore, comparing PRS across environmentally stratified cohorts, such as in some

biobanks with healthy volunteer effects versus disease study datasets or hospital-based

cohorts, requires careful consideration of technical differences, collider bias, as well as

variability in baseline health status among studies. It is also important to consider

differences in clinical definition of the phenotypes and heterogeneous constitution of

sub-phenotypes among countries.

Differences in environmental exposure, gene × environment interactions, historical

population size dynamics, and other factors will further limit generalizability for genetic

risk scores in an unpredictable, trait-specific fashion [40,41]. While non-linear genetic

factors explain little variation in complex traits beyond a purely additive model [42], some

unrecognized nonlinearities and gene × gene interactions can also induce genetic risk

prediction challenges, as pairwise interactions are likely to vary more across

populations than individual SNPs. Mathematically, we can simplistically think of this in

terms of a two-SNP model, in which the sum of two SNP effects is likely to explain more

phenotypic variance than the product of the same SNPs. Some machine learning

approaches may thus modestly improve genetic prediction accuracy for some

phenotypes [43], but these approaches are most likely to improve prediction accuracy for

atypical traits with simpler architectures, known interactions, and poor prediction

generalizability across populations, such as skin pigmentation [44].

Limited generalizability of genetic prediction across diverse populations

Previous work has assessed prediction accuracy across diverse populations in several

traits and diseases for which GWAS summary statistics are available. These

assessments are becoming increasingly feasible with the growth and public availability

of global biobanks for quantitative traits as well as diversifying priorities from funding

agencies [45,46]. As of yet, multi-ethnic work has been slow in most disease areas [47],

limiting even the opportunity to assess prediction utility in non-European cohorts.

Nonetheless, we have assembled prediction accuracy statistics from several studies

using the largest European GWAS to predict several phenotypes in target European

and non-European cohorts. For example, multiple schizophrenia studies consistently

predicted risk on average 2.2-fold worse in East Asians relative to Europeans, (i.e.

μ=0.46, σ=0.06), using summary statistics from a Eurocentric GWAS [11,14] (**Figure 3**),

despite the fact that there is no genetic heterogeneity in schizophrenia between the two

populations. This finding is even more pronounced in African Americans, where genetic

divergence from Europeans is greater than between Europeans and East Asians [26].

Across several phenotypes with a range of genetic architectures in which empirical

evaluations were available, including BMI, educational attainment, height, and

schizophrenia, prediction accuracy using European GWAS summary statistics was on

average 4.5-fold less accurate in African Americans than in Europeans (i.e. μ=0.22,

σ=0.09, **Figure 3**) [11,12,15-18]. By extension, prediction accuracy is expected to be even

lower in African Americans with higher than average African ancestry or among

populations with greater divergence from Europeans (e.g. some southern African

populations). These enormous disparities are not simply methodological issues, as

various approaches (e.g. pruning and threshold versus LDPred) and accuracy metrics

(R^2 for quantitative traits and various pseudo-R^2 metrics for binary traits) illustrate this

consistently poorer performance in populations distinct from the discovery sample

across a range of polygenic traits (**Table S2**).

Prioritizing diversity shows early promise for polygenic prediction

Early diversifying GWAS efforts have been especially productive for informing on these

questions surrounding risk prediction. Rather than varying the prediction target dataset,

some GWAS in diverse populations have increased the scale of non-European

summary statistics and also varied the study dataset in multi-ethnic PRS studies. For

example, a BioBank Japan GWAS study (N=158,284) showed that compared to a 2×

larger European GWAS (N=322,154), the variance in BMI explained in an independent

Japanese cohort with Japanese GWAS summary statistics was on average 1.5-fold

greater than with European GWAS summary statistics (R^2=0.154 vs 0.104 at p < 0.05,

respectively) [19]. Similarly, a Chinese schizophrenia study (N=7,699 cases and 18,327

controls) showed that compared to an effectively 5-fold larger European GWAS

(N=36,989 cases, 113,075 controls), prediction accuracy in an independent Chinese

cohort with GWAS summary statistics from China far surpassed prediction accuracy

from European summary statistics by 2.6-fold (2.3% versus 6.2%) [20]. Thus, even when

studies in non-European populations are only a fraction the size of the largest European

study, they are likely to have disproportionate value for predicting polygenic traits in

other individuals of similar ancestry.

Given this background, we performed a systematic evaluation of polygenic prediction

accuracy across 17 quantitative anthropometric and blood panel traits in British and

Japanese individuals [19,48,49] by performing GWAS with the exact same sample sizes in

each population. We symmetrically demonstrate that prediction accuracy is consistently

higher with GWAS summary statistics from ancestry-matched summary statistics

(**Figure 4**). Keeping in mind issues of comparability described above, we note that the

BioBank Japan (BBJ) is a hospital-based cohort, whereas UK Biobank (UKBB) is a

healthier than average population-based cohort, and that differences in observed

heritability among these cohorts (rather than among populations) likely explain lower

prediction accuracy from the BBJ GWAS summary statistics (**Table S3**). Some

statistical fluctuations in the relative differences in prediction accuracy across

populations are likely driven by differences in trans-ethnic genetic correlation (i.e.

comparing across ancestries the estimated correlation of common variant effect sizes at

SNPs common in both populations via Popcorn) and/or differences in heritability

measured in each population (**Figure S1**, **Table S3**). Prediction accuracy was far lower

in individuals of African descent in the UK Biobank (**Figure S5**) using GWAS summary

statistics from European or Japanese ancestry individuals (**Figure 4**). These population

studies demonstrate the power and utility of increasingly diverse GWAS for prediction,

especially in populations of non-European descent.

While many other traits and diseases have been studied in multi-ethnic settings, few

have reported comparable metrics of prediction accuracy across populations.

Cardiovascular research, for example, has led the charge towards clinical translation of

PRS [1]. This enthusiasm is driven by observations that a polygenic burden of LDL-

increasing SNPs can confer monogenic-equivalent risk of cardiovascular disease, with

polygenic scores improving clinical models for risk assessment and statin prescription

that can reduce coronary heart disease and improve healthcare delivery efficiency [2,3,5].

However, many of these studies have been conducted exclusively in European descent

populations, with few studies rigorously evaluating population-level applicability to non-

Europeans. Those existing findings indeed demonstrate a large reduction in prediction

utility in non-European populations [7], though often with comparisons of odds ratios

among arbitrary breakpoints in the risk distribution that make comparisons across

studies challenging. To better clarify how polygenic prediction will be deployed in a

clinical setting with diverse populations, more systematic and thorough evaluations of

the utility of PRS within and across populations for many complex traits are still needed.

These evaluations would benefit from rigorous polygenic prediction accuracy

evaluations, especially for diverse non-European patients [50-52].

Translational genetic prediction may uniquely exacerbate disparities

Our impetus for raising these statistical issues limiting the generalizability of PRS across

population stems from our concern that, while they are legitimately clinically promising

for improving health outcomes for many biomedical phenotypes, they may have a larger

potential to raise health disparities than other clinical factors for several reasons. The

opportunities they provide for improving health outcomes means they inevitably will and

should be pursued in the near term, but we urge that a concerted prioritization to make

GWAS summary statistics easily accessible for diverse populations and a variety of

traits and diseases is imperative, even when they are a fraction the size of the largest

existing European datasets. *Individual* clinical tests, biomarkers, and prescription drug

efficacy may vary across populations in their utility, but are fundamentally informed by

the same underlying biology [53,54] Currently, guidelines state that as few as 120

individuals define reference intervals for clinical factors (though often smaller numbers

from only one subpopulation are used) and there is no clear definition of who is "normal"

[53]. Consequently, reference intervals for biomarkers can sometimes deviate

considerably by reported ethnicity [55-57]. Defining ethnicity-specific reference intervals is

clearly an important problem that can provide fundamental interpretability gains with

implications for some major health benefits (e.g. need for dialysis and development of

Type 2 diabetes based on ethnicity-specific serum creatinine and hemoglobin A1C

reference intervals, respectively) [56]. Simply put, some biomarkers or clinical tests scale

directly with health outcomes independent of ancestry, and many others may have

distributional differences by ancestry but are equally valid after centering with respect to

a readily collected population reference.

In contrast, PRS are uniformly less useful in understudied populations due to

differences in genomic variation and population history [9,10]. No analogous solution of

defining ethnicity-specific reference intervals would ameliorate health disparities

implications for PRS or fundamentally aid interpretability in non-European populations.

Rather, as we and others demonstrate, PRS are unique in that even with multi-ethnic

population references, these scores are fundamentally less informative in populations

more diverged from GWAS study cohorts.

The clinical use and deployment of genetic risk scores needs to be informed by the

issues surrounding tests that currently would unequivocally provide much greater

benefit to the subset of the world's population which is already on the positive end of

healthcare disparities[*]. Conversely, African descent populations, which already endure

many of the largest health disparities across the globe, are often predicted marginally

better, if at all, compared to random (**Figure 4**). They are therefore least likely to benefit

---

[*] To maximally benefit all populations, the largest existing GWAS results should be used. Downsampling European GWAS for the sake of parity results in worse predictors for all individuals.

from improvements in precision healthcare delivery from genetic risk scores with

existing data due to human population history and study biases. This is a major concern

globally and especially in the U.S., which already leads other middle- and high-income

countries in both real and perceived healthcare disparities [58]. Thus, we would strongly

urge that any discourse on clinical use of polygenic scores include a careful,

quantitative assessment of the economic and health disparities impacts on

underrepresented populations that might be unintentionally introduced by the use of

PRS and raise awareness about how to eliminate these disparities.

How do we even the ledger?

What can be done? An equal investment in GWAS across all major ancestries and

global populations is the most obvious solution to truly generate a substrate for equally

informative risk scores, but is not likely to occur any time soon absent a dramatic priority

shift given the current imbalance and stalled diversifying progress over the last five

years (**Figure 1**). While it may be challenging or in some cases infeasible to acquire

sample sizes large enough for PRS to be equally informative in all populations, some

much-needed efforts towards increasing diversity in genomics that support open sharing

of GWAS summary data from multiple ancestries are underway. Examples include the

All of Us Research Program, the Population Architecture using Genomics and

Epidemiology (PAGE) Consortium, as well as some disease-focused consortia, such as

the T2D-genes and Stanley Global initiatives on the genetics of type II diabetes and

psychiatric disorders, respectively. The prerequisite data for dramatically increasing

diversity also hypothetically exist in several large-scale publicly funded datasets such as

the Million Veterans Project and Trans-Omics for Precision Medicine (TOPMed), but

with problematic data access issues in which even summary data from GWAS within

and across populations are not publicly shared. While there is an understandable

patient privacy balance to strike when sharing individual-level data, GWAS summary

statistics by population from all publicly funded and as many privately funded projects

as possible should be made easily and publicly accessible to improve global health

outcomes. Efforts to unify phenotype definitions, normalization approaches, and GWAS

methods among studies are also encouraged.

To enable progress towards parity, it will be critical that open data sharing standards be

adopted for all ancestries and for genetic studies of all sample sizes, not just the largest

European results. Locally appropriate and secure genetic data sharing techniques as

well as equitable technology availability will need to be adopted widely in Asia and

Africa as they are in Europe and North America, to ensure that maximum value is

achieved from existing and ongoing efforts that are being developed to help counter the

current imbalance. Methodological improvements that better define risk scores by

accounting for population allele frequency, LD, and/or admixture differences

appropriately are underway and may help considerably, but will not by themselves bring

equality. All of these efforts are important and should be prioritized not just for risk

prediction but more generally to maximize the use and applicability of genetics to inform

on the biology of disease. Given the acute recent attention on clinical use of PRS, we

believe it is paramount to recognize their potential to improve health outcomes for all

individuals and many complex diseases. Simultaneously, we as a field must address the

disparity in utility in an ethically thoughtful and scientifically rigorous fashion, lest we

inadvertently enable genetic technologies to contribute to, rather than reduce, existing

health disparities.

Acknowledgments

We thank Amit Khera for helpful discussions. We also thank Michiaki Kubo, Yoshinori

Murakami, and Masato Akiyama for their support in the BioBank Japan Project analysis.

This work was supported by funding from the National Institutes of Health

(K99MH117229 to A.R.M.). UK Biobank analyses were conducted via application

31063. The BioBank Japan Project was supported by the Tailor-Made Medical

Treatment Program of the Ministry of Education, Culture, Sports, Science, and

Technology (MEXT) and the Japan Agency for Medical Research and Development

(AMED). M.K. was supported by a Nakajima Foundation Fellowship and the Masason

Foundation.

Figures

Figure 1 – Ancestry of GWAS participants over time compared to the global

population. Cumulative data as reported by the GWAS catalog [23]. A notable caveat is

that because some cohorts are included in numerous studies, some individuals are

represented multiple times. This bias in multiple counting is especially likely for publicly

available cohorts, which are more likely to be of European or East Asian descent.

Individuals whose ancestry is "not reported" are not shown.

Figure 2 – Demographic relationships, allele frequency differences, and local LD patterns between population pairs. Data analyzed from 1000 Genomes, in which population labels are: AFR = continental African, EUR = European, and EAS = East Asian. A) Cartoon relationships among AFR, EUR, and EAS populations. B) Allele frequency distributions in AFR, EUR, and EAS populations of variants from the GWAS catalog. C-E) Color axis shows LD scale (r^2). LD comparisons between pairs of populations show the same region of the genome for each comparison (representative region is chr1, 51572kb-52857kb) among pairs of SNPs polymorphic in both populations, illustrating that different SNPs are polymorphic across some population pairs, and that these SNPs have variable LD patterns across populations.

Figure 3 – Empirical comparison of phenotypic variance explained across

populations using polygenic scores computed with European GWAS. All GWAS

studies included here were conducted in European ancestry populations, with PRS

calculated and evaluated in independent European, East Asian, and African American

target cohorts. The European study biases result in the highest prediction accuracies in

independent European cohorts, followed by declining accuracy with increased genetic

divergence from Europe. A) Proportion of variance explained in each of the original

studies. B) Relative proportion of variance explained in each population with respect to

an independent European target population in each study. The diminished proportion of

variance explained in East Asian and African American populations relative to

Europeans is remarkably consistent despite differing genetic architectures, prediction

methods, and accuracy metrics due to similar population histories within these cohorts.

BMI = body mass index, EA = educational attainment, and SCZ = schizophrenia.

Figure 4 – Polygenic risk prediction accuracy in Japanese, British, and African descent individuals using independent GWAS of equal sample sizes in the BioBank Japan and UK Biobank. All target prediction cohorts are withheld from the GWAS and thus independent. Sample sizes in each GWAS are identical between BBJ and UKBB (**Table S1**). To optimize signal to noise, each point shows the maximum R^2 (i.e. best predictor) across 10 p-value thresholds. R^2 values for all p-value thresholds are shown in **Figures S2-S4**. Prediction accuracy tends to be higher in the UK Biobank, likely because observed heritability tends to be higher than in the BioBank Japan (**Table S3**). A) Genetic prediction accuracy for 17 anthropometric and blood panel traits in

Japanese individuals using summary statistics from GWAS of independent BioBank

Japan versus UK Biobank samples. B) Genetic prediction accuracy for the same 17

anthropometric and blood panel traits in independent British individuals using summary

statistics from GWAS of independent BioBank Japan versus the UK Biobank samples.

C) Genetic prediction accuracy for 17 anthropometric and blood panel traits in African

descent individuals in the UK Biobank using summary statistics from GWAS of

independent BioBank Japan versus UK Biobank samples.

References

1. Knowles, J. W. & Ashley, E. A. Cardiovascular disease: The rise of the genetic risk score. *PLoS Med* **15,** e1002546–7 (2018).

2. Khera, A. V. *et al.* Genome-wide polygenic score to identify a monogenic risk-equivalent for coronary disease. (2017). doi:10.1101/218388

3. Kullo, I. J. *et al.* Incorporating a Genetic Risk Score Into Coronary Heart Disease Risk Estimates: Effect on Low-Density Lipoprotein Cholesterol Levels (the MI-GENES Clinical Trial). *Circulation* **133,** 1181–1188 (2016).

4. Paquette, M. *et al.* Polygenic risk score predicts prevalence of cardiovascular disease in patients with familial hypercholesterolemia. *Journal of Clinical Lipidology* **11,** 725–732.e5 (2017).

5. Natarajan, P. *et al.* Polygenic Risk Score Identifies Subgroup With Higher Burden of Atherosclerosis and Greater Relative Benefit From Statin Therapy in the Primary Prevention SettingClinical Perspective. *Circulation* **135,** 2091–2101 (2017).

6. Tikkanen, E., Havulinna, A. S., Palotie, A., Salomaa, V. & Ripatti, S. Genetic risk prediction and a 2-stage risk screening strategy for coronary heart disease. *Arterioscler. Thromb. Vasc. Biol.* **33,** 2261–2266 (2013).

7. Khera, A. V. *et al.* Genetic Risk, Adherence to a Healthy Lifestyle, and Coronary Disease. *N Engl J Med* **375,** 2349–2358 (2016).

8. Khera, A. V. & Kathiresan, S. Genetics of coronary artery disease: discovery, biology and clinical translation. *Nat Rev Genet* **18,** 331–344 (2017).

9. Martin, A. R. *et al.* Human Demographic History Impacts Genetic Risk Prediction across Diverse Populations. *Am. J. Hum. Genet.* **100,** 635–649 (2017).

10. Scutari, M., Mackay, I. & Balding, D. Using Genetic Distance to Infer the Accuracy of Genomic Prediction. *PLoS Genet* **12,** e1006288 (2016).

11. Vilhjálmsson, B. J. *et al.* Modeling Linkage Disequilibrium Increases Accuracy of Polygenic Risk Scores. *The American Journal of Human Genetics* **97,** 576–592 (2015).

12. Ware, E. B. *et al.* Heterogeneity in polygenic scores for common human traits. 1–13 (2017). doi:10.1101/106062

13. Curtis, D. Polygenic risk score for schizophrenia is more strongly associated with ancestry than with schizophrenia. *bioRxiv* (2018). doi:https://doi.org/10.1101/287136

14. Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511,** 421–427 (2014).

15. Belsky, D. W. *et al.* Development and Evaluation of a Genetic Risk Score for Obesity. *Biodemography and Social Biology* **59,** 85–100 (2013).

16. Domingue, B. W., Belsky, D. W., Conley, D., Harris, K. M. & Boardman, J. D. Polygenic Influence on Educational Attainment. *AERA Open* **1,** 233285841559997–13 (2015).

17. Wedow, R. *et al.* Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet* **92,** 109 (2018).

18.   Vassos, E. *et al.* An Examination of Polygenic Score Risk Prediction in Individuals
With First-Episode Psychosis. *Biological Psychiatry* **81,** 470–477 (2017).
19.   Akiyama, M. *et al.* Genome-wide association study identifies 112 new loci for
body mass index in the Japanese population. *Nat Genet* **49,** 1458–1467 (2017).
20.   Li, Z. *et al.* Genome-wide association analysis identifies 30 new susceptibility loci
for schizophrenia. *Nature Publishing Group* **49,** 1576–1583 (2017).
21.   Need, A. C. & Goldstein, D. B. Next generation disparities in human genomics:
concerns and remedies. *Trends in Genetics* **25,** 489–494 (2009).
22.   Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature* **538,**
161–164 (2016).
23.   Morales, J. *et al.* A standardized framework for representation of ancestry data in
genomics studies, with application to the NHGRI-EBI GWAS Catalog. 1–10
(2018). doi:10.1186/s13059-018-1396-2
24.   Rosenberg, N. A. *et al.* Genome-wide association studies in diverse populations.
Nat Rev Genet **11,** 356–366 (2010).
25.   Sham, P. C., Cherny, S. S., Purcell, S. & Hewitt, J. K. Power of linkage versus
association analysis of quantitative traits, by use of variance-components models,
for sibship data. *The American Journal of Human Genetics* **66,** 1616–1630
(2000).
26.   1000 Genomes Project Consortium *et al.* A global reference for human genetic
variation. *Nature* **526,** 68–74 (2015).
27.   Consortium, T. S. 2. D. Sequence variants in SLC16A11 are a common risk factor
for type 2 diabetes in Mexico. *Nature* **506,** 97–101 (2014).
28.   Estrada, K. *et al.* Association of a Low-Frequency Variant in HNF1A With Type 2
Diabetes in a Latino Population. *JAMA* **311,** 2305–2314 (2014).
29.   Haiman, C. A. *et al.* Genome-wide association study of prostate cancer in men of
African ancestry identifies a susceptibility locus at 17q21. *Nat Genet* **43,** 570–573
(2011).
30.   Genovese, G. *et al.* Association of trypanolytic ApoL1 variants with kidney
disease in African Americans. *Science* **329,** 841–845 (2010).
31.   Liu, J. Z. *et al.* Association analyses identify 38 susceptibility loci for inflammatory
bowel disease and highlight shared genetic risk across populations. *Nat Genet*
47, 979–986 (2015).
32.   Carlson, C. S. *et al.* Generalization and Dilution of Association Results from
European GWAS in Populations of Non-European Ancestry: The PAGE Study.
PLoS Biol **11,** e1001661 (2013).
33.   Easton, D. F. *et al.* Genome-wide association study identifies novel breast cancer
susceptibility loci. *Nature* **447,** 1087–1093 (2007).
34.   Mahajan, A. *et al.* Genome-wide trans-ancestry meta-analysis provides insight
into the genetic architecture of type 2 diabetes susceptibility. *Nat Genet* **46,** 234–
244 (2014).
35.   Waters, K. M. *et al.* Consistent Association of Type 2 Diabetes Risk Variants
Found in Europeans in Diverse Racial and Ethnic Groups. *PLoS Genet* **6,**
e1001078 (2010).
36.   Huang, H. *et al.* Fine-mapping inflammatory bowel disease loci to single-variant
resolution. *Nature* **547,** 173–178 (2017).

37.   Berg, J. J. *et al.* Reduced signal for polygenic adaptation of height in UK Biobank.
1–44 (2018). doi:10.1101/354951
38.   Sohail, M. *et al.* Signals of polygenic adaptation on height have been
overestimated due to uncorrected population structure in genome-wide
association studies. 1–44 (2018). doi:10.1101/355057
39.   Duncan, L. *et al.* Analysis of Polygenic Score Usage and Performance across
Diverse Human Populations. (2018). doi:10.1101/398396
40.   Novembre, J. & Barton, N. H. Tread Lightly Interpreting Polygenic Tests of
Selection. *Genetics* **208,** 1351–1355 (2018).
41.   Henn, B. M., Botigué, L. R., Bustamante, C. D., Clark, A. G. & Gravel, S.
Estimating the mutation load in human genomes. *Nat Rev Genet* **16,** 333–343
(2015).
42.   Zhu, Z. *et al.* Dominance Genetic Variation Contributes Little to the Missing
Heritability for Human Complex Traits. *The American Journal of Human Genetics*
96, 377–385 (2015).
43.   Paré, G., Mao, S. & Deng, W. Q. A machine-learning heuristic to improve gene
score prediction of polygenic traits. *Sci Rep* **7,** 12665 (2017).
44.   Martin, A. R. *et al.* An Unexpectedly Complex Architecture for Skin Pigmentation
in Africans. (2017). doi:10.1101/200139
45.   Hindorff, L. A. *et al.* Prioritizing diversity in human genomics research. *Nature*
Publishing Group 1–11 (2017). doi:10.1038/nrg.2017.89
46.   H3Africa Consortium *et al.* Research capacity. Enabling the genomic revolution in
Africa. *Science* **344,** 1346–1348 (2014).
47.   Duncan, L. E. *et al.* Largest GWAS of PTSD (N=20□070) yields genetic overlap
with schizophrenia and sex differences in heritability. *Molecular Psychiatry* (2017).
doi:10.1038/mp.2017.77
48.   Howrigan, D. Details and Considerations of the Uk Biobank GWAS.
http://www.nealelab.is/blog/2017/9/11/details-and-considerations-of-the-uk-
biobank-gwas (2017). Available at: (Accessed: 9 November 2017)
49.   Kanai, M. *et al.* Genetic analysis of quantitative traits in the Japanese population
links cell types to complex human diseases. *Nat Genet* 1–16 (2018).
doi:10.1038/s41588-018-0047-6
50.   Wray, N. R. *et al.* Pitfalls of predicting complex traits from SNPs. *Nature*
Publishing Group **14,** 507–515 (2013).
51.   Wray, N. R. *et al.* Research Review: Polygenic methods and their application to
psychiatric traits. *J Child Psychol Psychiatr* **55,** 1068–1087 (2014).
52.   Torkamani, A., Wineinger, N. E. & Topol, E. J. The personal and clinical utility of
polygenic risk scores. *Nat Rev Genet* 1–10 (2018). doi:10.1038/s41576-018-
0018-x
53.   Manrai, A. K., Patel, C. J. & Ioannidis, J. P. A. In the Era of Precision Medicine
and Big Data, Who Is Normal? *JAMA* **319,** 1981–1982 (2018).
54.   Plenge, R. M., Scolnick, E. M. & Altshuler, D. Validating therapeutic targets
through human genetics. *Nature Publishing Group* **12,** 581–594 (2013).
55.   Carroll, M. D., Kit, B. K., Lacher, D. A., Shero, S. T. & Mussolino, M. E. Trends in
lipids and lipoproteins in US adults, 1988-2010. *JAMA* **308,** 1545–1554 (2012).

56.   Rappoport, N. *et al.* Creating ethnicity-specific reference intervals for lab tests
570   from EHR data. *bioRxiv* (2017). doi:10.1101/213892
571   57.   Lim, E., Miyamura, J. & Chen, J. J. Racial/Ethnic-Specific Reference Intervals for
572   Common Laboratory Tests: A Comparison among Asians, Blacks, Hispanics, and
573   White. *Hawaii J Med Public Health* **74,** 302–310 (2015).
574   58.   Hero, J. O., Zaslavsky, A. M. & Blendon, R. J. The United States Leads Other
575   Nations In Differences By Income In Perceptions Of Health And Health Care.
576   *Health Aff* **36,** 1032–1040 (2017).
577