Current clinical use of polygenic scores will risk exacerbating health disparities

Alicia R. Martin[1,2,3], Masahiro Kanai[1,2,3,4,5], Yoichiro Kamatani[5,6], Yukinori Okada[5,7,8], Benjamin M. Neale[1,2,3], Mark J. Daly[1,2,3,9]

[1] Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA 02114, USA
[2] Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA
[3] Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA
[4] Department of Biomedical Informatics, Harvard Medical School, Boston, MA 02115, USA
[5] Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan
[6] Kyoto-McGill International Collaborative School in Genomic Medicine, Graduate School of Medicine, Kyoto University, Kyoto 606-8507, Japan
[7] Department of Statistical Genetics, Osaka University Graduate School of Medicine, Suita 565-0871, Japan
[8] Laboratory of Statistical Immunology, Immunology Frontier Research Center (WPI-IFReC), Osaka University, Suita 565-0871, Japan
[9] Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland

Correspondence: Alicia R. Martin, armartin@broadinstitute.org

Abstract

Polygenic risk scores (PRS) are poised to improve biomedical outcomes via precision

medicine. However, the major *ethical* and *scientific* challenge surrounding clinical

implementation is that they are many-fold more accurate in European ancestry

individuals than others. This disparity is an inescapable consequence of Eurocentric

genome-wide association study biases. This highlights that—unlike clinical biomarkers

and prescription drugs, which may individually work better in some populations but do

not ubiquitously perform far better in European populations—clinical uses of PRS today

would systematically afford greater improvement to European descent populations.

Early diversifying efforts show promise in levelling this vast imbalance, even when non-

European sample sizes are considerably smaller than the largest studies to date. To

realize the full and equitable potential of PRS, we must prioritize greater diversity in

genetic studies and public dissemination of summary statistics to ensure that health

disparities are not increased for those already most underserved.

Keywords: health disparities, genetic risk prediction, polygenic risk scores, diversity,

population genetics, statistical genetics

Polygenic risk scores (PRS), which predict complex traits using genetic data, are of

burgeoning interest to the clinical community as researchers demonstrate their growing

power to improve clinical care, genetic studies of a wide range of phenotypes increase

in size and power, and genotyping costs plummet to less than US$50. Many earlier

criticisms of limited prediction power are now recognized to have been chiefly an issue

of insufficient sample size, which is no longer the case for many outcomes[1]. For

example, polygenic risk scores alone already predict breast cancer, prostate cancer,

and type 1 diabetes risk in European descent patients more accurately than current

clinical models[2-4]. Additionally, integrated models of PRS together with other lifestyle

and clinical factors have enabled clinicians to more accurately quantify the risk of heart

attack for patients; consequently, they have more effectively targeted the reduction of

LDL cholesterol and by extension heart attack by prescribing statins to patients at the

greatest overall risk of cardiovascular disease[5-9]. Promisingly, return of genetic risk of

complex disease to at-risk patients does not induce significant self-reported negative

behavior or psychological function, and some potentially positive behavioral changes

have been detected[10]. While we share enthusiasm about the potential of PRS to

improve health outcomes through their eventual routine implementation as clinical

biomarkers, we consider the consistent observation that they are currently of far greater

predictive value in individuals of recent European descent than in others to be the major

ethical and *scientific* challenge surrounding clinical translation and, at present, the most

critical limitation to genetics in precision medicine. The scientific basis of this imbalance

has been demonstrated theoretically, in simulations, and empirically across many traits

and diseases[11-22].

All studies to date using well-powered genome-wide association studies (GWAS) to

assess the predictive value of PRS across a range of traits and populations have made

a consistent observation: PRS predict individual risk far more accurately in Europeans

than non-Europeans[15,16,18-24] . Rather than chance or biology, this is a predictable

consequence of the fact that the genetic discovery efforts to date heavily

underrepresent non-European populations globally. The correlation between true and

genetically predicted phenotypes decays with genetic divergence from the makeup of

the discovery GWAS, meaning that the accuracy of polygenic scores in different

populations is highly dependent on the study population representation in the largest

existing 'training' GWAS. Here, we document study biases that underrepresent non-

European populations in current GWAS, and explain the fundamental concepts

contributing to reduced phenotypic variance explained with increasing genetic

divergence from populations included in GWAS.

Predictable basis of disparities in PRS accuracy

Poor generalizability of genetic studies across populations arises from the

overwhelming abundance of European descent studies and dearth of well-powered

studies in globally diverse populations[25-28]. According to the GWAS catalog, ~79% of all

GWAS participants are of European descent despite making up only 16% of the global

population (**Figure 1**). This is especially problematic as previous studies have shown

that Hispanic/Latino and African American studies contribute an outsized number of

associations relative to studies of similar sizes in Europeans[27]. More concerningly, the

fraction of non-European individuals in GWAS has stagnated or declined since late

2014 (**Figure 1**), suggesting that we are not on a trajectory to correct this imbalance.

These numbers provide a composite metric of study availability, accessibility, and use—

cohorts that have been included in numerous GWAS are represented multiple times,

which may disproportionately include cohorts of European descent. However, whereas

the average sample sizes of GWAS in Europeans continue to grow, they have

stagnated and remain several-fold smaller in other populations (**Supplementary Figure**

1).

The relative sample compositions of GWAS result in highly predictable disparities in

prediction accuracy; population genetics theory predicts that genetic risk prediction

accuracy will decay with increasing genetic divergence between the original GWAS

sample and target of prediction, a function of population history[13,14]. This pattern can be

attributed to several statistical observations which we detail below: 1) GWAS favor the

discovery of genetic variants that are common in the study population; 2) linkage

disequilibrium (LD) differentiates marginal effect size estimates for polygenic traits

across populations, even when causal variants are the same; and 3) environment and

demography differ across populations. Notably, the first two phenomena degrade

prediction performance across populations substantially even when there exist no

biological, environmental, or diagnostic differences, whereas the environment and

demography may interact to drive differential forces of natural selection that in turn drive

differences in causal genetic architecture. (We define the causal genetic architecture as

the true effects of variants that impact a phenotype that would be identified in a

population of infinite sample size. Unlike effect size estimates, true effects are typically

modeled as invariant with respect to LD and allele frequency differences across

populations.)

Common discoveries and low-hanging fruit

First, the power to discover an association in a genetic study depends on the effect size

and frequency of the variant[29]. This dependence means that the most significant

associations tend to be more common in the populations in which they are discovered

than elsewhere[13,30]. For example, GWAS catalog variants are more common on

average in European populations compared to East Asian and African populations

(**Figure 2B**), an observation not representative of genomic variants at large.

Understudied populations offer low-hanging fruit for genetic discovery because variants

that are common in these groups but rare or absent in European populations could not

be discovered even with very large European sample sizes. Some examples include

SLC16A11 and *HNF1A* associations with type II diabetes in Latino populations, as well

as *APOL1* associations with end-stage kidney disease and associations with prostate

cancer in African descent populations[31-34]. If we assume that causal genetic variants

have an equal effect across all populations—an assumption with some empirical

support that offers the best case scenario for transferability[35-40]—Eurocentric GWAS

biases mean that variants associated with risk are disproportionately common and

discovered in European populations, accounting for a larger fraction of the phenotypic

variance there[13]. Furthermore, imputation reference panels share the same study

biases as in GWAS[41], creating challenges for imputing sites that are rare in European

populations but common elsewhere when the catalog of non-European haplotypes is

substantially smaller. These issues are insurmountable through statistical methods

alone[13], but rather motivate substantial investments in more diverse populations to

produce similar-sized GWAS of biomedical phenotypes in other populations.

Linkage disequilibrium

Second, LD, the correlation structure of the genome, varies across populations due to

demographic history (**Figure 2A,C-E**). These LD differences in turn drive differences in

effect size estimates (i.e. predictors) from GWAS across populations in proportion to LD

between tagging and causal SNP pairs, even when causal effects are the same [35,37-40]

(**Supplementary Note**). Differences in effect size estimates due to LD differences may

typically be small for most regions of the genome (**Figure 2C-E**), but PRS sum across

these effects, also aggregating these population differences. While it would be ideal to

use causal effects rather than correlated effect size estimates to calculate PRS, it may

not be feasible to fine-map most variants to a single locus to solve issues of low

generalizability, even with very large GWAS. This is because complex traits are highly

polygenic, meaning most of our prediction power comes from small effects that do not

meet genome-wide significance and/or cannot be fine-mapped, even in many of the

best-powered GWAS to date[42].

Complexities of history, selection, and the environment

Lastly, other cohort considerations may further worsen prediction accuracy differences

across populations in less predictable ways. GWAS ancestry study biases and LD

differences across populations are extremely challenging to address, but these issues

actually make many favorable assumptions that all causal loci have the same impact

and are under equivalent selective pressure in all populations. In contrast, other effects

on polygenic adaptation or risk scores such as long-standing environmental differences

across global populations that have resulted in differing responses of natural selection

can impact populations differently based on their unique histories. Additionally, residual

uncorrected population stratification may impact risk prediction accuracy across

populations, but the magnitude of its effect is currently unclear. These effects are

particularly challenging to disentangle, as has clearly been demonstrated for height,

where evidence of polygenic adaptation and/or its relative magnitude is under

question[43,44]. Comparisons of geographically stratified phenotypes like height across

populations with highly divergent genetic backgrounds and mean environmental

differences, such as differences in resource abundance during development across

continents, are especially prone to confounding from correlated environmental and

genetic divergence[43,44]. This residual stratification can lead to over-predicted differences

across geographical space[45].

Related to stratification, most PRS methods do not explicitly address recent admixture

and none consider recently admixed individuals' unique local mosaic of ancestry; further

methods development is needed. Additionally, comparing PRS across environmentally

stratified cohorts, such as in some biobanks with healthy volunteer effects versus

disease study datasets or hospital-based cohorts, requires careful consideration of

technical differences, collider bias, as well as variability in baseline health status among

studies. It is also important to consider differences in definitions of clinical phenotypes

and heterogeneity of sub-phenotypes among countries.

Differences in environmental exposure, gene-gene interactions, gene-environment

interactions, historical population size dynamics, statistical noise, some potential causal

effect differences, and/or other factors will further limit generalizability for genetic risk

scores in an unpredictable, trait-specific fashion[46-49]. Complex traits do not behave in a

genetically deterministic manner, with some environmental factors dwarfing individual

genetic effects, creating outsized issues of comparability across globally diverse

populations. Among psychiatric disorders for example, whereas schizophrenia has a

nearly identical genetic basis across East Asians and Europeans (r_g=0.98)[40],

substantially different rates of alcohol use disorder across populations are partially

explained by differences in availability and genetic differences impacting alcohol

metabolism[50]. While non-linear genetic factors explain little variation in complex traits

beyond a purely additive model[51], some unrecognized nonlinearities and gene-gene

interactions can also induce genetic risk prediction challenges, as pairwise interactions

are likely to vary more across populations than individual SNPs. Mathematically, we can

simplistically think of this in terms of a two-SNP model, in which the sum of two SNP

effects is likely to explain more phenotypic variance than the product of the same SNPs.

Some machine learning approaches may thus modestly improve PRS accuracy beyond

current approaches for some phenotypes[52], but most likely for atypical traits with simpler

architectures, known interactions, and poor prediction generalizability across

populations, such as skin pigmentation[53].

Limited generalizability of PRS across diverse populations

So far, multi-ethnic work has been slow in most disease areas[54], limiting even the

opportunity to assess PRS in non-European cohorts. Nonetheless, some previous work

has assessed prediction accuracy across diverse populations in several traits and

diseases for which GWAS summary statistics are available and identified large

disparities across populations (**Supplementary Note**). These disparities are not simply

methodological issues, as various approaches (e.g. pruning and thresholding versus

LDPred) and accuracy metrics (R^2 for quantitative traits and various pseudo-R^2 metrics

for binary traits) illustrate this consistently poorer performance in populations distinct

from discovery samples across a range of polygenic traits (**Supplementary Table 1**).

These assessments are becoming increasingly feasible with the growth and public

availability of global biobanks as well as diversifying priorities from funding

agencies[55,56]. We assessed how prediction accuracy decayed across globally diverse

populations for 17 anthropometric and blood panel traits in the UK Biobank (UKBB)

when using European-derived summary statistics (**Supplementary Note**). Consistent

with previous studies, we find that relative to European prediction accuracy, genetic

prediction accuracy was far lower in other populations: 1.6-fold lower in Hispanic/Latino

Americans, 1.7-fold lower in South Asians, 2.5-fold lower in East Asians, and 4.9-fold

lower in Africans on average (**Figure 3**).

Prioritizing diversity shows early promise for PRS

Early diversifying GWAS efforts have been especially productive for informing on

questions surrounding risk prediction. Rather than varying the prediction target dataset,

some GWAS in diverse populations have increased the scale of non-European

summary statistics and also varied the study dataset in multi-ethnic PRS studies[23,24,40].

These studies have shown that even when non-European cohorts are only a fraction the

size of the largest European study, they are likely to have disproportionate value for

predicting polygenic traits in other individuals of similar ancestry.

Given this background, we performed a systematic evaluation of polygenic prediction

accuracy across 17 quantitative anthropometric and blood panel traits and five disease

endpoints in British and Japanese individuals[23,57,58] by performing GWAS with the exact

same sample sizes in each population. We symmetrically demonstrate that prediction

accuracy is consistently higher with GWAS summary statistics from ancestry-matched

summary statistics (**Figure 4, Supplementary Figures 2-6**). Keeping in mind issues of

comparability described above, we note that BBJ is a hospital-based disease-

ascertained cohort, whereas UKBB is a healthier than average[59] population-based

cohort; thus, differences in observed heritability among these cohorts (rather than

among populations) due to differences in phenotype precision likely explain lower

prediction accuracy from the BBJ GWAS summary statistics for anthropometric and

blood panel traits, but higher prediction accuracy for five ascertained diseases

(**Supplementary Table 2**). Indeed, other East Asian studies have estimated higher

heritability for some quantitative traits than BBJ using the same methods, such as for

height (h^2 = 0.48±0.04 in Chinese women[60]). Some statistical fluctuations in the relative

differences in prediction accuracy across populations are likely driven by differences in

heritability measured in each population and/or trans-ethnic genetic correlation (i.e. of

common variant effect sizes at SNPs common in two populations, **Supplementary**

Figures 7-10, **Supplementary Tables 2–5**). These trans-ethnic correlation estimates

indicate that effect sizes were mostly highly correlated across ancestries, with a few

traits that were somewhat lower than excepted (e.g. height and BMI, with ρ_{ge}=0.69 and

0.75, respectively). Prediction accuracy was far lower in individuals of African descent in

the UK Biobank (**Supplementary Figures 4 and 11**) using GWAS summary statistics

from either European or Japanese ancestry individuals, consistent with reduced

prediction accuracy with increasing genetic divergence (**Figures 3 and 4**). These

population studies demonstrate the power and utility of increasingly diverse GWAS for

prediction, especially in populations of non-European descent.

While many other traits and diseases have been studied in multi-ethnic settings, few

have reported comparable metrics of prediction accuracy across populations.

Cardiovascular research, for example, has led the charge towards clinical translation of

PRS[1]. This enthusiasm is driven by observations that a polygenic burden of LDL-

increasing SNPs can confer monogenic-equivalent risk of cardiovascular disease, with

polygenic scores improving clinical models for risk assessment and statin prescription

that can reduce coronary heart disease and improve healthcare delivery efficiency[5-7].

However, many of these studies have been conducted exclusively in European descent

populations, with few studies rigorously evaluating population-level applicability to non-

Europeans. Those existing findings indeed demonstrate a large reduction in prediction

utility in non-European populations[11], though often with comparisons of odds ratios

among arbitrary breakpoints in the risk distribution that make comparisons across

studies challenging. To better clarify how polygenic prediction will be deployed in a

clinical setting with diverse populations, more systematic and thorough evaluations of

the utility of PRS within and across populations for many complex traits are still needed.

These evaluations would benefit from rigorous polygenic prediction accuracy

evaluations, especially for diverse non-European patients[61-63].

Clinical use of PRS may uniquely exacerbate disparities

Our impetus for raising these statistical issues limiting the generalizability of PRS across

population stems from our concerns that, while they are legitimately clinically promising

for improving health outcomes for many biomedical phenotypes, they may have a larger

potential to raise health disparities than other clinical factors for several reasons. The

opportunities they provide for improving health outcomes means they inevitably will and

should be pursued in the near term, but we urge that a concerted prioritization to make

GWAS summary statistics easily accessible for diverse populations and a variety of

traits and diseases is imperative, even when they are a fraction the size of the largest

existing European datasets. *Individual* clinical tests, biomarkers, and prescription drug

efficacy may vary across populations in their utility, but are fundamentally informed by

the same underlying biology[64,65]. Currently, guidelines state that as few as 120

individuals define reference intervals for clinical factors (though often smaller numbers

from only one subpopulation are used) and there is no clear definition of who is

"normal"[64]. Consequently, reference intervals for biomarkers can sometimes deviate

considerably by reported ethnicity[66-68]. Defining ethnicity-specific reference intervals is

clearly an important problem that can provide fundamental interpretability gains with

implications for some major health benefits (e.g. need for dialysis and development of

Type 2 diabetes based on ethnicity-specific serum creatinine and hemoglobin A1C

reference intervals, respectively)[67]. Simply put, some biomarkers or clinical tests scale

directly with health outcomes independent of ancestry, and many others may have

distributional differences by ancestry but are equally valid after centering with respect to

a readily collected population reference.

In contrast, PRS are uniformly less useful in understudied populations due to

differences in genomic variation and population history[13,14]. No analogous solution of

defining ethnicity-specific reference intervals would ameliorate health disparities

implications for PRS or fundamentally aid interpretability in non-European populations.

Rather, as we and others demonstrate, PRS are unique in that even with multi-ethnic

population references, these scores are fundamentally less informative in populations

more diverged from GWAS study cohorts.

The clinical use and deployment of genetic risk scores needs to be informed by the

issues surrounding tests that currently would unequivocally provide much greater

benefit to the subset of the world's population which is already on the positive end of

healthcare disparities. Conversely, African descent populations, which already endure

many of the largest health disparities globally, are often predicted marginally better, if at

all, compared to random (**Figure 4F**). They are therefore least likely to benefit from

improvements in precision healthcare delivery from genetic risk scores with existing

data due to human population history and study biases. This is a major concern globally

and especially in the U.S., which already leads other middle- and high-income countries

in both real and perceived healthcare disparities[69,70]. Thus, we would strongly urge that

any discourse on clinical use of PRS include a careful, quantitative assessment of the

economic and health disparities impacts on underrepresented populations that might be

unintentionally introduced, and raise awareness about how to eliminate these

disparities.

How do we even the ledger?

What can be done? The single most important step towards parity in PRS accuracy is

by vastly increasing the diversity of participants included and analyzed in genetic

studies, which will improve utility for all and most rapidly for underrepresented groups.

Regulatory protections against genetic discrimination are necessary to accompany calls

for more diverse studies; while some already exist in the U.S., including for health

insurance and employment opportunities via the Genetic Information Nondiscrimination

Act (GINA), stronger protections in these and other areas globally will be particularly

important for minorities and/or marginalized groups. An equal investment in GWAS

across all major ancestries and global populations is the most obvious solution to

generate a substrate for equally informative risk scores but is not likely to occur any

time soon absent a dramatic priority shift given the current imbalance and stalled

diversifying progress over the last five years (**Figure 1, Supplementary Figure 1**).

While it may be challenging or in some cases infeasible to acquire sample sizes large

enough for PRS to be equally informative in all populations, some much-needed efforts

towards increasing diversity in genomics that support open sharing of GWAS summary

data from multiple ancestries are underway. Examples include the *All of Us* Research

Program, the Population Architecture using Genomics and Epidemiology (PAGE)

Consortium, as well as some disease-focused consortia, such as the T2D-genes and

Stanley Global initiatives on the genetics of type II diabetes and psychiatric disorders,

respectively. Supporting data resources such as imputation panels, multi-ethnic

genotyping arrays, gene expression datasets from genetically diverse individuals, and

other tools are necessary to similarly empower these diverse studies for all populations.

The lack of supporting resources for diverse ancestries creates financial challenges for

association studies with limited resources, e.g. raising questions about whether to

genotype samples on GWAS arrays that may favor European allele frequencies versus

sequence samples, and how dense of an array to choose or how deeply to

sequence[71,72].

Additional leading global efforts also provide easy unified access linking genetic, clinical

record, and national registry data in more homogeneous continental ancestries, such as

the UK Biobank, BioBank Japan, China Kadoorie Biobank, and Nordic efforts (e.g. in

Danish, Estonian, Finnish, and other integrated biobanks). Notably, some of these

biobanks such as UK Biobank have participants with considerable global genetic

diversity that enables multi-ethnic comparisons; although minorities from this cohort

provide the largest deeply phenotyped GWAS cohorts for several ancestries, these

individuals are often excluded in current statistical analyses in favor of single ancestries,

large sample sizes, and the simplicity afforded by genetic homogeneity. These

considerations notwithstanding, there are critical needs and challenges for expanding

the scale of genetic studies of heritable traits in diverse populations; this is especially

apparent in Africa where humans originated and retain the most genetic diversity, as

Africans are understudied but disproportionately informative for genetic analyses and

evolutionary history[27,73]. The most notable investment here comes from the Human

Heredity and Health in Africa (H3Africa) Initiative, increasing genomics research

capacity in Africa through more than $216 million in funding from the NIH (USA) and

Wellcome Trust (UK) for genetics research led by African investigators[55,74]. The

increasing interest and scale of genetic studies in low- and middle-income countries

(LMICs) raises ethical and logistical considerations about data generation, access,

sharing, security, and analysis, as well as clinical implementation to ensure these

advances do not only benefit high-income countries. Frameworks such as the

H3ABioNet, a pan-African bioinformatics network designed to build capacity to enable

H3Africa researchers to analyze their data in Africa, provide cost-effective examples for

training local scientists in LMICs[75].

The prerequisite data for dramatically increasing diversity also hypothetically exist in

several large-scale publicly funded datasets such as the Million Veterans Project and

Trans-Omics for Precision Medicine (TOPMed), but with problematic data access issues

in which even GWAS summary data within and across populations are not publicly

shared. Existing GWAS consortia also need to carefully consider the granularity of

summary statistics they release, as finer scale continental ancestries and phenotypes in

large, multi-ethnic projects enable ancestry-matched analyses not possible with a single

set of summary statistics. While there is an understandable patient privacy balance to

strike when sharing individual-level data, GWAS summary statistics from all publicly

funded and as many privately funded projects as possible should be made easily and

publicly accessible to improve global health outcomes. Efforts to unify phenotype

definitions, normalization approaches, and GWAS methods among studies will also

improve comparability.

To enable progress towards parity, it will be critical that open data sharing standards be

adopted for all ancestries and for genetic studies of all sample sizes, not just the largest

European results. Locally appropriate and secure genetic data sharing techniques as

well as equitable technology availability will need to be adopted widely in Asia and

Africa as they are in Europe and North America, to ensure that maximum value is

achieved from existing and ongoing efforts that are being developed to help counter the

current imbalance. Simultaneously, ethical considerations require that research capacity

is increased in LMICs with simultaneous growth of diverse population studies to balance

the benefits of these studies to scientists and patients globally versus locally to ensure

that everyone benefits. Methodological improvements that better define risk scores by

accounting for population allele frequency, LD, and/or admixture differences

appropriately are underway and may help considerably but will not by themselves bring

equality. All of these efforts are important and should be prioritized not just for risk

prediction but more generally to maximize the use and applicability of genetics to inform

on the biology of disease. Given the acute recent attention on clinical use of PRS, we

believe it is paramount to recognize their potential to improve health outcomes for all

individuals and many complex diseases. Simultaneously, we as a field must address the

disparity in utility in an ethically thoughtful and scientifically rigorous fashion, lest we

inadvertently enable genetic technologies to contribute to, rather than reduce, existing

health disparities.

Author contributions

Competing interests

The authors declare no competing interests.

Acknowledgments

We thank Amit Khera for helpful discussions. We also thank Michiaki Kubo, Yoshinori

Murakami, Masato Akiyama, and Kazuyoshi Ishigaki for their support in the BioBank

Japan Project analysis. We are grateful to Steven Gazal for his help in calculating LD

scores. This work was supported by funding from the National Institutes of Health

(K99MH117229 to A.R.M.). UK Biobank analyses were conducted via application

31063. The BioBank Japan Project was supported by the Tailor-Made Medical

Treatment Program of the Ministry of Education, Culture, Sports, Science, and

Technology (MEXT) and the Japan Agency for Medical Research and Development

(AMED). M.K. was supported by a Nakajima Foundation Fellowship and the Masason

Foundation.

Figures

442 **Figure 1 – Ancestry of GWAS participants over time compared to the global**

443 **population**. Cumulative data as reported by the GWAS catalog[76]. Individuals whose

444 ancestry is "not reported" are not shown.

Figure 2 – Demographic relationships, allele frequency differences, and local LD

patterns between population pairs. Data analyzed from 1000 Genomes, in which

population labels are: AFR = continental African, EUR = European, and EAS = East

Asian. **a**) Cartoon relationships among AFR, EUR, and EAS populations. **b**) Allele

frequency distributions in AFR, EUR, and EAS populations of variants from the GWAS

catalog. **c-e**) Color axis shows LD scale (r^2). LD comparisons between pairs of

populations show the same region of the genome for each comparison (representative

region is chr1, 51572kb-52857kb) among pairs of SNPs polymorphic in both

populations, illustrating that different SNPs are polymorphic across some population

pairs, and that these SNPs have variable LD patterns across populations.

Figure 3 – Prediction accuracy relative to European ancestry individuals across 17 quantitative traits and 5 continental populations in UKBB. All phenotypes shown here are quantitative anthropometric and blood panel traits, as described in **Supplementary Table 6**, which includes discovery cohort sample sizes. Prediction target individuals do not overlap with the discovery cohort and are unrelated, with sample sizes shown in **Supplementary Table 7**. Violin plots show distributions of relative prediction accuracies, points show mean values, and error bars show standard errors of the means. Prediction R^2 for each trait and population are shown in **Supplementary Figure 12**.

Figure 4 – Polygenic risk prediction accuracy in Japanese, British, and African descent individuals using independent GWAS of equal sample sizes in the BioBank Japan (BBJ) and UK Biobank (UKBB). a) Explanatory diagram showing the different discovery and target cohorts/populations, and disease endpoints versus quantitative traits. **b-f**) Genetic prediction accuracy computed from independent BBJ and UKBB summary statistics with identical sample sizes (**Supplementary Tables 6 and 8**). Note that y-axes differ, reflecting differences in prediction accuracy. **b-c**) PRS accuracy for five diseases in: Japanese individuals in the BBJ (**b**) and British individuals in the UKBB. **d-f**) PRS accuracy for 17 anthropometric and blood panel traits in: Japanese individuals in the BBJ (**d**), British individuals in the UKBB (**e**), and African descent British individuals in the UKBB (**f**). Trait abbreviations are as in **Supplementary Table 6**. Each point shows the maximum R^2 (i.e. best predictor) across five p-value thresholds, and lines correspond to 95% confidence intervals calculated via bootstrap.

R^2 values for all p-value thresholds tested are shown in **Supplementary Figures 2-6**.

Prediction accuracy tends to be higher in the UKBB for quantitative traits than in BBJ

and vice versa for disease endpoints, likely because of concomitant phenotype

precision and consequently observed heritability for these classes of traits

(**Supplementary Tables 2-4**). Thalassemia and sickle cell disease are unlikely to

explain a significant fraction of prediction accuracy differences for blood panels across

populations, as few individuals have been diagnosed with these disorders via ICD-10

codes (**Supplementary Table 9**).

References

1. Knowles, J. W. & Ashley, E. A. Cardiovascular disease: The rise of the genetic risk score. *PLoS Med* **15,** e1002546–7 (2018).
2. Maas, P. *et al.* Breast Cancer Risk From Modifiable and Nonmodifiable Risk Factors Among White Women in the United States. *JAMA Oncol* **2,** 1295–8 (2016).
3. Schumacher, F. R. *et al.* Association analyses of more than 140,000 men identify 63 new prostate cancer susceptibility loci. *Nat Genet* **50,** 928–936 (2018).
4. Sharp, S. A. *et al.* Development and Standardization of an Improved Type 1 Diabetes Genetic Risk Score for Use in Newborn Screening and Incident Diagnosis. *Diabetes Care* **42,** 200–207 (2019).
5. Khera, A. V. *et al.* Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat Genet* **50,** 1219–1224 (2018).
6. Kullo, I. J. *et al.* Incorporating a Genetic Risk Score Into Coronary Heart Disease Risk Estimates: Effect on Low-Density Lipoprotein Cholesterol Levels (the MI-GENES Clinical Trial). *Circulation* **133,** 1181–1188 (2016).
7. Natarajan, P. *et al.* Polygenic Risk Score Identifies Subgroup With Higher Burden of Atherosclerosis and Greater Relative Benefit From Statin Therapy in the Primary Prevention Setting. *Circulation* **135,** 2091–2101 (2017).
8. Paquette, M. *et al.* Polygenic risk score predicts prevalence of cardiovascular disease in patients with familial hypercholesterolemia. *Journal of Clinical Lipidology* **11,** 725–732.e5 (2017).
9. Tikkanen, E., Havulinna, A. S., Palotie, A., Salomaa, V. & Ripatti, S. Genetic Risk Prediction and a 2-Stage Risk Screening Strategy for Coronary Heart Disease. *Arterioscler. Thromb. Vasc. Biol.* **33,** 2261–2266 (2013).
10. Frieser, M. J., Wilson, S. & Vrieze, S. Behavioral impact of return of genetic test results for complex disease: Systematic review and meta-analysis. *Health Psychol* **37,** 1134–1144 (2018).
11. Khera, A. V. *et al.* Genetic Risk, Adherence to a Healthy Lifestyle, and Coronary Disease. *N Engl J Med* **375,** 2349–2358 (2016).
12. Khera, A. V. & Kathiresan, S. Genetics of coronary artery disease: discovery, biology and clinical translation. *Nature Publishing Group* **18,** 331–344 (2017).
13. Martin, A. R. *et al.* Human Demographic History Impacts Genetic Risk Prediction across Diverse Populations. *Am. J. Hum. Genet.* **100,** 635–649 (2017).
14. Scutari, M., Mackay, I. & Balding, D. Using Genetic Distance to Infer the Accuracy of Genomic Prediction. *PLoS Genet* **12,** e1006288–19 (2016).
15. Vilhjálmsson, B. J. *et al.* Modeling Linkage Disequilibrium Increases Accuracy of Polygenic Risk Scores. *Am. J. Hum. Genet.* **97,** 576–592 (2015).
16. Ware, E. B. *et al.* Heterogeneity in polygenic scores for common human traits. *biorxiv.org* doi:10.1101/106062
17. Curtis, D. Polygenic risk score for schizophrenia is more strongly associated with ancestry than with schizophrenia. *Psychiatric Genetics* **28,** 85–89 (2018).

534   18.   Schizophrenia Working Group of the Psychiatric Genomics Consortium.
535   Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511,**
536   421–427 (2014).
537   19.   Belsky, D. W. *et al.* Development and Evaluation of a Genetic Risk Score for
538   Obesity. *Biodemography and Social Biology* **59,** 85–100 (2013).
539   20.   Domingue, B. W., Belsky, D. W., Conley, D., Harris, K. M. & Boardman, J. D.
540   Polygenic Influence on Educational Attainment. *AERA Open* **1,**
541   233285841559997–13 (2015).
542   21.   Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide
543   association study of educational attainment in 1.1 million individuals. *Nature*
544   *Publishing Group* 1–16 (2018). doi:10.1038/s41588-018-0147-3
545   22.   Vassos, E. *et al.* An Examination of Polygenic Score Risk Prediction in
546   Individuals With First-Episode Psychosis. *Biological Psychiatry* **81,** 470–477
547   (2017).
548   23.   Akiyama, M. *et al.* Genome-wide association study identifies 112 new loci for
549   body mass index in the Japanese population. *Nat Genet* **49,** 1458–1467 (2017).
550   24.   Li, Z. *et al.* Genome-wide association analysis identifies 30 new susceptibility loci
551   for schizophrenia. *Nat Genet* **49,** 1576–1583 (2017).
552   25.   Need, A. C. & Goldstein, D. B. Next generation disparities in human genomics:
553   concerns and remedies. *Trends in Genetics* **25,** 489–494 (2009).
554   26.   Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature* **538,**
555   161–164 (2016).
556   27.   Morales, J. *et al.* A standardized framework for representation of ancestry data in
557   genomics studies, with application to the NHGRI-EBI GWAS Catalog. 1–10
558   (2018). doi:10.1186/s13059-018-1396-2
559   28.   Rosenberg, N. A. *et al.* Genome-wide association studies in diverse populations.
560   *Nature Publishing Group* **11,** 356–366 (2010).
561   29.   Sham, P. C., Cherny, S. S., Purcell, S. & Hewitt, J. K. Power of linkage versus
562   association analysis of quantitative traits, by use of variance-components
563   models, for sibship data. *The American Journal of Human Genetics* **66,** 1616–
564   1630 (2000).
565   30.   The 1000 Genomes Project Consortium *et al.* A global reference for human
566   genetic variation. *Nature* **526,** 68–74 (2015).
567   31.   Consortium, T. S. 2. D. Sequence variants in SLC16A11 are a common risk
568   factor for type 2 diabetes in Mexico. *Nature* **506,** 97–101 (2014).
569   32.   Estrada, K. *et al.* Association of a Low-Frequency Variant in HNF1AWith Type 2
570   Diabetes in a Latino Population. *JAMA* **311,** 2305–10 (2014).
571   33.   Haiman, C. A. *et al.* Genome-wide association study of prostate cancer in men of
572   African ancestry identifies a susceptibility locus at 17q21. *Nature Publishing*
573   *Group* **43,** 570–573 (2011).
574   34.   Genovese, G. *et al.* Association of trypanolytic ApoL1 variants with kidney
575   disease in African Americans. *Science* **329,** 841–845 (2010).
576   35.   International Multiple Sclerosis Genetics Consortium *et al.* Association analyses
577   identify 38 susceptibility loci for inflammatory bowel disease and highlight shared
578   genetic risk across populations. *Nat Genet* **47,** 979–986 (2015).

36.   Carlson, C. S. *et al.* Generalization and Dilution of Association Results from
European GWAS in Populations of Non-European Ancestry: The PAGE Study.
PLoS Biol **11,** e1001661–11 (2013).
37.   Easton, D. F. *et al.* Genome-wide association study identifies novel breast cancer
susceptibility loci. *Nature* **447,** 1087–1093 (2007).
38.   DIAbetes Genetics Replication And Meta-analysis (DIAGRAM) Consortium *et al.*
Genome-wide trans-ancestry meta-analysis provides insight into the genetic
architecture of type 2 diabetes susceptibility. *Nat Genet* **46,** 234–244 (2014).
39.   Waters, K. M. *et al.* Consistent Association of Type 2 Diabetes Risk Variants
Found in Europeans in Diverse Racial and Ethnic Groups. *PLoS Genet* **6,**
e1001078–9 (2010).
40.   Lam, M. *et al.* Comparative genetic architectures of schizophrenia in East Asian
and European populations. 1–41 (2018). doi:10.1101/445874
41.   McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype
imputation. *Nat Genet* **48,** 1279–1283 (2016).
42.   Huang, H. *et al.* Fine-mapping inflammatory bowel disease loci to single-variant
resolution. *Nature* 1–19 (2017). doi:10.1038/nature22969
43.   Sohail, M. *et al.* Signals of polygenic adaptation on height have been
overestimated due to uncorrected population structure in genome-wide
association studies. 1–44 (2018). doi:10.1101/355057
44.   Berg, J. J. *et al.* Reduced signal for polygenic adaptation of height in UK
Biobank. 1–44 (2018). doi:10.1101/354951
45.   Kerminen, S. *et al.* Geographic variation and bias in polygenic scores of complex
diseases and traits in Finland:. 1–35 (2018). doi:10.1101/485441
46.   Novembre, J. & Barton, N. H. Tread Lightly Interpreting Polygenic Tests of
Selection. *Genetics* **208,** 1351–1355 (2018).
47.   Henn, B. M., Botigué, L. R., Bustamante, C. D., Clark, A. G. & Gravel, S.
Estimating the mutation load in human genomes. *Nature Publishing Group* **16,**
333–343 (2015).
48.   Brown, B. C., Asian Genetic Epidemiology Network Type 2 Diabetes Consortium,
Ye, C. J., Price, A. L. & Zaitlen, N. Transethnic Genetic-Correlation Estimates
from Summary Statistics. *Am. J. Hum. Genet.* **99,** 76–88 (2016).
49.   Galinsky, K. J. *et al.* Estimating cross-population genetic correlations of causal
effect sizes. *Genet. Epidemiol.* **200,** 1285 (2018).
50.   Li, D., Zhao, H. & Gelernter, J. Strong protective effect of the aldehyde
dehydrogenase gene (ALDH2) 504lys (*2) allele against alcoholism and alcohol-
induced medical diseases in Asians. *Human Genetics* **131,** 725–737 (2011).
51.   Zhu, Z. *et al.* Dominance Genetic Variation Contributes Little to the Missing
Heritability for Human Complex Traits. *The American Journal of Human Genetics*
96, 377–385 (2015).
52.   Paré, G., Mao, S. & Deng, W. Q. A machine-learning heuristic to improve gene
score prediction of polygenic traits. *Sci Rep* **7,** 12665 (2017).
53.   Martin, A. R. *et al.* An Unexpectedly Complex Architecture for Skin Pigmentation
in Africans. *Cell* **171,** 1340–1353.e14 (2017).

54. Duncan, L. E. *et al.* Largest GWAS of PTSD (N=20 070) yields genetic overlap with schizophrenia and sex differences in heritability. *Molecular Psychiatry* **23,** 666–673 (2018).

55. H3Africa Consortium *et al.* Research capacity. Enabling the genomic revolution in Africa. *Science* **344,** 1346–1348 (2014).

56. Hindorff, L. A. *et al.* Prioritizing diversity in human genomics research. *Nature Publishing Group* **19,** 175–185 (2018).

57. Kanai, M. *et al.* Genetic analysis of quantitative traits in the Japanese population links cell types to complex human diseases. *Nat Genet* **50,** 390–400 (2018).

58. Howrigan, D. Details and Considerations of the UK Biobank GWAS. (2017). Available at: http://scholar.google.comjavascript:void(0). (Accessed: 9 November 2017)

59. Fry, A. *et al.* Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. *Am. J. Epidemiol.* **186,** 1026–1034 (2017).

60. Liu, S. *et al.* Genomic Analyses from Non-invasive Prenatal Testing Reveal Genetic Associations, Patterns of Viral Infections, and Chinese Population History. *Cell* **175,** 347–359.e14 (2018).

61. Wray, N. R. *et al.* Pitfalls of predicting complex traits from SNPs. *Nature Publishing Group* **14,** 507–515 (2013).

62. Wray, N. R. *et al.* Research Review: Polygenic methods and their application to psychiatric traits. *J Child Psychol Psychiatr* **55,** 1068–1087 (2014).

63. Torkamani, A., Wineinger, N. E. & Topol, E. J. The personal and clinical utility of polygenic risk scores. *Nature Publishing Group* **19,** 581–590 (2018).

64. Manrai, A. K., Patel, C. J. & Ioannidis, J. P. A. In the Era of Precision Medicine and Big Data, Who Is Normal? *JAMA* **319,** 1981–3 (2018).

65. Plenge, R. M., Scolnick, E. M. & Altshuler, D. Validating therapeutic targets through human genetics. *Nature Publishing Group* **12,** 581–594 (2013).

66. Carroll, M. D., Kit, B. K., Lacher, D. A., Shero, S. T. & Mussolino, M. E. Trends in Lipids and Lipoproteins in US Adults, 1988-2010. *JAMA* **308,** 1545–1554 (2012).

67. Rappoport, N. *et al.* Comparing Ethnicity-Specific Reference Intervals for Clinical Laboratory Tests from EHR Data. *Jrnl App Lab Med* **3,** 366–377 (2018).

68. Lim, E., Miyamura, J. & Chen, J. J. Racial/Ethnic-Specific Reference Intervals for Common Laboratory Tests: A Comparison among Asians, Blacks, Hispanics, and White. *Hawaii J Med Public Health* **74,** 302–310 (2015).

69. Hero, J. O., Zaslavsky, A. M. & Blendon, R. J. The United States Leads Other Nations In Differences By Income In Perceptions Of Health And Health Care. *Health Affairs* **36,** 1032–1040 (2017).

70. Williams, D. R., Priest, N. & Anderson, N. B. Understanding associations among race, socioeconomic status, and health: Patterns and prospects. *Health Psychology* **35,** 407–411 (2016).

71. Gilly, A. *et al.* Very low depth whole genome sequencing in complex trait association studies. *Bioinformatics* (2018). doi:10.1093/bioinformatics/bty1032

72. Pasaniuc, B. *et al.* Extremely low-coverage sequencing and imputation increases power for genome-wide association studies. *Nat Genet* **44,** 631–635 (2012).

73.   Martin, A. R., Teferra, S., Möller, M., Hoal, E. G. & Daly, M. J. The critical needs
and challenges for genetic architecture studies in Africa. *Current Opinion in*
Genetics & Development **53,** 113–120 (2018).
74.   Coles, E. & Mensah, G. A. Geography of Genetics and Genomics Research
Funding in Africa. *Global Heart* **12,** 173–176 (2017).
75.   Mulder, N. J. *et al.* Development of Bioinformatics Infrastructure for Genomics
Research. *Global Heart* **12,** 91–98 (2017).
76.   MacArthur, J. *et al.* The new NHGRI-EBI Catalog of published genome-wide
association studies (GWAS Catalog). *Nucl. Acids Res.* **45,** D896–D901 (2017).
677