Phenotypic and genotypic resistance to commonly used insecticides in *Aedes aegypti* among four cities in southern Ecuador

Sadie J. Ryan[1,2†*] , Stephanie J. Mundis[1,2†], Alex Aguirre[3†], Catherine A. Lippi[1,2], Efraín Beltrán[5], Froilán Heras[5], Anna M. Stewart-Ibarra[4*], Marco Neira[3*]

[1]Quantitative Disease Ecology and Conservation (QDEC) Lab, Department of Geography, University of Florida, Gainesville, FL 32610 USA

[2]Emerging Pathogens Institute, University of Florida, Gainesville, FL 32610 USA

[3]Center for Research on Health in Latin America (CISeAL), Pontificia Universidad Católica del Ecuador, Quito, 170143 Ecuador

[4]Institute for Global Health and Translational Science and Department of Medicine, SUNY Upstate Medical University, Syracuse, NY 13210 USA

[5]Universidad Técnica de Machala, Machala, 070102 Ecuador

[†]These authors contributed equally to this work

*Corresponding authors:

Sadie J. Ryan: sjryan@ufl.edu

Anna M. Stewart-Ibarra: stewarta@upstate.edu

Marco Neira: mvneira@puce.edu.ec

Abstract

Insecticide resistance (IR) can undermine efforts to control vector species of public health importance. *Aedes aegypti* is the main vector of resurging diseases in the Americas such as yellow fever and dengue, as well as recently emerging chikungunya and Zika viruses, which have caused unprecedented epidemics in the region. Vector control remains the primary public health intervention to prevent outbreaks of *Aedes* transmitted diseases. In many high-risk regions, like southern Ecuador, we have limited information on IR in *Ae. aegypti*. In this study, IR status in *Ae. aegypti* was measured across four cities in El Oro Province in Ecuador using phenotypic assays and genetic screening for alleles associated with resistance to pyrethroid insecticides. Bottle bioassays showed significant inter-seasonal variation in resistance to deltamethrin, a pyrethroid insecticide commonly used by the Ministry of Health, and alpha-cypermethrin, as well as differences in deltamethrin resistance between cities. There was also a significant difference in phenotypic response to the organophosphate, Malathion, between two of the cities during the second sampling season. Genotyping showed moderate to high frequencies of the V1016I and F1534C resistance alleles in all four cites. Frequency of resistance genotypes varied significantly between cities in the first sampling season, and not in the later seasons, suggesting a possible selective response to vector control activity. Overall, resistance levels were highest in Machala, a city where dengue is hyperendemic and where insecticide use has historically been more intense than in the other cities included in the study. Despite statistically significant evidence that resistance alleles conferred phenotypic resistance, there was not precise correspondence between these indicators. We found that 17.6% of F1534C and 45.6% of V1016I mutant mosquitoes were susceptible in the bottle bioassays. This study shows there is spatiotemporal variability in IR in southern Ecuador, and serves as an initial examination of the genotypic and phenotypic indicators of IR in this region, providing important information to guide the vector control interventions by the public health sector.

Keywords: Insecticide resistance; *Aedes aegypti*; pyrethroid; organophosphate; Ecuador; arbovirus

Introduction

In Ecuador, dengue, chikungunya, and Zika viruses are present and transmitted to people by the *Aedes aegypti* mosquito, causing a high burden of febrile illness in susceptible populations. This species is a particularly effective vector of these viruses because it has evolved to live in urban environments, lay its eggs in small containers of water in and around human dwellings, and feed preferentially on humans [1]. There were 14,150 cases of dengue reported in Ecuador in the year 2016 [2], while chikungunya caused an estimated 2,025 cases [3] and Zika caused approximately 3,531 cases by the end of 2016 [4]. For all three of these diseases, actual incidence in Ecuador is likely much higher than the reported figures indicate because many cases are asymptomatic or mild, and access to laboratory diagnostics can be limited [5]. Furthermore, co-circulation of these viruses can lead to more severe disease outcomes and differential diagnosis is complicated [6]. To reduce the burden of disease, public health organizations rely heavily on vector control methods, particularly insecticide applications [7]. Additionally, individuals purchase products such as aerosols, repellents, and nets to reduce mosquito populations and prevent disease transmission at the household level, with low-income households in Machala, Ecuador, spending as much a 10% of their discretionary income on these items [5].

There are four main categories of chemical insecticides regularly used for disease vector control: organophosphates, pyrethroids, carbamates, and organochlorines, with organophosphates and pyrethroids being the most widely used in Ecuador for *Ae. aegypti* control [8]. For all four of these insecticide classes, regular deployment in locations around the world has been associated with the development of insecticide resistance (IR) in targeted vector populations, resulting in resurgences of mosquito-borne diseases [9–11]. While public health

organizations have made recommendations for monitoring and managing IR [12], these practices

are often resource-intensive, meaning many areas do not have the capacity to conduct regular IR

testing [13,14]. Additionally, local-scale IR can be driven by the largely unregulated application

of insecticide treatments at the household level by private individuals [15]. When resistance has

developed, but remains undetected, public health organizations may unintentionally spend

substantial time and resources applying insecticides that are ineffective [16] and may have

negative environmental and human health impacts [17].

A better understanding of the spatial and temporal variability in IR would greatly

enhance the ability of vector control groups to predict and mitigate IR in vector populations.

Research has shown that IR status can differ dramatically across cities within a country [18–20],

but further work needs to be conducted to understand the spatial scale of this variability,

particularly across cities with differing vector control needs and strategies. Furthermore, most

available reports in this area determine IR status at a single point in time, but IR exhibits

temporal fluctuations that should be taken into consideration [21,22]. This study addresses a

substantial knowledge gap regarding IR in *Ae. aegypti* by investigating differences in IR across

both space (with four cities included in the study area) and time (with sampling occurring in

three seasons), while also considering both genetic and phenotypic lines of evidence to determine

IR status. Furthermore, assays were conducted with multiple commonly-used insecticides,

allowing for comparisons of effectiveness that can inform local-scale vector-control decision-

making.

In Ecuador, vector control is conducted by field workers of the Ministry of Health (MoH)

in areas at risk of arboviral transmission, as well as focal control in and around homes with

suspected arboviral infections. Interventions to control adult mosquitoes include indoor residual

spraying (IRS) with deltamethrin (pyrethroid) and ultra-low volume (ULV) fumigation with

malathion (organophosphate). Interventions to control juvenile mosquitoes include application of

an organophosphate larvicide (temephos) to containers with standing water, as well as

community mobilization to eliminate larval habitats.

The objectives of this study were to evaluate overall IR status and describe seasonal and

inter-city variability in IR of Ecuadorian populations of *Ae. aegypti*, specifically testing for

susceptibility to insecticides commonly used in mosquito control campaigns by the public health

sector. In recent work in El Oro province, researchers demonstrated resistance to deltamethrin in

phenotypic assays and detected genetic mutations associated with resistance to pyrethroids in *Ae.*

aegypti (Neira, *unpublished data*). Beyond this small-scale study, however, IR monitoring is not

regularly performed and research in this area has been very limited, meaning the prevalence IR

in these *Ae. aegypti* populations is largely unknown. Due to the high morbidity and mortality

associated with the viruses transmitted by *Ae. aegypti*, as well as concerns regarding the cost and

efficacy of vector control efforts, leaders of the Ministry of Health (MoH) have indicated that

there is a critical need for operational research on IR in Ecuador.

Methods

Ethics Statement

This study was part of a longitudinal cohort study. The study protocol was reviewed and

approval by Institutional Review Boards (IRBs) at SUNY Upstate Medical University, the

Human Research Protection Office (HRPO) of the U.S. Department of Defense, the Luis

Vernaza Hospital in Guayaquil, Ecuador, and the Ecuadorean Ministry of Health. Prior to the

start of the study, all adult participants (18 years of age or older) engaged in a written informed

consent process.

Data collection for this project was also conducted with input from local vector control

and public health organizations. Sampling coordinates were stored in a secured database and all

resulting data were pooled to evaluate citywide trends in the analysis, meaning identifiable

individual sites were not shared.

Study sites and sampling period

Field samples of *Ae. aegypti* were collected in four cities in El Oro, a province located on

Ecuador's southern coast with a well-documented history of dengue transmission (Figure 1). The

cities included in this study historically range from high to low dengue case burden. Two sites

were located at sea level along the coast – Machala (3°15'09"S, 79°57'20"W; 6m elevation), a

port city and important center of agribusiness, and Huaquillas (3°28'33"S, 80°13'33"W; 15m

elevation), a town on the border with Peru. These cities have endemic transmission of dengue

and typically see high numbers of annual reported cases. In contrast, Portovelo (3°42'58"S,

79°37'08"W; 645 m elevation) and Zaruma (3°41'31"S, 79°36'47"W; 1,155m elevation) are

both mining towns located further inland and at higher elevations, and have limited

autochthonous transmission. The discrepancies in case burden between the cities translate into

differential mosquito control demands at the municipal level, making our selection of study sites

a diverse backdrop for investigating IR.

Collection of field samples was conducted in 2017 over three sampling periods: Season 1

(February 1 – April 30), Season 2 (May 1 – June 30), and Season 3 (July 1 – August 31).

Seasons were explicitly chosen to collect mosquito eggs at different phases of annual arbovirus

transmission in Ecuador, sampling during the peak (Season 1), decline (Season 2), and low

transmission (Season 3). These collection seasons correspond with historical trends in both

dengue transmission and mosquito densities in El Oro Province [23,24].

Egg collection and mosquito rearing

Ae. aegypti eggs were collected from households in the four cities using ovitraps lined with

oviposition paper [25]. Households were selected from ongoing surveillance study sites where

the ministry of health designated areas of the cities as having high historic risk of dengue. These

houses are distributed across each of the four cities to capture geographic variability, and are part

of a larger study looking at arboviral and vector dynamics in a cohort of homes over 3 years.

Ovitrapped houses for this study were purposely targeted to collect the greatest number of eggs,

so ovitraps were strategically placed in areas known to have a greater abundance of eggs,

representing a subset of the larger household cohort study. Ovitraps are a sensitive means of

identifying the presence of *Ae. aegypti*, especially in areas with low vector densities [26]. Papers

with eggs were collected in the field and transported to the Center for Research on Health in

Latin America (CISeAL) in Quito, Ecuador, where all insect rearing and handling was performed

under standard insectary conditions (28 ± 1 ºC, 80% +/- 10% relative humidity, 12h light; 12h

darkness photoperiod). Egg hatching was achieved by placing the papers in a plastic tray

containing distilled water. Larvae were fed finely ground fish flakes. Upon pupation, mosquito

specimens were placed in cages for adult emergence. To increase the number of adult specimens

available for experimentation, F0 adults were allowed to mate and adult female mosquitoes were

blood fed on mice and allowed to oviposit. Collected eggs (F1 generation) were hatched and

maintained under the aforementioned conditions. Upon pupation, F1 specimens were sorted by

sex and females were used for further experimentation. Males were killed by freezing and

discarded.

Resistance monitoring

We monitored phenotypic resistance to pesticides using the bottle bioassay method as described

by Brogdon and Chan [27]. Briefly, we coated glass bottles with ethanolic-insecticide solutions

at established diagnostic doses: 10 µg / bottle for both deltamethrin and alpha-cypermethrin

(both pyrethroids), and 50 µg / bottle malathion (an organophosphate). After the ethanol had

evaporated from the bottles, groups of 15-25 female mosquitoes were introduced into each bottle

and mortality was recorded at 15-minute intervals. Mosquitoes were considered dead when they

were incapable of flying or maintaining an upright posture on the surface of the bottle. Final

mortality counts were recorded after 30 minutes of exposure. Each bioassay consisted of four

pesticide-coated bottles and one control bottle, coated only with the ethanol diluent.

The number of F1 females procured from field-collected specimens allowed us to perform one to

three bioassay replicates for each city. The selection of pesticides (deltamethrin, alpha-

cypermethrin and malathion) corresponds to pesticides typically used for vector control in

Ecuador.

Genotyping

In addition to the monitoring of phenotypic resistance, we also performed genetic screening for

two mutations of the voltage-gated sodium channel gene (V1016I and F1534C) that are

associated with IR to pyrethroids in *Ae. aegypti*. Both mutations have been previously described

and are reported to alter the transmembrane domain of the voltage-gated sodium channel,

preventing pyrethroids from binding to these sites [28,29]. Previous genotyping work has shown

that *Aedes aegypti* that are homozygous for the isoleucine variant at position 1016 (I1016) are

resistant to pyrethroid treatments, while those that are homozygous for the valine variant

(V1016) allele are susceptible, and those that are heterozygous show intermediate resistance [30].

At position 1534, the cysteine variant (C1534) confers resistance, while the phenylalanine

variant (F1534) is susceptible, and the heterozygote shows intermediate resistance [31].

Following the aforementioned bioassays, DNA was extracted from specimens of known

phenotype (susceptible/resistant to each of the pesticides used for testing). Genomic DNA was

obtained using the Wizard ® Genomic DNA Purification Kit (Promega Corporation, Madison,

WI, USA), following the protocol established by the manufacturer. Quantification of the

concentration (ng/µL) and purity (absorbance index at 260nm / 280 nm) of the obtained DNA

was completed using a Nano-Drop 1000 V3.7 spectrophotometer (Thermo Fisher Scientific Inc.,

Wilmington, DE, USA).

Screening for the V1016I and F1534C mutations was performed as previously described

[28,29], using a Bio-Rad real-time thermocycler CFX96 (Bio-Rad, Hercules, CA, USA). Results

were visualized using CFX Manager Software (Version 3.1, Bio-Rad, Hercules, CA, USA).

Determination of the genotype was possible through the visualization of PCR product melting

curves. In the case of the V1016I mutation, a melting peak at 79ºC corresponded to isoleucine

(I/I – resistant mutant) and a melting peak at 85ºC corresponded to a valine (V/V – susceptible

wild type) (Figure S1). In the case of the F1534C mutation, a melting peak at 85ºC corresponded

to cysteine (C/C, resistant mutant) and a melting peak at 80ºC corresponded to a phenylalanine

(F/F– susceptible wild type) (Figure S2). Genotypic and allelic frequencies were calculated for

both mutations.

Seasonal differences in genotypic and phenotypic frequencies for all four cities were

mapped using ArcMap© version 10.6 (Environmental Systems Research Institute, Redlands,

California, USA) and tested for statistical significance ($\alpha = 0.05$) using Fisher's exact test of

independence in the R software for statistical computing, version 3.4.3 (The Foundation for

Statistical Computing, Vienna, Austria). When significant differences were detected, post-hoc

pairwise Fisher's tests with Bonferroni corrections were then performed to determine which

cities differ in the frequency of resistant genotypes. The same statistical methods were used to

examine phenotypic frequencies of resistance, as determined by bottle bioassays for

deltamethrin, alpha-cypermethrin, and malathion, between cities for each collection season and

in relation to corresponding genotypes.

Results

Resistance monitoring

High levels of IR to deltamethrin, alpha-cypermethrin, and malathion were detected in all

four cities, as determined by bottle bioassays. Machala had the lowest combined mortality

averaged across the three seasons (18.29%, SE±5.75), indicating the highest level of resistance,

followed by Portovelo (31.08%, SE±6.23) and Huaquillas (44.79%, SE±10.87). The mean

mortality rate for Zaruma was 23.67% (SE±17.93) in Season 1, which was the only time period

in which sufficient numbers of *Ae. aegypti* were collected for bioassays at this location.

Significant phenotypic differences in deltamethrin resistance between cities were

detected in all collection seasons (Fisher's exact test p-value<0.001) (Figure 2a). This pattern

was also seen in the post-hoc pairwise tests (Table S1). Mean mosquito mortality upon exposure

to deltamethrin in bottle bioassays ranged from 0.70% (SE±0.6) in Machala to 18.99% (SE±0) in

Huaquillas in Season 1, from 7.09% (SE±4.19) in Portovelo to 16.88% (SE±7.51) in Huaquillas

in Season 2, and from 0.37% (SE±0.37) in Machala to 56% (SE±2.49) in Huaquillas in Season 3,

with no mortality observed in control groups (Figure 2a).

Mean mosquito mortality upon exposure to alpha-cypermethrin in bottle bioassays ranged

from 59.41% (SE±1.49) in Machala to 59.24 (SE±13.86) in Zaruma in Season 1, from 12.1%

(SE±0.42) in Machala to 17.27 (SE±0.33) in Portovelo in Season 2, and from 0.38% (SE±0.38)

in Machala to 11.39 (SE±1.03) in Portovelo in Season 3, with no mortality observed in control

groups (Figure 2b). Significant differences in mean mortality rates between cities were detected

in seasons two and three (Fisher's exact test p value<0.001) (Figure 2b), with Machala having

the highest proportion of phenotypically resistant specimens in these two seasons (Table S2).

Collection counts were insufficient for alpha-cypermethrin bioassays in Portovelo for Season 1,

Huaquillas for Seasons 1 and 3, and Zaruma for Seasons 2 and 3.

Mean mosquito mortality when exposed to malathion in the bottle bioassays ranged from

5.07% (SE±0.35) in Machala to 3.03% (SE±1.52) in Zaruma in Season 1, from no mortality in

Machala to 1.45% (SE±1.45) in Portovelo in Season 2, and from 0.78% (SE±0.78) in Machala to

2.86% (SE±0.08) in Portovelo in Season 3, with no mortality observed in control groups.

Statistically significant differences in Malathion resistance were only detected in season two

between Machala and Portovelo (Fisher's exact test p value < 0.001). Due to low sample sizes,

bioassays could not be conducted in Portovelo for Season 1, Zaruma for Seasons 2 and 3, and

Huaquillas for all three seasons, and therefore further comparisons were not feasible.

Genotyping

Observed frequencies varied significantly between cities only in Season 1 for both

V1016I and F1534C alleles (Fisher's exact test p value < 0.001) (Figure 3). Pairwise post-hoc

analysis revealed significant differences in the frequencies of genotypes I/I (mutant) and V/I

(heterozygous) for the V1061I gene, with Huaquillas having a significantly higher frequency of

heterozygotes compared to Machala and Portovelo (Post-hoc Fisher's exact test p value < 0.001

and 0.03, respectively, Table S3). Portovelo and Zaruma differed in the frequency of I/I

(mutant) and V/V (wild type) genotypes, with Zaruma having a significantly higher frequency of

wild type mosquitoes during Season 1 (Post-hoc Fisher's exact test p value < 0.05, Table S3).

Genotypic frequencies of C/C (mutant) and F/C (heterozygous) genotypes of the F1534C

resistance gene differed across cities in Season 1, with Zaruma having a significantly higher

proportion of the heterozygous genotype than Huaquillas and Machala (Post-hoc Fisher's exact

test p value < 0.02 and < 0.001, respectively, Table S4). Although differences in the frequencies

of F1534C genotypes also varied significantly in the Season 3, conservative post-hoc analysis

did not reveal any significant pairwise relationships (Table S4).

There were statistically significant associations between the resistance genotypes and

phenotypic resistance results for 632 *Aedes aegypti* that were genotyped and subjected to the

pyrethroid (deltamethrin or alpha-cypermethrin) bottle bioassays. Due to low sample sizes, we

pooled pyrethroid assay results (Table 1), and present the separated analyses in supplemental

information (Tables S5, S6). For mutation V1016I, while the majority of the resistant

individuals (96.4%) presented the mutant or heterozygous genotype, 12 (3.6%) phenotypically

resistant individuals presented the wild type (V/V) genotype that is typically associated with

susceptibility to pyrethroids, and 40 (13.5%) susceptible individuals presented the genotype that

typically confers resistance. Similarly, at the F1534C locus, 276 (93.2%) individuals with the

mutant (C/C) genotype, typically associated with resistance, were susceptible to pyrethroid

treatments. Due to low sample sizes of the wildtype (F/F) for both pyrethroids, separate Fisher's

exact tests were not conducted (Table S6).

Discussion

In this study, exposure to diagnostic doses of deltamethrin, alpha-cypermethrin and

malathion resulted in mortality rates below 80% in all populations tested, regardless of collection

season. Based on the World Health Organization´s recommendations for assessing the

significance of detected resistance [27], our results suggest that these *Ae. aegypti* populations

should be considered as resistant to all the insecticides considered in our study. Furthermore, the

results of bioassays for malathion susceptibility are of particular interest, as they indicate that

these populations are extremely resistant to malathion, with no population showing more than

5.07% mortality when exposed to this pesticide, and populations from Machala reaching values

as low as 0% mortality during Season 2.

The molecular characterization shows that the resistant-associated mutant alleles V1016I

and F1534C are present in all the locations studied. In particular, allele F1534C was present at

very high frequencies (close to 1) in all the locations studied and across all three seasons, similar

to what was seen in recent work in Mexico [32]. This suggests that this gene has been subjected

to selective pressures in the past and is approaching or has reached fixation in these populations.

Both of these mutations typically confer resistance to dichlorodiphenyltrichloroethane (DDT) as

well as pyrethroids [33,34], meaning selection for these alleles likely began with earlier

widespread usage of DDT (which was used by the MoH until 1996 [35]) and has persisted as

pyrethroids became more commonly used [36]. Significant distinctions in genotypic and

phenotypic frequencies were not detected for many cities in later seasons.

It is worth noting that for some locations and seasons, periods of low vector densities in

the field caused the collection of low numbers of mosquito eggs. This scarcity of F0 individuals

translated into missing data for some cities and/or very low frequency counts in some categories,

thus providing a limited basis for making quantitative comparisons.

The detection of resistance to organophosphate and pyrethroid insecticides in Ecuador is

consistent with broader regional trends identified in recent years. In neighboring Peru, *Ae.*

aegypti strains have been shown to be resistant to multiple organophosphates and pyrethroids

[37], while resistance to commonly used pyrethroids has been detected in Colombia, though the

levels of resistance varied throughout the country [19]. However, Colombian populations remain

broadly susceptible to malathion, in contrast to the widespread resistance seen in the Ecuadorian

Ae. aegypti in this study [38]. Throughout the rest of the Americas, pyrethroid resistance appears

to be broadly distributed, particularly in Brazil and French Guiana, though some populations in

Costa Rica, Panama, and northern Colombia are still susceptible [8].

The temporal and spatial variability in the results from the bioassays highlight the

importance of regularly conducting IR monitoring across multiple locations to understand the

true extent of IR and make appropriate vector control decisions. For example, in Machala, the

mortality rate of *Ae. aegypti* treated with alpha-cypermethrin decreased from season one to

season three, indicating this population was becoming less susceptible throughout the course of

the study. By contrast, the mortality rate associated with the deltamethrin assays on populations

from Huaquillas increased from season one to season three, meaning this population was

becoming more susceptible over time. Continued work in this area could determine if these

trends are due to seasonal fluctuations, local-scale movements of *Ae. aegypti* populations with

varying levels of resistance, or long-term, inter-annual trends. There were also statistically

significant differences in mortality rates across the cities for the deltamethrin bioassays in all

three seasons and in two of the three seasons for the cypermethrin bioassays. Considering this

variability in the resistance phenotypes found within a single province of Ecuador, organizations

involved in decision-making about insecticide applications should be cautioned against inferring

the IR status of one *Ae. aegypti* population based on the status of populations in neighboring

municipalities.

To better contextualize this work for appropriate vector-control decision-making, the

relationships between genotypes, IR bioassay results, and actual IR status in the field should be

considered. While the V1016I and F1534C mutations are known markers of pyrethroid

resistance in *Ae. aegypti*, the results of this study showed that the genotypes were not perfectly

predictive of resistance phenotypes. In the recorded frequencies of genotypes versus bioassay

outcomes, resistant and susceptible phenotypes were observed for each genotype, although the

resistant phenotype was still statistically associated with the mutant genotypes for both genes.

This is likely due to other IR mechanisms, such as metabolic detoxification processes [16], which

were not considered in the current study, but could influence IR status in these populations.

Additionally, factors such as temperature, larval nutrition, larval density and age have been

shown to influence insecticide susceptibility in *Aedes* mosquitoes, leading to discrepancies

between bioassay results and the actual outcomes of insecticide treatments in the field [39].

Further work on IR in *Aedes* populations could identify and possibly reconcile differences

between results from the laboratory and the field. Additionally, to comprehensively evaluate IR

status, future studies should also investigate the effectiveness of larvicides, such as temephos,

which is commonly used in this study area, as well as *Bacillus thuringensis israelensis* (Bti).

Geographic methods, particularly spatial statistics and modelling, are well suited for

understanding the patterns and drivers of IR at meso- and local scales. Employing these

approaches can lead to more targeted, efficient, and sustainable vector control efforts. Future

research in this area should continue to explore the spatial and temporal variability in IR among

Ae. aegypti populations. Recent work in Yucatan, Mexico, demonstrated that IR levels could

vary significantly across neighborhoods within the same city [21]. Additionally, work on the

temporal dynamics of resistance could be beneficial for vector control decision-making. For

example, in a study on pyrethroid-resistant, field-derived *Ae. aegypti*, researchers demonstrated

that susceptibility to pyrethroids could be restored within ten generations when the selective

pressure of regular insecticide treatments was removed [22]. While this experiment was

conducted in a controlled environment, similar work within the context of urban environments,

such as the four cities included in this study, could help calibrate timing of insecticide class

rotations, allowing for better long-term management of susceptibility.

In conclusion, the *Ae. aegypti* collected in these four cities in Ecuador showed varying

levels of resistance to the insecticides tested, and these measures typically changed over the

course of the three seasons during which sampling took place. Regular IR monitoring should be

conducted as long as insecticide applications remain an integral component of vector control

activities, particularly in areas where these operations are deployed to control arbovirus

transmission. Beyond this monitoring process, appropriate alternative management strategies

should be deployed when IR is detected. These strategies can include biological control and

community mobilization to reduce *Ae. aegypti* breeding sites.

Acknowledgments:

We are grateful to Naveed Heydari, the Ministry of Health of Ecuador, and our colleagues who

facilitated data collection in the field.

References

1. Dzul-Manzanilla F, Ibarra-Lopez J, Bibiano Marin W, Martini-Jaimes A, Leyva JT, Correa-
Morales F, et al. Indoor Resting Behavior of Aedes aegypti (Diptera: Culicidae) in
Acapulco, Mexico. J Med Entomol. 2017;54: 501–504. doi:10.1093/jme/tjw203

2. Chatchen S, Sabchareon A, Sirivichayakul C. Serodiagnosis of asymptomatic dengue
infection. Asian Pac J Trop Med. 2017;10: 11–14. doi:10.1016/j.apjtm.2016.12.002

3. Pan American Health Organization / World Health Organization. Number of reported cases
of Chikungunya Fever in the Americas, by Country or Territory 2016. Washington, D.C.;

4. Pan American Health Organization / World Health Organization. Zika suspected and
confirmed cases reported by countries and territories in the Americas: Cumulative cases,
2015-2016. Washington, D.C.;

5. Heydari N, Larsen DA, Neira M, Beltrán Ayala E, Fernandez P, Adrian J, et al. Household
dengue prevention interventions, expenditures, and barriers to Aedes aegypti control in
Machala, Ecuador. Int J Environ Res Public Health. 2017;14: 196.

6. Rodriguez-Morales AJ, Villamil-Gómez WE, Franco-Paredes C. The arboviral burden of
disease caused by co-circulation and co-infection of dengue, chikungunya and Zika in the
Americas. Travel Med Infect Dis. 2016;14: 177–179.

7. Yakob L, Walker T. Zika virus outbreak in the Americas: the need for novel mosquito
control methods. Lancet Glob Health. 2016;4: e148–e149.

8.   Moyes CL, Vontas J, Martins AJ, Ng LC, Koou SY, Dusfour I, et al. Contemporary status
of insecticide resistance in the major Aedes vectors of arboviruses infecting humans. PLoS
Negl Trop Dis. 2017;11: e0005625.

9.   Hemingway J, Ranson H. Insecticide Resistance in Insect Vectors of Human Disease. Annu
Rev Entomol. 2000;45: 371.

10.   Emmanuel Chanda, Edward K. Thomsen, Mulenga Musapa, Mulakwa Kamuliwo, William
G. Brogdon, Douglas E. Norris, et al. An Operational Framework for Insecticide Resistance
Management Planning. Emerg Infect Dis J. 2016;22: 773. doi:10.3201/eid2205.150984

11.   Cohen JM, Smith DL, Cotter C, Ward A, Yamey G, Sabot OJ, et al. Malaria resurgence: a
systematic review and assessment of its causes. Malar J. 2012;11: 122.

12.   IRAC. Prevention and management of insecticide resistance in vectors and pests of public
health importance. Insecticide Resistance Action Committee; 2011.

13.   Zaim M, Guillet P. Alternative insecticides: an urgent need. Trends Parasitol. 2002;18:
161–163.

14.   Rose RI. Pesticides and public health: integrated methods of mosquito management. Emerg
Infect Dis. 2001;7: 17.

15.   Gray L, Florez SD, Barreiro AM, Vadillo-Sánchez J, González-Olvera G, Lenhart A, et al.
Experimental evaluation of the impact of household aerosolized insecticides on pyrethroid
resistant Aedes aegypti. Sci Rep. 2018;8: 12535.

16.   Liu N. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research
Directions. Annu Rev Entomol. 2015;60: 537–559. doi:10.1146/annurev-ento-010814-
020828

17.   Nicolopoulou-Stamati P, Maipas S, Kotampasi C, Stamatis P, Hens L. Chemical Pesticides
and Human Health: The Urgent Need for a New Concept in Agriculture. Front Public
Health. 2016;4: 148. doi:10.3389/fpubh.2016.00148

18.   Marcombe S, Mathieu RB, Pocquet N, Riaz M-A, Poupardin R, Sélior S, et al. Insecticide
resistance in the dengue vector Aedes aegypti from Martinique: distribution, mechanisms
and relations with environmental factors. PLoS One. 2012;7: e30989.

19.   Fonseca□González I, Quiñones ML, Lenhart A, Brogdon WG. Insecticide resistance status
of Aedes aegypti (L.) from Colombia. Pest Manag Sci. 2011;67: 430–437.

20.   Rawlins SC. Spatial distribution of insecticide resistance in Caribbean populations of Aedes
aegypti and its significance. Rev Panam Salud Pública. 1998;4: 243–251.

21.   Deming R, Manrique-Saide P, Medina Barreiro A, Cardeña EUK, Che-Mendoza A, Jones
B, et al. Spatial variation of insecticide resistance in the dengue vector Aedes aegypti

presents unique vector control challenges. Parasit Vectors. 2016;9: 67. doi:10.1186/s13071-
016-1346-3

22. Grossman MK, Uc-Puc V, Rodriguez J, Cutler DJ, Morran LT, Manrique-Saide P, et al.
Restoration of pyrethroid susceptibility in a highly resistant Aedes aegypti population. Biol
Lett. 2018;14: 20180022.

23. Stewart Ibarra AM, Lowe R. Climate and non-climate drivers of dengue epidemics in
southern coastal Ecuador. Am J Trop Med Hyg. 2013;88: 971–981. doi:10.4269/ajtmh.12-
0478

24. Stewart Ibarra AM, Ryan SJ, Beltrán E, Mejía R, Silva M, Muñoz Á. Dengue Vector
Dynamics (Aedes aegypti) Influenced by Climate and Social Factors in Ecuador:
Implications for Targeted Control. PLOS ONE. 2013;8: e78263.

25. Fay RW, Eliason DA. A preferred oviposition site as a surveillance method for Aedes
aegypti. Mosq News. 1966;26: 531–535.

26. Focks DA. A review of entomological sampling methods and indicators for dengue vectors.
Geneva WHO. 2003; Available: http://203.90.70.117/PDS_DOCS/B0219.pdf#page=233

27. Brogdon W, Chan A. Guidelines for evaluating insecticide resistance in vectors using the
CDC bottle bioassay/methods in Anopheles research. CDC Atlanta USA CDC Tech Rep.
2010;

28. Saavedra-Rodriguez K, Beaty M, Lozano-Fuentes S, Denham S, Garcia-Rejon J, Reyes-
Solis G, et al. Local Evolution of Pyrethroid Resistance Offsets Gene Flow Among Aedes
aegypti Collections in Yucatan State, Mexico. Am J Trop Med Hyg. 2015;92: 201–209.
doi:10.4269/ajtmh.14-0277

29. Yanola J, Somboon P, Walton C, Nachaiwieng W, Somwang P, Prapanthadara L.
High☐throughput assays for detection of the F1534C mutation in the voltage☐gated
sodium channel gene in permethrin☐resistant Aedes aegypti and the distribution of this
mutation throughout Thailand. Trop Med Int Health. 2011;16: 501–509.

30. Saavedra☐Rodriguez K, Urdaneta☐Marquez L, Rajatileka S, Moulton M, Flores AE,
Fernandez☐Salas I, et al. A mutation in the voltage☐gated sodium channel gene associated
with pyrethroid resistance in Latin American Aedes aegypti. Insect Mol Biol. 2007;16:
785–798.

31. Linss JGB, Brito LP, Garcia GA, Araki AS, Bruno RV, Lima JBP, et al. Distribution and
dissemination of the Val1016Ile and Phe1534Cys Kdr mutations in Aedes aegypti Brazilian
natural populations. Parasit Vectors. 2014;7: 25.

32. Lopez☐Monroy B, Gutierrez☐Rodriguez SM, Villanueva☐Segura OK, Ponce☐Garcia G,
Morales☐Forcada F, Alvarez LC, et al. Frequency and intensity of pyrethroid resistance
through the CDC bottle bioassay and their association with the frequency of kdr mutations
in Aedes aegypti (Diptera: Culicidae) from Mexico. Pest Manag Sci. 2018;

33.   Brito LP, Linss JG, Lima-Camara TN, Belinato TA, Peixoto AA, Lima JBP, et al.
465   Assessing the effects of Aedes aegypti kdr mutations on pyrethroid resistance and its fitness
466   cost. PloS One. 2013;8: e60878.

467   34.   Brengues C, Hawkes NJ, Chandre F, McCarroll L, Duchon S, Guillet P, et al. Pyrethroid
468   and DDT cross☐resistance in Aedes aegypti is correlated with novel mutations in the
469   voltage☐gated sodium channel gene. Med Vet Entomol. 2003;17: 87–94.

470   35.   Krisher LK, Krisher J, Ambuludi M, Arichabala A, Beltrán-Ayala E, Navarrete P, et al.
471   Successful malaria elimination in the Ecuador–Peru border region: epidemiology and
472   lessons learned. Malar J. 2016;15: 573. doi:10.1186/s12936-016-1630-x

473   36.   Dick OB, San Martín JL, Montoya RH, del Diego J, Zambrano B, Dayan GH. The history
474   of dengue outbreaks in the Americas. Am J Trop Med Hyg. 2012;87: 584–593.

475   37.   Rodríguez MM, Bisset JA, Fernández D. Levels of insecticide resistance and resistance
476   mechanisms in Aedes aegypti from some Latin American countries. J Am Mosq Control
477   Assoc. 2007;23: 420–429.

478   38.   Ocampo CB, Salazar-Terreros MJ, Mina NJ, McAllister J, Brogdon W. Insecticide
479   resistance status of Aedes aegypti in 10 localities in Colombia. Acta Trop. 2011;118: 37–
480   44.

481   39.   McAllister JC, Godsey MS, Scott ML. Pyrethroid resistance in Aedes aegypti and Aedes
482   albopictus from Port-au-Prince, Haiti. J Vector Ecol J Soc Vector Ecol. 2012;37: 325–332.
483   doi:10.1111/j.1948-7134.2012.00234.x

 Figures and Tables

Table 1. Occurrences of V1016I and F1534C Genotypes compared to pyrethroid-resistant

phenotype. Fisher's tests indicated significant associations between genotypes and phenotypes

for resistance.

Figure 1. Map of the four study site cities: Machala, Huaquillas, Portovelo, and Zaruma, in

southern Ecuador. Panel A shows the cities within southern Ecuador, on elevational relief;

panel B illustrates Ecuador's location along the northern Pacific coast of South America, and

panel C places panel A (red bounding box) within Ecuador's southern coastal region.

Figure 2. Bar plots of CDC bottle bioassay mortality for deltamethrin (A) alpha-

cypermethrin (B), and malathion (C). Note different axis height for (C). Asterisks indicate

Fisher's exact test showed statistically significant differences across cities for that season.

Insufficient numbers of field-collected mosquitos for bioassay within a city in a given season are

indicated by 'x'.

Figure 3. Genotypic frequencies of the V1016C and F1534C alleles at the four study site

locations during Season 1 (Feburary 1-April 30, 2017).

Supporting information

Table S1: Post-hoc Fisher's exact test p-values for Deltamethrin resistance between cities. Significant difference are denoted in bold.

Table S2: Post-hoc Fisher's exact test *p*-values for Alpha-Cypermethrin resistance between cities in the season two collection period. Significant difference are denoted in bold.

Table S3: Post-hoc Fisher's exact test *p*-values for genotype V1016I in season 1, with comparisons in genotype frequencies made between the cities of Huaquillas (H), Machala (M), Portovelo (P), and Zaruma (Z). Significant difference are denoted in bold.

Table S4: Post-hoc Fisher's exact test *p*-values for genotype F1534C in season 1, with comparisons in genotype frequencies made between the cities of Huaquillas (H), Machala (M), Portovelo (P), and Zaruma (Z). Significant difference are denoted in bold.

Table S5. V1016I and F1534C Genotypes compared to deltamethrin-resistant phenotype

Table S6. V1016I and F1534C Genotypes compared to alpha-cypermethrin resistant phenotype

Figure S1: Genotypic characterization of the V1016I mutation. Dissociation curves representing the susceptible (wild type) homozygous (A), heterozygous (B), and resistant (mutant) heterozygous (C) genotypes.

531 **Figure S2: Genotypic characterization of the F1534C mutation.** Dissociation curve

532 representing the wild type (susceptible) homozygous (A), heterozygous (B), and mutant (C)

533 genotypes.

535 **Table 1.**

V1016I		I/I (mutant)	V/I (heterozygote)	V/V (wild type)	*p*-value*
	Resistant	187	137	12	**<0.001***
	Susceptible	40	172	84	
F1534C		C/C (mutant)	F/C (heterozygote)	F/F (wild type)	
	Resistant	328	7	1	**0.02***
	Susceptible	276	17	3	

536 * Fisher's Exact Test

538 Figure 1

540   Figure 2

542   Figure 3

546   **Table S1.**

Season	City	Huaquillas	Machala	Portovelo
1	Machala	**< 0.001***		
	Portovelo	**0.002***	1	
	Zaruma	**0.008***	**< 0.001***	0.75
2	Machala	**0.006***		
	Portovelo	**< 0.001***	1	
3	Machala	**< 0.001***		
	Portovelo	**< 0.001***	**< 0.001***	

549   **Table S2.**

City	Huaquillas	Machala
Machala	**0.01***	
Portovelo	**< 0.001***	1

552   **Table S3.**

City:City*	I/I : V/I	I/I : V/V	V/I : V/V
H:M	**< 0.001***	1.00	1.00
H:P	**0.03***	0.94	1.00
H:Z	0.44	1.00	0.32
M:P	1.00	1.00	1.00
M:Z	0.09	0.08	1.00
P:Z	1.00	**0.05***	1.00

553   *H – Huaquillas, M-Machala, P-Portovelo, Z-Zaruma

555 **Table S4.**

City:City*	C/C : F/C	C/C : F/F	F/C : F/F
H:M	1.00	1.00	1.00
H:P	1.00	1.00	1.00
H:Z	**0.02***	1.00	1.00
M:P	1.00	1.00	1.00
M:Z	**< 0.001***	1.00	1.00
P:Z	0.18	1.00	1.00

556 *H – Huaquillas, M-Machala, P-Portovelo, Z-Zaruma

558 **Table S5.**

V1016I		I/I (mutant)	V/I (heterozygote)	V/V (wild type)	*p*-value*
	Resistant	107	79	10	**<0.001***
	Susceptible	17	88	52	
F1534C		C/C (mutant)	F/C (heterozygote)	F/F (wild type)	
	Resistant	190	5	1	0.10
	Susceptible	144	10	3	

559 *Fisher's Exact Test

561 **Table S6.**

V1016I		I/I (mutant)	V/I (heterozygote)	V/V (wild type)	*p*-value*
	Resistant	80	58	2	**<0.001***
	Susceptible	23	84	32	
F1534C		C/C (mutant)	F/C (heterozygote)	F/F (wild type)	
	Resistant	138	2	0	0.10
	Susceptible	132	7	0	

562 *Fisher's Exact Test

563   **Figure S1**

566　**Figure S2**