Principles of Meiotic Chromosome Assembly

*Stephanie A. Schalbetter[1], *Geoffrey Fudenberg[2], Jonathan Baxter[1], Katherine S. Pollard[2,3,4], and Matthew J. Neale[1]

* SAS & GF contributed equally to this work

1. Genome Damage and Stability Centre, School of Life Sciences, University of Sussex, UK

2. Gladstone Institutes for Data Science and Biotechnology, San Francisco, USA

3. Department of Epidemiology & Biostatistics, Institute for Human Genetics, Quantitative Biology Institute, and Institute for Computational Health Sciences, University of California, San Francisco, California, USA.

4. Chan-Zuckerberg Biohub, San Francisco, California, USA.

Abstract

During meiotic prophase, chromosomes organise into a series of chromatin loops emanating from a proteinaceous axis, but the mechanisms of assembly remain unclear. Here we elucidate how this elaborate three-dimensional chromosome organisation is underpinned by genomic sequence in *Saccharomyces cerevisiae*. Entering meiosis, strong cohesin-dependent grid-like Hi-C interaction patterns emerge, reminiscent of mammalian interphase organisation, but with distinct regulation. Meiotic patterns agree with simulations of loop extrusion limited by barriers, yet are patterned by convergent transcription rather than binding of the mammalian interphase factor, CTCF, which is absent in *S. cerevisiae*—thereby both challenging and extending current paradigms of local chromosome organisation. While grid-like interactions emerge independently of meiotic chromosome synapsis, synapsis itself generates additional compaction that matures differentially according to telomere proximity and chromosome size. Collectively, our results elucidate fundamental principles of chromosome assembly and demonstrate the essential role of cohesin within this evolutionarily conserved process.

Introduction and Results

During meiosis, eukaryotic chromosomes are broken, repaired and paired with their homologs followed by two rounds of segregation—a series of events accompanied by dynamic structural changes of the chromosomes. Most prominent is the paired arrangement of pachytene chromosomes into a dense array of chromatin loops emanating from proteinaceous axes linked by a central core, the synaptonemal complex (SC), which is highly conserved across eukaryotes[1,2]. In *S. cerevisiae*, structural components include the meiotic cohesin kleisin subunit, Rec8[3], the transverse filament, Zip1[4], the axial/lateral elements, Hop1 and Red1[5,6], and the pro-DSB factors Rec114-Mei4-Mer2 (RMM)[7,8]. Much of our understanding of meiotic chromosome structure has been deduced from a combination of electron microscopy, immunofluorescence microscopy, and the genome-wide patterns of protein localisation determined by ChIP. However, the link between key meiotic protein complexes, chromosome conformation, and genomic sequence remains uncharacterized.

Chromosome conformation capture (3C) techniques generate maps of pairwise contact frequencies that are snapshots of chromosome organisation. 3C methods were originally applied to assay chromosome conformation in *S. cerevisiae*, including during meiosis[9]. Now they are widely used across a range of organisms and cellular contexts to link 3D organisation directly with genomic sequence[10], revealing important roles of the Structural Maintenance of Chromosomes (SMCs) cohesin and condensin in genomic organization[11,12], where they likely mediate chromosome compaction via the process of loop extrusion[13]. Here we return to yeast meiosis to interrogate genome-wide chromosome organisation by Hi-C, elucidate mechanisms of chromosome assembly, and define the role of key meiotic chromosome components, including cohesin and the SC.

Starting with a synchronized G1 population we analysed timepoints encompassing DNA replication, meiotic prophase and both meiotic divisions (**Fig. 1a,b,c, Extended Data Fig. 1a,b,c**). In G1, we detect strong centromere clustering (**Fig. 1a,d**) and folding back of the arms at the centromeres (**Fig. 1a, Supplementary Fig. 2**), characteristic of a Rabl conformation[9,14]. During meiosis, centromere clustering is transiently dissolved (3-5h, **Fig. 1a,d, Extended Data Fig. 1a**); this coincides with a global decrease in inter-chromosomal contact frequency at mid-prophase, reflecting chromosome individualisation. Subtelomeric clustering also decreases during meiotic

prophase (**Fig. 1a,d, Extended Data Fig. 1a**), with no evidence for the transient telomeric

bouquet conformation, consistent with prior microscopic analyses[15].

Entering meiosis, contact frequency versus distance, *P(s),* curves display a shoulder, consistent

with the linear compaction of chromosome arms increasing due to *cis*-loop formation (2-4h, **Fig.**

1e, Extended Data Fig. 1d, e.g. as defined[16]; for review[13]). This change in *P(s)* is reminiscent of

the SMC-dependent changes observed via Hi-C during mitosis across species[17–21]. Compaction

coincides with meiotic prophase I and the formation of the SC at pachytene, and is lost at later

stages (**Fig. 1e, Extended Data Fig. 1d**).

To study meiotic chromosome conformation in more detail, and to eliminate cell-to-cell

heterogeneity (**Fig. 1b,c**), we enriched for pachytene cells in subsequent experiments by

inactivating Ndt80, a transcription factor required for exit from meiotic prophase[22]. *ndt80Δ* cells

entered meiosis synchronously, assessed by bulk DNA replication (**Fig. 1f**), but do not initiate

the first nuclear division[22]. Similar to the wild type prophase population (3-5h), but likely

accentuated by the increased homogeneity, Hi-C maps of pachytene-enriched cells displayed

total loss of centromere clustering (**Extended Data Fig. 2**) and dramatic chromosome arm

compaction (**Fig. 1e**). Analysing compaction in more detail, shorter chromosomes (**Extended**

Data Fig. 1e) and, in particular, shorter chromosome arms (**Fig. 1h, Extended Data Fig. 1f**),

displayed elevated contact frequency at short genomic separations, and an earlier shoulder,

apparently arising from distinct behavior of subtelomeric and subcentromeric regions (**Fig. 1h,**

Extended Data Fig. 1g). Moreover, distinct *P(s)* for chromosomes with different length arms

(**Extended Data Fig. 1h**) suggests that the centromere can insulate the process that leads to

differences between arms. In agreement with this, compaction is interrupted at centromeres in Hi-

C maps (**Fig. 1a**, **Extended Data Fig. 2b**).

Zooming in to consider within-arm organization revealed punctate grid-like Hi-C interactions

between pairs of loci during prophase (**Fig. 2a**), particularly prominent in *ndt80Δ* (**Fig. 2a,b**). Such

focal meiotic patterns are more prominent than reported previously[23]—resembling peaks between

CTCF sites[24] rather than topological domains[25,26] detected in mammalian interphase Hi-C maps—

and likely arise from a heterogeneous mixture of 'transitive' interactions and 'skipping' of peak

bases (**Fig. 2c**).

Genomic regions underlying the punctate Hi-C interactions display a remarkable visual (**Fig.**
2a,b), and quantitative (**Fig. 2d-g**), correspondence with previously characterized sites of high
Rec8 occupancy[27]. At pachytene, Rec8 sites display elevated cis/total contact frequencies (**Fig.**
2c), enriched contact frequency (**Fig. 2e,f**), and evidence of insulation (**Fig. 2g**)—features that
correlate with Rec8 occupancy measured by ChIP (**Fig. 2a, lower**). In wild type cells, Rec8-Rec8
interactions became visible in early prophase (2h), peaked at mid prophase (4h), and were
especially prominent in the homogenous *ndt80Δ* cell population (**Fig. 2a-g, Extended Data Fig.**
4b,c). Importantly, Rec8-Rec8 enrichments are strongest between adjacent sites, decrease
between non-adjacent sites with increasing genomic separation, and are absent in *trans*
(**Extended Data Fig. 4b,c**). As for enrichments between CTCF sites in mammalian interphase[28],
these observations argue that a cis-acting process generates such focal interactions in meiosis.
106
Rec8 is a central component of the meiotic chromosome axis[3]. Assaying a *rec8Δ* mutant enabled
us to determine that Rec8 is absolutely required for the emergence of the grid-like Hi-C patterns
present in meiosis (**Fig. 2a,b**). Moreover, *rec8Δ* cells completely lose the shoulder in *P(s)*,
indicative of a dramatic loss of arm compaction (**Fig. 2b, Extended Data Fig. 4a**), similar to that
caused by depletion of SMCs in diverse contexts[17,19,21,29–32]. Instead of assembling an axis of
loops, *rec8Δ* cells appear to be caught in a state with highly clustered telomeres (**Extended Data**
Fig. 4d, Extended Data Fig. 3), consistent with previous observations by microscopy[33,34].
Moreover, in *rec8Δ* cells *cis* contact frequency is reduced (**Fig. 2d**), similar to G1 cells, and
cis/total no longer correlates with Rec8 occupancy. Instead, *rec8Δ* cis/total displays a decreasing
trend along chromosome arms, likely due to persistent telomere clustering (**Extended Data Fig.**
4d).
118
To test how compaction and grid-like interaction patterns could jointly emerge in meiosis, we
developed polymer simulations (**Fig. 3a, Methods**) similar to those used to successfully describe
the assembly of TADs in mammalian interphase chromosomes[13]. Importantly, these simulations
employ the *cis*-acting process of loop extrusion, where extruders form progressively larger
chromatin loops, unless impeded by adjacent extruders or barrier elements (**Fig. 3a**). Extrusion
dynamics are controlled by parameters dictating the *processivity* (average loop size) and
separation (number of active extruders), as well as the *strength* of barriers (**Methods**). Because
the accumulation of Rec8 at ChIP-seq sites[27] concomitant with convergent transcription[35] is
indicative of barriers to extrusion[28], we positioned bi-directional barriers at Rec8 sites.
128

Simulations were used to explore variations in loop extrusion dynamics to determine whether
specific parameter combinations are able to generate Hi-C maps that agree with experimental
observation (**Fig. 3, Methods**). Models with excellent fits were identified in which ~65% of the
genome is covered by extruded loops (**Fig. 3b,c, Extended Data Fig. 5**)—a far denser array than
present in *S. cerevisiae* mitosis[21], but still less compact than human mitotic cells[20]. Even though
extrusion can generate compaction independently of barriers (**Fig. 3d**), an intermediate barrier
strength is essential to match the grid-like patterns observed experimentally (**Fig. 3b**). Despite
the simplifying assumptions, simulated chromosomes displayed many features observed
experimentally: (i) chromosomes fold into a loose polymer brush[3,36,37], with a Rec8-rich core[3] (**Fig.**
3f, Extended Data Fig. 5 a); (ii) a grid-like interaction pattern naturally emerges in simulated Hi-
C maps (**Fig. 3d**); (iii) importantly, because loop extrusion is a *cis*-acting process, pairs of Rec8
sites at increasing separations naturally have lower contact frequency (**Fig. 3e**).

Simulations also highlight the stochasticity of loop positions across the cell population, with most
barriers (73%) unoccupied by an extruder, and extruders paused with barrier elements on both
sides only a minority of the time (15%) in the best fitting models (**Extended Data Fig. 6 c**).
Because of this, the majority (65%) of extruded loops cross over Rec8 sites, consistent with an
average loop size roughly twice the average distance between Rec8 ChIP peaks (26 kb versus
12 kb, **Extended Data Fig. 6 d**), and remarkably consistent with estimates made using EM
(~20kb[36]). Most strikingly—despite the prominence of Rec8-dependent grid-like features in the
experimental data (**Fig. 2c**)—our simulations indicate that Rec8 sites are not always occupied by
extruding cohesins and thus are present at the meiotic chromosome core in only a subset of cells,
as inferred previously[38].

The range of loop extrusion parameters we explored encompasses the situation where Rec8 sites
always halt extrusion and *cis*-loops are formed between each consecutive Rec8 site. However,
simulations with these parameters have quantitatively poor fits with experimental maps (**Fig. 3d-**
e, ii): the bend in *P(s)* comes too early to recapitulate experimental *P(s)*, and Rec8-Rec8 contacts
are much too strong. The poor fit of such 'direct-bridging' simulations underscores the conclusion
that only a fraction of Rec8 sites are occupied in a given cell, and argues that cohesin-dependent
cis-loops must link regions that are not primary Rec8 binding sites in order to provide compaction
without making Rec8-Rec8 enrichments overly strong.

A crucial prediction of our loop extrusion simulations is that depletion of extruders in meiosis would
lead to both decompaction and loss of the grid-like pattern of Hi-C interactions. When we repeated
our fitting procedure for *rec8Δ*, the best fits were for simulations with either no, or very few,
extruded loops (**Extended Data Fig. 5e**). The lack of compaction in these simulations (**Extended
Data Fig. 5a**) is consistent with previous EM showing decompacted chromatids in *rec8Δ*[3]. Such
joint consistency between Hi-C and imaging data further supports loop extrusion as a mechanism
underlying assembly of the cohesin-rich core and contributing to chromosomal compaction in
meiosis. Our simulations also open the possibility that overly-shortened axes observed upon
Wapl[39,40] and Pds5[41] depletion may reflect heightened extruder processivity[42] upon which
shortened SCs are assembled, and predict that such perturbations would cause a rightward shift
in the *P(s)* shoulder measured via Hi-C (**Extended Data Fig. 5c**).

To investigate how homologue synapsis affects chromosome conformation, we assayed
pachytene cells in the absence of Zip1, the transverse filament of the SC[4], and Hop1, an axial
element required for Zip1 loading[6] (**Fig. 4a,b**). Both *zip1Δ* and *hop1Δ* Hi-C maps retained the
Rec8-dependent punctate interactions (**Fig. 4b, Extended Data Fig. 2b,c**), and displayed
compaction relative to G1 or *rec8Δ*, but with the *P(s)* shoulder shifted left relative to *ndt80Δ* (**Fig.**
4c). Attempts to model the known *zip1Δ* and *hop1Δ* defects in chromosome synapsis simply by
removing interhomologue crosslinks from best-fitting *ndt80Δ* simulations did not recapitulate the
P(s) shift observed experimentally (**Extended Data Fig. 5 f**), consistent with the suggestion that
interhomologue contacts make only a minor contribution within meiotic Hi-C maps[23]. Instead,
best-fitting simulations had shifts towards slightly lower processivity and larger separation,
consistent with less axial compaction relative to the *ndt80Δ* control (**Fig. 4e**). Interestingly,
subtelomeric regions no longer displayed a distinct *P(s)* in *zip1Δ* and *hop1Δ* (**Fig. 4d**), suggesting
that chromosome compaction at chromosome termini is regulated differently.

Discussion

Our analysis of meiotic chromosome organisation via Hi-C reconciles the function and localisation
of factors thought to shape meiotic chromosomes with their 3D organisation, revealing the
emergence of a punctate grid of interactions concomitant with initial stages of meiotic
chromosome compaction. Crucially, we formally demonstrate the link between preferential
positioning of meiotic cohesin along the genome[27,35] and the inference that these loci come into
close proximity based on the localization of Rec8 to the chromosomal axes[3]. Remarkably, the

punctate cohesin-dependent interactions in yeast meiosis emerge despite the absence of CTCF
in this organism; this challenges previous models where focal Hi-C peaks are strictly dependent
on CTCF[24,31,43], and indicates that alternative mechanisms of loop positioning must exist.

Notably, whilst much less prominent, locus-specific folding is evident in equivalent high-resolution
Hi-C maps of mitotic cells (**Extended Data Fig. 7**)—something that was hidden within lower-
resolution analyses[19,21]. The correspondence between cohesin positioning and convergent
transcription in both meiosis[35] and mitosis[44,45] argues that transcription may be a fundamental and
ubiquitous process capable of broadly patterning locus-specific chromosome organisation by
modulation of cohesin dynamics[46]. Indeed, the stronger meiotic patterns are particularly
reminiscent of the extended grid-like Hi-C patterns observed in interphase mammalian cells upon
depletion of the cohesion unloader, Wapl[31,47], wherein "vermicelli"-like chromatids arise with a
cohesin-rich backbone[48], emphasising the influence of cohesin dynamics on loop extrusion. We
favour the view that transcription acts as a barrier to cohesin-dependent loop extrusion, rather
than as a motive force as previously proposed[35,49,50], consistent with transcription-independent
compaction by cohesin in mammalian interphase[47] and direct observation of extrusion by the
related SMC condensin *in vitro*[51].

Consistent with the idea that large DNA-protein complexes, like a kinetochore, can impede
extrusion, we observe a paucity of Rec8-dependent loops spanning the centromere.
Nevertheless, whether barriers arise directly from Pol2 binding, or indirectly via other axis
proteins, remains to be determined. Indeed, the reason for the why loops are more strictly
positioned in meiosis compared to mitosis is intriguing. However, our observations enable us to
rule out the axial element, Hop1, the SC lateral element, Zip1, and the process of homologous
recombination mediated by Spo11, Sae2, and Dmc1 (unpub. obs.) as important for the generation
of such patterns.

Our simulations also reveal a nuanced picture of meiotic chromosome assembly: loops are, on
average, larger than the inter-Rec8 peak distance, and more than half of the loop bases are not
associated with preferred sites of Rec8 binding. Moreover, it is likely that loop sizes and positions
vary widely from one cell to another, making classifications of genomic regions as 'axis' or 'loop'
a great oversimplification. The agreement between our simulations and experimental data furthers
the case for loop extrusion as a general mechanism[19–21,23,28,52–55] that is flexibly employed and
regulated in interphase, mitosis, and meiosis.

Our results also reveal how the interplay between the synapsis components, Hop1 and Zip1, influences chromosome morphology. That Hop1 and Zip1 are both required to increase chromosome compaction at pachytene likely points at their joint role in promoting synapsis[4,6], and supports the view that synapsis itself modulates axial compaction. Interestingly, whilst Zip1 binds largely uniformly along the arms of pachytene chromosomes[56], subtelomeres and short chromosomes display an increase in short-range contacts and an earlier shoulder in *P(s)*, consistent with smaller loops or less compression of spacers between loops in these regions, and therefore less axial compaction. Because such differences correlate with disproportionate retention of Hop1 in these regions[56] and diminished efficiency of synapsis[57], it is possible that Hop1 impedes the pathway whereby Zip1 imposes additional compaction upon synapsis. Nevertheless, it is unclear whether Zip1 mediates this effect by modifying loop extrusion dynamics, or via a distinct process of axial compression, as has been argued for higher eukaryote mitotic chromosome compaction[20]. Given the influence that chromosome structure has over so many aspects of meiosis, teasing apart these mechanisms is of great future interest.

Acknowledgements & Contributions

SAS and MJN planned the study, performed wet-lab work and data analysis. GF and KSP developed polymer simulations and performed data analysis. SAS, GF, JB, KSP and MJN discussed results and wrote the manuscript. We thank Tim Cooper for deploying the HiC Pro installation, Scott Keeney and Franz Klein for sharing *S. cerevisiae* strains, Svetlana Lyalina for assistance with QB3 GPUs, Anton Goloborodko for suggesting the use of *looplib*, and Nezar Abdennur for feedback.

Data & Code Availability

All processed Hi-C matrices are publicly accessible via the interactive HiGlass viewer[58]. Live web links are provided with each figure. Raw sequence reads will be made publicly accessible via the SRA repository upon final publication. Code for data analysis is either already publicly available online or will be made available on request or via Github upon final publication.

Funding Statement

SS and MJN are supported by an ERC Consolidator Grant (#311336), the BBSRC (#BB/M010279/1) and the Wellcome Trust (#200843/Z/16/Z). KP and GF are supported by NIMH grant #MH109907, NHLBI grant #HL098179, and the San Simeon Fund.

Figure Legends

Figure 1. Chromosome conformation during yeast meiosis.

a. Cells were collected during meiosis at indicated timepoints and analysed by Hi-C. At 0h the cells are in G1. Representative Hi-C contact maps of chromosomes 6, 11, and 7 plotted at 5 kb resolution. Centromeres, telomeres and arm fold-back at the centromere are indicated by blue, red and grey arrows, respectively, and axial compaction by the width of the main diagonal relative to the fixed-width black clamp. For interactive HiGlass[58] views see: http://higlass.pollard.gladstone.org/app/?config=Z5iwKpjzQpePCXXyvuYGeQ

b. Meiotic entry assessed by FACS; at 4h, the majority of cells show a 4C peak indicating completion of DNA replication.

c. Meiotic progression was monitored by quantification of nuclear divisions determined by DAPI staining. Around 4h, cells start to undergo meiotic divisions I and II. The majority of cells undergo meiotic divisions between 4h and 8h, indicating the degree of heterogeneity within the cell population.

d. *Upper panels*: Average *trans* centromere-centromere contact maps. *Lower panels*: *trans* telomere-telomere contact maps. *Right*: ratio of *cis* to total contact frequency.

e. Intra-arm contact probability versus genomic distance, *P(s),* indicating the emergence (*left*) and disappearance (*right*) of chromosome arm compaction during meiosis. Shaded area bounded above and below by the two *ndt80Δ* 8h replicates.

f. Meiosis was induced in *ndt80Δ* cells for 8h and meiotic entry was checked by monitoring DNA replication by FACS.

g. *ndt80Δ* cells were grown for 8h in sporulation media and analysed by Hi-C (*left*). Log2 ratio of
ndt80Δ cells 8h over G1 (*right*). Centromeres and telomeres are indicated by blue and red arrows,
respectively, and axial compaction by a black clamp.

h. *Left*: Contact probability of individual chromosome arms stratified by length. *Right*: Contact
probability stratified by the distance from the telomere.

Figure 2. Emergence of a Rec8-dependent grid of punctate interactions in meiosis

a. Hi-C contact maps of chromosome 11 for the indicated genotypes plotted at 2 kb bin resolution,
showing near-diagonal interactions. Wild type timepoints as in **Fig. 1a**. **Lower panels**:
log2(insulation); cis/total ratio, Rec8 ChIP-seq[27], all binned at 2kb. Insulation and cis/total
calculated from *ndt80Δ* maps. Positions of Rec8 sites indicated as green circles. Genome-wide
cis/total (Spearman's R=0.62, P<1e-10) and insulation (R= -0.23, P<1e-10, insulation window =
20 kb) profiles are correlated with Rec8 occupancy.

b. Zoom-in into contact maps on chromosome 11 (0-200kb) of wt-4h and *ndt80Δ* (top) and *rec8Δ*
(*bottom left*). Contact probability versus genomic distance, *P(s)*, for G1(*ndt80Δ-0h*) and *ndt80Δ*
and *rec8Δ* (*bottom right*). Data shown is the average (n=2) except for wt-4h. Rec8 peak sites
called from ChIP-seq data[27] are indicated in green. For an interactive view see:
http://higlass.pollard.gladstone.org/app/?config=Twrh61jGT4SlxotaguTIJg

c. Simplified illustration of how a grid of peaks on a Hi-C map can emerge between Rec8 sites
either by transitive contacts between adjacent loops, or by loops that skip over adjacent sites.
Experimentally observed grids extend much further than separation=2 (**Extended Data Fig. 4c**)

d. Cis/total ratios for Rec8 (green) and nonRec8 (grey) sites for indicated datasets.

e. Contact frequency versus distance between Rec8-Rec8 sites (green), Rec8-nonRec8 sites
(light green) and nonRec8-nonRec8 sites (green).

f. Log2 ratio of contact frequency between adjacent Rec8-sites (separation=1) compared to
average cis interactions.

g. Log2 ratio of contact frequency centered at Rec8 sites compared to average cis interactions.
In *ndt80Δ*, Rec8 sites show: elevated cis/total frequency (0.85 versus 0.77), elevated pairwise
contact frequency (~2-fold at 20 kb), and mild insulation (**f-g**). These distinctions are similar in
wild type pachytene (4h) yet absent in G1 (*ndt80Δ-0h*) or in *rec8Δ*.

Figure 3. Modelling meiotic chromosome compaction.

a. In simulations, yeast chr13 was represented as a polymer fiber confined to the nucleus subject to additional meiosis-specific constraints. These include: extruded loops, sister crosslinks, and homolog crosslinks (**Methods**). Barriers to extruded loops were placed at Rec8 sites[27]. We imposed inter-sister and inter-homologue crosslinks at sites of extruded loop bases in order to approximate the paired arrangement of homologues at pachytene (**Extended Data Fig. 6**). For each set of extruded loop parameters (*processivity, separation,* and *barrier strength*), conformations were collected and used to generate simulated contact maps. Roughly, *processivity* dictates the size of an extruded loop unimpeded by collisions, *separation* controls the number of active extruders on the chromosome, and *barrier strength* controls the probability that an extruder gets paused when attempting to step past a barrier. Simulated and experimental contact maps were then compared via the combined average fold discrepancy between *P(s)* curves for Rec8-Rec8, Rec8-non, and non-non bin pairs at 2kb resolution.

b. Goodness-of-fit for indicated barrier strengths over coarse grids of processivity and separation demonstrate that intermediate barrier strengths are required to agree with experimental *ndt80Δ* Hi-C maps.

c. Goodness-of-fit for a fine grid of processivity versus separation at barrier strength 0.95. Best-fitting models had separation ~32kb and processivity ~76kb, corresponding to ~60% coverage of the genome by extruded loops of average length 26kb.

d. From *left* to *right*: contact maps for chr13 for *ndt80Δ*, and simulations with (i) best-fitting parameters, (ii) relatively stable loops between neighboring Rec8 sites, and (iii) no barriers.

e. P(s) split by Rec8-Rec8, Rec8-non, and non-non, as in **Fig. 2D**.

f. Conformations for best-fitting simulations, which highlight: (*left*) one chromatid colored from start (red) to end (blue); (*right*) extruders (yellow), extrusion barriers (red), and extruders paused at barriers (orange).

Figure 4. Hop1 and Zip1-dependent compaction of Rec8-dependent loops.

a. *Top*: Hi-C maps for *hop1Δ* and *zip1Δ* (plotted as in **Fig. 1a**). *Bottom*: Log2 ratio of *hop1Δ* over *ndt80Δ* (as in **Fig. 1g**). For interactive views of the full genome, see http://higlass.pollard.gladstone.org/app/?config=TTBGu5DDR0SHAa09zrjTXA

b. Hi-C contact maps of chromosome 11 for *hop1Δ* and *zip1Δ* plotted at 2kb bin resolution, showing near-diagonal interactions, as in **Fig. 2a**.

c. Contact probability versus genomic distance for G1, *ndt80Δ*, *hop1Δ*, *zip1Δ*. Shaded area
bounded above and below by *ndt80Δ* replicas. Average between two replicas for *zip1Δ* and one
sample for G1 and *hop1Δ* are shown.

d. Contact probability over genomic distance averaged over all chromosome arms stratified by
distance from the telomere.

e. Goodness-of-fit for simulations without homolog crosslinks with a fine grid of processivity
versus separation at barrier strength 0.95 *zip1Δ* and *hop1Δ*.

f. Model of meiotic chromosome compaction: Rec8-dependent loop formation leads to initial
chromosome arm compaction and emergence of a grid-like pattern of Hi-C interactions that jointly
agrees with a mechanism of loop extrusion including barrier elements. We suggest that
transcription could impose such barriers. Hop1 and Zip1 are dispensable for this step, but are
required for synapsis, where additional compaction occurs differentially along chromosome arms.
358

359 **Extended Data Fig. 1**
Temporal and chromosome length-specific analysis of meiotic chromatin conformation
a-d. Results from a replicate timecourse, collected and characterized independently of the
timecourse in Fig. 1.

a. Hi-C maps, plotted as in Fig. 1a.

b. FACS as in Fig. 1b.

c. DAPI as in Fig. 1c.

d. *P(s)* as in Fig. 1e.

e. *P(s)* for chromosomes stratified by size for *ndt80Δ*-0h, *ndt80Δ*-8h. Short chromosomes display
relatively elevated *P(s)* at short distances, and an earlier shoulder.

f. *Left*: *P(s)* for individual chromosome arms, stratified by size for wt-4h. Short arms display
relatively elevated *P(s)* at short distances, and an earlier roll-over. *Right*: Intra-arm *P(s)* stratified
by the distance from the telomere for wt-4h*,* averaged across all chromosomes. Telomere-
proximal regions display elevated *P(s)* at short distances.

g. Intra-arm *P(s)* stratified by the distance from the centromere for G1 (*ndt80Δ*-0h), *wt*-4h, *ndt80Δ*-
8h, averaged across all chromosomes.

h. Contact probability of single chromosome arms for *ndt80Δ*-8h*.*

Extended Data Fig. 2
Aggregate analysis of centromeric interactions in meiosis
376 **a.** Average *trans* centromere-centromere contact maps for indicated data sets.

377   **b.** Average *cis* centromere-centromere contact maps for indicated data sets. Note the loss of the

378   folding back in meiosis, and how the intra-arm enrichment is insulated at centromeres in meiosis.

379   **Extended Data Fig. 3**
       Aggregate analysis of telomeric interactions in meiosis

380   **a.** Average *trans* telomere-telomere contact maps for indicated datasets.

381   **b.** Average telomere-telomere contact maps between the two telomeres of the same

382   chromosome.

383   **c**. Average contact map around each telomere in *cis.*

384   **Extended Data Fig. 4**
       Preferred sites of Rec8 occupancy define sites of locus-specific interaction

a. *Left*: Hi-C contact maps of *rec8Δ ndt80Δ*. Chromosomes 6, 11 and 7 are shown as

representatives for the whole genome. *Right*: Log2 Hi-C ratio maps of *rec8Δ ndt80Δ / ndt80Δ*.

Plotted as in Fig. 1g.

b. Log2 observed over expected contact frequency at Rec8-Rec8 peak pairs as a function of

separation across datasets.

c. Log2 observed over expected contact frequency +/-8kb around Rec8-Rec8 peak pairs at the

indicated separations.

Together, **b-c** demonstrate that Rec8-Rec8 enrichments are strongest between adjacent sites,

decrease between non-adjacent sites with increasing genomic separation, and are absent in

trans. Equally important, these meiotic features are lost in *rec8Δ*. As for mammalian interphase,

this observation in meiosis argues for a *cis*-acting process underlying the formation of focal

interactions between Rec8 sites.

d. cis/total as a function of distance along the chromosomal arm, Rec8 sites marked in green.

Extended Data Fig. 5
       Polymer simulations of loop extrusion reveal best fitting parameters and conformations

a. Representative conformation for the indicated parameter sets. As in **Fig. 3F,** one chromatid

from a homologous quartet of chromatids colored from start to end according to the spectrum;

other three colored in grey.

b. For the same three conformations, positions of Rec8 sites indicated with red spheres, positions

of extruded loop bases in yellow, and extruders overlapping a Rec8 site in orange. Note the stable

loops between neighboring Rec8 sites creates a very elongated chromatid (ii). Also note the

majority of Rec8 sites are unoccupied in (iii), despite the self-assembly of two axial cores and a

strong brush. Finally, note very dispersed chromosomes in (iv), consistent with EM[3] for *rec8Δ*.

c. Contact frequency versus distance, *P(s),* for indicated simulations. Note that the loss of the

shoulder in *P(s)* in the case of full extruder depletion mirrors the difference between experimental

ndt80Δ and *rec8Δ* Hi-C maps. Simulations with increased processivity predict that *P(s)* would shift

rightward if unloading was impaired, as could happen in *waplΔ.* Conversely, if unloading was

enhanced, simulations with decreased processivity indicate a leftward shift in P(s), until the

absence of extruders.

d. Goodness-of-fit for a fine grid of processivity versus separation at barrier strength 0.90. The

best-fit occurs at similar processivity and separation as for barrier strength 0.95 shown in **Fig. 3c**,

but with slightly lower goodness-of-fit.

e. Goodness-of-fit to *rec8Δ* data for simulations with the indicated barrier strengths (in grey: 0.00,

0.75, 0.90, 0.95, 0.99, 1.00) over coarse grids of processivity and separation demonstrates that

the best fits have few if any extruded loops, regardless of barrier strength.

f. *P(s)* curves for simulations with sisters and homologs with the best-fitting parameters for

ndt80Δ-8h maps compared to P(s) for simulations with sisters only show that simply removing

homolog tethering does not recapitulate the sort of shifted P(s) seen experimentally in *zip1Δ* Hi-

C.

Extended Data Fig. 6
      Polymer simulations of loop extrusion elucidate meiotic barrier strength

a. Simulated contact maps for the indicated region of chr13 for: (i) best-fitting simulations, (ii)

simulations with relatively stable loops between neighboring Rec8 sites (barrier strength=1 and

high processivity), and (iii) no barriers, as in **Fig. 2d.**

b. Simulated ChIP-seq profiles for the indicated region of chr13. Best-fitting simulations (i) display

occupancy well below 100% at Rec8 sites. Simulations with stable loops (ii) display highly

occupied Rec8 sites. Simulations without barriers (iii) have homogenous Rec8 occupancy across

the genome.

c. Positions of extruded loops (arcs) sister crosslinks (solid black lines) and homolog crosslinks

(dashed lines) for four chromatids in two separate cells, showing how the simulated Hi-C maps

and ChIP-seq profiles emerge from the stochastic positioning of extruded loops from cell-to-cell.

For statistics, see **Supplementary Table S1**.

d. Histogram of extruded loop lengths for indicated parameters (i, ii, iii).

Extended Data Fig. 7
Cohesin and transcription patterns loops in meiosis and mitosis
a. Hi-C contact maps of chromosome 11 for meiotic (*ndt80Δ*, pachytene - top) and mitotic (wild

type, nocodazole arrest - bottom) plotted at 2 kb bin resolution, showing near-diagonal

interactions. Data shown is the average (n=2).

b. Zoom-in into contact maps on chromosome 11 (0-200kb) of *ndt80Δ* (top) and mitotic (*bottom*).

Data shown is the average (n=2). Rec8 peak sites called from ChIP-seq data[27] are indicated in

green. Arrowheads indicate sites of prominent focal interaction.

Tables

449 **Table 1. *S. cerevisiae* strains used in this study**

Strain name	genotype
MJ6	ho::LYS2/'', lys2/'', ura3/'', arg4-nsp/'', leu2::hisG/'', his4X::LEU2/'', nuc1::LEU2/''
SSY14	ho::LYS2/'', lys2/'', ura3/'', arg4-nsp/'', leu2::hisG/'', his4X::LEU2/'', nuc1::LEU2/'', ndt80Δ::LEU2/''
SSY20	ho::LYS2/'', lys2/'', ura3/'', arg4-nsp/'', leu2::hisG/'', rec8Δ::KanMX4/'', ndt80Δ::LEU2/''
SSY25	ho::LYS2/'', lys2/'', ura3/', arg4-nsp/'', leu2::hisG/'', his4X::LEU2/'', nuc1::LEU2/'', zip1::LEU2/'', ndt80Δ::LEU2/''
SSY49	ho::LYS2/'', lys2/'', ura3/'', arg4-nsp/'', leu2::hisG/'', nuc1::LEU2/'', his4X::LEU2/'', hop1::LEU2/'', ndt80Δ::LEU2/''
SSY58	ho::hisG/'', lys2/'', ura3/'', leu2::hisG/'', nuc1::LEU2/'', arg4-nsp/'', rec8::KanMX/'', ndt80Δ::LEU2/''

451 **Table 2. Hi-C Libraries**

Name	mutations	Sample name	valid pairs (M)
Main figures			
wt-0h/G1		HiC_MJ6_wt_2A_0h	14.5
wt-2h		HiC_MJ6_wt_2A1_2h	27.6
wt-3h		HiC_MJ6_wt_2A_3h	24.1
wt-4h		HiC_MJ6_wt_2A_4h	28
wt-5h		HiC_MJ6_wt_2A1_5h	27.6
wt-6h		HiC_MJ6_wt_2A1_6h	27.6
wt-8h		HiC_MJ6_wt_2A3_8h	19
rec8Δ	*rec8Δ ndt80Δ*	average	
rec8Δ replica 1	*rec8Δ ndt80Δ*	HiC_SSY20_ndt80Drec8D_1A2_8h	39.3
rec8Δ replica 2	*rec8Δ ndt80Δ*	HiC_SSY58_ndt80Drec8D_2A_8h	20.2
ndt80Δ	*ndt80Δ*	average, 8h	
G1	*ndt80Δ*	HiC_SSY14_ndt80D_1A2_0h	36
ndt80Δ-4h	*ndt80Δ*	HiC_SSY14_ndt80D_1A_4h	11.9
ndt80Δ replica 1	*ndt80Δ*	HiC_SSY14_ndt80D_1A1_8h	22.9
ndt80Δ replica 2	*ndt80Δ*	HiC_SSY14_ndt80D_2A2_8h	37
zip1Δ	*zip1Δ ndt80Δ*	average	
zip1Δ replica 1	*zip1Δ ndt80Δ*	HiC_SSY25_ndt80Dzip1D_1B2_8h	22.7
zip1Δ replica 2	*zip1Δ ndt80Δ*	HiC_SSY25_ndt80Dzip1D_2A_8h	28.6

hop1Δ	*hop1Δ ndt80Δ*	hop1 ndt80	
hop1Δ replica1	*hop1Δ ndt80Δ*	HiC_SSY49_ndt80Dhop1D_1A_8h	32.8
Supplementary figures			
wt-2h		HiC_MJ6_wt_3A_2h	22.5
wt-3h		HiC_MJ6_wt_3A_3h	19.8
wt-4h		HiC_MJ6_wt_3A_4h	16.7
wt-6h		HiC_MJ6_wt_3A_6h	37.6

453 **Table 3. Overview of proteins described in this study**

Protein	Description
Ndt80	Transcription factor required for exit from pachytene
Rec8	Meiosis-specific kleisin subunit of cohesin
Hop1	Axial element of the synaptonemal complex
Zip1	Transverse filament of the synaptonemal complex

Methods

Yeast strains and cell culture growth.

Strains used in this study were derived from SK1 and are listed in Table 1.

Monitoring meiotic progression by flow cytometry and quantification of nuclear divisions

Cells were fixed in 70% EtOH, digested with 1 mg/ml RNAse (10 mM Tris-HCl pH 8.0, 15 mM NaCl, 10 mM EDTA pH 8.0) for 2 h at 37 ˚C, 800 rpm and subsequently treated with 1 mg/ml Proteinase K in 50 mM Tris-HCl pH 8.0 at 50 ˚C, 800 rpm for 30 min for analysis by FACS. Cells were then washed in 50 mM Tris-HCl pH 8.0 and stained in the same buffer with 1 uM Sytox green overnight in the fridge. FACS profiles were plotted with R using the library *hwglabr2* (https://github.com/hochwagenlab/hwglabr2). Fixed cells were also used for quantification of nuclear divisions by spreading onto a microscope slide, mounting with Fluoroshield containing DAPI followed by analysis with a Zeiss Scope.A1 microscope.

Hi-C library preparation

The Hi-C protocol used was amended from[59] by ~5-fold reduction in all materials and volumes. Briefly, *S. cerevisiae* diploid cells were synchronised in G1 by growth at 30 ˚C for ~15 h in 30 ml YPA (1% Yeast extract, 2% Peptone, 1% K-acetate) to OD600 of ~4, harvested, washed, and resuspended in prewarmed sporulation medium (2% KAc with 0.2x nutritional supplements) before fixing 5 ml aliquots (20-30 ODs) of relevant timepoints with formaldehyde at 3% final concentration for 20 min at 30 °C, 250 rpm, then quenched by incubating with a final concentration of 0.35 M Glycine (2x the volume of Formaldehyde added) for an additional 5 minutes. Cells were washed with water split into two samples and stored at -80˚C ready for library preparation. Cells were thawed, washed in spheroplasting buffer (SB, 1 M Sorbitol, 50 mM Tris pH 7.5) and digested with 100 ug/ml 100T Zymolyase in SB containing 1% beta-Mercaptoethanol for 15-20 min at 35 ˚C. Cells were washed in restriction enzyme buffer, chromatin was solubilised by adding SDS to 0.1% and incubating at 65 ˚C for 10 minutes. Excess SDS was quenched by addition of Triton X100 to 1%, and chromatin was incubated with 2.07U/ul of *Dpn*II overnight at 37 ˚C. DNA ends were filled in with nucleotides, substituting dCTP for biotin-14-dCTP using Klenow fragment DNA polymerase I at 37 ˚C for 2 h followed by addition of SDS to 1.5% and incubation at 65 ˚C for 20 min to inactivate Klenow and further solubilise the chromatin. The sample volume was diluted 15-

fold, crosslinked DNA ends ligated at 16 ˚C for 8 h using 0.024U/ul of T4 DNA ligase, and

crosslinks reversed by overnight incubation at 65 ˚C in the presence of proteinase K. DNA was

precipitated with ethanol, dissolved in TE and passed through an Amicon 30 kDa column. DNA

was further purified by phenol:chloroform:isoamylalcohol extraction and precipitated again before

treating with RNaseA at 37 ˚C for 1 h. Biotin was removed from unligated ends by incubation with

T4 DNA polymerase at 20˚C for 4 h and at 75 ˚C for 20 min for inactivation of the enzyme. DNA

was subsequently fragmented using a Covaris M220 (Duty factor 20%, 200 cycles/burst, 350s,

20 ˚C), and DNA ends were repaired and A-tailed using T4 DNA polymerase, T4 Polynucleotide

Kinase and Klenow fragment DNA polymerase I before isolating fragments of 100-250 bp using

a Blue Pippin (Sage). Biotinylated fragments were enriched using streptavidin magnetic beads

(C1) and NextFlex (Bioo Scientific) barcoded adapters were ligated while the DNA was on the

beads. Resulting libraries were minimally amplified by PCR and sequenced using paired end 42

bp reads on a NextSeq500 (Ilumina; Brighton Genomics).

Hi-C data processing and analysis

Hi-C sparse matrices were generated at varying spatial resolutions using the Hi-C-pro pipeline[60],

using a customised S288c reference genome (*SK1Mod*, in which high confidence SK1-specific

polymorphisms were inserted in order to improve read alignment rates; manuscript in preparation)

and plotted using R Studio (version 1.0.44) after correcting for read depth differences between

samples. Raw read statistics are presented in **Table 2**. Repeat biological samples gave broadly

similar matrices and, unless indicated otherwise, were averaged to improve their expected

quantitative accuracy. As visual inspection indicated a number of potential translocations in the

SK1 strain as compared with the S288c reference genome, for conservative downstream

analyses, additional bins were masked if they contained potential translocations. Such bins were

identified if they either had values in *trans* at the level of the median of the third diagonal in *cis*, or

the maximum value in *trans* exceeded the maximum value in *cis* for SSY14 for bins displaying

these properties in either *ndt80D-0h* or in *ndt80D-8h* and for MJ6 in *wt-0h* or *wt-4h*. chr1 was

excluded from downstream analysis as few informative bins remained after filtering potential

translocations.

Average maps centered at centromeres and telomeres were calculated as in[61], ensuring that

collected patches for average centromere maps did not extend inter-chromosomally, and

collected patches for average telomere maps did not extend beyond centromeres or inter-

chromosomally. Contact frequency versus distance curves, *P(s),* were calculated from 2 kb

binned maps, with logarithmically-spaced bins in *s* (numutils.logbins, https://bitbucket.org /
mirnylab/mirnylib, start =2, end = max(binned arm lengths), N=50), and restricting the calculation
to bin pairs within chromosomal arms and excluding bins less than 20 kb from centromeres or
telomere (as in[61]), and normalized to the average value at 4 kb. *P(s)* stratified by distance to
telomeres was calculated using the combined distance to telomeres for each bin-pair (as in[62]),
and excluded bins-pairs where one bin was closer to a centromere than telomere along that arm.
Distance to centromeres, and *P(s)* stratified by this distance, was calculated similarly. Log2
insulation profiles were calculated using a sliding diamond window (as in[63]) with a +/-20 kb (+/-10
bins) extent; as in[43] downstream analyses were restricted to when there were zero or one filtered
bins in the sliding window. To calculate histograms of cis/total (**Fig. 2D**), bins were defined as
either Rec8 or non-Rec8. To calculate *P(s)* split by Rec8 bin-pair status, each bin-pair (i.e. entry
of the heatmap) was assigned as either Rec8-Rec8, Rec8-nonRec8, or non-non (e.g. **Fig. 2E**).
P(s) was then aggregated separately across chromosomes for these three categories, similar to
calculation of *P(s)* within and between TADs[28]. Average log2 observed/expected maps were
calculated by first dividing by intra-arm *P(s)* and then averaging together appropriate patches of
Hi-C maps. Correlations between Rec8 occupancy from[27] and insulation or cis/total profiles
excluded chromosome 12 because the rDNA locus greatly alters the insulation profile within the
right arm of the chromosome.

Polymer simulations

Meiotic loop extrusion simulations begin with a generic polymer representation of the yeast
chromatin fiber similar to that used in previous models of yeast mitotic chromosomes[21], where
each 20 nm monomer represents 640 bp (~4 nucleosomes). We simulated the chromatin fiber
with excluded volume interactions and without topological constraints, using Langevin dynamics
in OpenMM, as in[64,65]. Importantly, meiotic simulations remove the geometric constraints specific
to the Rabl conformation[66,67] because this is not visible in meiotic pachytene *ndt80Δ* Hi-C maps.
As our focus was to characterize the grids of intra-chromosomal interactions, we considered a
system with multiple copies of chromosome 13, equivalent to four copies of the haploid genome
in terms of total genomic content (4 x 13 copies of chromosome 13), to enable efficient
computational averaging of simulated Hi-C maps. Extruded loops were generated according to
parameters that describe the dynamics of loop extruders, using the simulation engine described
in[68]: extruder *separation*, extruder *processivity*, chromatin fiber *relaxation time* relative to extruder
velocity, and *barrier strength*. Because yeast chromosomes are short compared to higher
eukaryote chromosomes, relaxation time is relatively rapid and we focused on *separation,*

processivity, and *barrier strength*. At every given timepoint an extruded loop is realized as a bond
between monomers at the two bases of the loop (see *./src/examples/loopExtrusion* in
https://bitbucket.org/mirnylab/openmm-polymer/).

Upon encountering a barrier, a loop extruder is paused with probability according to the barrier
strength; barrier strength =1 indicates an impermeable barrier, barrier strength =0 indicates no
impediment to extrusion. We assume loop extrusion occurs independently on each chromatid,
and simulate loop extrusion dynamics on a 1D lattice (as in[28]) where the number of lattice sites
equals the total number of monomers (75,140). Bi-directional barriers were placed at monomers
with positions corresponding to Rec8 ChIP-seq sites[27], and pause extruders according the barrier
strength parameter. We assume a uniform birth probability, constant death probability, and that
all barriers have an equal strength; as additional data becomes available, these assumptions can
be relaxed and more detailed models can be built.

We investigated scenarios where chromatids are then either left individualized (52 copies),
crosslinked to sisters (26 pairs), or additionally paired with homologs (13 pairs-of-pairs). For
simulations with sister crosslinks, these were added (following[54]) when extruded loop bases were
present at cognate positions ±30 monomers (~20kb) on both chromatids (distance=20nm);
homolog crosslinks were added similarly when sister crosslinks were present on both chromatids
(distance=100nm); centromeres and telomeres were always paired, and both presented
impermeable (strength=1) boundaries to extruders. To avoid introducing pseudo-knots, if
extruded loops were nested only the outer cohesins were considered as possible bases for sister
crosslinks, sister crosslinks were only allowed between the same side of loop bases (i.e. left-to-
left arm or right-to-right arm), and sister crosslinks were only added between bases at the
reciprocal minimum distance.

For calculation of simulated Hi-C maps, contacts were recorded from conformations of the full
system, which includes intra- and inter-sister, and interhomologue contacts. Because
experimental Hi-C here does not distinguish either sisters or homologs, contacts were then
aggregated into one simulated map. For each model and parameter set we investigated, we
collected an ensemble of conformations, generated simulated chr13 Hi-C maps, and compared
their features and *P(s)* with those from experimental Hi-C maps. Each simulated chr13 map
represented an average over 5400 conformations. *P(s)* for chr13 was calculated from 2kb binned
simulated maps exactly as for experimental maps. Maps of goodness-of-fit between simulations

and experimental data (e.g. **Fig. 3b,c**) were computed as the geometric standard deviation of the

ratio of simulated to experimental *P(s)* combined across $P^{Rec8\text{-}Rec8}(s)$, $P^{Rec8\text{-}non}(s)$, and $P^{non\text{-}non}(s)$,

as was previously done for P(s) within TADs of multiple sizes and between TADs[28], for *s* from

10kb to 300kb. This measure reflects the typical fold-deviation for *P(s)*.

Simulated ChIP-seq profiles (**Extended Data Fig. 6b**) for Rec8 were generated by aggregating

the position of extruded loop bases (two per extruded loop) across conformations. Statistics of

extruded loop positioning relative to Rec8 sites was calculated with *loopstats.py* in *looplib*

(https://github.com/golobor/looplib), and arc diagrams (**Extended Data Fig. 6c**) with *loopviz.py*.

Conformations showing chromatids or positions of extruded loop bases were rendered in PyMOL

(https://pymol.org/sites/default/files/pymol.bib).

References

1. Zickler, D. & Kleckner, N. Meiotic chromosomes: integrating structure and function. *Annu Rev Genet* **33**,

603-754 (1999).

2. Gray, S. & Cohen, P. E. Control of Meiotic Crossovers: From Double-Strand Break Formation to

Designation. *Annu Rev Genet* (2016).

3. Klein, F. et al. A central role for cohesins in sister chromatid cohesion, formation of axial elements, and

recombination during yeast meiosis. *Cell* **98**, 91-103 (1999).

4. Sym, M., Engebrecht, J. A. & Roeder, G. S. ZIP1 is a synaptonemal complex protein required for meiotic

chromosome synapsis. *Cell* **72**, 365-378 (1993).

5. Smith, A. V. & Roeder, G. S. The yeast Red1 protein localizes to the cores of meiotic chromosomes. *J

Cell Biol* **136**, 957-967 (1997).

6. Hollingsworth, N. M., Goetsch, L. & Byers, B. The HOP1 gene encodes a meiosis-specific component of

yeast chromosomes. *Cell* **61**, 73-84 (1990).

7. Maleki, S., Neale, M. J., Arora, C., Henderson, K. A. & Keeney, S. Interactions between Mei4, Rec114,

and other proteins required for meiotic DNA double-strand break formation in Saccharomyces cerevisiae.

Chromosoma **116**, 471-486 (2007).

8. Li, J., Hooker, G. W. & Roeder, G. S. Saccharomyces cerevisiae Mer2, Mei4 and Rec114 form a

complex required for meiotic double-strand break formation. *Genetics* **173**, 1969-1981 (2006).

9. Dekker, J., Rippe, K., Dekker, M. & Kleckner, N. Capturing chromosome conformation. *Science* **295**,

1306-1311 (2002).

10. Bonev, B. & Cavalli, G. Organization and function of the 3D genome. *Nat Rev Genet* **17**, 661-678 (2016).

11. Merkenschlager, M. & Nora, E. P. CTCF and Cohesin in Genome Folding and Transcriptional Gene

Regulation. *Annu Rev Genomics Hum Genet* **17**, 17-43 (2016).

12.   Dekker, J. & Mirny, L. The 3D Genome as Moderator of Chromosomal Communication. *Cell* **164**, 1110-
1121 (2016).

13.   Fudenberg, G., Abdennur, N., Imakaev, M., Goloborodko, A. & Mirny, L. A. Emerging Evidence of
Chromosome Folding by Loop Extrusion. *Cold Spring Harb Symp Quant Biol* **82**, 45-55 (2017).

14.   Jin, Q., Trelles-Sticken, E., Scherthan, H. & Loidl, J. Yeast nuclei display prominent centromere
clustering that is reduced in nondividing cells and in meiotic prophase. *J Cell Biol* **141**, 21-29 (1998).

15.   Hayashi, A., Ogawa, H., Kohno, K., Gasser, S. M. & Hiraoka, Y. Meiotic behaviours of chromosomes and
microtubules in budding yeast: relocalization of centromeres and telomeres during meiotic prophase.
Genes Cells **3**, 587-601 (1998).

16.   Goloborodko, A., Marko, J. F. & Mirny, L. A. Chromosome Compaction by Active Loop Extrusion.
Biophys J **110**, 2162-2168 (2016).

17.   Gibcus, J. H. et al. A pathway for mitotic chromosome formation. *Science* **359**, (2018).

18.   Kakui, Y., Rabinowitz, A., Barry, D. J. & Uhlmann, F. Condensin-mediated remodeling of the mitotic
chromatin landscape in fission yeast. *Nat Genet* **49**, 1553-1557 (2017).

19.   Lazar-Stefanita, L. et al. Cohesins and condensins orchestrate the 4D dynamics of yeast chromosomes
during the cell cycle. *EMBO J* **36**, 2684-2697 (2017).

20.   Naumova, N. et al. Organization of the mitotic chromosome. *Science* **342**, 948-953 (2013).

21.   Schalbetter, S. A. et al. SMC complexes differentially compact mitotic chromosomes according to
genomic context. *Nat Cell Biol* **19**, 1071-1080 (2017).

22.   Xu, L., Ajimura, M., Padmore, R., Klein, C. & Kleckner, N. NDT80, a meiosis-specific gene required for
exit from pachytene in Saccharomyces cerevisiae. *Mol Cell Biol* **15**, 6572-6581 (1995).

23.   Muller, H. et al. Characterizing meiotic chromosomes' structure and pairing using a designer sequence
optimized for Hi-C. *Mol Syst Biol* **14**, e8293 (2018).

24.   Rao, S. S. P. et al. A 3D map of the human genome at kilobase resolution reveals principles of chromatin
looping. *Cell* **159**, 1665-1680 (2014).

25.   Nora, E. P. et al. Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* **485**,
381-385 (2012).

26.   Dixon, J. R. et al. Topological domains in mammalian genomes identified by analysis of chromatin
interactions. *Nature* **485**, 376-380 (2012).

27.   Ito, M. et al. Meiotic recombination cold spots in chromosomal cohesion sites. *Genes Cells* **19**, 359-373
(2014).

28.   Fudenberg, G. et al. Formation of Chromosomal Domains by Loop Extrusion. *Cell Rep* **15**, 2038-2049
(2016).

29.   Schwarzer, W. et al. Two independent modes of chromatin organization revealed by cohesin removal.
Nature **551**, 51-56 (2017).

30.   Rao, S. S. P. et al. Cohesin Loss Eliminates All Loop Domains. *Cell* **171**, 305-320.e24 (2017).

31.   Wutz, G. et al. Topologically associating domains and chromatin loops depend on cohesin and are
regulated by CTCF, WAPL, and PDS5 proteins. *EMBO J* **36**, 3573-3599 (2017).

32.   Gassler, J. et al. A mechanism of cohesin-dependent loop extrusion organizes zygotic genome
architecture. *EMBO J* **36**, 3600-3618 (2017).

33.   Conrad, M. N., Lee, C. Y., Wilkerson, J. L. & Dresser, M. E. MPS3 mediates meiotic bouquet formation in
Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A* **104**, 8863-8868 (2007).

34.   Trelles-Sticken, E., Adelfalk, C., Loidl, J. & Scherthan, H. Meiotic telomere clustering requires actin for its
formation and cohesin for its resolution. *J Cell Biol* **170**, 213-223 (2005).

35.   Sun, X. et al. Transcription dynamically patterns the meiotic chromosome-axis interface. *Elife* **4**, (2015).

36.   Møens, P. B. & Pearlman, R. E. Chromatin organization at meiosis. *Bioessays* **9**, 151-153 (1988).

37.   Marko, J. F. & Siggia, E. D. Polymer models of meiotic and mitotic chromosomes. *Mol Biol Cell* **8**, 2217-
2231 (1997).

38.   Blat, Y. & Kleckner, N. Cohesins bind to preferential sites along yeast chromosome III, with differential
regulation along arms versus the centric region. *Cell* **98**, 249-259 (1999).

39.   Crawley, O. et al. Cohesin-interacting protein WAPL-1 regulates meiotic chromosome structure and
cohesion by antagonizing specific cohesin complexes. *Elife* **5**, e10851 (2016).

40.   Challa, K., Lee, M.-S., Shinohara, M., Kim, K. P. & Shinohara, A. Rad61/Wpl1 (Wapl), a cohesin
regulator, controls chromosome compaction during meiosis. *Nucleic Acids Res* **44**, 3190-3203 (2016).

41.   Jin, H., Guacci, V. & Yu, H.-G. Pds5 is required for homologue pairing and inhibits synapsis of sister
chromatids during yeast meiosis. *J Cell Biol* **186**, 713-725 (2009).

42.   Petela, N. J. et al. Scc2 Is a Potent Activator of Cohesin's ATPase that Promotes Loading by Binding
Scc1 without Pds5. *Mol Cell* **70**, 1134-1148.e7 (2018).

43.   Nora, E. P. et al. Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains
from Genomic Compartmentalization. *Cell* **169**, 930-944.e22 (2017).

44.   Glynn, E. F. et al. Genome-wide mapping of the cohesin complex in the yeast Saccharomyces
cerevisiae. *PLoS Biol* **2**, E259 (2004).

45.   Lengronne, A. et al. Cohesin relocation from sites of chromosomal loading to places of convergent
transcription. *Nature* **430**, 573-578 (2004).

46.   Heinz, S. et al. Transcription Elongation Can Affect Genome 3D Structure. *Cell* **174**, 1522-1536.e22
(2018).

47.   Haarhuis, J. H. I. et al. The Cohesin Release Factor WAPL Restricts Chromatin Loop Extension. *Cell*
169, 693-707.e14 (2017).

48.   Tedeschi, A. et al. Wapl is an essential regulator of chromatin structure and chromosome segregation.
Nature **501**, 564-568 (2013).

49.   Busslinger, G. A. et al. Cohesin is positioned in mammalian genomes by transcription, CTCF and Wapl.
Nature **544**, 503-507 (2017).

50.   Ocampo-Hafalla, M., Muñoz, S., Samora, C. P. & Uhlmann, F. Evidence for cohesin sliding along
budding yeast chromosomes. *Open Biol* **6**, (2016).

51.   Ganji, M. et al. Real-time imaging of DNA loop extrusion by condensin. *Science* (2018).

52.   Sanborn, A. L. et al. Chromatin extrusion explains key features of loop and domain formation in wild-type
and engineered genomes. *Proc Natl Acad Sci U S A* **112**, E6456-65 (2015).

53. Nasmyth, K. Disseminating the genome: joining, resolving, and separating sister chromatids during
mitosis and meiosis. *Annu Rev Genet* **35**, 673-745 (2001).

54. Goloborodko, A., Imakaev, M. V., Marko, J. F. & Mirny, L. Compaction and segregation of sister
chromatids via active loop extrusion. *Elife* **5**, (2016).

55. Alipour, E. & Marko, J. F. Self-organization of domain structures by DNA-loop-extruding enzymes.
Nucleic Acids Res **40**, 11202-11212 (2012).

56. Subramanian, V. V. et al. Persistent DNA-break potential near telomeres increases initiation of meiotic
recombination on short chromosomes. *Biorxiv* 201889 (2017).

57. Thacker, D., Mohibullah, N., Zhu, X. & Keeney, S. Homologue engagement controls meiotic DNA break
number and distribution. *Nature* **510**, 241-246 (2014).

58. Kerpedjiev, P. et al. HiGlass: web-based visual exploration and analysis of genome interaction maps.
Genome Biol **19**, 125 (2018).

59. Belton, J. M. & Dekker, J. Hi-C in Budding Yeast. *Cold Spring Harb Protoc* **2015**, 649-661 (2015).

60. Servant, N. et al. HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol* **16**,
259 (2015).

61. Hsieh, T. S., Fudenberg, G., Goloborodko, A. & Rando, O. J. Micro-C XL: assaying chromosome
conformation from the nucleosome to the entire genome. *Nat Methods* **13**, 1009-1011 (2016).

62. Mizuguchi, T. et al. Cohesin-dependent globules and heterochromatin shape 3D genome architecture in
S. pombe. *Nature* **516**, 432-435 (2014).

63. Crane, E. et al. Condensin-driven remodelling of X chromosome topology during dosage compensation.
Nature **523**, 240-244 (2015).

64. Eastman, P. et al. OpenMM 7: Rapid development of high performance algorithms for molecular
dynamics. *PLoS Comput Biol* **13**, e1005659 (2017).

65. Eastman, P. & Pande, V. S. Efficient nonbonded interactions for molecular dynamics on a graphics
processing unit. *J Comput Chem* **31**, 1268-1272 (2010).

66. Wong, H. et al. A predictive computational model of the dynamic 3D interphase yeast nucleus. *Curr Biol*
22, 1881-1890 (2012).

67. Tjong, H., Gong, K., Chen, L. & Alber, F. Physical tethering and volume exclusion determine higher-order
genome organization in budding yeast. *Genome Res* **22**, 1295-1305 (2012).

68. Fudenberg, G. & Imakaev, M. FISH-ing for captured contacts: towards reconciling FISH and 3C. *Nat*
Methods **14**, 673-678 (2017).

Figure 1

Figure 1. Chromosome conformation during yeast meiosis.
a. Cells were collected during meiosis at indicated timepoints and analysed by Hi-C. At 0h the cells are in G1. Representative Hi-C contact maps of chromosomes 6, 11, and 7 plotted at 5 kb resolution. Centromeres, telomeres and arm fold-back at the centromere are indicated by blue, red and grey arrows, respectively, and axial compaction by the width of the main diagonal relative to the fixed-width black clamp. For interactive HiGlass[52] views see: http://higlass.pollard.gladstone.org/app/?config=Z5iwKpjzQpePCXXyvuYGeQ
b. Meiotic entry assessed by FACS; at 4 h, the majority of cells show a 4C peak indicating completion of DNA replication.
c. Meiotic progression was monitored by quantification of nuclear divisions determined by DAPI staining. Around 4 h, cells start to undergo meiotic divisions I and II. The majority of cells undergo meiotic divisions between 4 and 8 h, indicating the degree of heterogeneity within the cell population.
d. *Upper panels*: Average *trans* centromere-centromere contact maps. *Lower panels*: *trans* telomere-telomere contact maps. *Right*: ratio of *cis* to total contact frequency.
e. Intra-arm contact probability versus genomic distance, *P(s),* indicating the emergence (*left*) and disappearance (*right*) of chromosome arm compaction during meiosis. Shaded area bounded above and below by the two *ndt80Δ* 8h replicates.
f. Meiosis was induced in *ndt80Δ* cells for 8h and meiotic entry was checked by monitoring DNA replication by FACS.
g. *ndt80Δ* cells were grown for 8h in sporulation media and analysed by Hi-C (*left*). Log2 ratio of *ndt80Δ* cells 8h over G1 (*right*). Centromeres and telomeres are indicated by blue and red arrows, respectively, and axial compaction by a black clamp.
h. *Left*: Contact probability of individual chromosome arms stratified by length. *Right*: Contact probability stratified by the distance from the telomere.

Figure 2

Figure 2. Emergence of a Rec8-dependent grid of punctate interactions in meiosis

a. Hi-C contact maps of chromosome 11 for the indicated genotypes plotted at 2 kb bin resolution, showing near-diagonal interactions. Wild type timepoints as in **Fig. 1a**. **Lower panels**: log2(insulation); cis/total ratio, Rec8 ChIP-seq[28], all binned at 2kb. Insulation and cis/total calculated from *ndt80Δ* maps. Positions of Rec8 sites indicated as green circles. Genome-wide cis/total (Spearman's R=0.62, P<1e-10) and insulation (R= -0.23, P<1e-10, insulation window = 20 kb) profiles are correlated with Rec8 occupancy.

b. Zoom-in into contact maps on chromosome 11 (0-200kb) of wt-4h and *ndt80Δ* (top) and *rec8Δ* (*bottom left*). Contact probability versus genomic distance, *P(s)*, for G1(*ndt80Δ-0h*) and *ndt80Δ* and *rec8Δ* (*bottom right*). Data shown is the average (n=2) except for wt-4h. Rec8 peak sites called from ChIP-seq data[28] are indicated in green. For an interactiveview see: http://higlass.pollard.gladstone.org/app/?config=Twrh61jGT4SlxotaguTlJg

c. Simplified illustration of how a grid of peaks on a Hi-C map can emerge between Rec8 sites either by transitive contacts between adjacent loops, or by loops that skip over adjacent sites. Experimentally observed grids extend much further than separation=2 (**Extended Data Fig. 4c**)

d. Cis/total ratios for Rec8 (green) and nonRec8 (grey) sites for indicated datasets.

e. Contact frequency versus distance between Rec8-Rec8 sites (green), Rec8-nonRec8 sites (light green) and nonRec8-nonRec8 sites (green).

f. Log2 ratio of contact frequency between adjacent Rec8-sites (separation=1) compared to average cis interactions.

g. Log2 ratio of contact frequency centered at Rec8 sites compared to average cis interactions.

In *ndt80Δ*, Rec8 sites show: elevated cis/total frequency (0.85 versus 0.77), elevated pairwise contact frequency (~2-fold at 20 kb), and mild insulation. These distinctions are similar in wild type pachytene (4h) yet absent in G1 (*ndt80Δ-0h*) or in *rec8Δ*.

Figure 3

Figure 3. Model for meiotic chromosome compaction in prophase.

a. In simulations, yeast chr13 was represented as a polymer fiber confined to the nucleus subject to additional meiosis-specific constraints. These include: extruded loops, sister crosslinks, and homolog crosslinks (**Methods**). Barriers to extruded loops were placed at Rec8 sites[28]. We imposed inter-sister and inter-homologue crosslinks at sites of extruded loop bases in order to approximate the paired arrangement of homologues at pachytene (**Extended Data Fig. 6**). For each set of extruded loop parameters (*processivity, separation,* and *barrier strength*), conformations were collected and used to generate simulated contact maps. These were then compared with experimental contact maps via the combined average fold discrepancy with *P(s)* curves for Rec8-Rec8, Rec8-non, and non-non bin pairs at 2 kb resolution.

b. Goodness-of-fit for indicated barrier strengths over coarse grids of processivity and separation demonstrate that intermediate barrier strengths are required to agree with experimental *ndt80Δ* Hi-C maps.

c. Goodness-of-fit for a fine grid of processivity versus separation at barrier strength 0.95. Best-fitting models had separation ~32kb and processivity ~76kb, corresponding to ~60% coverage of the genome by extruded loops of average length 26kb.

d. From *left* to *right*: contact maps for chr13 for *ndt80Δ*, and simulations with (i) best-fitting parameters, (ii) relatively stable loops between neighboring Rec8 sites, and (iii) no barriers.

e. P(s) split by Rec8-Rec8, Rec8-non, and non-non, as in **Fig. 2d**.

f. Conformations for best-fitting simulations, which highlight: *(left)* one chromatid colored from start (red) to end (blue); *(right)* extruders (yellow), extrusion barriers (red), and extruders paused at barriers (orange).

Figure 4

Figure 4. Hop1 and Zip1-dependent compaction of Rec8-dependent loops.

a. *Top*: Hi-C maps for *hop1Δ* and *zip1Δ* (plotted as in **Fig. 1a)**. *Bottom*: Log2 ratio of hop1Δ over *ndt80Δ* (as in **Fig. 1g)**. For interactive views of the full genome, see http://higlass.pollard.gladstone.org/app/?config=TTBGu5DDR0SHAa09zrjTXA

b. Hi-C contact maps of chromosome 11 for *hop1Δ* and *zip1Δ* plotted at 2kb bin resolution, showing near-diagonal interactions, as in **Fig. 2a**.

c. Contact probability versus genomic distance for G1, *ndt80Δ*, *hop1Δ*, *zip1Δ*. Shaded area bounded above and below by *ndt80Δ* replicas. Average between two replicas for *zip1Δ* and one sample for G1 and *hop1Δ* are shown.

d. Contact probability over genomic distance averaged over all chromosome arms stratified by distance from the telomere.

e. Goodness-of-fit for simulations without homolog crosslinks with a fine grid of processivity versus separation at barrier strength 0.95 *zip1Δ* and *hop1Δ*.

f. Model of meiotic chromosome compaction: Rec8-dependent loop formation leads to initial chromosome arm compaction and emergence of a grid-like pattern of Hi-C interactions that jointly agrees with a mechanism of loop extrusion including barrier elements. We suggest that transcription could impose such barriers. Hop1 and Zip1 are dispensable for this step, but are required for synapsis, where additional compaction occurs differentially along chromosome arms.

Extended Data Figure 1

Extended Data Fig. 1 Temporal and chromosome length-specific analysis of meiotic chromatin conformation
a-d. Results from a replicate timecourse, collected and characterized independently of the timecourse in Fig. 1.
a. Hi-C maps, plotted as in Fig. 1a.
b. FACS as in Fig. 1b.
c. DAPI as in Fig. 1c.
d. *P(s)* as in Fig. 1e.
e. *P(s)* for chromosomes stratified by size for *ndt80Δ*-0h, *ndt80Δ*-8h. Short chromosomes display relatively elevated *P(s)* at short distances, and an earlier shoulder.
f. *Left*: *P(s)* for individual chromosome arms, stratified by size for wt-4h. Short arms display relatively elevated *P(s)* at short distances, and an earlier roll-over. *Right*: Intra-arm *P(s)* stratified by the distance from the telomere for wt-4h*,* averaged across all chromosomes. Telomere-proximal regions display elevated *P(s)* at short distances.
g. Intra-arm *P(s)* stratified by the distance from the centromere for G1 (*ndt80Δ*-0h), *wt*-4h, *ndt80Δ*-8h, averaged across all chromosomes.
h. Contact probability of single chromosome arms for *ndt80Δ*-8h*.*

Extended Data Figure 2

a Trans cen-cen

b Cis cen

Extended Data Fig. 2 Aggregate analysis of centromeric interactions in meiosis
a. Average *trans* centromere-centromere contact maps for indicated data sets.
b. Average *cis* centromere-centromere contact maps for indicated data sets. Note the loss of the folding back in meiosis, and how the intra-arm enrichment is insulated at centromeres in meiosis.

Extended Data Figure 3

a Trans tel-tel

b Cis tel-tel

c Cis-arm tel

Extended Data Fig. 3 **Aggregate analysis of telomeric interactions in meiosis**
a. Average trans telomere-telomere contact maps for indicated datasets.
b. Average telomere-telomere contact maps between the two telomeres of the same chromosome.
c. Average contact map around each telomere in cis.

Extended Data Figure 4

Extended Data Fig. 4 Preferred sites of Rec8 occupancy define sites of locus-specific interaction
a. *Left*: Hi-C contact maps of *rec8Δ ndt80Δ*. Chromosomes 6, 11 and 7 are shown as representatives for the whole genome.
Right: Log2 Hi-C ratio maps of *rec8Δ ndt80Δ / ndt80Δ*. Plotted as in Fig. 1g.
b. Log2 observed over expected contact frequency at Rec8-Rec8 peak pairs as a function of separation across datasets.
c. Log2 observed over expected contact frequency +/-8 kb around Rec8-Rec8 peak pairs at the indicated separations.
Together, **b-c** demonstrate that Rec8-Rec8 enrichments are strongest between adjacent sites, decrease between non-adjacent sites with increasing genomic separation, and are absent in *trans*. Equally important, these meiotic features are lost
in *rec8Δ*. As for mammalian interphase, this observation in meiosis argues for a *cis*-acting process underlying the formation
of focal interactions between Rec8 sites.
d. cis/total as a function of distance along the chromosomal arm, Rec8 sites marked in green.

Extended Data Figure 5

Extended Data Fig. 5 Polymer simulations of loop extrusion reveal best fitting parameters and conformations
a. Representative conformation for the indicated parameter sets. As in **Fig. 3F,** one chromatid from a homologous quartet of chromatids coloured from start to end according to the spectrum; other three coloured in grey.
b. For the same three conformations, positions of Rec8 sites indicated with red spheres, positions of extruded loop bases in yellow, and extruders overlapping a Rec8 site in orange. Note the stable loops between neighbouring Rec8 sites creates a very elongated chromatid (ii). Also note the majority of Rec8 sites are unoccupied in (iii), despite the self-assembly of two axial cores and a strong brush. Finally, note very dispersed chromosomes in (iv), consistent with EM[3] for *rec8Δ*.
c. Contact frequency versus distance, *P(s),* for indicated simulations. Note that the loss of the shoulder in *P(s)* in the case of full extruder depletion mirrors the difference between experimental *ndt80Δ* and *rec8Δ* Hi-C maps. Simulations with increased processivity predict that *P(s)* would shift rightward if unloading was impaired, as could happen in *waplΔ.* Conversely, if unloading was enhanced, simulations with decreased processivity indicate a leftward shift in P(s), until the absence of extruders.
d. Goodness-of-fit for a fine grid of processivity versus separation at barrier strength 0.90. The best-fit occurs at similar processivity and separation as for barrier strength 0.95 shown in **Fig. 3c**, but with slightly lower goodness-of-fit.
e. Goodness-of-fit to *rec8Δ* data for simulations with the indicated barrier strengths (in grey: 0.00, 0.75, 0.90, 0.95, 0.99, 1.00) over coarse grids of processivity and separation demonstrates that the best fits have few if any extruded loops, regardless of barrier strength.
f. *P(s)* curves for simulations with sisters and homologs with the best-fitting parameters for *ndt80Δ-8h* maps compared to P(s) for simulations with sisters only show that simply removing homolog tethering does not recapitulate the sort of shifted P(s) seen experimentally in *zip1Δ* Hi-C.

Extended Data Figure 6

Extended Data Fig. 6 Polymer simulations of loop extrusion elucidate meiotic barrier strength
a. Simulated contact maps for the indicated region of chr13 for: (i) best-fitting simulations, (ii) simulations with relatively stable loops between neighboring Rec8 sites (barrier strength=1 and high processivity), and (iii) no barriers, as in **Fig. 2d**.
b. Simulated ChIP-seq profiles for the indicated region of chr13. Best-fitting simulations (i) display occupancy well below 100% at Rec8 sites. Simulations with stable loops (ii) display highly occupied Rec8 sites. Simulations without barriers (iii) have homogenous Rec8 occupancy across the genome.
c. Positions of extruded loops (arcs) sister crosslinks (solid black lines) and homolog crosslinks (dashed lines) for four chromatids in two separate cells, showing how the simulated Hi-C maps and ChIP-seq profiles emerge from the stochastic positioning of extruded loops from cell-to-cell. For statistics, see **Supplementary Table S1**.
d. Histogram of extruded loop lengths for indicated parameters (i, ii, iii).

Extended Figure 7

Extended Data Figure 7. Cohesin and transcription patterns loops in meiosis and mitosis
a. Hi-C contact maps of chromosome 11 for meiotic (*ndt80Δ*, pachytene - top) and mitotic (wild type, nocodazole arrest - bottom) plotted at 2 kb bin resolution, showing near-diagonal interactions. Data shown is the average (n=2).
b. Zoom-in into contact maps on chromosome 11 (0-200kb) of *ndt80Δ* (top) and mitotic (*bottom*). Data shown is the average (n=2). Rec8 peak sites called from ChIP-seq data[27] are indicated in green. Arrowheads indicate sites of prominent focal interaction.