Word stress representations are language-specific: evidence from event-related brain

potentials

Ferenc Honbolygó[1,3], Andrea Kóbor[1], Borbála German[1,2], Valéria Csépe[1,2]

[1] Brain Imaging Centre, Research Centre for Natural Sciences, Hungarian Academy of

Sciences, Budapest, Hungary

[2] Department of Cognitive Science, Budapest University of Technology and Economics,

Budapest, Hungary

[3] Institute of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary

Running head: Word stress representations are language-specific

Corresponding author: Ferenc Honbolygó

Brain Imaging Centre, Research Centre for Natural Sciences, Hungarian Academy of Sciences;

Magyar tudósok körútja 2., H–1117, Budapest, Hungary.

E-mail: honbolygo.ferenc@ttk.mta.hu; Phone: + 36 13826615

22
23

Abstract

Understanding speech at the basic levels entails the simultaneous and independent

processing of phonemic and prosodic features. While it is well-established that phoneme

perception relies on language-specific long-term traces, it is unclear if the processing of

prosodic features similarly involves language-specific representations. In the present study,

we investigated the processing of a specific prosodic feature, word stress, using the method

of event-related brain potentials (ERPs) employing a cross-linguistic approach. Hungarian

participants heard disyllabic pseudowords stressed either on the first (legal stress) or on the

second (illegal stress) syllable, pronounced either by a Hungarian or a German speaker.

Results obtained using a data-driven ERP analysis methodology showed that all

pseudowords in the deviant position elicited an Early Differentiating Negativity (EDN) and a

Mismatch Negativity (MMN) component, except for the Hungarian pseudowords stressed on

the first syllable. This suggests that Hungarian listeners did not process the native legal stress

pattern as deviant, but the same stress pattern with a non-native accent was processed as

deviant. This implies that the processing of word stress was based on language-specific long-

term memory traces.

Keywords: ERP, MMN, speech, prosody, word stress

**Introduction**

Understanding speech requires the simultaneous processing of segmental and

suprasegmental (prosodic) information, suggested to be based on parallel and independent

processing mechanisms (Poeppel, 2014), and possibly relying on separate neural pathways

(Bornkessel-Schlesewsky & Schlesewsky, 2013). Among the available prosodic information,

word stress is a relative emphasis given to certain syllables within words or to certain words

in sentences (for review, see Kager, 2007). Word stress can emphasize or separate certain

parts of the speech stream; thus, it potentially contributes to the segmentation of

continuous speech into words (Cutler & Norris, 1988). However, languages differ in several

aspects regarding word stress: in the position of the stressed syllable within multisyllabic

words (initial, final, penultimate, etc.); in the variability of the stressed syllable's position

(free or fixed); and whether stress can distinguish lexical meaning (contrastive or non-

contrastive). Consequently, the language processing system of individuals has to adapt to

the specific features of word stress of a given language. In the present study, we investigated

whether stress related representations can be considered as language-specific, and whether

we find any evidence of language-specific processing at the neurocognitive level.

Previous studies investigating the neurocognitive basis of word stress processing studied

mostly the Mismatch Negativity (MMN) event-related brain potential (ERP) component. The

MMN is an auditory ERP component with a negative polarity and a frontocentral voltage

maximum, and has been an exceptionally useful tool in studying linguistic processing (for

review, see Näätänen, Paavilainen, Rinne, & Alho, 2007). The MMN is usually elicited in

passive oddball paradigms where frequently repeated standard stimuli are interspersed by

rarely repeated deviant stimuli differing from the standard in some discriminable features.

The MMN appears 100-250 ms after the onset of the change and can be elicited in the

absence of participants' attention. The MMN is currently interpreted as a brain electrical

correlate of the mainly pre-attentive detection of violation of simple or complex regularities

(Winkler, Denham, & Nelken, 2009).

A more complex model for the MMN's emergence is proposed in the AERS (Auditory Event

Representation System) predictive coding model suggested by Winkler and Schröger (2015).

According to the model, acoustic information reaching the auditory system undergoes an

initial sound analysis which extract the proto-features of sounds (for speech, these are

speech-related acoustic-phonetic information, such as voicing, formant transitions,

fundamental frequency, intensity, etc.). This is followed by binding the separate features to

unitary sensory memory representations (for speech, these are phonetic representations like

phonemes, prosodic features, etc.). These representations are then compared against the

predictions derived from the model of the acoustical environment. The predictive model

stores representations of the current regularities of the acoustical environment and it is also

affected by long-term experience, which is available for native linguistic representations

(speech sounds, prosodic patterns, words, etc.). The predictive model is the basis of the

mechanisms comparing incoming information with the regularity representations. If

predictions from the model fail, the model needs to be corrected via an updating process,

which is reflected by the MMN.

There are a number of studies that applied the MMN paradigm to study the processing of

word stress. Weber et al. (2004) found that German adults showed an MMN to the word

'baba' ('baby' in English) in the deviant position both in the trochaic form (stress on the first

syllable, i.e., /baꞎba/) and in the iambic form (stress on the second syllable, i.e., /babaꞎ/),

indicating that both stress patterns were equally well discriminated.

Ylinen et al. (2009) studied the effect of prosodic familiarity in the case of Finnish words and

pseudowords with familiar (trochaic, i.e., stress on the first syllable) and unfamiliar (iambic,

i.e., stress on the second syllable) stress patterns. They used the pseudowords with the

familiar stress pattern as standard and investigated the MMNs elicited by pseudowords with

unfamiliar stress pattern, words with familiar stress pattern, and words with unfamiliar

stress pattern. According to the results, the pseudowords with unfamiliar stress pattern

elicited two MMNs related to the first and second syllables of utterances and were

interpreted as the detection of the unexpected lack of stress on the first syllable and the

detection of unexpected presence of stress on the second syllable, respectively. Words with

familiar stress pattern elicited a single MMN in mid-latency time window, which was

interpreted as the detection of the change of the lexical status of the word, and words with

unfamiliar stress pattern elicited a single MMN related to the presence of stress on the

second syllable. These results demonstrated that the familiarity of the words modulated the

processing of stress pattern, and also that the familiarity of stress pattern modulated the

processing of words.

Zora, Heldner, & Schwarz (2016) investigated stress processing in Turkish. In the study, the

authors presented word and pseudoword stimuli differing in their stress pattern, with

stimuli with iambic stress pattern being always standards and stimuli with trochaic pattern

being deviants. They manipulated the acoustic features contributing to stress (f0, spectral

emphasis, duration, all features combined), in order to investigate to what extent they are

utilized for lexical access. Results showed that changes in prosodic features elicited ERPs

both in words and pseudowords but there were processing differences between the

prosodic features. The most prominent feature was the f0 eliciting an MMN in words but a

frontal P3a in pseudowords, and consequently, was claimed to be lexically specified in

Turkish by the authors. Results also indicated that the individual prosodic features rather

than combined features differentiated language-related effects, since the combined features

elicited similar MMN in both words and pseudowords.

The same authors (Zora, Schwarz, & Heldner, 2015) investigated also the contribution of two

specific features, f0 and intensity to stress processing in English. In the study, participants

heard stress minimal pairs versions (i.e., the same word with trochaic and iambic stress

pattern) of the word 'upset' and the pseudoword 'ukfet'. Deviants – having trochaic stress

compared to the standards having iambic stress – differed from the standards in f0,

intensity, or the combination of the two. According to the results, all deviants elicited the

MMN, although intensity change elicited a smaller MMNs in pseudowords than f0,

confirming the importance of both f0 and intensity to determine words stress and contribute

to lexical access.

Finally, in a study comparing the processing of duration-related stress in speech and music in

English (Peter, Mcarthur, & Thompson, 2012), the authors found that in the case of speech,

only the stress on the first syllable condition elicited an MMN, while in the case of music

stimuli, both long-short and short-long patterns (the musical equivalent of stress on the first

and stress on the second syllable) elicited an MMN, indicating a fundamental difference

between speech and non-speech related processing.

The above studies established convincingly that the MMN can be reliably used to investigate

the neurocognitive background of stress processing and demonstrated the sensitivity of this

particular ERP component to acoustic, phonetic, and lexical effects. In a series of studies, we

investigated word stress processing in Hungarian. Hungarian is an excellent target language

for studying word stress representations because the stress pattern of disyllabic word is

highly regular: stress always falls on the first syllable. Therefore, any changes in the stress

pattern is considered as violating the regularity, leading to illegal stress patterns. First, it was

demonstrated that a word with stress on the second syllable elicited two MMN components

when contrasted with a word with stress on the first syllable (Honbolygó, Csépe, & Ragó,

2004). In a subsequent study (Honbolygó and Csépe, 2013), it was also shown that

pseudowords with stress on the second syllable elicited two consecutive MMN components,

while pseudowords with a familiar stress pattern in a deviant position did not elicit an MMN,

suggesting that stress processing is modulated by top-down processes, supporting the

importance of familiarity in word stress processing. This study was replicated using

meaningful words (Garami, Ragó, Honbolygó, & Csépe, 2017), demonstrating that lexicality

facilitates the processing of familiar stress patterns but not the unfamiliar ones. These

studies provided further evidence that Hungarian speakers rely on long-term

representations of the fixed Hungarian stress pattern that appear to contain a description of

the expected word stress pattern characteristic for a given language at the phonological

level. However, the language-specific nature of these representations has not been directly

investigated in a cross-linguistic experiment so far; therefore, the present study specifically

aimed to clarify this aspect.

The MMN paradigm has also been used to investigate language-specific processing of

linguistic representations. It has been demonstrated in several studies (Dehaene-Lambertz,

1997; Kraus & Cheour, 2000; Näätänen et al., 1997; Winkler et al., 1999) that native

language phonemes elicit enhanced MMN as compared to non-native phonemes, indicating

the modulation of speech processing mechanisms by the exposure to native language.

Chandrasekaran, Krishnan, & Gandour (2007) used the MMN paradigm to study the effect of

language experience on the processing of pitch contour in Mandarin Chinese in a cross-

linguistic study. According to the results, native Chinese speakers showed a larger MMN

component to highly dissimilar tones than non-native (English) speakers, demonstrating that

the MMN is modulated by the long-term experience with the native language.

In the present study, we also used the MMN paradigm to investigate the language-specific

nature of word stress representations in native speakers of Hungarian. In the study, we used

naturally produced, meaningless pseudowords having stress either on the first or the second

syllable. Moreover, we created a native and a non-native (German) version of both types of

pseudowords in order to study the ERP correlates of non-native stress processing. We

assumed that listeners would detect preattentively if a pseudoword was pronounced in a

native or non-native way, and only the native-like stimuli will activate the long-term stress

representation. We expected that the activation of the long-term representation would be

indicated by not recognizing the native legal stress patterns as deviant (as they do not differ

from the expected stress pattern specified in the long-term representation). Based on these

assumptions, we hypothesized that pseudowords stressed on the second syllable (illegal

stress pattern in Hungarian) would elicit the MMN component irrespective of the language.

However, we expected that pseudowords stressed on the first syllable (legal stress pattern in

Hungarian) would elicit the MMN component only when the non-native version is

presented, and no MMN would be elicited by the native legal pseudoword, because listeners

would not detect this pattern as irregularity.

**Methods**

**Participants**

Thirty-two healthy young adults took part in the experiment. Two of them were excluded by

reason of excessive artifacts (see EEG Recording and Pre-Processing). Therefore, thirty

participants remained in the final sample (19 females).  The participants, aged between 18

and 25 years ($M_{Age}$ = 21.7 years, $SD$ = 1.6 years), were native speakers of Hungarian who

have not  studied German as L2 according to a questionnaire screening their linguistic

background (Kóbor et al., 2018). All of them were right-handed according to the Edinburgh

Handedness Inventory revised version (Dragovic, 2004) ($M_{LQ}$ = 90.4, $SD$ = 12.9[1]). Most of the

participants were undergraduate students enrolled in different universities in Budapest

($M_{Years\ of\ education}$ = 15.1 years, $SD$ = 1.6 years). They had normal or corrected-to-normal vision

and normal hearing level according to the audiometry measurement. Prior to their inclusion

in the study, participants provided informed consent to the procedure as approved by the

Ethical Review Committee for Research in Psychology, Hungary. The study was conducted in

accordance with the Declaration of Helsinki. Participants received payment (3300 HUF, ca.

10 Euros) or course credit for taking part in the experiment.

**Stimuli**

Four pseudowords were used as stimuli: two spoken by a Hungarian native speaker and two

spoken by a German native speaker (both were female). The same reiterative pseudoword

'bébé' (/be□□be□/) was recorded by each speaker with two different stress patterns: one

with stress on the first syllable (/be□□be□/ trochaic stress pattern, henceforth referred to as

S1), and one with stress on the second syllable (/be□be□□/ iambic stress pattern, henceforth

referred to as S2). Speakers were asked to produce the pseudowords as naturally as they

could, and no further software editing was done on the stimuli, except for equalizing the

overall loudness (RMS equalization) and applying a 10 ms long onset and offset ramp. The

acoustical properties of stimuli are illustrated in Figure 1 and summarized in Table 1.

---

[1] LQ = Laterality Quotient; LQ = -100 means complete left-handedness, LQ = 100 means complete right-handedness.

INSERT TABLE 1 AROUND HERE

INSERT FIGURE 1 AROUND HERE

**Procedure**

Stimuli were presented in a *passive oddball paradigm* with a stimulus onset asynchrony

(SOA) varying randomly between 850 and 1050 ms. The duration of the S1 stimulus was 746

ms in Hungarian and 573 ms in German, and the duration of the S2 stimulus was 747 ms in

Hungarian and 574 ms in German. In our design, we varied language (Hungarian, German),

stress position (S1, S2) and role (standard, deviant) yielding *eight different conditions*: the S1

and S2 as a standard stimulus and the S1 and S2 as a deviant stimulus in Hungarian and

German, respectively. Stimuli were presented in *four experimental blocks* according to the

following pairings: (1) Hungarian S1 standard with Hungarian S2 deviant; (2) Hungarian S2

standard with Hungarian S1 deviant; (2) German S1 standard with German S2 deviant; (3)

German S2 standard with German S1 deviant. Participants were presented 666 stimuli in

each block with 566 standard stimuli and 100 deviant stimuli (i.e., the probability of the

deviant was 15%). Each block began with at least fifteen standard stimuli, and then deviants

were separated by at least two standard stimuli. We created a different pseudorandom

sequence of stimuli for each block, and within each block, the same sequence of stimuli was

used for all participants. The order of the four experimental blocks was counterbalanced

across participants. During the EEG experiment, participants were seated in a comfortable

chair in an acoustically shielded, dimly lit room. They were instructed to ignore the stimulus

sequence presented via headphones (Sennheiser PX 200) while they were watching a silent

movie.

After the EEG experiment and the removal of the electrode net, a behavioral discrimination

task was administered to check whether the variations in stress position were

distinguishable within a language. Namely, a *same-different paradigm* with two blocks was

used: one block tested the Hungarian stimuli, another tested the German stimuli. Stimuli

were presented pairwise with a SOA of 1000 ms. Forty stimulus pairs were presented in each

block: The four possible combinations of stimulus pairs (S1-S1, S2-S2, S1-S2, S2-S1) were

tested ten times. The order of blocks was counterbalanced across participants. Participants

were required to decide as fast and accurate as possible whether the two stimuli presented

as a stimulus pair were the same or different. They had unlimited time to initiate a response

for each stimulus pair. The next stimulus pair was presented after the participant gave

behavioral response. We used a Cedrus RB-740 response pad (Cedrus Corporation, San

Pedro, CA) to record participants' responses in this task. Assignment of response keys to

same/different responses was counterbalanced across participants. The entire experimental

procedure lasted about 2 hours, including the application of the electrode net. Both the EEG

and the behavioral experiment were written in and controlled by the Presentation software

(v. 17.0, Neurobehavioral Systems).

**EEG Recording and Pre-processing**

The continuous EEG activity was recorded using the Electrical Geodesics system (GES 400;

Electrical Geodesics, Inc., Eugene, OR) and Net Station 4.5.1 software. We used a 128-

channel HydroCel Geodesic Sensor Net with saline electrolyte solution (see Figure 2).

Electrode Cz was used as a reference and a sampling rate of 500 Hz was applied.

INSERT FIGURE 2 AROUND HERE

Data were analyzed using BrainVision Analyzer software (Brain Products GmbH, Munich,

Germany) and Matlab 8.5 (MathWorks Inc.). Spline interpolation of bad electrodes was

performed if necessary. Zero – three ($M$ = 0.61) channels per participant were interpolated.

As the first step of pre-processing, EEG data were band-pass filtered offline between 0.3 – 30

Hz (48 dB/oct) and notch filtered at 50 Hz to remove additional electrical noise. Second, we

corrected horizontal and vertical eye-movement artifacts and heartbeats with independent

component analysis (ICA; Delorme, Sejnowski, & Makeig, 2007). Using ICA, 1-6 ICA

components ($M$ = 2.93) were removed, then, the channel-based EEG data were recomposed.

Third, EEG data were re-referenced to the average activity of all electrodes. After that, the

continuous EEG was segmented into epochs. Epochs extended from -100 to 1000 ms relative

to the onset of the standard and deviant stimuli, respectively, in each condition. An equal

number of epochs (n = 100) were selected for both the standard and deviant stimuli in each

condition by analyzing only the standards preceding the deviants. To remove artifacts still

present in the data after ICA corrections, we used an automatic artifact rejection algorithm

implemented in BrainVision Analyzer software. This algorithm rejected epochs where the

activity exceeded +/- 100 μV at any of the electrode sites. Only those participant's data were

included in further analysis where the percentage of rejected epochs was below 30% in any

of the conditions. Accordingly, two participants' data were excluded. The mean number of

retained epochs collapsed across all conditions was 95 out of the possible 100 ($SD$ = 5.4;

range: 72 – 100). After artifact rejection, artifact-free epochs were baseline corrected based

on the mean activity from -100 to 0 ms. Finally, the remaining epochs were averaged.

**Data/ERP Analysis**

ERP analysis involved a permutation based non-parametric topographic analysis of variance

(TANOVA, Murray, Brunet, & Michel, 2008) conducted in Ragu (Koenig, Kottlow, Stein, &

Melie-García, 2011) on the individual ERP waveforms. The aim of the analysis was to

determine those latency ranges where there are significant differences between the ERP

curves related to the effect of interest (see below). The TANOVA allows the quantification of

ERP components without selecting time windows arbitrarily (i.e., where the ERP

components' amplitude seems to be maximal or the difference between particular

conditions seems to be maximal) or by following previous literature; therefore, it is and

unbiased approach (see Brooks, Zoumpoulaki, & Bowman, 2017).

The analysis was conducted separately for the Hungarian and German stimuli. The reason for

this was that differences in duration between the stimuli (see Figure 1 and Table 1) made the

point-by-point comparison of the ERPs problematic across the languages, as the shorter first

syllables in German could potentially elicit ERP components with shorter latencies.

We calculated the TANOVA for topographic dissimilarity (TD) (Koenig & Melie-Garcia, 2009;

Wirth et al., 2008), which is a single measure of the distance between two electric field

topographies. We assumed that the ERPs to standard and deviant stimuli and to different

stress patterns would demonstrate slightly different topographic distributions. The TANOVA

was performed with Role (Standard vs. Deviant) and Stress Position (S1 vs. S2) as within-

subjects factors. Standard and deviant responses for the same stimuli (i.e., either S1 or S2)

taken from different conditions were used in the analysis. This approach enabled us to

measure genuine MMN effects as we compared physically identical stimuli within each

language differing only in their role (standard vs. deviant) in the experimental blocks (see

also Honbolygó & Csépe, 2013; Jacobsen & Schröger, 2003).

We performed 1000 randomization runs and applied a 5% significance threshold on the

individual averages during the entire epoch length (no normalization was used). Regarding

the main question of this study, we focused on the Stress Position * Role interaction effect,

which would show the time range where the difference between deviants and standards is

modulated by the stress pattern. As mentioned above, we did not compare effects directly

between languages.

**Results**

**Behavioral results**

One participant's data in the same-different paradigm are missing because of technical

reasons. Thus, behavioral data were analyzed in 29 participants. As Hungarian and German

stimuli differed in duration, we did not compare the RT of responses across the different

contrasts. Instead, accuracy, defined as the percentage of correct responses, was used as a

measure of performance. Responses slower that 2000 ms were excluded from further

analysis, thus, accuracy was calculated without these trials. We found ceiling effect on this

task as the mean accuracy for the same pairs was 99.48% ($SD$ = 1.55%) in the Hungarian

block and 98.79% ($SD$ = 2.18%) in the German block. For the different pairs, this was 98.45%

($SD$ = 3.02%) in the Hungarian block and 99.12% ($SD$ = 2.38%) in the German block.

Therefore, participants were able to discriminate pseudowords with different stress pattern

both in Hungarian and German.

**ERP results**

Grand average ERP waveforms to the standard and deviant stimuli and the difference curves

split by language and stress position are presented in Figure 3. As it can be seen in the figure,

stimuli elicited a complex set of ERP waveforms, starting with positive ERP deflections that

went in the negative direction from around 200-300 ms. ERPs to the stimuli in the standard

and deviant role started to deviate from each other already from around 100 ms.

INSERT FIGURE 3 AROUND HERE

Difference curves calculated by subtracting the ERPs to the standard from the ERPs to the

deviant elicited by the same stimuli are depicted on Figure 4. Difference waves were

calculated as follows: (S1 HU) Hungarian S1 deviant minus Hungarian S1 standard; (S2 HU)

Hungarian S2 deviant minus Hungarian S2 standard; (S1 GE) German S1 deviant minus

German S1 standard; (S2 GE) German S2 deviant minus German S2 standard. The heat maps

under the ERPs in Figure 4 show those latency ranges where the TANOVA indicated

significant effects (main effect or interaction) for scalp topographies (colors denote the *p*

values).  We found a significant Stress position * Role interaction effect only for the

Hungarian stimuli between 176-254 ms and 374-418 ms. There was no significant interaction

effect for the German stimuli. These two latency windows correspond to two negative ERP

deflections, and the significant interaction is due to S1 HU not eliciting or eliciting smaller

deflections. As for the German stimuli, these two deflections are also present, but they are

not different for the S1 and S2 stimuli. A more detailed presentation of the TANOVA results,

and the ANOVA results of the ERP data in the above time windows for both languages

separately are presented in the supplementary material.

INSERT FIGURE 4 AROUND HERE

**Discussion**

This study investigated the language-specific nature of word stress representations in a

fixed-stress language. We studied the ERP correlates of the processing of disyllabic

pseudowords having stress on the first or the second syllable produced by either a native

Hungarian or a native German speaker. Using a data-driven analysis approach, we found two

time windows that demonstrated stimulus-related effects.

The ERP activity in the first (176-254 ms) and second (374-418 ms) time windows showed a

difference between the standard and the deviant stimuli in all conditions except for the

Hungarian stress on the first syllable (S1) stimulus. This means that Hungarian listeners

detected the difference between the native stress patterns when the stimulus had stress on

the second syllable (S2) but not when the stress was on the first syllable. At the same time,

they detected the difference between the German stress patterns in both S1 and S2. We

interpret these results as evidence for language-specific processing of stress patterns. We

suggest that the lack of Hungarian S1 effect means that the legal stress patterns was

processed as not being deviant. Based on our earlier results (Garami et al., 2017; Honbolygó

& Csépe, 2013; Honbolygó, Kolozsvári, & Csépe, 2017), we assume that the legal stress

pattern is processed in relation to long-term stress representations, so called stress-

templates. As we argued previously (Honbolygó & Csépe, 2013), the lack of MMN in S1

cannot be explained simply with reference to acoustic differences between the stimuli, as in

the case of the trochaic stress pattern, the deviant actually contains more salient acoustic

features than the standard.

We suggest that the ERP activity in the first time window is not an MMN component, we

rather consider it to be an Early Differentiating Negativity (EDN). The sensitivity of the EDN

to feature changes was investigated by Kuuluvainen, Alku, Makkonen, Lipsanen, and Kujala

(2016) who studied the discrimination of five syllable features (consonant, vowel, vowel

duration, fundamental frequency, and intensity), including both segmental and prosodic

phonetic changes in pre-school children. They found that the EDN was elicited around 100-

200 ms after stimulus onset by all feature changes except for the vowel duration, and non-

speech consonant and f0 changes. Because the EDN was larger for speech than for non-

speech stimuli, the authors suggested that this component was related to speech-specific

processing mechanisms. They left open the possibility that the EDN could reflect a difference

in the neural refractoriness resulting from enhanced N2s for the novel deviant features. In

another study investigating stress discrimination (Cunillera, Toro, Sebastián-Gallés, &

Rodríguez-Fornells, 2006), the authors found an increased N2 component for the initially

stressed stimuli compared with the initially unstressed ones. However, in our study, the EDN

component was larger for the initially unstressed syllable compared with the initially

stressed one, which suggests that the negative deflection appearing in our study is different

from the N2 components. Since data on the EDN component are still scarce, we have to be

cautious about the functional interpretation.  However, as our data suggest, the EDN is

sensitive to both the acoustical information and the regularity of the stimuli.

We considered the ERP deflection in the second time window (374-418 ms) as a genuine

MMN component. This component has been previously found in relation to stress

processing both in our studies investigating Hungarian language  (Garami et al., 2017;

Honbolygó & Csépe, 2013; Honbolygó et al., 2017) and in other studies investigating Finnish

(Ylinen et al., 2009), German (Weber et al., 2004), and English (Peter et al., 2012) languages.

As described above, the MMN component was elicited in the Hungarian S2 and German S1

and S2 conditions but not in the Hungarian S1 condition, indicating that stress-related

linguistic information was processed in a language-specific way. The functional difference

between the EDN and MMN is currently unclear and requires further investigation.

However, based on the temporal relations between the syllables and the ERP deflections

(see Figure 3) we can speculate that the EDN and MMN reflect the processing of different

information: because its latency is very short, the EDN is probably related to processing the

onset of the first syllable, while the MMN may be related to the processing of acoustic-

phonetic changes inside the first syllable. Because the second syllable starts around 360 ms,

it is unlikely that the MMN would be related to the processing of that.

These results can be interpreted in the AERS model described in the Introduction (Winkler &

Schröger, 2015). In our case, the lack of ERP difference in the two time windows for the

Hungarian S1 can be explained by assuming that the native stress pattern in the deviant

position did not start the updating process because it was fully predicted by the model based

on long-term stress-related representations. Thus, in the model, the long-term

representations overwrote or overweighed the short-term representations based on the

stimulus sequence, in which the deviant stimuli still differed from the standard. In all other

cases, the incoming deviant stimuli did start the updating process, because they violated

both the short-term (standard vs. deviant) and the long-term (irregular native stress, regular

and irregular non-native stress) regularities.

**Summary and conclusion**

In summary, we found evidence for the language-specific processing of word stress patterns

indicated by the lack of early ERP (EDN and MMN) activities to the native stress pattern in

the deviant position. Our results are compatible with a predictive coding model of auditory

event representations and suggest that the word stress is processed predictively, based on

long-term memory traces activated only by native language input. Further studies are

required to extend these findings to languages with variable stress, and to show the

limitations imposed by the native memory traces when learning a second language with

different stress pattern.

**Acknowledgment**

The authors are thankful for Gabriella Baliga, Renáta Szűcs, Lívia Elek for their help in EEG

recording and data collection, and Dorottya Gyarmathy and Mária Gósy for their help in

recording the stimuli. The study was supported by the Hungarian Scientific Research Fund

(OTKA K 119365, PI: V.Cs.), the János Bolyai Research Fellowship of the Hungarian Academy

of Sciences (F.H.), and the Postdoctoral Fellowship of the Hungarian Academy of Sciences (to

A.K.)

**References**

Bornkessel-Schlesewsky, I., & Schlesewsky, M. (2013). Reconciling time, space and function:

A new dorsal-ventral stream model of sentence comprehension. *Brain and Language*,

*125*(1), 60–76. https://doi.org/10.1016/j.bandl.2013.01.010

Brooks, J. L., Zoumpoulaki, A., & Bowman, H. (2017). Data-driven region-of-interest selection

without inflating Type I error rate. *Psychophysiology*, *54*(1), 100–113.

https://doi.org/10.1111/psyp.12682

Chandrasekaran, B., Krishnan, A., & Gandour, J. T. (2007). Mismatch negativity to pitch

contours is influenced by language experience. *Brain Research*, *1128*(1), 148–156.

https://doi.org/10.1016/j.brainres.2006.10.064

Cunillera, T., Toro, J. M., Sebastián-Gallés, N., & Rodríguez-Fornells, A. (2006). The effects of

stress and statistical cues on continuous speech segmentation: An event-related brain

potential study. *Brain Research*, *1123*(1), 168–178.

Cutler, A., & Norris, D. (1988). The Role of Strong Syllables in Segmentation for Lexical

Access. *Journal of Experimental Psychology: Human Perception and Performance*, *14*(1),

113–121.

Dehaene-Lambertz, G. (1997). Electrophysiological correlates of categorical phoneme

perception in adults. *NeuroReport*, *8*(4), 919–924.

Delorme, A., Sejnowski, T., & Makeig, S. (2007). Enhanced detection of artifacts in EEG data

using higher-order statistics and independent component analysis. *NeuroImage*, *34*,

1443–1449.

Dragovic, M. (2004). Towards an improved measure of the Edinburgh Handedness Inventory:

A one-factor congeneric measurement model using confirmatory factor analysis.

*Laterality: Asymmetries of Body, Brain and Cognition*, *9*(4), 411–419.

https://doi.org/10.1080/13576500342000248

Garami, L., Ragó, A., Honbolygó, F., & Csépe, V. (2017). Lexical influence on stress processing

in a fixed-stress language. *International Journal of Psychophysiology*, *117*, 10–16.

https://doi.org/10.1016/j.ijpsycho.2017.03.006

Honbolygó, F., & Csépe, V. (2013). Saliency or template? ERP evidence for long-term

representation of word stress. *International Journal of Psychophysiology*, *87*(2), 165–

172. https://doi.org/10.1016/j.ijpsycho.2012.12.005

Honbolygó, F., Csépe, V., & Ragó, A. (2004). Suprasegmental speech cues are automatically

processed by the human brain: a mismatch negativity study. *Neuroscience Letters*,

*363*(1), 84–88. https://doi.org/10.1016/j.neulet.2004.03.057

Honbolygó, F., Kolozsvári, O., & Csépe, V. (2017). Processing of word stress related acoustic

information: A multi-feature MMN study. *International Journal of Psychophysiology*,

*118*, 9–17. https://doi.org/https://doi.org/10.1016/j.ijpsycho.2017.05.009

Jacobsen, T., & Schröger, E. (2003). Measuring duration mismatch negativity. *Clinical*

*Neurophysiology*, *114*(6), 1133–1143. https://doi.org/10.1016/S1388-2457(03)00043-9

Kager, R. (2007). Feet and metrical stress. In P. de Lacy (Ed.), *The Cambridge Handbook of*

*Phonology* (pp. 195–228). Cambridge University Press, Cambridge.

Kóbor, A., Honbolygó, F., Becker, A. B. C., Schild, U., Csépe, V., & Friedrich, C. K. (2018). ERP

evidence for implicit L2 word stress knowledge in listeners of a fixed-stress language.

*International Journal of Psychophysiology*, *128*, 100–110.

https://doi.org/10.1016/j.ijpsycho.2018.04.006

Koenig, T., Kottlow, M., Stein, M., & Melie-García, L. (2011). Ragu: A free tool for the analysis

of EEG and MEG event-related scalp field data using global randomization statistics.

*Computational Intelligence and Neuroscience*, *2011*, 1–15.

https://doi.org/10.1155/2011/938925

Koenig, T., & Melie-Garcia, L. (2009). Statistical analysis of multichannel scalp field data. In

*Electrical Neuroimaging* (pp. 169–189). Cambridge University Press New York.

Kraus, N., & Cheour, M. (2000). Speech sound representation in the brain. *Audiology &*

*Neuro-Otology*, *5*(3–4), 140–150. https://doi.org/13876

Kuuluvainen, S., Alku, P., Makkonen, T., Lipsanen, J., & Kujala, T. (2016). Cortical speech and

non-speech discrimination in relation to cognitive measures in preschool children.

*European Journal of Neuroscience*, *43*(6), 738–750. https://doi.org/10.1111/ejn.13141

Murray, M. M., Brunet, D., & Michel, C. M. (2008). Topographic ERP analyses: A step-by-step

tutorial review. *Brain Topography*, *20*(4), 249–264. https://doi.org/10.1007/s10548-

008-0054-5

Näätänen, R., Lehtokoski, A., Lennes, M., Cheour, M., Huotilainen, M., Iivonen, A., … Alho, K.

(1997). Language-specific phoneme representations revealed by electric and magnetic

brain responses. *Nature, 385*, 432–434.

Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in

basic research of central auditory processing: a review. *Clin Neurophysiol, 118*(12),

2544–2590.

Peter, V., Mcarthur, G., & Thompson, W. F. (2012). Discrimination of stress in speech and

music: A mismatch negativity (MMN) study. *Psychophysiology, 49*(12), 1590–1600.

https://doi.org/10.1111/j.1469-8986.2012.01472.x

Poeppel, D. (2014). The neuroanatomic and neurophysiological infrastructure for speech and

language. *Current Opinion in Neurobiology, 28*, 142–149.

https://doi.org/10.1016/j.conb.2014.07.005

Weber, C., Hahne, A., Friedrich, M., & Friederici, A. D. (2004). Discrimination of word stress

in early infant perception: Electrophysiological evidence. *Cognitive Brain Research*,

*18*(2), 149–161.

Winkler, I., Denham, S. L., & Nelken, I. (2009). Modeling the auditory scene: predictive

regularity representations and perceptual objects. *Trends in Cognitive Sciences, 13*(12),

532–540. https://doi.org/10.1016/j.tics.2009.09.003

Winkler, I., Lehtokoski, A., Alku, P., Vainio, M., Czigler, I., Csépe, V., … Näätänen, R. (1999).

Pre-attentive detection of vowel contrasts utilizes both phonetic and auditory memory

representations. *Cognitive Brain Research, 7*(3), 357–369.

Winkler, I., & Schröger, E. (2015). Auditory perceptual objects as generative models: Setting

the stage for communication by sound. *Brain and Language, 148*, 1–22.

https://doi.org/10.1016/j.bandl.2015.05.003

Wirth, M., Horn, H., Koenig, T., Razafimandimby, A., Stein, M., Mueller, T., … Strik, W. (2008).

The early context effect reflects activity in the temporo-prefrontal semantic system:

Evidence from electrical neuroimaging of abstract and concrete word reading.

*NeuroImage*, *42*(1), 423–436. https://doi.org/10.1016/j.neuroimage.2008.03.045

Ylinen, S., Strelnikov, K., Huotilainen, M., & Näätänen, R. (2009). Effects of prosodic

familiarity on the automatic processing of words in the human brain. *International*

*Journal of Psychophysiology*, *73*(3), 362–368.

Zora, H., Heldner, M., & Schwarz, I. C. (2016). Perceptual correlates of turkish word stress

and their contribution to automatic lexical access: Evidence from early ERP

components. *Frontiers in Neuroscience*, *10*(JAN), 1–11.

https://doi.org/10.3389/fnins.2016.00007

Zora, H., Schwarz, I.-C., & Heldner, M. (2015). Neural correlates of lexical stress.

*NeuroReport*, *26*, 791–796. https://doi.org/10.1097/WNR.0000000000000426

537                                          **Tables**

539     Table 1. Acoustical properties of stimuli.

| | | Hungarian | | German | |
|---|---|---|---|---|---|
| | | 1st syllable | 2nd syllable | 1st syllable | 2nd syllable |
| intensity (dB)[1] | S1 | 82.13 | 66 | 78.79 | 70.8 |
| | S2 | 75.57 | 80.96 | 74.35 | 78.78 |
| f0 (Hz) [2] | S1 | 225.32 | 202.65 | 213.61 | 215.44 |
| | S2 | 211.32 | 206.36 | 217.17 | 209.41 |
| duration (ms) | S1 | 360 | 386 | 170 | 403 |
| | S2 | 350 | 396 | 140 | 433 |

540     *Note*. [1]: maximum amplitude measured in the whole length of syllable; [2]: maximum f0 measured in the whole

541     length of syllable

544    Figure captions
545
546    Figure 1. Acoustic characteristics of the stimuli. The figures show the spectrogram, pitch (red
547    line) and intensity (blue line) of the four stimuli. Note the different time axes of Hungarian
548    and German stimuli. *Note*: S1: stimuli with stress on the first syllable; S2: stimuli with stress
549    on the second syllable; HU: Hungarian stimuli; GE: German stimuli.
550
551    Figure 2. The electrode montage with 128 electrodes used in the experiment. The blue
552    ellipsis shows those electrodes (5, 6, 11, 12) that were pooled together in the fronto-central
553    electrode pool used for statistical analysis.
554
Figure 3. Grand average ERP waveforms split by language and stress position to the standard
(Std) and deviant (Dev) stimuli together with the difference wave (Diff) calculated by
subtracting the ERPs to standard stimuli from that to the corresponding deviant stimuli.
Negativity is plotted upwards. ERPs are plotted for the fronto-central electrode pool. The
waveform of the respective stimuli below the ERPs illustrates the temporal relation between
the stimuli and ERP responses. *Note*: S1: stimuli with stress on the first syllable; S2: stimuli
with stress on the second syllable; HU: Hungarian stimuli; GE: German stimuli.
562
Figure 4. Difference waves in the four different conditions split by language (Hungarian: left
side; German: right side) and the TANOVA results. ERPs are plotted for the fronto-central
electrode pool. The results of the TANOVA are depicted below the ERPs. The colored
intervals show those time windows where the TANOVA indicated significant main effects of
Stress Position, Role, and Stress Position * Role interaction. Only those time windows are
shown where the effect was significant for at least 10 consecutive time points. The *p*-value
of the significance is shown as a colored scale in the range of 0-0.05. Shaded areas show the
time windows on the ERPs selected for later analysis (presented in the supplementary
material). *Note*: S1: stimuli with stress on the first syllable; S2: stimuli with stress on the
second syllable; HU: Hungarian stimuli; GE: German stimuli. EDN: Early Differentiating
Negativity.
574
575
576
577