Combining transcriptomic and fitness data reveals adaptive and additive plastic responses of gene expression to multiple stressors in *Tribolium castaneum*.

Eva L. Koch[1] and Frédéric Guillaume[1]

[1] Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstr. 190, 8057 Zurich, Switzerland

Corresponding author:
Prof. Dr. Frédéric Guillaume
tel: ++41 44 635 6623
frederic.guillaume@ieu.uzh.ch

Abstract

Gene expression is known to be highly responsive to the environment and important for

adjustment of metabolism but there is also growing evidence that differences in gene regulation

contribute to species divergence and differences among locally adapted populations. However,

most studies so far investigated populations when divergence already occurred. Not much is

known about the evolutionary processes acting on expression levels when populations first

encounter environmental change. Understanding the mechanisms is further complicated by the

fact that environmental change is often multivariate, meaning that organisms are exposed to

multiple stressors simultaneously, with interactive effects, which can constrain adaptation when

multiple responses cannot be optimized at the same time. Here we use a novel approach by

combining fitness and whole-transcriptome data in a large-scale experiment to investigate

responses to drought, heat and their combination in *Tribolium castaneum*. We found that fitness

was reduced by both stressors and their combined effect was almost additive. Expression data

showed that both stressors were acting independently and did not interfere physiologically.

Since we measured expression and fitness within the same individuals, we were able to estimate

selection on single gene expression levels. We found that variation in fitness can be attributed to

gene expression variation and that dependent on the environment different selection pressures

are acting, which were opposite between control and stress conditions. We could further show

that plastic responses of expression are largely adaptive, meaning they changed in the direction

that should increase fitness.

Keywords: gene expression, plasticity, multiple stressors, selection, *Tribolium castaneum*,

fitness

Introduction

One of the major goals of evolutionary biology is to understand the genetic basis of phenotypic

variation and how it is shaped by natural selection. The mapping of genetic to phenotypic

variation depends on many cellular processes, of which RNA transcript abundance, or gene

expression, has been shown to play a central role (Abzhanov et al. 2004; Prud 'homme et al.

2007; Carroll 2008; Fay and Wittkopp 2008; Romero et al. 2012). For variation in expression to

be relevant for evolution it needs a heritable genetic basis and a link with fitness variation. While

the heritability of expression variation has been established in many cases (Cui et al. 2006;

Ayroles et al. 2009; Skelly et al. 2009; Long et al. 2012; Morris et al. 2014; McCairns et al. 2016),

we still lack direct estimates of the strength of selection acting on transcript level abundance.

The link between expression levels and fitness variation is not obvious, if not doubted, since

mRNA abundance must be translated into protein abundance, enzyme activity and ultimately phenotypic variation (Feder and Walser 2005; Evans 2015). So far, the evidence for a link between fitness and gene expression variation is mixed and mostly indirect. For instance, in yeast, the knocking-out of many genes first had mostly non-consequential effects on fitness (Giaever 2002; Benoit et al. 2009). However, more recent evidence in yeast show that variation in expression can significantly affect fitness (Keren et al. 2016). Unfortunately, data in more complex organisms are still scarce, especially on a transcriptome-wide scale. Indirect evidence supporting the importance of gene expression in evolution comes from studies showing differences in expression levels between adaptively divergent populations in yeast (Townsend et al. 2003), humans (Fraser 2013), *Drosophila* (Hutter et al. 2008) or fish (Whitehead and Crawford 2006; McCairns and Bernatchez 2009; Morris et al. 2014; Dayan et al. 2015; Ghalambor et al. 2015; Leder et al. 2015). In such cases, further support can be brought when evolved differences in regulatory DNA sequences are found (Zheng et al. 2011; Sato et al. 2016). Additionally, experimental evolution approaches were successful to detect altered expression levels within a few generations in multiple organisms that adapted to different environmental conditions (Riehle et al. 2003; Telonis-Scott et al. 2009; Yampolsky et al. 2012; Ghalambor et al. 2015; Huang and Agrawal 2016).

In contrast to expression differentiation caused by genetic divergence (Brem et al. 2002; Fay and Wittkopp 2008; Romero et al. 2012), phenotypic plasticity is also expected to play an important role in expression differentiation, especially at the onset of adaptation to novel environments (Yeh and Price 2004; Lande 2009; Chevin et al. 2010; Ehrenreich and Pfennig 2016). Expression is a highly plastic trait and is involved in the immediate response of organisms to changes in their environment (Gasch et al. 2000; Chen et al. 2003; Gibson 2008; López-Maury et al. 2008; Des Marais et al. 2013). The role of plasticity in evolution is, however, contentious. It is often argued that if plasticity is adaptive, it should impede evolution since it can hide genetic variance on which selection would act, thus weakening selection (Price et al. 2003). Yet, plasticity is also key for population persistence in a changing environment because it can keep populations at higher sizes, or buffer novel variants against purifying selection (Chevin et al. 2010; Fitzpatrick 2012), thus facilitating long-term adaptation (Yeh and Price 2004; Lande 2009; Ehrenreich and Pfennig 2016).Studies comparing plastic and evolved responses of gene expression in natural populations (Sørensen et al. 2001; Dayan et al. 2015; Ghalambor et al. 2015) repeatedly found that plasticity was non-adaptive because opposed in direction to the evolutionary response. It may thus be that non-adaptive plasticity facilitates evolutionary divergence by increasing the strength of selection (Ghalambor et al. (2007, 2015)).

However, these studies and others compared expression responses of non-adapted individuals to adapted populations or selection lines in common garden experiments (Whitehead and Crawford 2006; McCairns and Bernatchez 2009; Morris et al. 2014; Ghalambor et al. 2015; Leder et al. 2015), thereby examining patterns when divergence has already occurred. They provide little information on how evolutionary forces have acted in the past and shaped expression but show only the current state after divergence. Studying organisms that have been exposed to new conditions recently can give us more insight into the initial processes leading to divergence between populations experiencing novel environmental conditions and how changes in transcription may contribute to it. In particular, it is still unknown how short term selection pressures are linked to long term optimum expression levels. The two may differ because organisms may first activate immediate stress responses that are beneficial and thus adaptive when they appear, but will not persist because costly to maintain on the long term resulting in quite different responses (Moya et al. 2012, 2015), especially for negative stress effects like slowing down of cell cycle and protein synthesis, and protein damage (Enzor and Place 2014; Huth and Place 2016). In contrast to a first-step stress response, long term adaptation necessitates to optimally re-allocate energy resources between reproduction and stress protection. Optimal solutions for this trade-off between reproduction and survival might also differ between conditions (Schlichting and Smith 2002) and because environmental changes in nature are often multivariate, responses may be constrained by trade-offs among multiple stressors. A beneficial response elicited by one environmental factor may be overridden by a negative effect in presence of a second factor and generate a pattern of non-adaptive plasticity. The joint effects of stress factors may not be simply deduced from single responses and may result in complex interactions when in combination (Crain et al. 2008; Holmstrup et al. 2010; Byrne and Przeslawski 2013; Gobler et al. 2014; Gunderson et al. 2016). It is thus crucial to understand the trade-offs faced by organisms when adapting to new environments (DeBiasse and Kelly 2016; Kelly et al. 2016). Transcriptomics can give us insight into the mechanisms underlying such trade-offs. It can potentially show which physiological processes are activated, thereby giving us information about how resources are used. These trade-offs in stress responses can not only limit plastic responses but may also constrain evolution and future adaptation (Etterson and Shaw 2001; Pörtner et al. 2006; Kelly et al. 2016). Therefore, linking the immediate plastic responses of gene expression with their fitness effects at the onset of adaptive divergence should allow a better understanding of their adaptive value and evolutionary consequences.

In this study we asked how *Tribolium castaneum* (the red flour beetle) was affected by heat and drought in single stressor treatments and in a combination treatment. We combined a fitness assay with measuring gene expression using RNA-seq. This allowed us to get more insight into the physiological processes affected by different stressors, to examine whether the transcriptomic responses to heat and drought overlapped and if gene expression was changed in the same direction by both stressors. Based on this, we can make predictions about how stressors will act in combination and test this in the combined stressor treatment. Since we measured expression and fitness in the same individuals in sufficient sample size, we were able to estimate transcriptome-wide selection intensities on single-gene expression levels giving us an unprecedented overview of selection pressures on gene expression in novel environments. Further, we were able to examine whether plastic responses were adaptive or non-adaptive, thus contributing to our understanding of how plasticity will affect future adaptation.

Results

We used a *T. castaneum strain* (Cro1) (Milutinović et al. 2013), which was collected from a wild population in 2010 and adapted to standard control conditions (33 °C, 70% relative humidity (r.h.)) since then. We measured the number of adult offspring as well as gene expression changes relative to control conditions when the beetles were exposed to a drought, a heat and a combined heat-drought treatment (conditions: Dry: 33°C, 30% r.h.; Hot: 37 °C, 70 % r.h; Hot-Dry: 37°C, 30% r.h.). Individuals were transferred to treatments in the egg stage and stayed there during their whole life time. Fitness and expression were measured in females at the age of eleven weeks. Gene expression was measured by RNA-seq using mRNA of the whole body.

Fitness assay

The proportion of reproducing females was significantly different between conditions (χ^2= 627.35, df=3, P< 2.2e-16). The highest proportion of non-reproducing females was found in Hot (Fig. 1A, Tab. 1). Offspring number of reproducing females decreased with increasing temperature ($F_{1,5157}$ = 1981.07 , P < 2.2e-16) and decreasing humidity ($F_{1,5184}$ = 262.05, P < 2.2e-16), with a stronger effect of heat (-15.98 ± 0.57 SE) than of drought (-5.12 ± 0.50 SE). The lowest offspring number was found when heat and drought were combined (Fig. 1B). Interaction between temperature and humidity was also significant ($F_{1,5128}$= 8.37 , P = 0.003835) and led to an additional decrease of -2.22 ± 0.77 compared to purely additive effects.

Figure 1 Effect of temperature and humidity on reproduction. Proportion of reproducing females in four different conditions: Control (33°C, 70 % r. h.), Dry (33°C, 30 % r. h.), Hot (37°C, 70% r. h.), Hot-Dry (37°C, 30% r. h.) (A). Offspring number of reproducing females. Females could lay eggs for one week and adult offspring was counted five weeks later (B).

Table 1: Results of the fitness assay. Number of adult offspring that females produced within one week of egg-laying in different conditions (Control: 33°C, 70% relative humidity (r.h.).; Dry: 33°C, 30% r.h.; Hot: 37°C, 70% r.h.; Hot-Dry: 37°C, 70% r.h.). For calculating offspring number per female only reproducing females were used

Condition	N	females with offspring	proportion of reproducing females (± SE)	offspring per reproducing female (±SE)
Control	1575	1514	97.62 ± 0.50 %	48.99 ± 0.44
Dry	1642	1603	98.66 ± 0.25 %	43.87 ± 0.43
Hot	1401	1005	76.38 ± 2.70 %	33.01 ± 0.52
Hot-Dry	1567	1396	92.66 ± 1.01 %	25.68 ± 0.45

Gene expression response to heat is stronger than to drought, with little functional overlap

The number of differentially expressed (DE) genes relative to Control was lowest in Dry and largest in Hot (Fig. 2, Tab. 2). Drought induced up-regulation of 52 and down-regulation of 48 genes. In contrast, the response to heat was much more pronounced, with a significantly higher number of DE genes than in Dry (up: 1594, down: 1255; permutation test $P < 0.004$). The overlap between heat and drought responses included only 26 genes (Fig. 2). To investigate whether the DE genes were involved in specific biological processes, we performed pathway, protein

domains, and Gene Ontology (GO) enrichment tests. Because of a low number of DE genes in Dry, only few enrichments could be detected (Supplementary Material S1, S2). Genes up-regulated by drought showed the strongest enrichment in active ion transmembrane transporter activity (GO:0022853), down-regulated genes in hydrolase activity, hydrolyzing O-glycosyl compounds (GO:0004553) and protein family Thaumatin (IPR001938). Analysis of the *Tribolium* genome had revealed a high number of genes thought to be involved in endocrine regulation of diuresis, including several that encode putative neuroendocrine peptides like antidiuretic factors (Aikins et al. 2008; Hauser et al. 2008; Li et al. 2008; Park et al. 2008). Nevertheless, none of these genes was found to respond to the Dry treatment.

Genes up-regulated in Hot were enriched in many metabolic processes e.g. carbohydrate metabolic process (GO:0005975), Citrate cycle (KEGG 00020), Pyruvate metabolism (KEGG 00620) (Supplementary Material S1, S2). The most strongly enriched category was chitin metabolic process (GO:0006030). A protein domain analysis also showed significant enrichment of heat shock proteins (IPR031107, IPR018181, IPR008978) (Supplementary Material S2). Down-regulated genes were enriched in pathways for DNA replication (KEGG 03030), nucleotide excision repair (KEGG 03420) and Ubiquitin mediated proteolysis (KEGG 04120).

Table 2: Number of differently expressed genes (FDR< 5 %) when comparing different conditions. Positive: higher expression in second condition. Negative: lower expression in second condition. Differential expression analysis was conducted in edgeR (Robinson et al. 2010).

	positive	negative	total
Dry vs Control	52	48	100
Hot vs Control	1594	1255	2849
Hot-Dry vs Control	1866	1705	3571
Hot vs Dry	1553	1290	2843
Hot-Dry vs Dry	1928	1813	3741

Figure 2 Venn-Diagram showing the number of differentially expressed genes in three treatments relative to control conditions. Overlapping regions represent genes that were found in more than one treatment and changed expression levels in the same direction. Sizes of circles as well as of overlapping regions are proportional to number of genes.

Response to stressor combination is dominated by heat response, single stress responses are mainly not modified in combination

When both stressors were combined in Hot-Dry, we found 3571 DE genes (up: 1866, down: 1705). Among them, 1515 (42.4%) were not found in single stressor treatments. However, only 69 of those genes (up: 30, down: 39) were found significantly DE between Hot-Dry and Dry, or Hot-Dry and Hot. This indicates that in most cases the combined stress did not induce expression changes in a different set of genes but modified their expression levels over and above their responses to single stressors. Compared to Hot, the Hot-Dry response had a significantly higher magnitude of expression change (permutation test: $P < 0.0001$), a higher number of down-regulated genes ($P = 0.007$), but a similar number of up-regulated and total number of DE genes ($P = 0.29$ and, $P = 0.054$, respectively). The functional response to Hot-Dry resembles the response to Hot, but more enriched GO categories and pathways could be found (see Supplementary Material S1, S2).

To further examine how single stress responses are modified during combination, we classified the top 1000 most responding transcripts (based on p-values) of each treatment into different response categories following Rasmussen et al. (2013). Only 8% of the genes showed a *similar* response mode, with same response to Dry, Hot and Hot-Dry (Fig. 3). Most responding genes (56 %) were classified as *independent*, with a response that is not altered in presence of a second

stressor. Most of those genes showed the same response in Hot and Hot-Dry (44.5 %) but no

response in Dry, in agreement with our DE analysis. Few genes (6 %) had a *combinatorial*

response mode in which they did not respond to heat and drought alone, but to their

combination. These represent cases in which presence of an additional stressor magnifies the

effect of another. Especially interesting are cases with opposite responses to both stressors but

with one response *prioritized* when combined, which can help to identify trade-offs. We found

13 % of DE genes falling into that category, most of them showing prioritization of the Hot

response in Hot-Dry (Supplementary Material S4), also in agreement with our DE analysis. But

more common than *prioritized* is the *cancelled* mode with 18 % of the genes, which show

opposite responses to both stressors or respond to one stressor only but return to control levels

in combination, showing no response to stress combination.

Figure 3: Response modes of the 1000 most significantly responding transcripts in the stress

treatments (Dry, Hot, Hot-Dry. *Combinatorial*: Similar levels in the two individual stresses but a

different response to combined stresses; *cancelled*: transcript response to either or both

individual stresses returned to control levels; *prioritized*: opposing responses to the individual

stresses and one stress response prioritized in response to combined stresses; *independent*:

response to only one single stress and a similar response to combined stresses; *similar*: similar

responses to both individual stresses and to combined stresses. Subcategories of different

response modes giving more details about the most prevalent patterns in supplementary

material (S4).

Immune response: Interaction between drought and heat

We noticed that among the genes up-regulated in Hot many are involved in immune response (Oppert et al. 2012; Behrens et al. 2014). These transcripts encode putative chitin-interacting proteins (binding, deacetylase), allergens, thaumatins, glucosidases, and serine proteases, whereas these genes are down-regulated at low humidity. For instance, Thaumatin protein family, which is involved in antimicrobial response (Altincicek et al. 2008) showed the strongest enrichment in down-regulated genes (Supplementary Material S2) and a pathogenesis related protein and several antimicrobial peptides are among those genes that showed the strongest response in Hot. We applied a gene set enrichment analysis to test for an overrepresentation of immune genes in responses to the different treatments. To avoid including immune genes also involved in heat stress response (Behrens et al. 2014), we used a test set of 15 antimicrobial peptides, which should be specific to a pathogen response, based on Zou et al. 2007, Oppert et al. 2012, Altincicek et al. 2013 (Supplementary Material S3A). We found that antimicrobial peptides were significantly up-regulated in Hot (p-value = 0.001) and down-regulated in Dry (p-value = 0.001) (Tab. 3). When we compared Hot-Dry with Hot, these genes became down-regulated (p-value = 0.002), indicating that it is mainly the combination of high temperature with relatively high humidity (i.e. 70% r.h.) that activates this response, pointing to a possible higher parasite load under these conditions.

Table 3 Results of enrichment analysis of antimicrobial peptides. Gene set enrichment test was conducted in edgeR (Robinson et al. 2010). Prop.Down and Prop.Up give the proportion of genes that are down- and up-regulated. The direction of change is determined from the significance of changes in each direction, and is shown in the Direction column. The p-value provides evidence for whether the majority of genes in the set are DE in the specified direction

Contrast	Prop. Down	Prop. Up	Direction	p-value
Dry-Control	0.5333	0	Down	0.001
Hot-Control	0	0.4667	Up	0.001
Hot-Dry–Control	0.1333	0.2667	Up	0.832
Hot-Dry–Hot	0.4	0.0667	Down	0.002

Treatment effects on reproduction related processes

To get further insights into the molecular processes that link the observed decline in offspring number with transcriptomic data, we focused on the juvenile hormone (JH), 20-hydroxecdysone (20 E), and insulin/insulin-like peptides (IIS) target of rapamycin (TOR) signaling pathways (IIS-

TOR), which are important mediators in the trade-off controlling stress response and

reproduction (Schwenke et al. 2016), along with vitellogenin, which is the main nutrient source

of eggs, and vitellogenin receptors. We found that heat and heat-drought stress, led to a

significant down-regulation of all pathways and repression of vitellogenin as well as vitellogenin

receptors (Supplementary Material S3B). Drought did not show any significant effect. A gene set

test based on a selection of key reproduction genes within these pathways (Supplementary

Material S3C) confirmed that the reproduction genes were mainly down-regulated in Hot and

Hot-Dry (Tab. 4).

Table 4: Results of gene set enrichment analysis of genes involved in reproduction. Gene set

enrichment test was conducted in edgeR (Robinson et al. 2010). Prop.Down and Prop.Up give

the proportion of genes that are down- and up-regulated. The direction of change is determined

from the significance of changes in each direction, and is shown in the Direction column. The *P*-

value provides evidence for whether the majority of genes in the set are DE in the specified

direction. The genes were selected based on (Parthasarathy, Sheng, et al. 2010; Parthasarathy,

Sun, et al. 2010; Xu et al. 2010; Parthasarathy and Palli 2011).

Contrast	PropDown	PropUp	Direction	*P*-value
Dry–Control	0.054	0.036	Up	0.912
Hot–Control	0.464	0.071	Down	0.001
Hot-Dry–Control	0.589	0.071	Down	0.001

Selection on expression levels is environment specific

Since we measured offspring number and transcription within the same individuals, we could

estimate selection intensity on single gene expression levels in each condition separately by

performing a linear regression of relative fitness on standardized expression levels (z-score). In

control conditions, expression levels of 2158 genes showed a negative and 21 a positive and

significant correlation with fitness. The two genes under strongest positive selection coded for

vitellogenin. In Dry, Hot and Hot-Dry we could not detect any significant selection after

correcting p-values for multiple comparisons. To gain a better overview of the variation in

selection pressures among treatments and to avoid stringent significance thresholds on single-

gene fitness-expression correlations, we then compared selection intensities on all genes

between conditions. We found significant negative correlations of selection intensities between

Control and all stress treatments, and positive correlations among stress treatments (p-values <

2.2e-16) (Fig 4). Control and Dry had the strongest negative correlation (-0.24), while Hot and

Hot-Dry had the highest positive correlation (0.34). Furthermore, significantly DE genes

responding to Hot and Hot-Dry were overrepresented among those that were negatively

selected in control conditions (Hot: $\chi2$= 158.62, df = 1, p-value < 2.2e-16, Hot-Dry: $\chi2$= 177.97, df

= 1, p-value < 2.2e-16), with 361 (22.6 %) up-regulated genes in Hot and 417 (22.3 %) in Hot-Dry.

The magnitudes of selection intensities were also significantly different between conditions

(median (and SD) of absolute values in Control: 0.09 (0.06), Dry: 0.03 (0.03), Hot: 0.09 (0.07),

Hot-Dry: 0.05 (0.04); Kruskal-Wallis rank sum test: $\chi2$= 9333.3, df = 3, p-value < 2.2e-16), with the

majority of genes negatively selected in Control, and similar proportions of genes under positive

and negative selection in stress treatments (Fig. 5). Estimated selection intensities were rather

low compared to those that were reported for morphological and life-history traits in natural

populations (Kingsolver et al. 2001; Hereford et al. 2004), but within the range that was

estimated for selection on expression levels (McCairns and Bernatchez 2009). Although the Dry

treatment had the largest negative correlation with Control expression levels and largest number

of genes switching sign relative to Control (9591), it had the lowest magnitude of change in

selection intensity of those genes (median: 0.14). In contrast, the Hot environment had the

strongest magnitude of change in selection intensity compared to Control (median: 0.19, 6644

genes) followed by Hot-Dry (median: 0.16, 7412 genes). Overall, a greater proportion of genes

switched from negative to positive selection (Dry: 0.86, Hot: 0.76, Hot-Dry: 0.8).

Figure 4: Pairwise correlations of selection intensities on single gene expression levels in

different conditions. Blue indicates a positive and red a negative correlation. Values are given in

the lower triangle. Confidence intervals for correlations: Control-Dry: -0.25 ,-0.22; Control-Hot: -

0.11, -0.08; Control–Hot-Dry: -0.23, -0.20; Dry-Hot: 0.08, 0.11; Dry–Hot-Dry: 0.00, 0.03; Hot–Hot-

Dry: 0.33, 0.36.

Figure 5: Distribution of selection intensities on gene expression levels under different

conditions. Linear regression of relative fitness (number of adult offspring) on normalized RNA-

seq read counts (z-score) was used to estimate selection intensity (slope of selection gradient). 0

is marked by the vertical line.

Is response in gene expression adaptive?

To examine whether the plastic responses in gene expression are adaptive, we tested whether

significantly up-regulated genes were more positively selected and significantly down-regulated

genes more negatively selected than non-responding genes. We found that the response to Hot-

Dry was mainly adaptive with up- and down-regulated genes significantly more positively and

negatively selected than non-responding genes, respectively (permutation tests: $P < 0.0001$) (Fig.

6C). In contrast, some parts of the response in Dry seemed maladaptive: Down-regulated genes

were not under significantly different selection, but up-regulated genes were more negatively

selected (Fig. 6A). In Hot, the response is partly adaptive since down-regulated genes were

significantly more negatively selected, while up-regulated genes were not under significantly

different selection (Fig. 6B).

Figure 6: Distribution of selection intensities on gene expression levels under different conditions: Dry (A), Hot (B), Hot-Dry (C). Linear regression of relative fitness (number of adult offspring) on normalized RNA-seq read counts (z-score) was used to estimate selection intensity (slope of selection gradient). Genes that became significantly up-regulated (FDR< 5%) in the respective condition relative to control are represented in green, significantly down-regulated genes in blue and genes that did not show a significant response in grey. Insets show the distribution of differences in the mean selection intensity between down-regulated and not responding genes (left) and up-regulated and not responding genes (right) from 10,000 permutations. Arrows indicate the observed difference.

Discussion

Physiological trade-offs caused by single stressors and their combination

We found that the effects of heat and drought on expression differed in the processes affected and the magnitude of their response. Heat stress triggered a response in a larger number of genes and affected a larger number of biological processes than drought, with little overlap between the two responses. These results are in agreement with many studies showing that heat is a major driver of expression change, especially in ectotherms (Neven 2000; Nguyen et al. 2009; Levine et al. 2011; Chen and Stillman 2012; Morris et al. 2014). The heat-induced expression changes were characteristic of stress responses: Up-regulation of heat shock proteins and metabolic pathways, whereas genes involved in cell growth and cell cycle, and reproduction were repressed (Gasch et al. 2000; Kültz 2005; Kassahn et al. 2009; Enders et al. 2015). This pattern of change in gene expression together with the overall reduction of the reproductive output of the females assayed suggests a shift of resource utilization from growth to stress protection, indicative of a physiological trade-off between stress response and reproduction

(Salmon et al. 2001; Fedorka et al. 2004; Harshman and Zera 2007). We can further confirm that

this physiological trade-off is at play because the decrease of females' reproductive output was

mainly driven by a loss of fecundity (egg number) in the Hot and Hot-Dry treatments and not a

decrease of larvae survival (see Supplementary Material S5), which matches well with the down-

regulation of vitellogenin and the other reproduction-related genes we monitored (JH, the IIS-

Tor and 20E signaling pathways). It has been suggested that those genes indeed play an

important role in mediating the trade-offs between stress-responses and cellular growth and

maintenance (Flatt et al. 2005; Gruntenko and Rauschenbach 2008; Schwenke et al. 2016). The

fitness costs associated with expression of heat shock proteins are also known (Feder and

Hoffman 1999; Silbermann and Tatar 2000; Kassahn et al. 2009). Finally, a trade-off between

immune response and reproduction may also have affected our results. Indeed, we found a

larger proportion of non-reproducing females (Fig.1B, Tab. 1) and an up-regulation of immune

response genes in the Hot treatment. These two results may point to a higher pathogen load

when high temperature (37°C) and relatively high humidity (70% r.h.) are combined, as described

before (Sokoloff 1972). However, the similar or even stronger down-regulation of reproduction-

related genes in the Hot-Dry treatment in which antimicrobial peptides were not up-regulated,

would suggest that the trade-off between reproduction and immune response was not the main

cause of reduction in fecundity in Hot, although it may have plaid a role in the number of

reproducing females. Overall, our analysis shows that different physiological trade-offs between

reproduction and multiple stressors exist in complex environments and elicit gene expression

responses that overlap with each other, resulting in a general decrease of population fitness.

However, precisely separating the responses of multiple stressors is difficult because of the large

pleiotropic effects of the core stress response genes part of the pathways (JH, IIS-TOR, 20E) and

stress response mechanisms (e.g. Hsp) mentioned above.

In contrast to the above energy allocation trade-offs, there was a lack of a strong trade-off

between heat and drought responses of gene expression. We investigated this trade-off by

comparing the single stressor transcriptomic responses to their combined response in the Hot-

Dry treatment. As expected from the small functional and DE genes overlap between the Hot

and Dry treatments, the top responding genes in the Hot-Dry treatment showed an independent

action mode of the individual stress responses as the most common response mode. When

stressors require different protection mechanisms and affect different pathways, it is unlikely

that they interfere with each other during combination and the effect of combination should

thus be additive (Folt and Chen 1999). Accordingly, we found that the observed reproductive

output in the combined Hot-Dry treatment was close to what would be expected for purely

additive effects between Hot and Dry. Detection of a significant interaction between heat and

drought on offspring number was due to high sample size and accordingly high statistical power

and it should be noted that the effect was small (-2.22 ± 0.77). Together with a larger overall

expression response, including stronger down-regulation of reproduction-related genes, this

points to a larger shift of resource allocation from growth to stress protection in the Hot-Dry

compared to the Hot treatment caused by the addition of a milder drought stress to the large

heat stress.

Evolutionary trade-offs in gene expression across environments

Our experiment revealed the fitness effects of the gene expression changes in the new

environments from which evolutionary trade-offs can be uncovered. A shift of the direction of

selection on gene expression variation from one environment to the next is symptomatic to an

evolutionary trade-off and can lead to phenotypic changes between environments when

selection acts on existing genetic variation. We found strong evidence of such switches of the

direction of selection between Control and test environments. The distribution of selection

intensities in Control showed that most expression levels were under negative selection,

probably because gene expression is costly (Castillo-Davis et al. 2002; Wagner 2005; Lang et al.

2009; Frumkin et al. 2017), and selection should therefore favor low expression levels under mild

laboratory conditions to which the beetles were adapted. The distribution changed when

conditions became stressful with a shift towards more transcripts under positive selection (about

80% of the shifting genes, see Results) in the three stress treatments. This translated into strong

negative correlations of selection intensities between Control and treatments. The protection

provided by stress response genes might thus have outweighed the costs of expression. Yet,

more importantly, although stress response trades-off negatively with reproduction and growth,

leading to an overall decrease of fitness in stress treatments, the gene expression change of the

underlying genes is mainly in the direction favored by selection (i.e. negative selection intensities

for down-regulated genes and vice-versa for positive selection, see Fig. 6). This means that the

physiological trade-off in energy allocation may not constrain the evolutionary response of those

genes because their responses were aligned with the direction of selection. There is thus a

strong potential for the evolution of gene expression, mostly in the Hot-Dry and to a lesser

extent in the Hot environments depending on the amount of genetic variance in expression

levels, which we haven't attempted to estimate here.

Plasticity and the evolutionary potential for adaptation

Based on our findings, we could expect divergence in gene expression to evolve during adaptation to stressful conditions compared to the baseline expression in Control, with some convergence of the evolutionary responses in the Hot and Hot-Dry conditions because of the similarity of their expression responses and selection pressures (evident by positive correlation). However, we have here documented the first-step plastic response of the adaptive process to a new stressful environment. Although there is some evidence for a relationship between plastic and evolved responses in gene expression (Riehle et al. 2003; Landry et al. 2007; Swindell et al. 2007; Tirosh et al. 2009; Ragland et al. 2015; Mäkinen et al. 2016), the changes observed may not indicate an evolutionary trend. A common observation is that adaptation reverses first-step plastic responses, as shown in experimental evolution studies (Fong et al. 2005; Rodŕiguez-Verdugo et al. 2016; Ho and Zhang 2018). Stress response is costly and long-term adaptation might involve different genes that are not part of the immediate stress response processes. After adaptation has occurred, stress protection mechanisms are not required anymore and expression reverses to pre-stress levels after rebalancing of the energy allocation trade-off in favor of growth and reproduction (Rodŕiguez-Verdugo et al. 2016). Therefore, previous studies comparing evolved with plastic responses inferred that plastic responses were mainly maladaptive because in a direction opposed to the evolved responses (Ghalambor et al. 2015; Huang and Agrawal 2016; Ho and Zhang 2018). Here, we showed that the first-step plastic responses were adaptive and increased fitness, thereby helping populations to persist, at least in Hot-Dry and partially in Hot. Heat and drought are stressors that *T. castaneum* has certainly encountered in the past (Sokoloff 1972) meaning that there has probably been selection on plastic responses to be adaptive. The results in Dry are less clear. Changes in expression were small and included only a few genes for which the response was not in the direction that should increase fitness. This is consistent with previous studies showing that well-adapted populations have weaker responses in expression (Heckel et al. 2016; Levis et al. 2017). Special anatomical adaptations allow *T. castaneum* to cope with dry conditions by reducing water loss dramatically (King and Denholm 2014), and selection on expression levels was weak in Dry, indicating that modifying transcription will play a minor role during potential future adaptation. More generally, adaptation can occur by either changing the mean of the expression levels or by modifying the extent of plasticity to match the optimal expression level in a new environment. It was found that most genes change rather mean expression than plasticity during adaptation (Yampolsky et al. 2012; Morris et al. 2014), but it was also shown that plasticity in expression could evolve during adaptation to novel conditions (Levine et al. 2011; Morris et al. 2014; Huang and Agrawal

2016; Levis et al. 2017; Passow et al. 2017). With our data it is not yet clear, which of these

mechanisms will be more prevalent or whether they are of equal importance.

Conclusions

We contend that current approaches of comparing already divergently adapted populations

cannot reveal the selection pressures acting at the beginning of divergence when populations

are exposed to different environments. They thus cannot say whether the first-step plastic

changes were adaptive or non-adaptive in the population where they appeared. In contrast, our

experimental approach gives a more accurate account of the selection pressures acting on gene

expression variation at the early onset of adaptive divergence by integrating fitness and

expression data in a large-scale stress-response experiment. Our approach also shows how

transcriptomics can be used to get information about the relative importance of different

stressors, their interaction and the potential constraints acting on plastic and evolutionary

responses when several environmental variables change at the same time. We were thus able to

evaluate the immediate adaptiveness of the plastic changes in gene expression. However,

further work is needed to clarify how we can use plastic responses to predict long-term

evolutionary outcomes, for instance by using long-term evolution experiments.

Material and Methods

Animal rearing and stress treatments

We used the *Tribolium castanuem* Cro1 strain (Milutinović et al. 2013), collected from a wild

population in 2010 and adapted to lab standard conditions (32°C, 70% relative humidity) for

more than 20 generations. Beetles were kept in 24h dark on organic wheat flour mixed with 10%

organic baker's yeast. We sterilized the flour and yeast by heating them for 12h at 80°C before

use. We tested the response of fitness and gene expression of the beetles to heat, drought, and

a combination of both stressors. The conditions in the treatments were: Hot: 37°C and 70% r. h.,

Dry: 33°C and 30% r. h., Hot-Dry: 37°C and 30% r. h. Parents of the studied beetles grew up and

mated in control conditions at the age of four weeks in 15 mL tubes with 1 g of medium. Each

virgin male was mated with a virgin female. After four days, in which the beetles could mate and

lay eggs, each mating pair was transferred to a new vial. We repeated this three times, resulting

in four vials per mating pair containing medium and eggs. Vials of each mating pair were

randomly assigned to the four different conditions, resulting in full-sib families split across all

conditions. Male and female offspring (four females and four males per family and condition)

were separated at the pupal stage and transferred to 10 mL tubes with 1 g of medium and
remained there until they were used for the fitness assay eight weeks later. After the fitness
assay, males and females were transferred to 1 mL tubes, frozen in liquid nitrogen and stored at
-80°C.
489

Fitness assay
To test the effects of the different conditions on fitness, we measured reproduction in 6183
virgin females (ca. 1500 per condition, Table 1). We mated each virgin female with one unrelated
male from the same condition in 15 mL tube with 1 g medium. The male was removed after 24 h.
Females were removed from the tubes after one week of egg laying, and 9 g medium was added
to provide food for the developing offspring. After five weeks the number of offspring was
counted. At this time, all offspring had reached the adult stage. Some females did not produce
any offspring, in proportions that differed between conditions. To test whether there was an
effect of treatment on the number of reproducing and non-reproducing females, we used a
generalized linear mixed model with reproduction success (binomial: offspring/no offspring) as
response and condition as fixed effect. Since some of the tested females and males were full-sibs
and developed within the same tube, we used male and female families as random factors to
account for non-independence due to relatedness and a shared environment during
development. To test how offspring number of reproducing females was influenced by
conditions we used a linear mixed model with offspring number as response, temperature,
humidity and their interaction as fixed effects and female and male family as random factors.
Denominator degrees of freedom were estimated using Satterthwaite approximation. Statistical
analyses were performed using the Lme4 package (Bates et al. 2015) version 1.1-17 in R (R Core
Team 2015).
509

RNA extraction, library preparation and sequencing
183 female beetles with known fitness (ca. 45 per condition), which had been stored at -80°C,
were homogenized in Tri-Reagent® (Zymo Research, California, USA) using an electric bead mill.
RNA was extracted with the RNA Mini Prep kit (Zimo Research, California, USA) following the
instructions of the manufacturer. RNA-quality was checked with the Bioanalyzer 2100 (Agilent,
Waldbronn, Germany) and concentrations were measured with aQubit® Fluorometer (Life
Technologies, California, USA). Libraries were created with 500 ng RNA for each individual
separately with the LEXOGEN mRNA-Seq Library Kit following the manual (LEXOGEN GmbH,
Vienne, Austria). Library quality was checked on a TapeStation (Agilent, Waldbronn, Germany)

and concentrations were determined by qPCR. Libraries were diluted to the same molarity.

Concentrations of dilutions were checked again by qPCR and libraries were pooled (36 libraries

per pool). All treatments were randomized during RNA-extraction, library preparation, and

sequencing. The single-end sequencing was performed in five runs on the Illumina NextSeq 500

(Illumina, Inc, California, USA) using the 75 cycles High Output Kit. Each run resulted in 550-600

million reads that passed the internal sequencer filter. After quality control using FastQC

(www.bioinformatics.bbsrc.ac.uk/projects/fastqc), reads were mapped against the reference

genome(ftp://ftp.ensemblgenomes.org/pub/release30/metazoa/gtf/tribolium_castaneum/Tribo

lium_castaneum.Tcas3.30.gtf.gz) with STAR v.2.5 (Dobin et al. 2013) (adaptors were trimmed

and the first 10 bases were hard trimmed, minimum average quality Q10, minimum tail quality

10, minimum read length 20). We then used FeatureCounts (Liao et al. 2014) to count the

number of reads that mapped to each gene in the reference genome. Mapping as well as read

counting was performed within the data analysis framework SUSHI (Hatakeyama et al. 2016).

Differential expression and enrichment analysis

We conducted a differential expression analysis using the R package edgeR (Robinson et al.

2010). We tested for differently expressed genes between the treatments (Dry, Hot, Hot-Dry)

relative to the control as well as to each other. A gene is classified as DE with a FDR ≤5% after

adjusting for multiple testing (Benjamini and Hochberg 1995). To test whether the number of DE

genes (relative to Control) was significantly different between two environmental conditions a

permutation tests was used. For each permutation entire RNA-seq samples of the two groups

were randomly assigned to conditions and the edgeR analysis repeated. Significance was

assessed by number of times the observed DE number was higher than the DE number obtained

by permutations. To test whether the magnitude of change in expression levels relative to

control was significantly different between Hot and Hot-Dry, we performed a permutation test.

Absolute log2-fold changes of each transcript were randomly assigned to the two groups and

differences in the mean were calculated. We then compared the distribution of differences

obtained by permutations to the observed difference between mean absolute log2-fold changes

in expression. Gene set enrichment analyses for immune response genes and reproduction

related genes were conducted using edgeR. A GO enrichment analysis of DE genes was

performed with gProfiler Version: r1622_e84_eg31 (Reimand et al. 2011) and pathway and

protein domain enrichment analysis with STRING v.10.0 (Szklarczyk et al. 2015).

Classification of response mode

Following Rasmussen et al. (2013), we created 20 predefined expression profiles each

representing a potential response when two single stressors are combined: *Combinatorial*:

similar expression levels in single stress treatments but a different level in stress combination,

cancelled: response to one or both single stressors but expression levels similar to control

conditions when both are combined, *prioritized*: opposite responses to single stressors and

expression levels in combination similar to one of them, *independent*: response to only one

single stressor and the same response in combination, *similar*: same response in single stressor

treatments and combination. For creating predefined expression profiles we used 0 as control

level, 1 and -1 as expression levels for up- and down-regulation. We then tested the correlation

of normalized read counts of the top 1000 genes that showed the most significant responses

(based on significance of the DE analysis) in Dry, Hot and Hot-Dry conditions with each of these

expression profiles and assigned the genes to the category to which they showed the highest

correlation (Pearson correlation).

Selection

We measured selection intensity on gene expression separately for each treatment using

univariate linear regression methods (Lande and Arnold 1983; Brodie III et al. 1995). Fitness

(number of adult offspring) was normalized by dividing each individual value by the mean (w' =

w_i/mean(w)). For each gene, expression levels were first normalized to cpm (counts per million,

TMM normalization) using edgeR and then transformed to standardized z-scores by subtracting

the mean and dividing by the standard deviation ($z = (x_i - mean(x))/ SD(x)$). Resulting regression

coefficients of relative fitness on standardized expression levels give an estimate of the selection

intensity. P-values were corrected for multiple comparisons. To test whether up- and down-

regulated genes were under significantly different selection compared to genes without a

significant response, we used permutation tests. For each permutation (10,000 for each test) we

randomly assigned the categories "not DE" and "up" (or "down" respectively) to each estimated

selection intensity and calculated the difference in mean selection intensity between both

groups. Significance was tested by counting the number of permutations that showed a

difference higher or equal to the observed one.

Acknowledgements

This work was supported by the Swiss National Science Foundation, grants PP00P3_1144846 and

PP00P3_176965 to FG. We thank Sonja Sbilordo, Julian Bauer, Valérian Zeender for helping

during the fitness assay, Maria Domenica Moccia for her help during the fitness assay and RNA

extraction and Lucy Poveda for advice in library preparation and sequencing.

References

Abzhanov A, Protas M, Grant BR, Grant PR, Tabin CJ. 2004. Bmp4 and morphological variation of
beaks in Darwin's finches. *Science* 305(5689):1462–1465.

Aikins MJ, Schooley DA, Begum K, Detheux M, Beeman RW, Park Y. 2008. Vasopressin-like
peptide and its receptor function in an indirect diuretic signaling pathway in the red flour
beetle. *Insect Biochem Mol Biol* 38(7):740–748.

Altincicek B, Elashry A, Guz N, Grundler FMW, Vilcinskas A, Dehne HW. 2013. Next Generation
Sequencing Based Transcriptome Analysis of Septic-Injury Responsive Genes in the Beetle
Tribolium castaneum. *PLoS One* 8(1).

Altincicek B, Knorr E, Vilcinskas A. 2008. Beetle immunity: Identification of immune-inducible
genes from the model insect Tribolium castaneum. *Dev Comp Immunol* 32(5):585–595.

Ayroles JF, Carbone MA, Stone EA, Jordan KW, Lyman RF, Magwire MM, Rollmann SM, Duncan
LH, Lawrence F, Anholt RRH, et al. 2009. Systems genetics of complex traits in Drosophila
melanogaster. *Nat Genet* 41(3):299–307.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using **lme4**. *J
Stat Softw* 67(1):1–48.

Behrens S, Peuß R, Milutinović B, Eggert H, Esser D, Rosenstiel P, Schulenburg H, Bornberg-Bauer
E, Kurtz J. 2014. Infection routes matter in population-specific responses of the red flour
beetle to the entomopathogen Bacillus thuringiensis. *BMC Genomics* 15(1).

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing Testing. *J R Stat Soc Ser B* 57:289–300.

Benoit JB, Lopez-martinez G, Elnitsky MA, Lee RE, Denlinger DL. 2009. Comparative Biochemistry
and Physiology , Part A Dehydration-induced cross tolerance of Belgica antarctica larvae to
cold and heat is facilitated by trehalose accumulation. *Comp Biochem Physiol Part A*
152(4):518–523.

Brem RB, Yvert G, Clinton R, Kruglyak L. 2002. Genetic dissection of transcriptional regulation in
budding yeast. *Science* 296(5568):752–5.

Brodie III ED, Moore AJ, Janzen FJ. 1995. Visualizing and quantifying natural selection. *Trends
Ecol Evol* 10(8):313–318.

Byrne M, Przeslawski R. 2013. Multistressor impacts of warming and acidification of the ocean
on marine invertebrates' life histories. *Integr Comp Biol* 53(4):582–596.

Carroll SB. 2008. Evo-Devo and an Expanding Evolutionary Synthesis: A Genetic Theory of
Morphological Evolution. *Cell* 134(1):25–36.

Castillo-Davis CI, Mekhedov SL, Hartl DL, Koonin E V., Kondrashov FA. 2002. Selection for short
introns in highly expressed genes. *Nat Genet* 31(4):415–418.

Chen D, Toone WM, Mata J, Lyne R, Burns G, Kivinen K, Brazma A, Jones N. 2003. Global
transcriptional responses of fission yeast to environmental stress. *Mol Biol Cell* 14:214–229.

Chen X, Stillman JH. 2012. Multigenerational analysis of temperature and salinity variability
affects on metabolic rate , generation time , and acute thermal and salinity tolerance in
Daphnia pulex. *J Therm Biol* 37(3):185–194.

Chevin LM, Lande R, Mace GM. 2010. Adaptation, plasticity, and extinction in a changing
environment: Towards a predictive theory. *PLoS Biol* 8(4).
Crain CM, Kroeker K, Halpern BS. 2008. Interactive and cumulative effects of multiple human
stressors in marine systems. *Ecol Lett* 11(12):1304–1315.
Cui X, Affourtit J, Shockley KR, Woo Y, Churchill GA. 2006. Inheritance patterns of transcript
levels in F1 hybrid mice. *Genetics* 174(2):627–637.
Dayan DI, Crawford DL, Oleksiak MF. 2015. Phenotypic plasticity in gene expression contributes
to divergence of locally adapted populations of Fundulus heteroclitus. *Mol Ecol*
24(13):3345–3359.
DeBiasse MB, Kelly MW. 2016. Plastic and evolved responses to global change: What can we
learn from comparative transcriptomics? *J Hered* 107(1):71–81.
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. 29(1):15–21.
Ehrenreich IM, Pfennig DW. 2016. Genetic assimilation: A review of its potential proximate
causes and evolutionary consequences. *Ann Bot* 117(5):769–779.
Enders LS, Bickel RD, Brisson JA, Heng-moss TM, Siegfried BD, Zera AJ, Miller NJ. 2015. Abiotic
and Biotic Stressors Causing Equivalent Mortality Induce Highly Variable Transcriptional
Responses in the Soybean Aphid. 5(February):261–270.
Enzor LA, Place SP. 2014. Is warmer better? Decreased oxidative damage in notothenioid fish
after long-term acclimation to multiple stressors. *J Exp Biol* 217(18):3301–3310.
Etterson JR, Shaw RG. 2001. Constraint to adaptive evolution in response to global warming.
Science 294(5540):151–154.
Evans TG. 2015. Considerations for the use of transcriptomics in identifying the "genes that
matter" for environmental adaptation. *J Exp Biol* 218(12):1925–1935.
Fay JC, Wittkopp PJ. 2008. Evaluating the role of natural selection in the evolution of gene
regulation. *Heredity* 100(2):191–199.
Feder ME, Hoffman GE. 1999. Heat-shock proteins, molecular chaperones, and the stress
response: evolutionary and ecological physiology. *Annu Rev Physiol* 61:243–282.
Feder ME, Walser JC. 2005. The biological limitations of transcriptomics in elucidating stress and
stress responses. *J Evol Biol* 18(4):901–910.
Fedorka KM, Zuk M, Mousseau TA. 2004. Immune suppression and the cost of reproduction in
the ground cricket, *Allonemobius socius*. *Evolution* 58(11):2478–2485.
Fitzpatrick BM. 2012. Underappreciated consequences of phenotypic plasticity for ecological
speciation. *Int J Ecol* 2012:32–37.
Flatt T, Tu MP, Tatar M. 2005. Hormonal pleiotropy and the juvenile hormone regulation
ofDrosophila development and life history. 27(10):999–1010.
Folt C, Chen C. 1999. Synergism and antagonism among multiple stressors. *Limnol Oceanogr*
44(February):864–877.
Fong SS, Joyce AR, Palsson BØ. 2005. Parallel adaptive evolution cultures of Escherichia coli lead
to convergent growth phenotypes with different gene expression states. *Genome Res*
15(10):1365–1372.
Fraser HB. 2013. Gene expression drives local adaptation in humans Gene expression drives local
adaptation in humans. :1089–1096.

Frumkin I, Schirman D, Rotman A, Li F, Zahavi L, Mordret E, Asraf O, Wu S, Levy SF, Pilpel Y. 2017.
Gene Architectures that Minimize Cost of Gene Expression. *Mol Cell* 65(1):142–153.

Gasch AP, Spellman PT, Kao CM, Carmel-Harel O, Eisen MB, Storz G, Botstein D, Brown PO. 2000.
Genomic expression programs in the response of yeast cells to environmental changes. *Mol*
Biol Cell 11(12):4241–4257.

Ghalambor CK, Hoke KL, Ruell EW, Fischer EK, Reznick DN, Hughes KA. 2015. Non-adaptive
plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature*
525(7569):372–375.

Ghalambor CK, McKay JK, Carroll SP, Reznick DN. 2007. Adaptive versus non-adaptive phenotypic
plasticity and the potential for contemporary adaptation in new environments. *Funct Ecol*
21(3):394–407.

Giaever EA. 2002. Functional profiling of the Saccharomyces cerevisiae genome. *Nature*
418(6896):387–91.

Gibson G. 2008. The environmental contribution to gene expression profiles. *Nat Rev Genet*
9(8):575.

Gobler CJ, DePasquale EL, Griffith AW, Baumann H. 2014. Hypoxia and acidification have additive
and synergistic negative effects on the growth, survival, and metamorphosis of early life
stage bivalves. *PLoS One* 9(1).

Gruntenko NE, Rauschenbach IY. 2008. Interplay of JH, 20E and biogenic amines under normal
and stress conditions and its effect on reproduction. *J Insect Physiol* 54(6):902–908.

Gunderson AR, Armstrong EJ, Stillman JH. 2016. Multiple Stressors in a Changing World: The
Need for an Improved Perspective on Physiological Responses to the Dynamic Marine
Environment. *Ann Rev Mar Sci* 8(1):357–378.

Harshman LG, Zera AJ. 2007. The cost of reproduction: the devil in the details. *Trends Ecol Evol*
22(2):80–86.

Hatakeyama M, Opitz L, Russo G, Qi W, Schlapbach R, Rehrauer H. 2016. SUSHI: An exquisite
recipe for fully documented, reproducible and reusable NGS data analysis. *BMC*
Bioinformatics 17(1):1–9.

Hauser F, Cazzamali G, Williamson M, Park Y, Li B, Tanaka Y, Predel R, Neupert S, Schachtner J,
Verleyen P, et al. 2008. A genome-wide inventory of neurohormone GPCRs in the red flour
beetle Tribolium castaneum. *Front Neuroendocrinol* 29(1):142–165.

Heckel K Von, Stephan W, Hutter S. 2016. Canalization of gene expression is a major signature of
regulatory cold adaptation in temperate Drosophila melanogaster. *BMC Genomics*:1–14.

Hereford J, Hansen T, Houle D. 2004. Comparing strenghts of directional selection: how strong is
strong? *Evolution* 58(10):2133–2143.

Ho WC, Zhang J. 2018. Evolutionary adaptations to new environments generally reverse plastic
phenotypic changes. *Nat Commun* 9(1):1–11.

Holmstrup M, Bindesbøl A mette, Janneke G, Duschl A, Scheil V, Köhler H r, Loureiro S, Soares
AMVM, Ferreira ALG, Kienle C, et al. 2010. Science of the Total Environment Interactions
between effects of environmental chemicals and natural stressors : A review. *Sci Total*
Environ 408(18):3746–3762.

Huang Y, Agrawal AF. 2016. Experimental Evolution of Gene Expression and Plasticity in
Alternative Selective Regimes. *PLoS Genet* 12(9):1–23.

Huth TJ, Place SP. 2016. Marine Genomics RNA-seq reveals a diminished acclimation response to
the combined effects of ocean acidi fi cation and elevated seawater temperature in
Pagothenia borchgrevinki. *Mar Genomics* 28:87–97.

Hutter S, Saminadin-Peter SS, Stephan W, Parsch J. 2008. Gene expression variation in African
and European populations of Drosophila melanogaster. *Genome Biol* 9(1):R12.

Kassahn KS, Crozier RH, Pörtner HO, Caley MJ. 2009. Animal performance and stress: Responses
and tolerance limits at different levels of biological organisation. *Biol Rev* 84(2):277–292.

Kelly MW, Debiasse MB, Villela VA, Hope L, Cecola CF. 2016. Adaptation to climate change :
trade-offs among responses to multiple stressors in an intertidal crustacean. :1147–1155.

Keren L, Hausser J, Lotan-Pompan M, Vainberg Slutskin I, Alisar H, Kaminski S, Weinberger A,
Alon U, Milo R, Segal E. 2016. Massively Parallel Interrogation of the Effects of Gene
Expression Levels on Fitness. *Cell* 166(5):1282–1294.e18.

King B, Denholm B. 2014. Malpighian tubule development in the red flour beetle (Tribolium
castaneum). *Arthropod Struct Dev* 43(6):605–613.

Kingsolver JG, Hoekstra HE, Hoekstra JM, Berrigan D, Vignieri SN, Hill CE, Hoang A, Gibert P,
Beerli P. 2001. The Strength of Phenotypic Selection in Natural Populations. *Am Nat*
157(3):245–261.

Kültz D. 2005. Molecular and Evolutionary Basis of the Cellular Stress Response. *Annu Rev Physiol*
67(1):225–257.

Lande R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic plasticity
and genetic assimilation. *J Evol Biol* 22(7):1435–1446.

Lande R, Arnold SJ. 1983. The Measurement of Selection on Correlated Characters. *Evolution*
37(6):1210.

Landry CR, Lemos B, Rifkin SA, Dickinson WJ, Hartl DL. 2007. Genetic properties influencing the
evolvability of gene expression. *Science* 317(5834):118–121.

Lang GI, Murray AW, Botstein D. 2009. The cost of gene expression underlies a fitness trade-off
in yeast. *Proc Natl Acad Sci* 106(14):5755–5760.

Leder EH, McCairns RJS, Leinonen T, Cano JM, Viitaniemi HM, Nikinmaa M, Primmer CR, Merilï¿½
J. 2015. The evolution and adaptive potential of transcriptional variation in sticklebacks -
Signatures of selection and widespread heritability. *Mol Biol Evol* 32(3):674–689.

Levine MT, Eckert ML, Begun DJ. 2011. Whole-genome expression plasticity across tropical and
temperate Drosophila melanogaster populations from eastern Australia. *Mol Biol Evol*
28(1):249–256.

Levis NA, Serrato-Capuchina A, Pfennig DW. 2017. Genetic accommodation in the wild: evolution
of gene expression plasticity during character displacement. *J Evol Biol* 30(9):1712–1723.

Li B, Predel R, Neupert S, Hauser F, Tanaka Y, Cazzamali G, Williamson M, Arakane Y, Verleyen P,
Schoofs L. 2008. Genomics, transcriptomics, and peptidomics of neuropeptides and protein
hormones in the red flour beetle Tribolium castaneum. *Genome Res* 18(1):113–122.

Liao Y, Smyth GK, Shi W. 2014. FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. 30(7):923–930.

Long Y, Li L, Li Q, He X, Cui Z. 2012. Transcriptomic characterization of temperature stress
responses in larval zebrafish. *PLoS One* 7(5).

López-Maury L, Marguerat S, Bähler J. 2008. Tuning gene expression to changing environments:

From rapid responses to evolutionary adaptation. *Nat Rev Genet* 9(8):583–593.

Mäkinen H, Papakostas S, Vøllestad LA, Leder EH, Primmer CR. 2016. Plastic and evolutionary
gene expression responses are correlated in European grayling (Thymallus thymallus)
subpopulations adapted to different thermal environments. *J. Hered.* 82–89.

Des Marais DL, Hernandez KM, Juenger TE. 2013. Genotype-by-Environment Interaction and
Plasticity: Exploring Genomic Responses of Plants to the Abiotic Environment. *Annu Rev*
Ecol Evol Syst 44(1):5–29.

McCairns RJS, Bernatchez L. 2009. Adaptive divergence between freshwater and marine
sticklebacks: insights into the role of phenotypic plasticity from an integrated analysis of
candidate gene expression. *Evolution* 64(4):1029–1047.

McCairns RJS, Smith S, Sasaki M, Bernatchez L, Beheregaray LB. 2016. The adaptive potential of
subtropical rainbowfish in the face of climate change: Heritability and heritable plasticity
for the expression of candidate genes. *Evol Appl* 9(4):531–545.

Milutinović B, Stolpe C, Peuß R, Armitage SAO, Kurtz J. 2013. The Red Flour Beetle as a Model for
Bacterial Oral Infections. *PLoS One* 8(5).

Morris MRJ, Richard R, Leder EH, Barrett RDH, Aubin-Horth N, Rogers SM. 2014. Gene expression
plasticity evolves in response to colonization of freshwater lakes in threespine stickleback.
Mol Ecol 23(13):3226–3240.

Moya A, Ganot P, Furla P, Sabourault C. 2012. The transcriptomic response to thermal stress is
immediate, transient and potentiated by ultraviolet radiation in the sea anemone
Anemonia viridis. *Mol Ecol* 21(5):1158–1174.

Moya A, Huisman L, Forêt S, Gattuso JP, Hayward DC, Ball EE, Miller DJ. 2015. Rapid acclimation
of juvenile corals to CO2-mediated acidification by upregulation of heat shock protein and
Bcl-2 genes. *Mol Ecol* 24(2):438–452.

Neven LG. 2000. Physiological responses of insects to heat. *Postharvest Biol Technol* 21(1):103–
111.

Nguyen TTA, Michaud D, Cloutier C. 2009. A proteomic analysis of the aphid Macrosiphum
euphorbiae under heat and radiation stress. *Insect Biochem Mol Biol* 39(1):20–30.

Oppert B, Dowd SE, Bouffard P, Li L, Conesa A, Lorenzen MD, Toutges M, Marshall J, Huestis DL,
Fabrick J, et al. 2012. Transcriptome profiling of the intoxication response of Tenebrio
molitor larvae to Bacillus thuringiensis Cry3Aa protoxin. *PLoS One* 7(4).

Park Y, Aikins J, Wang LJ, Beeman RW, Oppert B, Lord JC, Brown SJ, Lorenzen MD, Richards S,
Weinstock GM, et al. 2008. Analysis of transcriptome data in the red flour beetle, Tribolium
castaneum. *Insect Biochem Molec Biol* 38(4):380–386.

Parthasarathy R, Palli SR. 2011. Molecular analysis of nutritional and hormonal regulation of
female reproduction in the red flour beetle, Tribolium castaneum. *Insect Biochem Mol Biol*
41(5):294–305.

Parthasarathy R, Sheng Z, Sun Z, Palli SR. 2010. Ecdysteroid regulation of ovarian growth and
oocyte maturation in the red flour beetle, Tribolium castaneum. *Insect Biochem Mol Biol*
40(6):429–439.

Parthasarathy R, Sun Z, Bai H, Palli SR. 2010. Juvenile hormone regulation of vitellogenin
synthesis in the red flour beetle, Tribolium castaneum. *Insect Biochem Mol Biol* 40(5):405–
414.

Passow CN, Henpita C, Shaw JH, Quackenbush CR, Warren WC, Schartl M, Arias-Rodriguez L, Kelley JL, Tobler M. 2017. The roles of plasticity and evolutionary change in shaping gene expression variation in natural populations of extremophile fish. *Mol Ecol* 26(22):6384–6399.

Pörtner HO, Bennett AF, Bozinovic F, Clarke A, Lardies MA, Lucassen M, Pelster B, Schiemer F, Stillman JH. 2006. Trade-Offs in Thermal Adaptation: The Need for a Molecular to Ecological Integration. *Physiol Biochem Zool* 79(2):295–313.

Price TD, Qvarnström A, Irwin DE. 2003. The role of phenotypic plasticity in driving genetic evolution. *Proc R Soc B Biol Sci* 270(1523):1433–1440.

Prud 'homme B, Gompel N, Carroll SB. 2007. Emerging principles of regulatory evolution. *Proc Natl Acad Sci* 104 (suppl:8605–8612.

R Core Team. 2015. *R: A Language and Environment for Statistical Computing*

Ragland GJ, Almskaar K, Vertacnik KL, Gough HM, Feder JL, Hahn DA, Schwarz D. 2015. Differences in performance and transcriptome-wide gene expression associated with Rhagoletis (Diptera: Tephritidae) larvae feeding in alternate host fruit environments. *Mol Ecol* 24(11):2759–2776.

Rasmussen S, Barah P, Suarez-Rodriguez MC, Bressendorff S, Friis P, Costantino P, Bones AM, Nielsen HB, Mundy J. 2013. Transcriptome Responses to Combinations of Stresses in Arabidopsis. *PLANT Physiol* 161(4):1783–1794.

Reimand J, Arak T, Vilo J. 2011. g:Profiler—a web server for functional interpretation of gene lists (2011 update). *Nucleic Acids Res* 39(suppl_2):W307–W315.

Riehle MM, Bennett AF, Lenski RE, Long AD. 2003. Evolutionary changes in heat-inducible gene expression in lines of *Escherichia coli* adapted to high temperature. *Physiol Genomics* 14(1):47–58.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. 26(1):139–140.

Rodŕiguez-Verdugo A, Tenaillon O, Gaut BS. 2016. First-Step mutations during adaptation restore the expression of hundreds of genes. *Mol Biol Evol* 33(1):25–39.

Romero IG, Ruvinsky I, Gilad Y. 2012. Comparative studies of gene expression and the evolution of gene regulation. *Nat Rev Genet* 13(7):505–516.

Salmon AB, Marx DB, Harshman LG. 2001. a Cost of Reproduction in Drosophila Melanogaster : Stress Susceptibility a Cost of Reproduction in Drosophila Melanogaster : 55(8):1600–1608.

Sato MP, Makino T, Kawata M. 2016. Natural selection in a population of Drosophila melanogaster explained by changes in gene expression caused by sequence variation in core promoter regions. *BMC Evol Biol* 16(1):1–12.

Schlichting CD, Smith H. 2002. Phenotypic plasticity: linking molecular mechanisms with\revolutionary outcomes. *Evol Ecol* 16:189–211.

Schwenke RA, Lazzaro BP, Wolfner MF. 2016. Reproduction–Immunity Trade-Offs in Insects. *Annu Rev Entomol* 61(1):239–256.

Silbermann R, Tatar M. 2000. Reproductive costs of heat shock protein in transgenic *Drosophila melanogaster*. *Evolution* 54(6):2038–2045.

Skelly DA, Ronald J, Akey JM. 2009. Inherited Variation in Gene Expression. *Annu Rev Genomics Hum Genet* 10(1):313–332.

Sokoloff A. 1972. The biology of Tribolium with special emphasis on genetic aspects I. Oxford.
Oxford University Press.

Sørensen JG, Dahlgaard J, Loeschcke V. 2001. Genetic variation in thermal tolerance among
natural populations of Drosophila buzzatii: down regulation of Hsp70 expression and
variation in heat stress resistance traits. *Funct Ecol* 15(3):289–296.

Swindell WR, Huebner M, Weber AP. 2007. Plastic and adaptive gene expression patterns
associated with temperature stress in Arabidopsis thaliana. *Heredity* 99(2):143–150.

Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth A,
Santos A, Tsafou KP, et al. 2015. STRING v10: protein–protein interaction networks,
integrated over the tree of life. *Nucleic Acids Res* 43(D1):D447–D452.

Telonis-Scott M, Hallas R, McKechnie SW, Wee CW, Hoffmann AA. 2009. Selection for cold
resistance alters gene transcript levels in Drosophila melanogaster. *J Insect Physiol*
55(6):549–555.

Tirosh I, Barkai N, Verstrepen KJ. 2009. Promoter architecture and the evolvability of gene
expression. *J Biol* 8(11):1–6.

Townsend JP, Cavalieri D, Hartl DL. 2003. Population genetic variation in genome-wide gene
expression. *Mol Biol Evol* 20(6):955–963.

Wagner A. 2005. Energy constraints on the evolution of gene expression. *Mol Biol Evol*
22(6):1365–1374.

Whitehead A, Crawford DL. 2006. Neutral and adaptive variation in gene expression. *Proc Natl*
Acad Sci United States Am 103(14):5425–5430.

Xu J, Tan A, Palli SR. 2010. The function of nuclear receptors in regulation of female reproduction
and embryogenesis in the red flour beetle, Tribolium castaneum. *J Insect Physiol*
56(10):1471–1480.

Yampolsky LY, Glazko G V., Fry JD. 2012. Evolution of gene expression and expression plasticity in
long-term experimental populations of Drosophila melanogaster maintained under
constant and variable ethanol stress. *Mol Ecol* 21(17):4287–4299.

Yeh PJ, Price TD. 2004. Adaptive Phenotypic Plasticity and the Successful Colonization of a Novel
Environment. *Am Nat* 164(4):531–542.

Zheng W, Gianoulis TA, Karczewski KJ, Zhao H, Snyder M. 2011. Regulatory Variation Within and
Between Species. *Annu Rev Genomics Hum Genet* 12(1):327–346.

Zou Z, Evans JD, Lu Z, Zhao P, Williams M, Sumathipala N, Hetru C, Hultmark D, Jiang H. 2007.
Comparative genomic analysis of the Tribolium immune system. *Genome Biol* 8(8):1–16.