The relative influence of niche versus neutral processes on Ediacaran communities

*Emily G .Mitchell[1], Simon Harris[2], Charlotte G. Kenchington[1,] Philip Vixseboxse[3], Lucy

Roberts[4], Catherine Clark[1], Alexandra Dennis[1], Alexander G. Liu[1], Philip R. Wilby[2].

[1]Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge CB2

3EQ, UK.

[2]British Geological Survey, Nicker Hill, Keyworth, Nottingham NG12 5GG, United

Kingdom.

[3]School of Earth Sciences, University of Bristol, Wills Memorial Building, Queens Road,

Bristol, BS8 1RJ, United Kingdom.

[4]Department of Zoology, University of Cambridge…

*Correspondence: ek338@cam.ac.uk.

Keywords: Ediacaran, neutral theory, spatial point process analysis, paleoecology, ecology,

paleontology, rangeomorph

Abstract

A fundamental question in community ecology is the relative influence of niche versus

neutral processes in determining ecosystem dynamics. The extent to which these processes

structured early animal communities is yet to be explored. Here we use spatial point process

analyses (SPPA) to determine the influence of niche versus neutral processes on early total-

group metazoan paleocommunities from the Ediacaran Period ~565 million years in age.

Preservation of these sessile organisms in large in-situ populations on exposed bedding

planes enables inference of the most likely underlying processes governing their spatial

distributions by SPPA. We conducted comprehensive spatial mapping of six of the largest

Ediacaran paleocommunities in Newfoundland, Canada and Charnwood Forest, UK using

LiDAR, photogrammetry and a laser-line probe. For each paleocommunity we determined

the best-fit spatial model for each univariate and bivariate species distribution, comparing

four sets of spatial models (complete spatial randomness, dispersal, habitat, and combined

dispersal with habitat) using goodness-of-fit tests. Random and dispersal models are

considered neutral processes while habitat and combined models are considered niche

processes. We find the dynamics of these paleocommunities to be dominated by neutral

processes, with limited influence from niche processes. Our findings are consistent with

community model predictions of when neutral dynamics dominate, but are in stark contrast to

the niche-dominated communities of the modern marine realm. Thus, while the underlying

processes determining metazoan community assembly appear to have been in place since the

appearance of the first macroscopic, complex animals, the dynamics of these early metazoan

communities were fundamentally different to those of extant communities.

Significance statement

The extent to which habitat and dispersal processes structure the earliest animal communities found during the Ediacaran Period ~565 million years ago, is unknown. In this study we analyse six of the largest and most diverse fossil assemblages from Newfoundland, Canada and Charnwood Forest, UK, using spatial point processes analyses to determine the relative influence of habitat processes (niche) and dispersal processes (neutral) on the spatial distribution of taxa. The vast majority of Ediacaran taxon distributions were controlled by neutral processes, in striking contrast to niche-dominated modern marine ecosystems, but consistent with model predictions of when neutral dynamics dominate communities. Thus the underlying processes determining metazoan community assembly have been in place since the appearance of the first macroscopic, complex animals.

Two opposing theories lie at the centre of debate regarding the fundamental dynamics that

govern ecosystem structure and biodiversity: niche and neutral. Niche theory is a central

tenet of classical ecological theory, whereby species avoid competitive exclusion by

occupying different niches within the ecosystem (MacArthur 1984). The smaller the niche

overlap, the less competition occurs between taxa, allowing more taxa to exist in an area

without driving each other to extinction. Species are able to co-exist because they are

different. Niche models describe selection-dominated ecosystems, whereby species

dynamics operate deterministically as a series of inter-specific interactions, which act as

stabilizing mechanisms for the ecosystem (Adler et al. 2007).

Neutral processes are often referred to as the null model of niche processes: instead of species

differences enabling co-existence, their *similarity* drives high diversity (Hubbell 2001).

Within neutral models, species fitness is constant, and so different taxa can co-exist because

none has a significant competitive advantage over the other. Despite this seemingly highly

unrealistic assumption, neutral theories have been able to accurately reproduce certain

species-area-distributions (SADs; Hubbell 2001); sometimes better than niche theories;

(MacArthur 1984), as well as SAD and beta diversity patterns (Condit et al. 2002, Chisholm

et al. 2010).

Unified or continuous theories, whereby niche and neutral processes combine to generate

species coexistence (Grave et al. 2004, Adler et al. 2007), have emerged in recent years. In

these combined models, species can exhibit strong differences and strong stabilizations

(niche-type), or weak stabilizations because of similar fitness (neutral-type), with classic

niche and neutral models the extreme endmembers of this continuum model. However, it is

often not possible to analytically determine whether the niche or neutral model fits the data

better, making it hard to untangle the relative influence of niche and neutral-type processes

within modern complex ecosystems.

In order to investigate whether niche or neutral processes were the most important during the

first establishment of modern-style ecosystems, we focus on some of the oldest known total-

group metazoan communities: those comprising the Ediacaran macrobiota, dated to ~571-560

million years in age (Droser et al., 2017). The evolution of macroscopic metazoans was

coupled with a transformation in ecosystem dynamics. Paleocommunities evolved from the

assumed simple community structure of pre-Ediacaran microbial populations (Butterfield

2007), through the Ediacaran paleocommunities that exhibited both simple and complex

community structures (Darroch et al. 2018), and on into the Cambrian 'modern' metazoan

ecosystems with a similar ecosystem structure to the present (Dunne et al. 2008).

The oldest metazoan-dominated communities form part of the Avalonian Assemblage of the

Ediacara Biota (Waggoner 2003), and are known primarily from Newfoundland, Canada and

Leicestershire, UK. Avalonian soft-bodied organisms were preserved in-situ in deep-water

strata dated to ~572-560 Ma (Pu et al. 2016, Noble et al. 2016), beneath volcanic ash-rich

event beds (Wood et al. 2003, Narbonne 2005). As such, exposed bedding-plane surfaces

preserve near-complete censuses of the communities (Wood et al. 2003, Clapham et al.

2003); though the impact of erosion of these surfaces needs to be taken into account

(Matthews et al. 2017), cf. Mitchell et al. 2015). Since they were soft bodied, dead organisms

could not accumulate over long time periods, removing problems associated with time-

averaging of the paleocommunity. Furthermore, Avalonian ecosystems pre-date macro-

predation and vertical burrowing, such that upon death, organisms were not eaten and

remained in place.   As a result, and because the organisms are considered to have been non-

mobile, each individual bedding plane is interpreted to record organisms that lived

contemporaneously, with recently deceased and decaying organisms being the primary

recognised record of identifiable time-averaging (Liu et al. 2011; Mitchell and Butterfield

2018; see also Wilby et al. 2015).

Consequently, the position and size of each fossil specimen can be interpreted to capture the

life history of the organism (i.e. the dispersal, habitat and community interactions it was

subject to in life). As a result, population studies using spatial point process analyses (SPPA)

can infer the most likely underlying ecological and biological processes recorded within the

spatial distributions of the fossils (Illian et al. 2008). For sessile organisms, community-scale

spatial distributions depend on the interplay of a limited number of different factors: physical

environment, which manifests as habitat associations of a taxon or taxon-pairs (Wiegand et

al. 2007), organism dispersal/reproduction (Seidler and Plotkin 2006), competition for

resources (Getzin et al. 2006), facilitation between taxa (Lingua et al. 2008), and differential

mortality (Getzin et al. 2008).

To assess the relative influence of niche and neutral processes for these sessile communities,

niche processes are identified as intra- or inter-specific habitat associations, and/or intra- and

inter-specific competition (Lin et al. 2011). Density-dependent competition typically

generates a segregated spatial distribution which, when broken down into taxon population

size-classes, exhibits segregated largest specimens, and either random or aggregated

distributions of small specimens (Diggle 2013). Such segregation is intra-specific for

univariate distributions, and inter-specific for bivariate distributions, between two taxa.

Neutral processes are identified where univariate distributions exhibit complete spatial

randomness (CSR), and by dispersal processes that are independent of local environment (i.e.

habitat heterogeneities; Gunatilleke et al. 2006; Comita et al. 2007; Wiegand et al. 2007a; Lin

et al. 2011). Intra-specific habitat associations are best-modelled as a heterogeneous Poisson

model (HP), or when combined with dispersal-limitations, ITC (Harms et al. 2001; Lin et al.

2011) and inter-specific habitat associations are best modelled by a shared parents models

(SPM). Dispersal patterns are indicated by a best-fit model of a Thomas Cluster (TC) or

Double Thomas Cluster (DTC) model (Lin et al. 2011). Where dispersal processes are

coupled to a habitat association, this process is best modelled by a Thomas cluster model

combined with an inhomogeneous Poisson model (ITC; Lin et al. 2011). Therefore, for

univariate distributions, neutral processes are indicated by CSR and HP, and niche processes

by segregation and HP and ITC models (Fig. 2). CSR is considered a neutral process because

there are no biologically or ecologically significant intrinsic or extrinsic influences on the

spatial distribution. TC and DTC aggregations are also considered neutral since they describe

dispersal processes, whereby aggregations arise from propagules only traveling a limited

distance (thus being unable to reach all suitable substrates regardless of underlying habitat

heterogeneities or species requirements; Hubbell et al. 1999; Harms et al. 2001; Seidler and

Plotkin 2006). Species associations with habitat heterogeneities lead to spatial aggregations

(or segregations) corresponding to the underlying habitat variations on which the species

depend. Density-dependent competition, as indicated by size-dependent spatial segregation

(Kenkel 1988), indicates a lack of sufficient resources, and is therefore also a niche-based

process.

Figure 1: Locality Map of study sites, showing: A, the relative location of sites within the

micro-continent of Avalonia. B, the Newfoundland sites of the Bristy Cove (BR5), 'D' and

'E' surface, Mistaken Point Ecological Reserve, the St Shott's (Sword Point) surface, and the

H14/Johnson surface, Bonavista Peninsula (modified from Liu 2016). Associated spatial

maps for each locality show the positions of the fossil specimens (indicated by a circle).

Black scale bar = 1m, grey scale bar = 0.1m. Different colors indicate different taxa as

follows: Thectardis *navy;* Fractofusus *light blue;* Charnia *bright yellow;* Charniodiscus *dark*

red; Aspidella *light green;* Bradgatia *dark green ; Feather Dusters light orange;*

Primocandelabrum *dark orange;* Trepassia *dark purple;* Beothukis *bright pink;* Pectinifrons

dark blue; Brushes brown; Avalofractus *navy;* Hylaecullulus *light yellow.*

In this study we assessed the univariate and bivariate spatial distributions of taxa from six

Avalonian communities: the 'D', 'E' and Bristy Cove surfaces in the Mistaken Point

Ecological Reserve; the St Shott's surface at Sword Point; the H14 (Johnson) surface at

Little Catalina in Newfoundland, Canada; and Bed B, Charnwood Forest, UK.   We mapped

the bedding-plane assemblages of fossils at a coarse scale using LiDAR, and then captured

the fine-scale fossil details using photogrammetry and high-resolution laser scanning (SI

Methods). Combination of these datasets enabled full maps of the paleocommunities to be

reconstructed (Fig 1).

For each surface, we tested for erosional biases and tectonic deformation, taking these factors

into account if they were considered to have significantly affected specimen density

distributions (SI Methods). Non-abundant taxa (< 30 specimens) and taphomorphs (such as

organ taxa or the decaying and/or poorly preserved remains of carcasses, ivesheadiomorphs)

were excluded from analyses, leaving 10 abundant taxa (see SI Methods for full taxonomic

descriptions), of which three (*Charniodiscus*, *Charnia*, *Bradgatia*) are found on two bedding

planes and one (*Fractofusus*) on four bedding planes. The univariate spatial distributions of

each taxon on individual bedding planes were described using pair correlation functions

(PCFs). A PCF = 1 indicates a distribution that was completely spatially random (CSR); PCF

> 1 indicates aggregation; and PCF < 1 indicates segregation (Diggle 2003, 2015; Illian et al.

2008). Monte Carlo simulations and Diggle's goodness-of-fit tests (p_d) were used to indicate

significantly non-CSR distributions where the observed PCF deviated outside the simulation

envelope and p_d << 1 (Diggle 2013). Where a non-CSR distribution was observed, HP, TC

and ITC models were fit to the data, with the highest p_d value indicating the best-fit model

(Fig. 2). Identifying the processes behind spatial patterns is not straightforward (Levin 1992;

Murrell and Law 2003; Wiegand et al. 2007; McIntier and Fajardo 2009; Wiegand and

Moloney 2013), but the best-fit model is interpreted to indicate the most likely underlying

processes, with CSR and TC reflecting neutral processes and HP and ITC reflecting niche

processes (Lin et al. 2011). Bivariate distributions were assessed to ascertain inter-specific

interactions and associations between taxa (cf. Mitchell and Butterfield 2018). Niche

processes are bivariate habitat associations indicated by shared parents models (SP) and

density-dependent competition. Density-dependent competition has the spatial signature of a

segregated distribution, which, when broken down into PCFs of size-classes, has segregated

largest specimens and either CSR or aggregated small specimens (Diggle 2013).

Figure 2. Univariate PCF for 'D' surface Fractofusus *and 'E' surface* Beothukis *under four*

different spatial models: CSR, HP, DTC, ITC. The model lines are dashed, the solid lines are

the observed spatial distributions and the grey area represents the simulation envelope of

999 Monte Carlo simulations.

Results and Discussion

Figure 3 Univariate PCF for surfaces A) 'E' surface, B) 'D' surface, C) Bristy Cove, D) St.

Shott's, E) H14/Johnson and F) Bed B. Where a niche model is the best-fit model for the

distribution, it is drawn as a dashed line. Neutral models are drawn as solid lines. Different

205 *colors indicate different taxa as follows:* Thectardis *navy;* Fractofusus *light blue;* Charnia

206 *bright yellow;* Charniodiscus *dark red;* Aspidella *light green;* Bradgatia *dark green; 'Feather*

207 *Duster', light orange;* Primocandalebrum *dark orange;* Trepassia *dark purple;* Beothukis

208 *bright pink;* Pectinifrons *dark blue; Brushes brown;* Avalofractus *navy;* Hylaecullulus *light*

209 *yellow.*

	CSR	HP	TC/DTC	ITC
BED B	0.00%	0.00%	100.00%	0.00%
H14/JOHNSON	0.00%	0.00%	100.00%	0.00%
ST SHOTTS	50.00%	0.00%	50.00%	0.00%
BRISTY COVE	100.00%	0.00%	0.00%	0.00%
D SURFACE	66.67%	0.00%	33.33%	0.00%
E SURFACE	16.67%	16.67%	66.67%	0.00%

Table 1. Proportion of best-fit univariate models by surface, showing the percentage of taxa

with univariate spatial distributions that are best described by CSR, HP, TC (or DTC) and

ITC models.

Across the six surfaces and the 16 taxon univariate distributions examined, 5 taxon

distributions were best modelled by CSR (Fig. 3, Table 1. SI Table 2). Of the non-CSR taxa

distributions 10 were best modelled by TC (or DTC). No taxa were best modelled by ITC

models (Table 1). Only *Beothukis* on the 'E' surface had a HP best-fit model univariate

spatial distribution (Fig. 2, Table 1). None of the 16 univariate taxon distributions exhibited

intra-specific spatial segregation, meaning there is no evidence of intra-specific competition.

Our results reveal only one univariate taxon distribution ('E' surface *Beothukis*) that was

best-modelled by a niche process.

The studied paleocommunities only have spatial scales in the order of decameters. However,

by comparing the univariate spatial distributions of the same taxa found at different sites

representing communities separated by large spatial and temporal scales, we can gain some

indication of whether taxa behave differently over large-spatial scales. Four taxa are abundant

across multiple bedding planes (*Bradgatia*, *Charnia*, *Charniodiscus* and *Fractofusus*), and

these taxa all exhibit the same type of best-fit model (CSR, TC, TC and TC/DTC) on all the

surfaces**.** Previous work has demonstrated how *Fractofusus* shows the same consistent

spatial distributions across multiple surfaces (Mitchell et al. 2015) in different geological

formations (potentially indicating different environments and ages; cf. H14 and 'E' surface).

Bradgatia, *Charnia* and *Charniodiscus* also show consistent spatial distributions (Fig. 3),

even when the taxa inhabit ecosystems originally located in different basins (e.g.

Charniodiscus from the 'E' surface and Bed B) and separated by ~6 million years (Noble et

al. 2015, Pu et al. 2016). The consistency of these results suggests that the small-spatial scale

ecological behaviour of these taxa did not change over large spatial and temporal scales.

Figure 4: Multi-surface univariate PCFs of A) Fractofusus *(from Mitchell et al. 2015). B)*

Non-CSR bivariate distributions of the 'E' surface (from Mitchell and Butterfield 2018) and

C) the non-CSR bivariate distributions Bed B, Charnwood Forest and). The x-axis is the

inter-point distance between organisms in metres. On the y-axis, PCF=1 indicates complete

spatial randomness (CSR), <1 indicates segregation, and >1 indicates aggregation.

Of the four communities with more than one abundant (> 30 specimens) taxon present, two

surfaces exhibited only CSR bivariate distributions ('D' surface (Mitchell and Butterfield

2018) and St Shotts). The 'E' surface (Mitchell and Butterfield 2018) and Bed B exhibit non-

CSR bivariate distributions (Fig 3). The three non-CSR bivariate distributions on Bed B

indicate shared habitat associations (Fig 4c, SI Table 1) as do the three bivariate distributions

on the 'E' surface (Mitchell and Butterfield 2018, Mitchell and Kenchington 2018). Habitat

associations are niche processes (Lin et al. 2011), thus these results demonstrate that both

niche and neutral processes operated within these communities. These bivariate associations

are much weaker in PCF magnitude than the univariate distributions (Figs 3 and 4), showing

that niche processes had less impact on spatial distributions than neutral processes.

Furthermore, competition is rare, and is also relatively weak in magnitude when observed

(Fig 4b,c, Mitchell and Butterfield 2018, Mitchell and Kenchington 2018). These results

demonstrate that environmental associations had a relatively small impact on the community

structure of these Ediacaran communities. On the 'E' surface, segregations reduced specimen

density by 25%, and aggregations increased taxon density by 56% (Fig. 4). In contrast, intra-

specific aggregations are large, reflecting an increase in taxon density of 250–600%. The two

habitat associations on the 'E' surface (Feather Dusters–*Fractofusus* and Feather Dusters–

Charniodiscus) exhibited small scale specimen aggregations (increase of 34% under 0.2m

and 56% under 1.2m respectively) with large scale reduction in specimen density (11% over

1m and 13% over 2.1m respectively). Similarly on Bed B, habitat association between

Charnia and *Primocandelabum* increased taxon density by 87%, whereas segregations

reduced taxon density by 10%. Univariate dispersal-generated aggregations increased taxon

density by 180–500% (Figs 3 and 4). These results demonstrate that habitat associations were

present in some paleocommunties, but had a relatively small impact on organism

distributions.

Our results support combined theories of community assembly whereby niche and neutral

theories are not mutually exclusive; they act along a continuum or spectrum, with differing

extents of niche and neutral processes present in different circumstances (Gravel et al. 2006;

Fisher and Mehta 2014). The dominance of univariate niche best-fit models, repetition of

best-fit univariate models across different communities, and the rarity and weakness of

bivariate niche best-fit models, all combine to provide strong evidence that neutral processes

dominated Avalonian communities, with only limited niche-based influence. These neutral-

dominated community dynamics contrast with those observed in the modern marine realm,

where neutral models do not provide good descriptions of the community dynamics

(Dornelas et al. 2006, Connolly et al. 2014).

The striking difference in dominance of niche versus neutral processes raises the intriguing

question as to whether the community dynamics of these oldest known macroscopic animal

communities had fundamentally different dynamics to the present.   Previous work on the

balance of niche–neutral influences on community assemblage in Quaternary fossil

assemblages provides strong model and empirical support for environment-led models of

assembly (Jackson and Blois 2015). However, Avalonian communities appear to differ from

the majority of existing marine systems in that the maximum number of generations currently

documented for these Avalonian paleocommunities is three generations (Mitchell et al.

2015), though some communities show rare survivors (Wilby et al. 2015) or evidence of

secondary community succession (Liu et al., 2012). These population structures suggest that

the preserved paleocommunities are not always mature, being culled prematurely by the

frequent event beds that killed and preserved them, limiting their maturity (Wilby et al.

2015). Frequent sedimentary disturbances could also explain why there is limited intra-

specific competition, with the lack of intra-specific competition demonstrating that

populations did not reach their carrying capacities. Recent models show that community

dynamics in small populations in fluctuating environments are dominated by neutral

processes (Fisher and Mehta 2014).

While some Avalonian taxa were global, with wide dispersal ranges (indicated here by CSR),

the majority of the specimens within each community consisted of taxa with limited dispersal

ranges on the order of decimetres (Mitchell et al. 2015. Mitchell and Kenchington 2018).

Such limited dispersal range have been shown to decrease ecological selection and decrease

effective community size, while increasing the dominance of neutral-processes (Ron et al.

2018). Therefore, while the dominance of neutral-based processes within these

paleocommunities differs significantly to the modern marine realm, the underlying dynamics

can still be described by established models of community dynamics. Therefore, we propose

that the oldest known macroscopic metazoan paleocommunities operated under different

processes to the present, but that underlying dynamics of community assembly have existed

unchanged over the last ~570 million years.

References

Adler, P. B., HilleRisLambers, J., & Levine, J. M. (2007). A niche for neutrality. Ecology

letters, 10(2), 95-104.

Boag, T. H., Darroch, S. A., & Laflamme, M. (2016). Ediacaran distributions in space and

time: testing assemblage concepts of earliest macroscopic body fossils. Paleobiology, 42(4),

574-594.

Baddeley, A., E. Rubak, and R. Turner. (2015). Spatial point patterns: methodology and

applications with R. CRC Press.

Butterfield, N. J. (2007). Macroevolution and macroecology through deep time.

Paleontology, 50(1), 41-55.

Chave, J. (2004). Neutral theory and community ecology. Ecology letters, 7(3), 241-253.

Chisholm, R. A., & Pacala, S. W. (2010). Niche and neutral models predict asymptotically

equivalent species abundance distributions in high-diversity ecological communities.

Proceedings of the National Academy of Sciences, 201009387.

Clapham, M. E., G. M. Narbonne, and J. G. Gehling. (2003). Paleoecology of the oldest

known animal communities: Ediacaran assemblages at Mistaken Point, Newfoundland.

Paleobiology 29:527–544.Comita et al. 2007;

Condit, R., Pitman, N., Leigh, E. G., Chave, J., Terborgh, J., Foster, R. B., ... & Muller-

Landau, H. C. (2002). Beta-diversity in tropical forest trees. Science, 295(5555), 666-669.

Connolly, S. R., MacNeil, M. A., Caley, M. J., Knowlton, N., Cripps, E., Hisano, M., ... &

Brandt, A. (2014). Commonness and rarity in the marine biosphere. Proceedings of the

National Academy of Sciences, 201406664.

Darroch, S. A., Laflamme, M., & Wagner, P. J. (2018). High ecological complexity in

benthic Ediacaran communities. Nature ecology & evolution, 2(10), 1541.

Diggle, P. Statistical Analysis of Spatial Point Patterns 2nd edn (Arnold, 2003).

Diggle, P., Zheng, P. and Durr, P., (2005). Nonparametric estimation of spatial segregation in

a multivariate point process: bovine tuberculosis in Cornwall, UK. Journal of the Royal

Statistical Society: Series C (Applied Statistics), 54(3), pp.645-658.

Dornelas, M., Connolly, S. R., & Hughes, T. P. (2006). Coral reef diversity refutes the neutral

theory of biodiversity. Nature, 440(7080), 80.

Droser, M. L., Tarhan, L. G., & Gehling, J. G. (2017). The rise of animals in a changing

environment: global ecological innovation in the late Ediacaran. Annual Review of Earth

and Planetary Sciences, 45, 593-617.

Dunn, F.S., Liu, A.G. & Donoghue, P.C. (2018) Ediacaran developmental biology. Biological

Reviews 93, 914-932 .

Dunne, J. A., Williams, R. J., Martinez, N. D., Wood, R. A., & Erwin, D. H. (2008).

Compilation and network analyses of Cambrian food webs. PLoS biology, 6(4), e102.

Fisher, C. K., & Mehta, P. (2014). The transition between the niche and neutral regimes in

ecology. Proceedings of the National Academy of Sciences, 111(36), 13111-13116.

Getzin, S., Ch. Dean, F. He, J. A Trofymow, K. Wiegand, and T. Wiegand. (2006). Spatial

patterns and competition of tree species in a Douglasfir chronosequence on Vancouver

Island. Ecography 29:671–682.

Getzin, S., T. W., K. Wiegand, and F. He. (2008). Heterogeneity influences spatial patterns

and demographics in forest stands. Journal of Ecology 96:807–820.

Gravel, D., Canham, C. D., Beaudet, M., & Messier, C. (2006). Reconciling niche and

neutrality: the continuum hypothesis. Ecology letters, 9(4), 399-409.

Gunatilleke. (2009). Ecological information from spatial patterns of plants: insights from

point process theory. Journal of Ecology 97:616–628.harms et al. 2001

Hubbell, S. P. (2001). The unified neutral theory of biodiversity and biogeography (MPB-32)

(monographs in population biology).

Hubbell, S. P., Foster, R. B., O'Brien, S. T., Harms, K. E., Condit, R., Wechsler, B., & De

Lao, S. L. (1999). Light-gap disturbances, recruitment limitation, and tree diversity in a

neotropical forest. Science, 283(5401), 554-557.

Illian, J., A. Penttinen, H. Stoyan, and D. Stoyan. (2008). Statistical analysis and modelling of

spatial point patterns. Vol. 70. John Wiley & Sons.

Jackson, S. T., & Blois, J. L. (2015). Community ecology in a changing environment:

Perspectives from the Quaternary. Proceedings of the National Academy of Sciences,

201403664.

Kenkel, N. C. (1988). Pattern of self thinning in jack pine: testing the random mortality

hypothesis. Ecology, 69(4), 1017-1024.

Levin, S. A. in Ecological Time Series Vol.2 (edsPowell, T. M.&Steele, J. H.) 277–326

(Springer, 1995).

Levin, S. A. (1992). The problem of pattern and scale in ecology: the Robert H. MacArthur

award lecture. Ecology, 73(6), 1943-1967.

Lin, Y. C., Chang, L. W., Yang, K. C., Wang, H. H., & Sun, I. F. (2011). Point patterns of

tree distribution determined by habitat heterogeneity and dispersal limitation. Oecologia,

165(1), 175-184.

Lingua, E., Cherubini, P., Motta, R., & Nola, P. (2008). Spatial structure along an altitudinal

gradient in the Italian central Alps suggests competition and facilitation among coniferous

species. Journal of Vegetation Science, 19(3), 425-436.

Liu, A. G., D. Mcilroy, J. B. Antcliffe, and M. D. Brasier. (2011). Effaced preservation in the

Ediacara biota and its implications for the early macrofossil record. Paleontology 54:607–

630.

Liu, A. G., D McIlroy, J. J. Matthews, and M. D. Brasier. (2012). A new assemblage of

juvenile Ediacaran fronds from the Drook Formation, Newfoundland. Journal of the

Geological Society 169:395–403.

Liu, A. G., C. G. Kenchington, and E. G. Mitchell. (2015). Remarkable insights into the

paleoecology of the Avalonian Ediacaran macrobiota. Gondwana Research 27:1355–

1380.MacArthur 1979

MacArthur, R. H. (1984). Geographical ecology: patterns in the distribution of species.

Princeton University Press.

Matthews, J. J., A. G. Liu, and D. McIlroy. (2017). Post–fossilization processes and their

implications for understanding Ediacaran macrofossil assemblages. Geological Society,

London, Special Publications 448:251–269.

McIntire, E. J., & Fajardo, A. (2009). Beyond description: the active and effective way to

infer processes from spatial patterns. Ecology, 90(1), 46-56.

Murrell, D.J. and Law, R., (2003). Heteromyopia and the spatial coexistence of similar

competitors. Ecology letters, 6(1), pp.48-59.

Mitchell, E. G., & Kenchington, C. G. (2018). The utility of height for the Ediacaran

organisms of Mistaken Point. Nature Ecology and Evolution, 2, 1218-1222.

Mitchell, E. G., Kenchington, C. G., Liu, A. G., Matthews, J. J., & Butterfield, N. J. (2015).

Reconstructing the reproductive mode of an Ediacaran macro-organism. Nature, 524(7565),

343.

Mitchell, E. G & Butterfield, N. J. (2018) Spatial analyses of Ediacaran communities at

Mistaken Point. Paleobiology, doi.org/10.1017/pab.2017.35.

Murrell, D. J., & Law, R. (2003). Heteromyopia and the spatial coexistence of similar

competitors. Ecology letters, 6(1), 48-59.

Narbonne, G. M. (2005). The Ediacara biota: Neoproterozoic origin of animals and their

ecosystems. Annu. Rev. Earth Planet. Sci., 33, 421-442.

Noble, S. R., Condon, D. J., Carney, J. N., Wilby, P. R., Pharaoh, T. C., & Ford, T. D.

(2015). U-Pb geochronology and global context of the Charnian Supergroup, UK:

constraints on the age of key Ediacaran fossil assemblages. Bulletin, 127(1-2), 250-265.

Pu, J.P., Bowring, S.A., Ramezani, J., Myrow, P., Raub, T.D., Landing, E., Mills, A.,

Hodgin, E. and Macdonald, F.A., (2016). Dodging snowballs: Geochronology of the

Gaskiers glaciation and the first appearance of the Ediacaran biota. Geology, 44(11),

pp.955-958.

Ron, R., Fragman-Sapir, O., & Kadmon, R. (2018). Dispersal increases ecological selection

by increasing effective community size. Proceedings of the National Academy of Sciences,

115(44), 11280-11285.

Seidler, T. G., and J. B. Plotkin. (2006). Seed dispersal and spatial pattern in tropical trees.

PLoS Biology 4:e344.

Waggoner, B. (2003). The Ediacaran biotas in space and time.Integrative and Comparative

Biology 43:104–113.

Wan, B., Yuan, X., Chen, Z., Guan, C., Pang, K., Tang, Q. and Xiao, S. (2016) Systematic

description of putative animal fossils from the early Ediacaran Lantian Formation of South

China. Paleontology 59(4), 1-18 .

Wiegand, T. & Moloney, K. A. Handbook of Spatial Point-Pattern Analysis in Ecology (CRC

Press, Boca Raton, 2013).

Wiegand, T., S. Gunatilleke, and N. Gunatilleke. (2007A). Species associations in a

heterogeneous Sri Lankan dipterocarp forest. The American Naturalist 170:E77–E95.

Wiegand, T., S. Gunatilleke, N. Gunatilleke, and T. Okuda. (2007B). Analyzing the spatial

structure of a Sri Lankan tree species with multiple scales of clustering. Ecology 88:3088–

3102

Wilby, P. R., J. N. Carney, and M. P.A. Howe. (2011). A rich Ediacaran assemblage from

eastern Avalonia: evidence of early widespread diversity in the deep ocean. Geology

39:655–658.

Wilby, P. R., Kenchington, C. G., & Wilby, R. L. (2015). Role of low intensity

environmental disturbance in structuring the earliest (Ediacaran) macrobenthic tiered

communities. Paleogeography, Paleoclimatology, Paleoecology, 434, 14-27.

Wood, D. A., Dalrymple, R. W., Narbonne, G. M., Gehling, J. G. & Clapham, M. E.

(2003).Paleoenvironmental analysis of the late Neoproterozoic Mistaken Point and

Trepassey formations, southeastern Newfoundland. Can. J. Earth Sci. 40, 1375–1391.

Yuan, X., Chen, Z., Xiao, S., Zhou, C. & Hua, H. (2011) An early Ediacaran assemblage of

macroscopic and morphologically differentiated eukaryotes. Nature 470 (7334), 390–393.

23