An application of higher order multivariate cumulants in modelling of myoelectrical

activity of porcine uterus during early pregnancy.

Malgorzata Domino[a], Krzysztof Domino[b], Zdzislaw Gajewski[a,*]

[a] Department of Large Animal Diseases with Clinic, Veterinary Research Centre and Center

for Biomedical Research, Faculty of Veterinary Medicine, Warsaw University of Life

Sciences (WULS – SGGW), Warsaw, Poland

[b] Institute of Theoretical and Applied Informatics, Polish Academy of Sciences, Gliwice,

Poland

[*]Corresponding authors E-mail address: zgajewski@supermedia.pl (Z. Gajewski)

ABSTRACT

The analysis of the uterine contraction have become a general practice in an effort to improve the clinical management of uterine contractions during pregnancy and labour in human beings. The fluctuations in uterine activity may occur without affecting progress of gestation, however the painful and fashion contractions may be the first threat of misscarage. While pigs were considered as an referential preclinical model, the computational modelling of spontaneous myoelectrical activity of complex systems of porcine myometrium in peri-fertilization period has been proposed. The higher order statistic, multivariate cumulants and Joint Skewness Band Selection method, have been applied to study the dependence structure of electromyographic (EMG) signal with an effective EMG feature. Than the model of recognition of multivariate, myoelectricaly changes according to crucial stages for successful fertilization and early pregnancy maintenance has been estimated. We found that considering together time and frequency features of EMG signal was extremely non-Gaussian distributed and the higher order multivariate statistics such as cumulants, have to be used to determine the pattern of myoelectrical activity in reproductive tract. We confirmed the expectance that the probabilistic model changes on a daily base. We demonstrated the changes in proposed model at the crucial time points of in peri-fertilization period. We speculate the activity of the middle of uterine horn and the power (minimum and maximum) and pauses between myoelectrical burst features are essential for the functional role of uterine contractility in peri-fertilization period.

Keywords:

EMG activity, higher order cumulant tensors, features selection, myometrium, early pregnancy, sow

1. Introduction

The analysis of the uterine contraction have become a general practice in an effort to improve the clinical management of uterine contractions during pregnancy and labour (Figueroa et al., 1987; Gajewski and Faundez, 1992; Maul et al., 2003; Rabotti et al., 2008; Jacod et al., 2010; Lammers, 2013; Rooijakkers et al., 2014; Pawlinski et al., 2017). In healthy, no pregnant uteri specific contractile patterns evolve during the estrus cycle both in humans and animals (Fanchin and Ayoubi, 2009; Kuijsters et al., 2017; Domino et. al, 2018a). Significant attention has been dedicated to evolving patterns being in line with the hypothesis of a functional role of uterine activity in promotion of fertilization (Pusey et al., 1980; Sammali et al., 2018). Uterine contractility is known to affect embryo implantation (Rogers et al., 1983; Ziecik et al., 2011; Lammers, 2013). Furthermore, the quiescence of the uterus during pregnancy is required to maintain pregnancy and allow adequate nourishment and development of the foetus (Gajewski and Faundez, 1992; Rabboti and Mischi, 2015; Markiewicz et al., 2016). During early pregnancy, some fluctuations in uterine activity can always be present without affecting progress of gestation, however the painful uterine contractions occurring in a coordinated and forceful fashion may be the first threat of a preterm delivery (Norwitz and Robinson, 2001).

The animal and *in vitro* experiments have been found to provide valuable results and methods for parameter estimation, undertaking up to humans clinical studies (Devedeux et al., 1993; Rabotti et al., 2010). While pigs were considered as an ideal preclinical model due to many anatomical and functional similarities with humans, the investigations carried out on the porcine reproductive tract may be treated as a referential (Ziecik et al., 1993; Gajewski et al, 2001; Kobayashi et al., 2012). The electrical activity of the porcine uterus as the primary cause of contractions, was established both during estrus cycle (Gajewski et al., 2001; Domino et al., 2018a) and early pregnancy (Maner et al., 2003; Markiewicz et al., 2016; Pawlinski et al., 2017). In those cases smooth muscle cells (SMCs) contract, action potentials reach a depolarization threshold and generate an electromagnetic field (Eswaran et al., 2004). The billions of SMCs are coupled electrically by the gap junctions into the complex biological system (Garfield et al., 1977) within numerous electrochemical events generate electric currents as a sum of action potential differences between SMCs (Alkan and Günay, 2012). Those myometrial electromagnetic field is possible to be measured as voltage and quantified with different sensitivity by electromyography (EMG) (Gajewski et al., 2001; Wolinski et al., 2003; Eswaran et al., 2004) and electrohysterography (EHG) (Rabotti et al., 2008; Jacod et al., 2010; Rabotti et al., 2010; Sammali et al., 2018). The EHG was reported as a feasible

option to evaluate noninvasively and objectively the myometrial activity in humans, both of

the non-pregnant uterus (Sammali et al., 2018) as well as during late gestation and labour

(Jacod et al., 2010; Rabotti et al., 2010). In early gestation in humans, EHG method is

insufficient to register the subtle changes in spectral content of uterine EMG signal, whereas

invasive EMG method is unacceptable for obvious, ethical reasons (Rabotti et al., 2010). The

needle EMG gathered the bioelectrical signal directly associated with the contractile activity

of myometrium, while EHG collected signal abundant in noise generated between uterus and

implanted superficial electrodes positioned on the surface of abdomen (Devedeux et al.,1993;

Jacod et al., 2010). Therefore, the electrical activity recorded more invasively, directly from

myometrium, may provide crucial information about uterine activity. For this reason the

animal models are essential for understanding the sophisticated mechanisms behind the

phenomena of contribution of uterine activity into the fertilization and early pregnancy

maintenance.

In the case of all electroencephalograms (EEG) (Singh et al., 2015; Chen et al., 2016),

EMG (Domino et al., 2016; 2017a; 2018a) and EHG (Farina and Negro, 2007; Garfield and

Maner, 2007a; Domino et al., 2017a), considering spectral content in addition to studying

changes in its time domain features is widely used. Moreover a variety of features have been

extracted by different feature selection methods in the time and frequency domains. The time

domain features of EMG include duration (D), amplitude (A) and root mean square (RMS) of

bursts as well as duration of pauses between bursts (P) (Devedeux et al.,1993; Gajewski et al.,

2004; Garfield et al., 2007a; Pawlinski et al., 2016). It has been established that the uterine

EMG signals can be quantified sufficiently with mathematical functions and transforms such

as power spectral analysis (Maner et al., 2003; Garfield and Maner, 2007b). In frequency

domain we receive features corresponding to each frequency bands. However in our work for

simplicity and informative significance we concentrate on: dominant frequency (DF),

maximum power (Max P) and the minimum power (Min P) which can be estimated by Fast

Fourier transform (FFT) method (Oczeretko et al., 2007; Domino et al., 2016; 2017a; 2018a).

Recent methods used in a detailed analysis of myoelectrical signals are divided in four major

categories: cross-coherence function (Farina and Merletti, 2004; Domino et al., 2016), phase

difference (Hunter et al., 1987), maximum likelihood (Farina et al., 2001; Rabotti et al., 2010)

and the detection of spectral relations (Farina and Negro, 2007; Domino et al., 2017; Garfield

and Maner 2007b). Garfield and Maner (2007b) suggested the features of power density

spectrum as a more effective than the conventional time dependent features in reflecting the

actual function of the uterus. The decomposition of myoelectrical signals into their individual

frequency or 'scale' subcomponents provided much more advantageous data, that can be used as indicators of changes in uterine status such as pre-term delivery (Maner et al., 2003; Garfield and Maner, 2007b). In our research we have acquired both recent features (D, A, RMS, P) and following frequency domain subcomponents (DF, Max P, Min P) of EMG signal, from porcine uterus in relation to early pregnancy phenomena. Further we speculate that those features are non-Gaussian distributed due to complex biochemical processes that take place in myometrium, which represents contractile element of the uterine wall composing of SMCs (Rabboti and Mischi, 2015). Observe that higher order multivariate cumulants can be used to analyse non-Gaussian distributed multivariate data (Domino et al., 2018b) such as considering together time and frequency features in our case. Observe as well, that higher order cumulants approach was successfully applied in EEG (Becker et al., 2014) and superficial EMG (Ju and Liu, 2011) data analysis. However, none of recent researchers has introduced the computational modelling of spontaneous myoelectrical activity of complex SMCs systems in peri-fertilization period. An additional novelty of this paper resides in the porcine EMG signal analyzing methodology based on the use of higher order multivariate cumulants. We have applied those cumulants to study the dependence structure of EMG patterns with an effective EMG feature, and then have built up the myoelectrical activity templates for recognizing complex uterine contraction according to crucial stages for successful fertilization and early pregnancy maintenance.

2. Materials and methods

2.1. Experimental design

The spontaneous myoelectrical activity of uterus has been recorded from 8 Polish Landrace sows according to protocol approved by the III Local Ethical Committee on Animal Testing in Warsaw (Permit Number: 71/2009, from 19.11.2009) on behalf of the National Ethical Committees on Animal Testing. The implantable telemetry method has been applied to obtain the EMG signal from subsequent parts of corpus uteri and unilateral uterine horn. The experiment has been started on non-pregnant sows during diestrus, subsequently the induction and synchronization of estrus was performed to manage an artificial insemination (AI) according to Pawlinski et al. (2017) (day -4: eCG, 750 IU/animal, i.m., Werfaser, Fa; Alvetra, Austria; day -1: hCG, 500 IU/animal, i.m., Werfaser, Fa; Alvetra, Austria; day 0: AI). Indeed, spontaneous uterine activity was registered during long-term registration both in non-pregnant and pregnant state since day 12 after AI.

The surgery under general anesthesia has been performed in order to position the combination of three silver, bipolar, needle electrodes connected to 3-channel telemetry transmitter TL10M3-D70-EEE (DSI, USA) according to Gajewski et al. (2004). Animals were premedicated with an azaperone (Stresnil, 3.0 mg/kg b.w., i.m., Fa; Janssen Pharmaceutica, Belgium). General anesthesia was conducted using combined administration of medetomidine (Cepetor, 1.0 mg/kg b.w., i.v., Fa; CP-Pharma HgmbH, Germany), butorphanol (Butomidor, 0.2 mg/kg b.w., i.v., Fa; Richter Pharma AG, Austria), ketamine (Bioketan, 3.0 mg/kg b.w., i.v., Fa: Vetoquinol Biowet, Poland) and propofol (Propofol, 2.0-4.0 mg/kg b.wt., i.v., Fa; Pfizer, USA). After surgery meloxicam (Metacam, 0.4 mg/kg b.w., i.m., Fa; Boehringer Ingelheim, Germany) and cefquinom (Cobactan, 2.0 mg/kg b.w., i.m., Fa. Intervet, Poland) had been administered for five days. The electrodes were sutured unilaterally onto the surfaces of uterus: the tip of right uterine horn, (channel 1), the middle of right uterine horn (channel 2) and corpus uteri (channel 3). Channel 1 was positioned at a distance of 2 cm from oviduct, whereas distance between the other two channels was fixed at 30 cm. The 3-channel telemetry transmitter was positioned between abdominal muscles directly under the skin (Fig. 1A). Finally, the silicone cannula was inserted into the brachial vein in order to daily blood sample collection.

2.1.1 Time-line determination

The experimental time-line was determined based on the hormonal and ovarian status as well as the time of AI (day 0) (Fig. 2). The daily blood samples were used for hormones concentrations: luteinizing hormones (LH) and progesterone (P_4), determination. Blood samples (7.0 ml every 4 h) collected into vacutainer tubes (Fa; KABE, Poland) were centrifuged (5.590 g; 10 min) and the serum samples were frozen until assaying. The hormones concentrations was determined using radioimmunoassay: a high sensitivity, non-commercial for LH (intra-assay coefficient of variation <6.7% and inter-assay coefficient of variation <12.5%, Ziecik et al., 1978) and a commercial for P_4 (intra-assay coefficient of variation <5.6% and inter-assay coefficient of variation <8.8%; KIP 1458; Fa; DIAsource ImmunoAssays SA, Belgium).

Day -6 and -5 correspond to the diestrus stage (hormone concentration: LH <1.0 ng/ml; P4 >1.0 ng/ml), whereas days from -4 to 1 show changes in the proestrus (day -4 to -2; LH <1.0 ng/ml; P4 <1.0 ng/ml) and estrus (day -1 to 1; LH >2.0 ng/ml; P4 <1.0 ng/ml) stage of the estrus cycle, respectively. During day -1 the concentration of LH increased (LH >4.0 ng/ml; P4 <1.0 ng/ml) concurrently and the AI were conducted twice (day 0 and 1). Finally,

the P4 concentration gradually increased (LH <1.0 ng/ml; P4 >1.0 ng/ml) from day 2 and

remained at a high level (LH <1.0 ng/ml; P4 >4.0 ng/ml) since day 12 after AI. At day 12

sows were euthanized (sodium pentobarbital; Morbital 100.0 or more mg/kg b.w., i.v., Fa;

Biowet Pulawy, Poland) and the pregnancy has been confirmed after flushing the reproductive

tract in order to collect embryos according to Pawlinski et al. (2017). Good quality, viable

embryos have been collected from each animal. In both uterine horns 12 - 19 embryos in size

6 - 8 mm were observed under a dissecting microscope (Fa; Olympus SZ51, Japan).

2.2. EMG data acquisition

From day 4. after surgery the EMG signal was measured continuously before and

during estrus synchronization as well as during AI and early pregnancy. The EMG recording

was continued for 12 days after first AI. The EMG signal was acquired with a 3-channel

transmitter (TL10M3-D70-EEE; DSI, USA) with sampling frequency - 100 Hz. Obtained

analogue signal was digitalized and sent by radio waves to the telemetric receiver (DL10

analogue output; Fa; DSI, USA). The band-pass digital filter (5-50 Hz) and the notch filter

were applied for lowering the power line interference from the EMG raw data (PowerLab 4e;

Fa; ADInstruments, Australia). Mean and linear trends were removed and the EMG signal

was analyzed using LabChart 8 Reader (Fa; ADInstruments, Australia) software. The

electrical activity of myometrium reflecting uterine contraction was defined as burst when

amplitude exceeded 5 μV for longer than 3 s, the individual bursts were separated with non

active periods from longer than 0.5 s, according to Gajewski et al. (2004). In the time domain

bursts parameters were determined using duration (D, s), amplitude (A, mV) and root mean

square (RMS, mV) of bursts as well as duration of pauses between bursts (P, s). In the

frequency domain EMG signal spectral content was calculated after signal multiplication by

Hamming window as in Muthuswamy and Thakor (1998) and Domino et al. (2016). By

means of the FFT: the dominant frequency (DF, Hz; the frequency of the maximum power

within the selected frequency range), maximum power (Max P, V^2 x10^{-11}; the maximum

power within the selected frequency range) and minimum power (Min P, V^2 x10^{-24}; the

minimum power within the selected frequency range) was assessed for each burst (Fig. 1B).

In recent papers the action potentials measured *in vivo* were determined mainly by one

statistical outcome variable at a time (A, RMS, D or their univariate statistic) (Gajewski et al.,

2004; Garfield and Manner, 2007a; Domino et al., 2016; Pawlinski et al., 2016). Since

Garfield and Maner (2007b) and Govindan et al. (2015) suggested analysis of the uterine

electrical activity in frequency domain as a potential indicators of predicting uterine

contraction, the methods taking into account the effects of all features on the recent responses
of interest has become necessary. Therefore the following probabilistic model was introduced
including both timer domain and frequency domain features and their mutual
interdependency.

2.3. EMG data processing

Obtained data are represented the form of matrix $X \in \mathbb{R}^{t \times n}$ with elements $x_{j,i}$ where j
index correspond to realizations and i index corresponds to features being univariate
marginals. In our approach we took under consideration 7 features (Duration; Pause;
Amplitude; RMS; MaxP; MinP; DF), which order is consistent with indexing. The single
realisation vector was $x_j = \left[x_{j,1}, ..., x_{n,j} \right] \in \mathbb{R}^7$.

While introducing the probabilistic model, supposed data are modelled by the
multivariate distribution with the following multivariate probability density function PDF
$f(x): \mathbb{R}^n \to \mathbb{R}$. We assumed that probabilistic model does not change during a given day, but
may change between days. The characteristic function (Kendall and Stuart, 1946; Lukacs,
1970) of $f(x)$ was given by:

$$\tilde{\phi}: \mathbb{R}^n \to \mathbb{C} \qquad \tilde{\phi}(\tau) = \int_{\mathbb{R}^n} \exp(i\tau^T x) f(x) dx \quad (1)$$

where τ is an argument vector $\tau = \left[\tau_1, ..., \tau_n \right]$ and i is an imaginary unit. The d^{th} multivariate
cumulant is an object (a tensor) indexed by $i_1, ... i_d$, where each $i_k \in \left[1, 2, ..., n \right]$, with following
elements:

$$c_{i_1, ..., i_d} = (-i)^d \frac{\partial^d}{\partial \tau_{i_1}, ..., \partial \tau_{i_d}} \log\left(\tilde{\phi}(\tau) \right) \Big|_{\tau = [0, ..., 0]} \quad (2).$$

It was easy to show that for the probabilistic model of the multivariate Gaussian distribution
$\log\left(\tilde{\phi}(\tau) \right)$ is quadratic in τ, hence for $d \geq 3$ Equation (2) gives zeros. In consistency with a
standard notation for $d \geq 3$ we have higher order cumulants. Observe that for $d = 1$ we have a
mean vector and for $d = 2$ a covariance matrix, we call them the standard cumulants.
Following Domino and Gawron (2018c) one can use the Frobenius norm on the higher
order multivariate cumulant to measure how far data are from hypothetical multivariate
Gaussian model. Importantly multivariate Gaussian distribution model does not allow for
simultaneous extreme values of many features (Domino et al., 2018b), hence the appearance

of such extreme simultaneous events should be reflected in higher order cumulants values.

The Frobenius norm of the d^{th} cumulant tensor can be defined as follows:

237
$$\|C_d\| = \sqrt{\sum_{i_1=1}^{n}...\sum_{i_d=1}^{n}\left(c_{i_1,...,i_d}\right)^2} \quad (3).$$

To calculate higher order cumulants we use the efficient algorithm introduced in Domino et

al. (2018b) and implemented in Julia programming language (Domino et al., 2017b).

2.3.1 Information extraction and hierarchy of features

For each day of the peri-fertilization period we have collected data in the following

matrix form: $X \in \mathbb{R}^{400x7}$. Due to the specifics of the signal collected, values of various features

differ by many orders of magnitude. Hence we are rather interested in information carried by

the whole features set. To overcome an impact of different orders of magnitudes of features

we normalise them by dividing each by its standard deviation. Normalized data $\hat{X}$ are defined

in element wise notation:

248
$$\hat{x}_{j,i} = \frac{x_{j,i}}{\sigma_i} \quad (4)$$

where σ_i is a standard deviation of the i^{th} column (i^{th} marginal) of X.

It would be interesting to determine the hierarchy of features including information

they are caring. For this purpose we use the Joint Skewness Band Selection (JSBS) method

that was primary introduced to analyse information stored in non-Gaussian features of

hyperspectral images (Geng et al., 2015). Moreover the JSBS was used for information

analysis concerning artificially generated data by means of copulas (Domino and Glos,

2018d). Recall that the application of the JSBS to analyse biomedical data is pioneer. Let us

move to the formal introduction of the JSBS.

To introduce formally the JSBS we need to refer to Equation (2) again. We can

introduce the 3^{rd} cumulant's tensor $C_3 \in \mathbb{R}^{n \times n \times n}$ being an 3 dimensional Array and containing

all cumulant's elements c_{i_1,i_2,i_3}, for the formal tensor definition see (Kolda and Bader, 2009).

Observe that:

261
$$c_{i_1,i_2,i_3} = c_{i_1,i_3,i_2} = c_{i_2,i_1,i_3} = c_{i_2,i_3,i_1} = c_{i_3,i_1,i_3} = c_{i_3,i_2,i_1} \quad (5)$$

262 since differentiation is commutative, this observation is called the super-symmetry of the

263 cumulant's tensor (Schatz et al., 2014). Following (Kolda and Bader, 2009) the first mode

unfolding of 3^{rd} cumulant's tensor, noted as $(C_3)_{(1)} \in \mathbb{R}^{n \times n^2}$ is the matrix composed of following elements:

$$c_{(1)_{i,j}} = c_{i_1, i_2, i_3} \, , \quad (6)$$

where its second index is:

$$j = i_2 + (i_3 - 1)n \, . \quad (7)$$

in our case. Reffer to Kolda and Bader (2009) for the general definition. Recall that, there exists other possible methods to unfold a tensor into a matrix. However due to the supper-symmetry of C_3, all those methods are equivalent Kolda and Bader (2009). This is an argument for taking a first mode unfold as its mathematical notation is simple. Having defined a tensor unfold into matrix, we can define the "joint skewness" matrix:

$$\mathbb{R}^{n \times n} \ni \mathbf{M}_3 = (C_3)_{(1)} ((C_3)_{(1)})^{\mathrm{T}} \, . \quad (8)$$

As argued in Geng et al. (2015) such matrix measures information extracted by the 3^{rd} cumulant about a non-Gaussian distribution of data. Finally to measure the relative information about the non-Gaussianity, we use the following target function:

$$f_{JSBS} = \frac{1}{2} \log(\det(\mathbf{M}_3)) - \frac{3}{2} \log(\det \Sigma) \, , \quad (9)$$

which is a logarithm of the target function introduced in Geng et al. (2015). The log values are better to present, and logarithm is strictly increasing function, hence it does not affect the information hierarchy. Since $\Sigma \in \mathbb{R}^{n \times n}$ is a covariance matrix, we may think about f_{JSBS} as the measure of information carried by the 3^{rd} cumulant, but not by a covariance matrix. Referring to Figure 3 the covariance matrix is little affected by the biological process, hence its information is redundant here. Importantly determinant is the product of eigenvalues of the matrix and as such it is proportional to the information hyper-ellipsoid volume (Sheffield, 1985) of the matrix.

Given a target function we perform the following features elimination scheme:

1. we compute the f_{JSBS} for all data $\hat{\mathrm{X}}$,

2. we remove one marginal of $\hat{\mathrm{X}} \in \mathbb{R}^{t \times n}$ in such a way that the target function of new $\hat{\mathrm{X}}' \in \mathbb{R}^{t \times (n-1)}$ is maximises, removed feature (marginal) is less informative,

3. we repeat $n-1$ times the procedure in point 2 to obtain the order of features by their information significance.

3. Results and discussion

The clear, free of artefacts EMG signal was recorded from all electrodes positioned along one side of uterus from day -6 to day 12 in respect to AI. We used the Frobenius norm on the higher order multivariate cumulant and found that considering together time and frequency features of EMG signal was extremely non-Gaussian distributed as following $\|C_3\| \gg 0$ and $\|C_4\| \gg 0$ in all points of designed time-line.

Beginning with the standard cumulants the Frobenious norm of the mean vector (for $d = 1$) and the covariance matrix (for $d = 2$) of $\hat{X}$ are presented in Figure 3. One can observe that there are no sharp changes as expected for those values in our experimental system. No significant changes are visible and repeatable in all three measurement sites (Ch1, Ch2, Ch3). Such sharp changes would be informative for our system analysis, since we expect the probabilistic model to change between days due to the change of the complex dynamics of EMG signal pattern. Hence such standard multivariate statistics are not informative here.

Following on higher order multivariate statistics the Froebenious norm of the 3^{rd} cumulant's (for $d = 3$) and the 3^{rd} cumulant's (for $d = 4$) of $\hat{X}$, measuring of the distance from the Gaussian multivariate distribution, are presented in Figure 4 and 5, respectively. There black dash lines gives the 0.975 quantile if particular date were from the Gaussian multivariate distribution. The examined features are high above black dash line indicating that the pattern of EMG activity is significantly far from hypothetical multivariate Gaussian model, in all measurement sites repetitively. Our results indicate than bioelectrical signal, such as uterine EMG, brings within much more information than has been recently achieved from univariate statistic (Jedruch et al., 1989a; Gajewski et al., 2004; Pawlinski et al., 2016; Pawlinski et al., 2017) or multivariate Gaussian distribution model (Fig. 3). Secondly there are high jumps (a value of the norm may double) yielding a change of the probabilistic model and suggesting transformations of EMG signal pattern.

To explain such behaviour we suggest the appearance of simultaneous extreme values of EMG signal features after estrus cycle synchronization beginning in corpus uteri (day -4, Fig. 4C, Fig. 5C) and spreading along uterine horn (days -3 to 0, Fig. 4B, Fig. 5B) into the tip of uterine horn (day -3, Fig. 4A, Fig. 5A). In the middle of uterine horn the most changes in myoelectrical activity pattern between days -3 and 0 have been observed. Garfield and Yallampalli (1994) suggested in uterine muscle signal parameters and its propagation depend on properties of wave propagation medium, which differed significantly witch changes of reproductive status (Jedruch et al., 1989a; Jedruch et al., 1989b; Andersen and Barclay, 1995;

Kitazawa et al., 2001; Mueller et al., 2006). Those differences in the properties of the medium

apply components of intracellular systems at the level of the SMCs plasma membrane both

the number of gap junction and expression of hormone (including reproductive hormones)

receptors (Garfield and Maner, 2007a). The changes in hormone (and their receptors)

concentration and ion channel distribution interact in dynamical manner to modulate the

membrane potential, the type of action potential, and the ionic conductance in the

myometrium (Wray et al., 2015). Therefore the changes in probabilistic model may also be

the most visible in estrus, when the endogenous regulatory impact is the highest. Our results

indicate the largest changes in EMG signal pattern during estrus in the uterine horn, which

coincides with the highest receptors for ovarian hormones (Kitazawa et al., 2001; Mueller et

al., 2006) and gap junction (Garfield et al., 1977; Andersen and Barclay, 1995) output in this

part of reproductive tract.

The second high jump occur shortly after AI again in the first in corpus uteri (day 1,

Fig. 4C, Fig. 5C) and then in the middle (day 2, Fig. 4B, Fig. 5B) and in the tip (day 3, Fig.

4B, Fig. 5B) of uterine horn. After AI due to sperm cells motion and myometrium

contractions, a small subpopulation of sperm are very quickly (within 1-2 hours) transported

from the uterus towards the utero-tubal junction where they colonize sperm reservoir (Rath et

al., 2008). The fate of the majority of sperm is elimination and the rapid removal of sperm is

thought to prevent acquired immune response against sperm (Hansen, 2011; Katila, 2012).

The myometral contractions takes active part in the elimination of sperm, which starts

simultaneously with semen transport. Frequent uterine contractions carry semen back and

forth in the uterus (Katila et al., 2000; Sumransap et al., 2007). Because the elimination of

sperm and post insemination inflammatory products is important for fertility, the efficient

myometrial contractions are a prerequisite for uterine cleaning and preparing the intrauterine

environment for the development of the embryo (Katila, 2012). The crucial changes in

probabilistic model may also be observed. The fate of the majority of sperm is elimination.

Sperm cells are antigenic to females, and immunization against sperm can result in infertility.

Rapid removal of sperm is thought to prevent acquired immune responses against sperm

(Hansen, 2011).

The third high jump was detected only in uterine horn, both in the tip (days 8 to 10,

Fig. 4C, Fig. 5C) and the middle (days 6 to 8, Fig. 4B, Fig. 5B) corresponding to the descent

of embryos into the uterine lumen. That intrauterine migration of porcine embryos occurs

between day 6. and 10(11). (day 0 = fertilization) (Ziecik et al., 2011). In this period the

changes in uterine activity during may be expected. Pope et al. (1982) showed that

myometrial contractility increased concomitantly with embryo migration. Similarly

Markiewicz et al. (2016) described an increase of contractile activity in the porcine uterus

with embryos. The changes of probabilistic model which we are demonstrated are in line with

the recent studies.

Interestingly, at days 11 and 12 the evidences of the largest order were observed in

corpus and horns of uterus at once (Figs. 4-5). At the day 11 after AI, the embryos starting

secretion of estrogen, which stimulates luteoprotective prostaglandin E_2 (PGE_2) synthesis in

the process of maternal recognition of pregnancy (Ziecik et al., 2011). In this process embryo

signals its presence in the uterus and mother receives and accepts this signal (Short, 1969).

When embryo signals are systemically recognized by mother, the corpus luteum function and

also pregnancy are maintained, therefore the changes in uterine activity pattern during this

period should be also expected. The complex system in which billions of cells comprising

myometrium, responds similarly and accordingly to biological signals transmitted through

uterine luminal epithelium and underlying smooth muscle layers (Dantzer, 1985; Franczak

and Bogacki, 2009). The interaction between embryos and uterine wall provides the

quiescence of the uterus requiring to maintain pregnancy.

Referring to Figure 4 one can conclude that the 3-order multivariate cumulant gives

optimal information about biological phenomena that we are investigate in. The large

estimation error of higher order cumulants, especially the 4^{th} order, should be associated with

the relatively small data samples $t = 400$. This disadvantage of our method may be improved

by increasing radically data sample size as one discussed in Appendix A (Domino et al.,

2018b). Nevertheless, in our case higher order multivariate cumulants are much more

informative that classical statistics. The following cumulants based tools find application to

analyse our data. The norm of the higher order cumulants tensors is used to analyse how far

the probabilistic model of data is from a Gaussian multivariate distribution. Moreover the 3^{rd}

cumulant's tensor based Joint Skewness Band Selection (JSBS) is used to order features

according to information they carry (Geng et al., 2015). The hierarchy of features has been

successfully determined using the pioneer application of the JSBS to analyse biomedical

features. An application of f_{JSBS} allowed to determinate the dependence structure of EMG

patterns with an effective EMG feature. Some of examples of target function values during

features elimination process are presented in Figure 6. Sometimes we have a maximum if all

features are included, and sometimes after removing one feature as in Table 1. The

explanation is not straight forward, since the f_{JSBS} is rather used to compare equally large

subsets of features. Recall that the JSBS is searching for a subset of features that has significantly non-Gaussian joint distribution, while highly non-Gaussian data are informative for our model. We have shown that Min P, Max P and P features carries highest information according to our model, while DF, D and RMS are little informative. Hence we have a combination of features both in time domain and frequency domain that are meaningful.

Our results obtained in animal *in vivo* model supporting the hypothesis of a functional role of uterine activity in promotion of fertilization both in estrus (days -4 to 0) and after AI (days 1), preparation of uterine environment to embrace the embryos (days 2 to 3) and support of embryos migration during early pregnancy (days 6 to 10). We suspect the middle of uterine horn as the most heavily involved area affecting the phenomena of embryo development and implantation and the Min P, Max P and P features as the most informative in the model of spontaneous myoelectrical activity of complex SMCs systems in peri-fertilization period. Due to the fact that we obtained good quality, viable embryos at day 12, the studied development of early pregnancy should be considered proper. The applied approach is suitable in recognition of crucial stages in peri-fertilization processes providing successful fertilization and early pregnancy maintenance.

5. Conclusions

In conclusion, due to non-Gaussian joint distribution of features such as, higher order multivariate statistics such as cumulants, have to be used to determine the pattern of myoelectrical activity in reproductive tract. Using those tools, we confirmed the expectance that the probabilistic model changes on a daily base, while analyzing the complex biological system of porcine SMCs and the functional role of uterine activity in peri-fertilization period. We speculate the middle of uterine horn and the Min P, Max P and P features are crucial elements of the uterine activity model, whereas further information is needed to elucidate the coordination and propagation of EMG signal. Concluding, the higher order multivariate cumulants modelling offers a unique opportunity to understand the mechanisms underlying uterine contractility and development of early pregnancy in animals. The authors belief that similar dynamics may appear in human beings due to high applicability of porcine preclinical models.

Conflict of interest

None.

Acknowledgements

This work was conducted in the Veterinary Research Centre WULS (WCB) and the Center

for Biomedical Research (CBB) supported by EFRR RPO WM 2007-2013.

References

Alkan, A. & Günay, M., Identification of EMG signals using discriminant analysis and SVM

classifier. *Expert Syst Appl*. 39: 44–47, DOI: 10.1016/j.eswa.2011.06.043 (2012).

Andersen H.F. & Barclay M., A computer model of uterine contractions based on discrete

contractile elements. *Obstet Gynecol*. 86(1): 108–11, DOI: 10.1016/0029-

7844(95)00111-4 (1995).

Becker, H., Albera, L., Comon, P., Haardt, M., Birot, G., Wendling, F., Gavaret, M., Bénar,

C.-G. & Merlet, I., EEG extended source localization: tensor-based vs. conventional

methods. *NeuroImage*. 96: 143–157, DOI: 10.1016/j.neuroimage.2014.03.043 (2014).

Chen, S.J., Peng, Ch.J., Chen, Y.Ch., Hwang, Y.R., Lai, Y.S., Fan, S.Z. & Jen, K.K.,

Comparison of FFT and marginal spectra of EEG using empirical mode decomposition

to monitor anesthesia. *Comput Methods Programs Biomed*. 137: 77–85, DOI:

10.1016/j.cmpb.2016.08.024 (2016).

Dantzer, V., Electron microscopy of the initial stages of placentation in the pig. *Anat

Embryol*. 172: 281–293 (1985).

Devedeux, D., Marque, C., Mansour, S., Germain, G. & Duchene, J., Uterine

electromyography: a critical review. *Am J Obstet Gynecol*. 169: 1636–1653, DOI:

10.1016/0002-9378(93)90456-S (1993).

Domino, K. & Gawron P., Sliding window high order cumulant tensors calculation algorithm.

arXiv preprint arXiv:1701.06446 (2018c).

Domino, K. & Glos, A., Hiding higher order cross-correlations of multivariate data using

archimedean copulas. arXiv preprint arXiv:1803.07813 (2018d).

Domino, K., Gawron P. & Pawela L., Cummulants.jl.,

https://github.com/kdomino/Cumulants.jl, DOI: 10.5281/zenodo.1254364 (2017b).

Domino, K., Gawron, P. & Pawela, L. Efficient Computation of Higher-Order Cumulant

Tensors. *SIAM J Sci Comput*. 40(4): A1590–A1610, DOI: 10.1137/17M1149365

(2018b).

Domino, M., Pawlinski, B. & Gajewski, Z., Biomathematical pattern of EMG signal

propagation in smooth muscle of the non-pregnant porcine uterus. *PLoS One*. 12(3):

e0173452, DOI: 10.1371/journal.pone.0173452 (2017a).

Domino, M., Pawlinski, B. & Gajewski, Z., The linear synchronization measures of uterine
EMG signals: Evidence of synchronized action potentials during propagation.
Theriogenology. 86(8): 1873–1878, DOI: 10.1016/j.theriogenology.2016.03.036 (2016).

Domino, M., Pawlinski, B., Gajewska, M., Jasinski, T., Sady, M. & Gajewski, Z. Uterine
EMG activity in the non-pregnant sow during estrous cycle. *BMC Vet Res.* 14(1): 176–
185, DOI: 10.1186/s12917-018-1495-z (2018a).

Eswaran, H., Preissl, H., Wilson, J.D., Murphy, P. & Lowery, C.L., Prediction of labor in
term and preterm pregnancies using non-invasive magnetomyographic recordings of
uterine contractions. *Am J Obstet Gynecology.* 190(6): 1598–1603, DOI:
10.1016/j.ajog.2004.03.063 (2004).

Fanchin, R. & Ayoubi, J.M., Uterine dynamics: impact on the human reproduction process.
Reprod Biomed Online. 18(suppl 2): 57–62, DOI: 10.1016/S1472-6483(10)60450-6
(2009).

Farina, D. & Merletti, R., Methods for estimating muscle fibre conduction velocity from
surface electromyographic signals. *Med Biol Eng Comput.* 42: 432–445 (2004).

Farina, D. & Negro, F., Estimation of muscle fiber conduction velocity with a spectral
multidip approach. *IEEE Trans Biom Eng.* 54: 1583–1589, DOI:
10.1109/TBME.2007.892928 (2007).

Farina, D., Muhammad, W., Fortunato, E., Meste, O., Merletti, R. & Rix, H., Estimation of
single motor unit conduction velocity from surface electromyogram signals detected
with linear electrode arrays. *Med Biol Eng Comput.* 39: 225–236 (2001).

Figueroa, J., Massmann, A., Pimentel, G. & Nathanielsz, P., Characteristics of the
electromyogram recorded from the mesometrium of the pregnant ewe from 106 days'
gestation to delivery: similarities with and differences from the electromyogram
obtained from the myometrium. *Am J Obstet Gynecol* 157: 991–998, DOI:
10.1016/S0002-9378(87)80102-3 (1987).

Franczak, A. & Bogacki, M., Local and systemic effects of embryos on uterine tissues during
early pregnancy in pigs. *J Reprod Dev.* 55(3): 262–72, DOI: 10.1262/jrd.20148 (2009).

Gajewski, Z. & Faundez, R. Characteristics and analysis of the uterus electrical activity in the
pregnant cows. *Theriogenology.* 37: 1133–1145, DOI: 10.1016/0093-691X(92)90110-D
(1992).

Gajewski, Z., Blitek, M., Klos, J., Gromadzka-Hliwa, K., Pawlinski, B., Andrzejczak, A. &
Ziecik, A., Oviductal and uterine myometrial activity during periovulatory period in the
pig. *Reprod. Dom. Anim.* 1: 41–47 (2004).

Gajewski, Z., Wolinski, J., Korczynski, W., Ziecik, A., Babelewska, M. & Zabielski, R.,
Application of telemetry to long-term electromyography recordings of the reproductive
tract in the pig. Newsletter 5th ESDAR, Viena. 1: 36–41 (2001).

Garfield, R.E. & Maner, W.L., Biophysical methods of prediction and prevention of preterm
labor: uterine electromyography and cervical light-induced fluorescence–new
obstetrical diagnostic techniques. In: Preterm Birth: Mechanisms, Mediators, Prediction,
Prevention, and Interventions. eds. Petraglia, F., Strauss, J.F., Gabbe, S.G., Weiss, G.,
pp.131–145, (2007b).

Garfield, R.E. & Maner, W.L., Physiology and electrical activity of uterine contractions.
Semin Cell Dev Biol 18: 289–295, DOI: 10.1016/j.semcdb.2007.05.004 (2007a).

Garfield, R.E. & Yallampalli, C., Structure and Function of Uterine Muscle. The Uterus. In:
Cambridge Reviews in Human Reproduction. eds. Chard, T., Grudzinskas, J.G., pp. 54–
93 (1994).

Garfield, R.E., Sims, S. & Daniel, E.E., Gap junctions: Their presence and necessity in
myometrium during parturition. *Science* 198: 958–960, DOI: 10.1126/science.929182
(1977).

Geng, X., Sun, K., Lujan, J., Tang, H. & Zhao Y., Joint Skewness and Its Application in
Unsupervised Band Selection for Small Target Detection. *Scientific reports* 5(9915): 1–
9, DOI: 10.1038/srep09915 (2015).

Govindan, R.B., Siegel, E., Mckelvey, S., Murphy, P., Lowery, C.L. & Eswaran, H., Tracking
the changes in synchrony of the electrophysiological activity as the uterus approaches
labor using magnetomyographic technique. *Reprod Sci.* 22(5): 595–601, DOI:
10.1177/1933719114556484 (2015).

Hansen, P.J., The immunology of early pregnancy in farm animals. *Reprod Domest Anim*. 46
(Suppl 3): 18–30, DOI: 10.1111/j.1439-0531.2011.01850.x (2011).

Hunter, I., Kearney, R. & Jones, L., Estimation of the conduction velocity of muscle action
potentials using phase and impulse response function techniques. *Med Biol Eng.* 25:
121–126 (1987).

Jacod, B.C., Graatsma, E.M., van Hagen, E. & Visser, G.H. A validation of
electrohysterography for uterine activity monitoring during labour. *J Matern Fetal
Neonatal Med*. 23: 17–22, DOI: 10.3109/14767050903156668 (2010).

Jedruch, J., Gajewski, Z., Kuussaari, J., The effect of detomidine hydrochloride on the
electrical-activity of uterus in pregnant mares. *Acta Vet Scand.* 30(3): 307–311 (1989a).

Jedruch, J., Gajewski, Z., Ratajska-Michalczak, K., Uterine motor-responses to an alpha-2-
adrenergic agonist medetomidine hydrochloride in the bitches during the end of
gestation and the post-partum period. *Acta Vet Scand Suppl.* 85: 129–134 (1989b).

Ju, Z. & Liu, H., Empirical Copula Based Templates to Recognize Surface EMG Signals of
Hand Motions. *Int J HR.* 8(4): 725-741, DOI: 10.1142/S0219843611002630 (2011).

Katila, T., Post-mating Inflammatory Responses of the Uterus. *Reprod Dom Anim.* 47 (Suppl.
5), 31–41 DOI: 10.1111/j.1439-0531.2012.02120.x (2012).

Katila, T., Sankari, S. & Makela, O., Transport of spermatozoa in the mare's genital tract
studied by a scintigraphic method. J Reprod Fertil Suppl 56: 571– 578 (2000).

Kendall, M.G. & Stuart, A., The advanced theory of statistics. In: The advanced theory of
statistics., no. 2nd Ed (1946).

Kitazawa, T., Kajiwara, T., Kiuchi, A., Hatakeyama, H. & Taneike, T., Muscle layer- and
region-dependent distributions of oxytocin receptors in the porcine myometrium.
Peptides. 22(6): 963–974, DOI: 10.1016/S0196-9781(01)00402-8 (2001).

Kobayashi, E., Hishikawa, S., Teratani, T. & Lefor, A.T., The pig as a model for translational
research: overview of porcine animal models at Jichi Medical University. *Transplant
Res.* 1(1): 8, DOI: 10.1186/2047-1440-1-8 (2012).

Kolda, T.G., & Bader, B.W., Tensor decompositions and applications. *SIAM review.* 51(3):
455–500, DOI: 10.1137/07070111X (2009).

Kuijsters, N.P.M., Methorst, W.G., Kortenhorst, M.S.Q., Rabotti, C., Mischi, M., Schoot,
B.C., Uterine peristalsis and fertility: current knowledge and future perspectives: a
review and meta-analysis. *Reprod Biomed Online.* 35(1): 50-71, DOI:
10.1016/j.rbmo.2017.03.019 (2017).

Lammers, W.J.E.P., The electrical activities of the uterus during pregnancy. *Reprod Sci.* 20:
182–189, DOI: 10.1177/1933719112446082 (2013).

Lukacs E., Characteristics functions. Griffin, London (1970).

Maner, W., Garfield, R., Maul, H., Olson, G. & Saade, G., Predicting term and preterm
delivery with transabdominal uterine electromyography. *Obstet Gynecol.* 101: 1254–
1260, DOI: 10.1016/S0029-7844(03)00341-7 (2003).

Markiewicz, W., Bogacki, M., Blitek, M. & Jaroszewski, J.J., Comparison of the porcine
uterine smooth muscle contractility on days 12–14 of the estrous cycle and pregnancy.
Acta Vet Scand. 58: 20–30, DOI: 10.1186/s13028-016-0201-z (2016).

Maul, H., Maner, W.L., Saade, G.R. & Garfield, R.E., The physiology of uterine contractions.
Clin Perinatol. 30: 665–676, DOI: 10.1016/S0095-5108(03)00105-2 (2003).

Mueller, A., Siemer, J., Schreiner, S., Koesztner, H., Hoffmann, I., Binder, H., Beckmann,
M.W. & Dittrich, R., Role of estrogen and progesterone in the regulation of uterine
peristalsis: results from perfused non-pregnant swine uteri. *Hum Reprod*. 21(7): 1863–
1868, DOI: 10.1093/humrep/del056 (2006).

Muthuswamy, J. & Thakor, N.V., Spectral analysis methods for neurological signals. *J*
Neurosci Methodes. 83: 1–14, DOI: 10.1016/S0165-0270(98)00065-X (1998).

Norwitz, E.R. & Robinson, J.N., A systematic approach to the management of preterm labor.
Semin Perinatol 25: 223–235, DOI: 10.1053/sper.2001.26417 (2001).

Oczeretko, E., Kitlas, A., Borowska, M., Swiatecka, J. & Laudanski, T., Uterine contractility:
visualization of synchronization measures in two simultaneously recorded signals. *Ann*
N Y Acad Sci 101: 49–60, DOI: 10.1196/annals.1389.007 (2007).

Pawlinski, B., Domino, M., Aniolek, O., Ziecik, A. & Gajewski, Z., Bioelectrical activity of
porcine oviduct and uterus during spontaneous and induced estrus associated with cyclic
hormone changes. *Theriogenology*. 86: 2312–2322, DOI:
10.1016/j.theriogenology.2016.07.029 (2016).

Pawlinski, B., Domino, M., Zabielski, R., Siewruk, K., Polanska-Plachta, M. & Gajewski, Z.,
Characteristics of bioelectrical activity of oviducts and uterus during early pregnancy in
sows recorded by telemetry method. *Exp Physiol*. 102(12): 1672–1682, DOI:
10.1113/EP086491 (2017).

Pope, W.F., Maurer, R.R. & Stormshak, F., Intrauterine migration of the porcine embryo-
interaction of embryo, uterine flushings and indomethacin on myometrial function in
vitro. *J Anim Sci*. 55: 1169–1178 (1982).

Pusey, J., Kelly, W.A., Bradshaw, J.M. & Porter, D.G., Myometrial activity and the
distribution of blastocysts in the uterus of the rat: interference by relaxin. *Biol Reprod*.
23(2): 394–397 (1980).

Rabotti, C. & Mischi, M., Propagation of electrical activity in uterine muscle during
pregnancy: a review. *Acta Physiol*. 213: 406–416, DOI: 10.1111/apha.12424 (2015).

Rabotti, C., Mischi, M., Oei, G. & Bergmans, J.W.M., Noninvasive Estimation of the
Electrohysterographic Action-Potential Conduction Velocity *IEEE Eng Med Biol Soc*.
57(9): 2178–2186, DOI: 10.1109/TBME.2010.2049111 (2010).

Rabotti, C., Mischi, M., van Laar, J.O.E.H., Oei, G.S. & Bergmans, J.W.M. Estimation of
internal uterine pressure by joint amplitude and frequency analysis of
electrohysterographic signals. *Physiol Meas*. 29: 829–841, DOI: 10.1088/0967-
3334/29/7/011 (2008).

Rath, D., Schuberth, H.J., Coy, P. & Taylor, U., Sperm interactions from insemination to
fertilization. *Reprod Domest Anim.* 43(Suppl 5): 2–11, DOI: 10.1111/j.1439-
0531.2008.01250.x (2008).

Rogers, P., Murphy, C.R., Squires, K.R. & MacLennan, A,H., Effects of relaxin on the
intrauterine distribution and antimesometrial positioning and orientation of rat
blastocysts before implantation. *J Reprod Fertil.* 68(2): 431-435, DOI: doi:
10.1530/jrf.0.0680431 (1983).

Rooijakkers, M.J., Rabotti, C., Oei, S.G., Aarts, R.M. & Mischi, M., Low-complexity
intrauterine pressure estimation using the Teager energy operator on
electrohysterographic recordings. *Physiol Meas.* 35: 1215–1228, DOI: 10.1088/0967-
3334/35/7/1215 (2014).

Sammali, F., Kuijsters, N.P.M., Schoot, B.C., Mischi, M. & Rabotti, C., Feasibility of
Transabdominal Electrohysterography for Analysis of Uterine Activity in Nonpregnant
Women. *Reprod Sci.* 25(7): 1124–1133, DOI: 10.1177/1933719118768700 (2018).

Schatz M.D., Low, T.M., van de Geijn, R.A. & Kolda T.G., Exploiting symmetry in tensors
for high performance: Multiplication with symmetric tensors. *SIAM J Sci Comput.*
36(5): C453–C479, DOI: 10.1137/130907215 (2014).

Sheffield, C., Selecting band combinations from multispectral data. Photogramm Eng Remote
Sensing. 51(): 681– 687 (1985).

Short, R.V., Implantation and the maternal recognition of pregnancy. In: Wolstenholme
GEW, eds. O'Connor, M., Ciba Foundation Symposium on Foetal Autonomy, eds.
Churchil, A., Churchil, J., London. pp. 2–26 (1969).

Singh, Y., Singh, J., Ratna, S. & Talwar, A., FFT transformed quantitative EEG analysis of
short term memory load. *Ann Neurosci.* 22(3): 176–179, DOI:
10.5214/ans.0972.7531.220308 (2015).

Sumransap, P., Tummaruk, P. & Kunavongkrit, A., Sperm distribution in the reproductive
tract of sows after intrauterine insemination. *Reprod Domest Anim.* 42: 113–117, DOI:
10.1111/j.1439-0531.2006.00696.x (2007).

Wolinski, J., Ziecik, A., Gajewski, Z., Korczynski, W., Zabielski R., A method of recording
the oviduct and uterus myoelectrical activity in gilts using implantable telemetry. *J
Anim Feed Sci.* 12: 359–367, DOI: 10.22358/jafs/67715/2003 (2003).

Wray, S., Burdyga, T., Noble, D., Noble, K., Borysova, L. & Arrowsmith, S., Progress in
understanding electro-mechanical signalling in the myometrium. *Acta Physiol (Oxf).*
213: 417–431, DOI: 10.1111/apha.12431 (2015).

Ziecik, A., Gajewski, Z., Jedruch, J. & Barcikowski, B. Effect of human chorionic
gonadotropin on myometrial electrical activity in the pig. *J Anim Sci*. 31: 131–139,
DOI: 10.1016/0378-4320(93)90033-N (1993).

Ziecik, A., Goralska, M., Krzymowski, T. & Pogorzelski, K., Isolation and purification of
porcine LH for radioimmunoassey and radioreceptor assay. *Bull Acad Pol Sci Biol*. 26:
734–744 (1978).

Ziecik, A.J., Waclawik, M.M., Kaczmarek, A., Blitek, B., Moza Jalali, B. & Andronowska,
A., Mechanisms for the establishment of pregnancy in the pig. *Reprod Domest Anim*.
46: 31–41, DOI: 10.1111/j.1439-0531.2011.01843.x (2011).

Figures and tables

Fig. 1. A. The unilateral arranged of the electrodes (Ch1, Ch2, Ch3) in the different parts of
uterus. B. The samples of EMG signal in time and frequency domain obtained at day 0 in
Ch1, Ch2, Ch3, respectively.

Fig. 2. The time-line of considering in: A. procedures of fixed-time artificial insemination,

B. hormonal status, C. ovarian status, D. content of uteri lumen.

Fig. 3. Standard statistics; mean vector ($\|\mu\|$) and covariance matrix ($\|\Sigma\|$) for seven features

of EMG signal obtained in the different parts of uterus from: A. channel 1, B. channel 2, C.

channel 3.

Fig. 4. Higher order multivariate statistics; The 3^{rd} multivariate cumulant ($\|C_3\|$) for seven

features of EMG signal obtained in the different parts of uterus from: A. channel 1, B.

channel 2, C. channel 3. The black dash lines gives the 0.975 quantile if particular date were from the Gaussian multivariate distribution.

Fig. 5. Higher order multivariate statistics; The 4^{th} multivariate cumulant ($\|C_4\|$) for seven features of EMG signal obtained in the different parts of uterus from: A. channel 1, B. channel 2, C. channel 3. The black dash lines gives the 0.975 quantile if particular date were from the Gaussian multivariate distribution.

Fig. 6. Higher order multivariate statistics; 3^{rd} cumulant's tensor based JSBS (f_{JSBS}); The examples of target function values during features elimination process for seven features of EMG signal obtained in the different parts of uterus from: A. channel 1, B. channel 2, C. channel 3 at day -6, 0, 6 of experiment.

Fetures / day	Ch	- 6	- 5	- 4	- 3	- 2	- 1	0	1	2	3	4	5	6	7	8	9	10	11	12
D	1	IV	VI	VI	V	VI	VI	IV	VI	VII	VI	VI	VI	V	IV	VI	VI	VI	IV	III
Duration	2	IV	IV	IV	VI	VI	VI	VI	VI	VII	V	VI	VI	VI	VI	VII	IV	VI	III	VI
[s]	3	VI	VI	VI	IV	VI	VII	VI	VII	VII	V	II	VI	IV	VI	VI	VII	VI	VI	VI
P	1	II	I	II	VI	V	V	VI	II	V	IV	III	I	I	I	II	II	III	II	IV
Pause	2	I	II	II	IV	I	I	I	I	VI	I	II	I	I	I	I	III	I	II	III
[s]	3	III	III	I	VI	V	I	I	I	III	II	VI	I	II	I	II	II	V	I	II
A	1	VI	V	IV	III	IV	IV	II	V	III	II	II	V	VI	VI	V	V	V	VI	VI
Amplitude	2	V	V	VI	III	V	IV	V	V	III	VI	IV	IV	V	IV	V	VI	IV	V	IV
[mV]	3	IV	IV	IV	III	III	V	V	V	IV	IV	IV	IV	V	III	IV	IV	II	IV	V
RMS	1	VII	VI	V	IV	III	III	III	VI	II	V	IV	IV	IV	V	IV	IV	IV	V	V
Root mean square	2	VI	VI	V	V	IV	III	VI	VI	VI	III	V	V	IV	V	IV	V	V	VI	V
[mV]	3	V	V	V	V	II	IV	IV	IV	VII	VI	III	III	VI	IV	V	V	I	V	IV
Max P	1	III	III	III	II	I	II	V	III	I	III	V	III	III	III	III	III	II	III	II
Maximum power	2	III	III	III	II	III	V	III	II	II	VI	III	III	III	III	II	III	IV	II
[V^2]	3	II	II	III	II	IV	III	III	III	II	III	V	II	III	II	III	III	III	III	I
Min P	1	I	II	I	I	II	I	I	I	IV	I	I	II	II	II	I	I	I	I	I
Minimum power	2	II	I	I	I	II	II	II	III	I	II	I	II	II	II	II	I	II	I	I
[V^2]	3	I	I	II	I	I	II	II	II	I	I	I	V	I	V	I	I	IV	II	III
DF	1	V	VII	VII	VII	VII	VII	VII	VII	VI	VII	VII	VII	VII	VII	VII	VII	VII	VII	VII
Dominant frequency	2	VII	VII	VII	VII	VII	VII	VII	VII	V	VII	VII	VII	VII	VII	VII	VII	VII	VII	VII
[Hz]	3	VII	VII	VII	VII	VII	VI	VII	VI	V	VII	VII	VII	VII	VII	VII	VI	VII	VII	VII

672   Footnotes: D, P, A, RMS, Max P, Min P, DF - the EMG signal features considered in experiment; Ch - the different parts of uterus from

673   (channel 1, channel 2, channel 3); day - the days considered in experiment; I - VII - the hierarchy of features indicated by 3rd cumulant's tensor

674   based JSBS.