Programmed Delayed Splicing: A Mechanism for Timed Inflammatory Gene Expression

Jacob S. Dearborn[1], **Luke Frankiw**[2], **Damas W. Limoge**[3], **Christian H. Burns**[2], **Logan Vlach**[2], **Patricia Turpin**[2], **Tylar Kirch**[1], **Zachary D. Miller**[1], **William Dowell**[1], **Sylvester Languon**[1], **Yvette Garcia-Flores**[1], **Robert C. Cockrell**[1], **David Baltimore**[4], **Devdoot Majumdar**[1,2]

*For correspondence:
dev.majumdar@uvm.edu
(Devdoot Majumdar)

[1]Department of Surgical Research, Larner College of Medicine, University of Vermont, Burlington, United States; [2]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, United States; [3]Nanotronics, Brooklyn, New York, United States; [4]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, United States

Abstract   Inflammation involves timed gene expression, suggesting that the fine-tuned onset, amplitude, and termination of expression of hundreds of genes is of critical importance to organismal homeostasis. Recent study of post-transcriptional regulation of inflammatory gene expression led to the suggestion of a regulatory role for pre-mRNA splicing. Here, using a hybrid capture approach to purify incompletely spliced, chromatin-associated pre-mRNAs, we use deep sequencing to study pre-mRNA splicing of the NF-κB transcriptome. By freezing transcription and examining subsequent splicing of complete transcripts, we find many introns splice tens to hundreds of times slower than average. Investigating the basis of these delays, we focused on evolutionarily conserved introns with suboptimal splice donor sequences and found that strengthening these donor sites by as few as two nucleotides in minigene reporter assays markedly increased gene expression for several targets. This suggests that such sites can act as timing elements that both delay mRNA production and limit expression amplitude. To broaden this mechanistic view, we applied deep learning sequence-to-function models with feature attribution to identify additional regulatory sequences—both intronic and exonic—that may contribute to delayed splicing through mechanisms independent of donor site strength. This integrated approach revealed non-canonical motifs enriched in slow-splicing introns, pointing to a broader repertoire of cis-elements that can fine-tune transcript maturation during inflammation. Together, these findings support a model in which the temporal regulation of pre-mRNA splicing serves as a layer of control in inflammatory gene expression, and raise the possibility that similar timing mechanisms operate in other rapid-response transcriptional programs.

Keywords: inflammation, innate immunity, macrophage, gene expression, splicing

Introduction

Gene expression in response to an inflammatory stimulus begins rapidly and is tightly controlled by conventional means (transcription and protein turnover; (*Chen and Chen, 2013*; *Gautier et al., 2012*; *Smale et al., 2014*)) and by an expanding list of modalities that have gained in appreciation as being general regulatory strategies (RNA stabilization, RNA deadenylation, ribosomal regulation, microRNA regulation, as examples; (*Hao and Baltimore, 2009*; *Leppek et al., 2013*; *O'Connell et al., 2012*; *Wan et al., 2007*)). We and others have recently investigated the role of RNA splicing kinetics—independently of alternative splicing—in gene expression (*Hao and Baltimore, 2013*; *Pandya-Jones et al., 2013*; *Rabani et al., 2011*, *2014*). In macrophages, an inflammatory stimulus leads to upregulation of expression of pre-mRNAs from hundreds of genes, providing an experimentally favorable system to investigate whether differential kinetics of pre-mRNA splicing may control the timing of gene expression following an inducing stimulus.

Pre-mRNA conversion to mRNA has been implicated in regulation of gene expression in diverse systems. As part of the cellular response to various environmental stressors, mRNAs for ribosomal proteins were shown to be downregulated due to decreased splicing efficiency in yeast (*Bergkessel et al., 2011*). Global changes in efficiency of pre-mRNA splicing have been shown to be a developmental prerequisite for *Drosophila* early embryonic development (*Guilgur et al., 2014*). The developing vertebrate embryo obeys a "segmentation clock" determining body segment length whose very timing relies on delays attributable to control of the splicing rate of the *Hes7* transcriptional repressor (*Takashima et al., 2011*).

In certain well-studied cases, as with the cytokine TNFα, regulatory mechanisms modulating RNA levels exert significant physiological effects (*Eissa et al., 1996*; *Hargreaves et al., 2009*). The insight that TNFα contains AU-rich elements in its 3′ untranslated region that act as mRNA degradation signals (*Han et al., 1990*), and subsequent observations that a mouse in which these AU-rich elements were removed resulted in a robust autoimmune phenotype (*Kontoyiannis et al., 1999*), was an early indication of the importance of precisely tuned mRNA levels in the regulation of inflammation to avoid autoimmunity. Given the role of pre-mRNA splicing in biogenesis of mature mRNA, we and others (*Hao and Baltimore, 2013*; *Bhatt et al., 2012*; *Cho et al., 2014*; *Davis-Turak et al., 2015*; *Grabherr et al., 2011*) suggested that regulation of splicing kinetics may influence the gene expression kinetics that define the inflammatory cascade. Consistent with this idea, *Braunschweig et al.* (*2014*) demonstrated that intron retention is a widespread and regulated feature of mammalian transcriptomes, often associated with reduced mRNA abundance and subject to tissue-specific control.

Deep learning models have rapidly advanced our ability to interpret noncoding regions of the genome by learning complex regulatory logic directly from DNA sequence. Early models such as DeepSEA demonstrated that convolutional neural networks could predict the effects of noncoding variants on transcription factor binding and chromatin state, laying a foundation for sequence-to-function inference (*Zhou and Troyanskaya, 2015*). Building on this, SpliceAI adapted deep convolutional architectures to model splic-

ing dynamics directly from primary sequence, achieving high accuracy in predicting splice site usage and the effects of sequence variants on splicing (*Jaganathan et al., 2019*). More recent models such as Enformer and Borzoi have expanded this paradigm further by predicting RNA-based outputs, including gene expression levels, transcript isoform usage, and transcription start site precision (*Avsec et al., 2021*; *Linder et al., 2025*). These models also support feature attribution through interpretable machine learning approaches such as saliency mapping and deepSHAP, allowing inference of sequence elements—such as transcription factor binding sites and splicing regulators—that most influence RNA output. Together, these approaches provide an unprecedented modeling context—spanning hundreds of kilobases—that enables the systematic dissection of sequence-level determinants of intron retention and the broader landscape of RNA processing.

To examine the timing of intron removal from 230 different transcripts induced by TNFα in macrophages, we have developed a method for highly enriching transcript populations for mRNAs of interest, which is followed by deep sequencing of the largely pre-mRNA populations we purify. The induction of transcripts and removal of introns can be quantified with precision, with lifetimes of introns determined by blocking transcription early after induction. Among genes whose mRNAs appear more slowly after induction, we identify ones containing introns with poor binding sites for splicing factor U1, usually finding one per transcript. We call these "bottleneck" introns. Among the most rapidly induced genes we find no such introns. We show that in these pre-mRNAs, the sequence of the U1 binding site is critical for the speed of intron removal by building mini-genes with "repaired" introns and showing that these splice at the canonical rate (identified as about 20 seconds after polymerase has passed that point). To complement these findings and explore regulatory determinants at larger scale, we apply a deep learning model trained on genomic sequence to predict splicing dynamics. Feature attribution methods identified sequence features associated with the splicing delays. We observe and model fine-tuning improved detection of patterns associated with intron retention. We propose that bottleneck introns are important for determining either the rate of degradation of pre-mRNAs or the rate of appearance of mature mRNA or both.

Results

Hybrid Capture of Chromatin-Associated Transcripts

It has been established that pre-mRNA is highly enriched in the chromatin-associated, polyadenylated RNA fraction (*Tilgner et al., 2012*). To examine splicing events in pre-mRNAs, we performed time-course experiments with TNF-stimulated bone marrow-derived macrophages (BMDMs), isolating RNA after biochemical separation of chromatin-associated material (*Bhatt et al., 2012*). Using a hybrid-capture approach, we targeted sequencing toward transcripts of 230 genes previously identified as TNF-induced inflammatory mRNAs (*Ramirez-Carrozzi et al., 2009*).

The hybrid capture strategy, based on a published approach (*Engreitz et al., 2013*), involved: (1) microarray printing of 12,000 150-bp ssDNAs designed from tiled fragments of the last exon of each gene of interest; (2) conversion to a pool of biotinylated ssRNA probes using PCR followed by in vitro transcription with biotinylated ribonucleotides; (3) hybridization of ssRNA pools to cDNA from each biological experiment; and (4) streptavidin-

coated bead-mediated capture of the transcripts of interest (Fig. S1).

This approach resulted in a ~30-fold enrichment of genes of interest, with 70% of the sequenced reads corresponding to the genes of interest; RNA submitted only to poly(A) selection contained only 2% of such reads (Fig. S2A).In this way we could analyze 1098 introns (after TPM filter >100) from genes induced by an inflammatory stimulus (Fig. S2B).

The selected chromatin-associated transcripts collected across time points after induction are shown as read-density histograms, displaying sequencing read abundance along each gene to reveal exon–intron structure. In Fig. 1A, *Nfkbia* (encoding IκBα) is shown over 1 hour after TNF induction, with $\log_{10}$-scaled densities normalized within each time point to highlight transcriptional kinetics. New transcription becomes evident by 6 minutes, especially on an unnormalized linear scale (Fig. S3). The corresponding log-scaled histograms permit visualization of the intronic signal as a function of time after induction.

Furthermore, from Fig. 1A it is evident that at all time points following induction, the 5′-proximal introns have been totally removed from the sequenced transcripts, indicating that the selection against partial transcripts is quite complete. Whereas excision of the first intron is always observed, the middle three introns are seen at intermediate states of excision at all time points such that intron definition for these introns is readily observable from read density histograms. We attribute this first exon excision largely to co-transcriptional splicing, consistent with other genome-wide splicing studies (*Bentley, 2014*). Strikingly, the final intron deviates significantly in its kinetic trajectory, as its read density does not obey a similar relative reduction. This might be due to a lag in terminal intron splicing (*Carrillo Oesterreich et al., 2010*) or a feature of splicing that accompanies transcript release from chromatin.

Quantifying Splice Completion Across the Transcriptome

To better quantify the observed dynamics, we adapted the Coefficient of Splicing (CoSI) (Fig. 1B), which quantifies the extent of splicing as a ratio of spliced to total (spliced and unspliced) junction reads such that CoSI values of ~1 and ~0 imply near-complete splicing and virtually unspliced states, respectively (*Tilgner et al., 2012*). Though we observed a decrease in read density as a function of distance from the 3′ end of the gene (Fig. S3), presumably as a consequence of premature termination of the reverse transcriptase during copying of the pre-mRNA, the use of CoSI allows for an intron-specific splicing score regardless of read densities at neighboring introns. Using the CoSI metric, a time course plotting of the extent of splicing showed a dip in CoSI at ~6 minutes (Fig. 1C) corresponding to the aforementioned accumulation of new, unspliced transcripts. The splicing dynamics of each *Nfkbia* intron can be inferred from the CoSI dynamics, and the notable difference in splicing between the 5′ and 3′ introns (Fig. 1D) is demonstrated by the amplitude of the "dip" at 6 minutes and the time required for each intron to return to CoSI of 1.

A surprising heterogeneity in CoSI was observed among all inflammatory introns (Fig. 2A), implying diversity in their propensity to be spliced (Fig. 2). When considering all 1,024 introns in the chromatin-associated TNF time course we find most introns very near to CoSI ~1 relatively soon after induction, indicating that most introns do not remain unspliced long after induction begins between 4–8 minutes (Fig. 2A). The observation that many TNF-responsive transcripts initiate within 4–8 minutes is consistent with that of other re-

lated studies (*Bhatt et al., 2012*). Unexpectedly, although the median CoSI value remains high, we identified considerable heterogeneity among introns, some at and remaining near CoSI values < 0.5, indicating relatively poor splicing, found very late into the time course.

As an example of a poorly spliced intron, chemokine *Cxcl10* intron 2 (Fig. 2A) is notable as it remains poorly spliced despite clear excision of neighboring introns, remaining quite unspliced even ~30 minutes after induction. It is possible that this intron undergoes splicing after its nascent chromatin-associated state, as is likely the case for the 3′-terminal intron of *Nfkbia*. It is also possible that this intron targets *Cxcl10* transcripts for degradation and the relatively fixed nature of intron 2's splicing status throughout the time course is a function of a constant rate of degradation. Introns with low CoSI at late time points post-TNF induction were considered putative "bottleneck introns"—borrowing from the language that accompanied the discovery and characterization of slowly splicing U12-type introns (*Patel et al., 2002*). These introns were so slow to splice that they may intrinsically delay gene expression. Notably, the distribution of CoSI values of the entire dataset (Fig. 2A) was very broad, and though most introns spliced immediately, many introns showed evidence of splicing bottlenecks, noticeable by their significant deviation from mean CoSI. At 10 and 60 minutes post-induction, 14% and 11% of introns, respectively, had CoSI values below one standard deviation of the mean (0.86 ± 0.25 and 0.91 ± 0.19, respectively). Shown as examples are *Cd40*, *Cxcl10*, *Daxx*, and *Irf7* (Fig. 2A–B), genes whose immunological and inflammatory importance is well-established in studies with knock-out mice (*Honda et al., 2005*; *Lei et al., 1998*; *Michaelson et al., 1999*).

Measurements of Intron Splicing Half-Lives

To quantify splicing kinetics, we used Actinomycin D (actD) to freeze transcription and followed the loss of intron and accumulation of splice junctions. In these experiments, splicing was analyzed at many time points immediately following actD treatment on the same 230 transcripts of interest and selected using hybrid capture from the total pool of cellular RNA rather than chromatin-associated RNA. Fitting the accumulation of spliced transcripts (as measured by CoSI) with an exponential distribution, we were able to extrapolate intron excision half-lives (Fig. 3). Because total cellular RNA was used, observed rates were independent of chromatin localization. We find intron half-lives that range from 20–40 s (56% of introns splicing in this timing window) to several minutes, reflecting the considerable heterogeneity that is observed from CoSI differences.

A Multifaceted Approach for Identification of Determinants of Slow Splicing

Building on our observation that certain introns exhibit markedly delayed splicing by both CoSI dynamics and direct half-life measurements, we sought to uncover the underlying sequence features responsible for these delays. To this end, we employed a multifaceted strategy: (1) computational analysis of 5′ splice donor strength using MaxEntScan, (2) experimental testing of splicing efficiency with a minigene reporter assay, and (3) interpretable deep learning to identify additional cis-regulatory elements—beyond canonical splice sites—that may modulate splicing kinetics.

Computational Analysis of Splice Donor Strength

A delay in splicing at certain sites could simply confer a delay in gene expression (by ~5 minutes in the slow case shown in Fig. 3, IκBε), or, as is seen in yeast studies, it could result in both gene expression delay and gene expression diminution due to degradation of slowly splicing pre-mRNA (*Koodathingal et al., 2010*). Prior studies placed IκBε in a delayed splicing category (*Hao and Baltimore, 2013*), suggesting a pronounced splicing delay relative to rapidly induced genes. To understand a potential mechanistic basis for these differences in splicing time, each intron within our dataset was assessed computationally for the concurrence of its 5′ splice donor sequence to a consensus sequence (*Pessa et al., 2006*). The 5′ splice donor is a highly conserved sequence that directly base pairs with splicing factor U1 (*Freund et al., 2005*); deviation from the consensus sequence confers a significantly reduced ability to engage the splicing machinery.

A maximum entropy model (*Yeo and Burge, 2004*) was used to calculate an intron quality score measuring extent of deviation from the consensus splice sequence (Fig. S4). Among the inflammatory transcripts studied, many examples of introns with poor 5′ donor scores were identified such as *Irf7* and *Il12b*, where lower scores indicate significant deviation from consensus. We suggest that having non-consensus splice sites may be a regulatory mechanism affecting gene expression. We considered that splicing might show profound differences in the previously defined categories of induction (immediate/early/intermediate/late) characteristic of the inflammatory gene expression kinetics (*Bhatt et al., 2012*). We found that the "immediate" genes showed consistently fast splicing (highest CoSI values) at all of their introns except the most 3′ intron, whereas the other three groups shared similar CoSI distributions (Fig. S6). Using the bioinformatics "intron quality score" we also found that the introns of the later gene classes have significantly lower scoring 5′ splice donor sequences (Fig. S4). Therefore, from experimental measurement of splicing and sequence-based prediction, genes expressed immediately following inflammatory stimulus are spliced faster, whereas all other inflammatory genes have a complex and heterogeneous distribution of splicing efficiency that does not stratify cleanly into the later kinetic categories (early/intermediate/late). Slowly splicing introns are found throughout these later kinetic categories in similar abundance, perhaps playing very gene-specific roles in diverse kinetic categories.

Experimental Validation with Minigene Fluorescent Reporter Assay

To test whether delays in splicing result in changes to gene expression, we identified a set of introns with the following criteria: (1) introns that splice poorly as defined by RNA-Seq, (2) introns that contain a low-scoring (non-consensus) 5′ splice donor, and (3) introns whose weak 5′ splice donor is evolutionarily conserved across many mammalian species. These introns were tested in the context of a splicing reporter expressed on a bidirectional promoter (*Mukherji et al., 2011*). For each intron of interest, the reporter construct consists of a single transcript containing: (1) the 5′ neighboring exon from the gene of interest, the intron of interest, and the 3′ neighboring exon; (2) a 2A "self-cleaving" peptide; and (3) the GFP gene. In the opposite orientation but from the same promoter, a blue fluorescent protein (BFP) mRNA is made in equal amounts to the intron-GFP construct. GFP fluorescence of cells transfected with this reporter is a readout of splicing efficiency of the intron-GFP construct, when normalized to BFP fluorescence levels. Transfected into HEK293T cells and expressed for 24 hours, this bidirectional reporter enabled

us to understand, at steady state, whether gene expression is affected by slow splicing introns. Measured using flow cytometry, the slope of the line corresponding to BFP:GFP ratios provides a relative metric of splicing efficiency, where slopes ~0.9 and ~0.1 imply efficient and inefficient splicing, respectively.

To test the effect of a poor splice donor, we "rescued" some poorly splicing introns in the context of the reporter by mutation to consensus splice donor "GTAAG." For instance, for *IL12* intron 3, the splice donor sequence of "GTAAT" that is conserved among many mammalian species was altered to "GTAAG" (Fig. S5). Expression levels of the reporter construct with the wild-type *IL12* intron were found to be about half (57%) of the levels of the same construct with a single base pair alteration to make the stronger splice donor. *IRF7* intron 5 was tested against a "fixed" intron as well as a wild-type intron from an actin gene, both resulting in two-fold improvements of gene expression. In one case, expression of TFEC (transcription factor EC) was not altered by splice site repair, suggesting that other mechanisms beyond 5'–splice site deficiencies may be involved in mediating slow splicing. Generally, when the BFP:GFP slopes of wild-type and mutated introns were compared by taking a ratio of their slopes, a change of ~2 nucleotides dramatically altered the slope of the line (Fig. 4).

Deep Learning–Based Discovery of Non-Canonical Regulatory Motifs

To extend our understanding of regulatory features beyond canonical splice donor sequences, we applied a supervised regulatory sequence model (Borzoi) to investigate sequence determinants associated with slow splicing. Borzoi is a model originally trained on steady-state RNA profiles across diverse cell types and tissues, which may limit its ability to capture stimulus-specific regulatory dynamics at play during TNFα-induced intron retention. Fine-tuning of this model offers a way to adapt a large, pre-trained sequence model to a specialized context by continuing training on domain-specific datasets, leveraging the general regulatory knowledge already encoded in the model weights while learning new patterns relevant to the task at hand. Fine-tuning has been shown to enhance Borzoi's performance in other specialized contexts, including tissue-specific expression, transcription factor knockdowns, and cellular aging (*Yuan et al., 2025*). We fine-tuned it using RNA-seq coverage data from BMDMs at 18, 20, and 30 minutes post-TNFα induction—timepoints that capture dynamic intron retention. Fine-tuning was performed over 20 epochs using the Adam optimizer (learning rate = 1×10^{-6}, MSE loss). To evaluate generalization, chromosomes 10 and 11 were held out for training and used for validation and testing, respectively. Model loss steadily decreased over the training period for both datasets (Fig. 5A), and Pearson correlation (reflecting the strength of their linear relationship) between predicted and observed expression improved from $r = 0.51$ ($p = 4.44 \times 10^{-16}$) for the pretrained model to $r = 0.61$ ($p = 4.84 \times 10^{-24}$) after fine-tuning (Fig. 5B).

To interpret the model's predictions and identify candidate regulatory elements, we applied a widely adopted backpropagation-based technique, DeepLIFT (*Shrikumar et al., 2019*), to compute nucleotide-resolution importance scores, capturing the contribution of each base to splicing of the retained intron. Attribution scores were computed for each intron along with sequence within the model input window (524 kb). In these scores, larger positive values indicate nucleotides that the model predicts as contributing to intron retention (i.e., delaying splicing), whereas negative values indicate nucleotides pre-

dicted to be associated with splice completion. We analyzed these scores using modisco-lite, which clusters recurring high-importance patterns into position weight matrices (PWMs) representing putative regulatory motifs (*Shrikumar et al., 2020*). To assess motif enrichment among delayed introns, we focused on the 50 slowest-splicing introns as ranked by area under the CoSI-time curve (Fig. S7C). Motif occurrences were identified using a conventional motif scanning approach, Finding Individual Motif Occurrences (FIMO) (*Grant et al., 2011*), and enrichment was calculated relative to either the remaining introns in our dataset or all introns in the transcriptome.

We compared the frequency of putative regulatory motifs in the 50 slowest-splicing introns to their frequency across all introns in the murine transcriptome, highlighting both the most enriched motifs overall and specific examples of their genomic locations within delayed introns. Among these, a GA-repeat–like motif showed enrichment in the slow-splicing set (FIMO, 80.0% vs 55.2%) (Fig. 6A). Representative instances of this motif were also recovered from the original source seqlets used to generate the PWM by modisco-lite, providing further support for its regulatory relevance (Fig. 6B–D). These findings illustrate how sequence-to-expression modeling, combined with attribution-based motif discovery, can uncover non-canonical cis-regulatory elements that may influence splicing efficiency during inflammatory gene activation.

Discussion

In this study we sought to understand splicing kinetics of the large number of genes that comprise the inflammatory response and to assess whether splicing itself might play a regulatory role in inflammation. We developed a targeted sequencing strategy, purifying transcripts containing each gene's terminal exon. This approach allowed us to sequence the 1,024 introns within inflammatory genes and permitted direct assessment of the structures of nearly-completed transcripts. We found considerable heterogeneity in splicing efficiency among these introns. In studying evolutionarily conserved weak 5′ splice donors, we have isolated one cause of slow appearance of mRNA following a pulse of stimulus; many other slowly spliced introns without such sequences were also identified in this study and suggest other regulatory mechanisms may be responsible.

Crucially, the hybrid capture approach averts a common ambiguity in analyzing splicing kinetics of not being able to differentiate completed pre-mRNA from nascent transcripts during an induction pulse—this often leads to an overestimation of the unspliced status of early introns and complicates quantification of splicing kinetics. To the contrary, we rarely found genes containing unspliced first introns. This was true of chromatin-associated RNA and of whole-cell RNA after inhibition of transcription with actD. These effects are consistent with the emerging model of co-transcriptional pre-mRNA splicing, where the splicing machinery has been suggested to lag 3–5 kb behind the polymerase. Indeed, several recent global studies of RNA splicing bolster the claim that much pre-mRNA is spliced co-transcriptionally: 74% (yeast) or 75–84% (human) of introns are found to be at least 50% spliced by the time of transcription termination in several other studies (*Tilgner et al., 2012*; *Carrillo Oesterreich et al., 2010*; *Ameur et al., 2011*; *Brugiolo et al., 2013*; *Girard et al., 2012*; *Khodor et al., 2011*). Surprisingly, this ~80% figure remains constant whether total RNA or chromatin-associated RNA is measured, implying that our choice to analyze chromatin-associated RNA does not significantly overrepresent splicing intermediates.

We found that most introns are spliced very efficiently, appearing and disappearing as a rapid dip of CoSI immediately following induction, returning to a CoSI of ~0.95 within minutes after induction. Notably, the distribution of CoSI values of the entire dataset (Fig. 2) was very broad. Though most introns spliced immediately, there were several "bottleneck introns." In order to determine more specifically the rates of slowly splicing introns, studies employing actD to stall transcription and examine intron splicing half-lives corroborated the idea that there is tremendous intron-to-intron heterogeneity. Most delayed introns ultimately reached higher CoSI values over the time course, consistent with completion of splicing rather than stable, long-term retention (see Figs. 2–3). Whereas most introns spliced within 20–40 s, some were delayed significantly (upwards of 5 minutes). Of note, however, is that our 20–40 s rate of splicing is somewhat at odds with other figures in the literature of 8–10 minutes for intron excision (*Pandya-Jones et al., 2013*) after a washout of the drug D-ribofuranosylbenzimidazole. There is some debate as to the perturbative role of actD in splicing, with one report observing that splicing intermediates in the context of the MS2 reporter system are prematurely liberated from chromatin upon actD treatment. Even in this case, the rapid actD-based rates are likely underestimating even faster kinetics if one considers that the co-transcriptional splicing machinery targets chromatin mRNA faster than released mRNA (*Martin et al., 2013*). However, even in the absence of actD, stimulation revealed that most of *IκBα*'s introns are spliced in less than two minutes when one takes into account the "dip" in CoSI due to induction and the time to reach steady CoSI levels (Fig. 2). While the terminal intron of *IκBα* appears to have a longer half-life, this unique feature of terminal introns is consistent with prior studies (*Carrillo Oesterreich et al., 2010*).

In testing gene expression differences in bottleneck introns among introns with poor splice sites that are also evolutionarily conserved, we found that steady-state levels of reporter proteins were upregulated when the 5′ splice donor sequence was mutated to the consensus sequence 'GTAAG' in all cases but one. Attenuated U1 binding provides a mechanistic insight for bottleneck introns that were chosen for their weak 5′ splice donors. This implies that at the level of splicing, either due to delays in expression or perhaps degradation due to delayed expression, significant differences in gene expression arise from small differences in nucleotide sequence. These reporter assays measure steady-state expression influenced by intron sequence; while consistent with changes in splicing efficiency, other post-transcriptional mechanisms may be at work to preclude these introns from efficient splicing. We also find many slowly spliced introns not explained by weak 5′ splice donors. In some cases, we find multiple bottleneck introns per gene, as is the case of *IRF7*, where one bottleneck (intron 5) was attributable to an evolutionarily conserved weak 5′ splice donor while another (intron 7) was observed experimentally but of unknown cause. This may be due to any of a number of potential mechanisms that may also serve in tuning the speed of splicing: cis-regulatory protein recruitment, 3′ splice acceptor sequence or other sequence elements, or alterations of RNA polymerase speed or chromatin marks or three-dimensional gene structure.

Central to our inquiry is the enigmatic nature of these bottlenecks remaining in physiologically critical genes, often evolutionarily conserved, and yet intrinsically mediating an inefficiency in gene expression. Importantly, the conservation of these weak donor sites suggests they confer a regulatory advantage—such as fine-tuning of expression timing or transcript abundance—rather than representing neutral or deleterious features merely

tolerated by selection. We posit that the gene expression changes that are shown in bone marrow-derived macrophages offer a regulatory strategy to slow up and maybe restrict expression of genes in a manner dependent on the composition of mRNA processing factors in the cell ("the splicing landscape"), the cell type, or the stimulus type in question. Recent studies have demonstrated global changes in intron retention preferences in B cell lymphomagenesis and granulocyte differentiation (*Koh et al., 2015*). In a similar manner, we suggest that selection of splicing and kinetics of splicing might allow a previously unappreciated level of specificity to gene expression decisions in cells presented with an inflammatory stimulus (57–63).

Induction with TNF is a particularly favorable situation because many of the genes we examined were up-regulated in their transcription within 4–6 minutes of adding inducer (Fig. S2B), allowing examination of large numbers of pre-mRNA transcripts. This, in concert with the hybrid capture approach that provides a large number of junctional sequencing reads, has permitted unique insight into the kinetics of splicing of mature transcripts and revealed surprising heterogeneity. We suggest that this methodology and analysis could have wider applicability for other gene induction situations.

Despite the promise of deep learning models, challenges remain in translating their predictions into biological insight. While performance on benchmark tasks continues to improve—often surpassing traditional motif-finding tools—generalization across conditions and cell types is still constrained by the scope of training data. Interpretability is another major hurdle, particularly for genomic sequences. Interpretable machine learning methods can highlight sequence elements that influence model predictions, but connecting these regions to underlying biological mechanisms often demands extensive experimental validation. In the case of RNA splicing, complexity of overlapping regulatory layers—such as RNA secondary structure, co-transcriptional dynamics, and nuclear export—can make it difficult to disentangle causal sequence features from correlated signals. Nonetheless, sequence-to-expression modeling provides a powerful framework for identifying sequence features correlated with delayed splicing and for guiding experimental discovery of novel regulators through fine-tuning and perturbation-informed learning.

Our deep learning–based analysis further supports the notion that splicing efficiency is shaped by a broad array of cis-regulatory features, many of which fall outside canonical splice site motifs. By leveraging a sequence-to-expression transformer trained on multimodal regulatory data, we identified non-canonical sequence motifs enriched in the slowest-splicing introns—motifs that may act as silencers or delay elements involved in splicing. These results offer a complementary, agnostic perspective to our mutational analyses, and suggest that intronic bottlenecks can arise from diverse sequence architectures. The enrichment of these motifs in bottleneck introns—independent of donor site strength—points to additional layers of splicing control that may be particularly relevant in rapid-response transcriptional programs like inflammation. We propose that such regulatory elements may encode a form of temporal tuning, modulating transcript availability through fine control of intron excision. More broadly, our work demonstrates that interpretable machine learning can uncover latent regulatory features that elude traditional sequence analysis, advancing efforts to decode the logic of splicing regulation in a cell-type or stimulus-specific context.

Methods

Cells

C56BL6/J mice were sacrificed via CO_2 euthanasia and sterilized with 70% ethanol. Femur and tibia bones were harvested and stripped of muscle tissue. Bone marrow cells were resuspended in 20 mL of fresh DMEM. 2.5×10^6 bone-marrow cells were plated in a 15-cm dish in 20 mL of BMDM media (DMEM, 20% FBS, 30% L929 conditioned media, and 1% Pen/Strep) and grown at 5% CO_2 and 37°C. BMDM media was completely replaced on day 3 as well as a supplemental addition of 5 mL L929 conditioned media on day 5.

RNA fractionation

RNA was fractionated into cytoplasmic, nucleoplasmic, and chromatin-associated pools as previously described (*Bhatt et al., 2012*) with modifications. Confluent 15 cm dishes of mature BMDMs were scraped into 400 μL cold NP-40 lysis buffer (10 mM Tris-HCl pH 7.5, 0.08% NP-40, 150 mM NaCl) and layered onto a 1 mL sucrose cushion (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 24% w/v sucrose). Samples were centrifuged at 13,000 rpm for 10 min at 4°C. The supernatant (cytoplasmic fraction) was mixed with 3 volumes of 100% ethanol and 2 volumes of buffer RLT (4 M GuSCN, 0.1 M β-mercaptoethanol, 0.5% N-lauroylsarcosine, 25 mM Na-citrate, pH 7.2) and stored at –80°C.

The nuclear pellet was resuspended in 200 μL cold glycerol buffer (20 mM Tris-HCl pH 7.5, 75 mM NaCl, 0.5 mM EDTA pH 8.0, 50% glycerol, 0.85 mM DTT) and lysed with an equal volume of nuclear lysis buffer (20 mM HEPES pH 7.5, 7.5 mM $MgCl_2$, 0.2 mM EDTA pH 8.0, 1 M urea, 1% NP-40). After vortexing and centrifugation (14,000 rpm, 5 min, 4°C), the supernatant (nucleoplasmic fraction) was processed as above.

The remaining chromatin pellet was hydrated in 1× PBS and dissolved in 500 μL TRIzol reagent at 50°C with intermittent vortexing. After phase separation with 100 μL chloroform, the aqueous phase was recovered and processed as above. All RNA fractions were purified using the Qiagen RNeasy protocol and eluted in nuclease-free water. Typical yields were 300–500 ng/μL for cytoplasmic RNA, 100–250 ng/μL for nucleoplasmic RNA, and 300–500 ng/μL for chromatin-associated RNA. RNA was DNase-treated (TURBO DNase, Thermo Fisher) and stored at –80°C.

Template-switch reverse transcription

One microgram of RNA was mixed with 1 μM oligo(dT)$_{30}$ (5′-AAGCAGTGGTATCAACGCA GAGTACT$_{30}$-3′), heated to 80°C for 2.5 min, and snap-cooled on ice. A 10 μL reverse-transcription mix containing 10 μM template-switch oligo (5′-AAGCAGTGGTATCAACGCAG AGTACACArGrGrG-3′), 20 mM DTT, 2× First-Strand Buffer (Invitrogen), 1 mM dNTPs, 40 U Murine RNase Inhibitor (NEB), and 200 U SuperScript II (Invitrogen) was added. Reactions were incubated sequentially at 42°C (30 min), 45°C (30 min), and 50°C (10 min), then heat-inactivated at 80°C (10 min).

RNA templates were degraded by adding NaOH (final 0.1 M) and EDTA (5 mM) and heating to 70°C for 10 min, followed by neutralization with HCl. cDNA was purified using 2× Sera-Mag carboxylate-modified magnetic beads (GE Healthcare) with standard PEG/ethanol washes and eluted in nuclease-free water.

Hybrid capture probe design and synthesis

Biotinylated RNA probes were designed against the terminal exons of inflammatory genes (see Supplemental Table 2). For each gene, 100-bp overlapping oligonucleotides were synthesized (CustomArray Inc.) and pooled into nine expression-matched subgroups. Each subpool was PCR-amplified to append a T7 RNA polymerase promoter and transcribed in vitro using the AmpliScribe T7 Biotin Flash Kit (Epicentre). Purified RNA probe subpools were combined in weighted ratios (A–I) to normalize capture representation across expression levels as below:

Pool	Pool composition
A	0.01113
B	0.02110
C	0.03235
D	0.05514
E	0.10542
F	0.21908
G	0.46740
H	1.00000
I	1.00000

cDNA hybrid capture and elution

Biotinylated probes were hybridized to cDNA at 74°C for 4.5 min, followed by addition of 2× hybridization buffer (1 M LiCl, 40 mM Tris-HCl pH 7.5, 20 mM EDTA pH 8.0, 4 M urea, 0.5% Triton X-100, 1% SDS, 0.2% Na-deoxycholate). Hybridization continued 30 min at 70°C. Streptavidin BioMag beads (0.3 mg) were washed and incubated 20 min at 70°C to capture cDNA–probe complexes. Beads were washed twice with 1× HYB, once each with Wash 4 and Wash 5 buffers, and eluted in 35 μL base elution buffer (125 mM NaOH, 10 mM EDTA pH 8.0, 10 mM Tris-HCl pH 7.5) at 74°C for 5 min. The eluate was neutralized and purified with 1× Sera-Mag beads and eluted in 45 μL and stored at –80°C.

Determining the efficiency of cDNA pulldown

Pulldown efficiency was evaluated by qPCR to quantify enrichment of target transcripts and depletion of background RNA. qPCR reactions (KAPA SYBR 2× Master Mix) compared pre- and post-pulldown cDNA at a 2:1 ratio. Primers:

- L32 (background control): F 5′-AAGCGAAACTGGCGGAAAC-3′; R 5′-TAACCGATGTTG GGCATCAG-3′
- NF-κBIA (spliced exon 5–6 junction): F 5′-ACGGAGTCAGAATTCACAGAGG-3′; R 5′-CA CAAAGACAACAGCCGAATC-3′

Cycle-threshold (Ct) values were used to calculate ΔCt between L32 and NF-κBIA. Successful pulldowns typically showed ΔL32 = –7 to –9 cycles and ΔNF-κBIA = –2 to –4, corresponding to ΔL32/ΔNF-κBIA > 2.0.

Post-pulldown cDNA amplification

Pulldown cDNA was amplified prior to tagmentation. PCR reactions contained Q5 High-Fidelity 2× Master Mix (NEB), 1 μM primer (5′-AAGCAGTGGTATCAACGCAGAGTACT-3′),

and ~5% of the pulldown reaction. Cycling: 95°C (2 min) → 20–25 cycles of 95°C (30 s), 62.5°C (30 s), 72°C (150 s) → 72°C (5 min). PCR products were purified (0.9× Sera-Mag) and eluted in 25 μL H_2O. Concentrations were determined using a Qubit HS dsDNA Assay.

Tagmentation of cDNA libraries with Tn5 transposase

Tn5 transposase was purified as in Picelli et al., *Genome Res.* (2014) and pre-loaded with hybridized adapter oligos (Tn5MErev 5′-[phos]CTGTCTCTTATACACATCT-3′; Tn5ME-A 5′-T CGTCGGCAGCGTCAGATGTGTATAAGAGACAG-3′; Tn5ME-B 5′- GTCTCGTGGGCTCGGAGAT GTGTATAAGAGACAG-3′). Tagmentation reactions contained ~40 ng amplified cDNA, 0.2 μL Tn5, 5% PEG8000, 10 mM TAPS (pH 8.5), 5 mM $MgCl_2$, incubated 10 min at 55°C. SDS (0.02%) was added and incubation continued 10 min to inactivate Tn5. Reactions were 1.4× Sera-Mag purified and eluted in 20 μL H_2O.

Library barcoding and sequencing

Tagmented libraries were PCR-amplified with paired barcode oligos using Q5 High-Fidelity 2× Master Mix (NEB). Cycling: 72°C (3 min), 98°C (2 min), then 25 cycles of 98°C (10 s), 63°C (30 s), 72°C (30 s), and 72°C (5 min). Libraries were double-purified (1.0×/1.4× Sera-Mag), quantified by Qubit HS dsDNA assay, pooled equimolarly, repurified, and analyzed with an Agilent Bioanalyzer 2100. Sequencing was performed on an Illumina HiSeq 2500 (50 bp, single-end mode).

RNA-seq alignment and analysis

Single-end 50 bp reads were aligned to the mm10 genome using STAR. Junctions were retained only if both exonic and intronic segments were ≥3 bp. Using `pysam`, splice and intron junctions were classified as types a, b, or c, and Completion of Splicing Index (CoSI) values were computed using custom Python scripts. All analyses were performed in Python and R. Sequencing data are being deposited to NCBI GEO, and all analysis code will be made available on GitHub.

Deep Learning Analysis of Intron Regulatory Features

Model architecture and initialization

We used Borzoi, a transformer-based sequence-to-function model trained to predict steady-state RNA abundance from genomic DNA sequence. The pretrained model captures general patterns of cis-regulatory architecture by integrating multi-omic training targets (CAGE, ATAC-seq, and ChIP-seq tracks). However, because Borzoi is optimized for baseline expression rather than stimulus-responsive contexts, we adapted it to our macrophage TNFα time course through targeted fine-tuning.

Fine-tuning on TNFα-stimulated macrophage data

We generated genome-wide RNA-seq coverage tracks from bone marrow–derived macrophages (BMDMs) collected at 18, 20, and 30 minutes post-TNFα induction—timepoints representative of dynamic intron retention. These coverage values served as supervised training targets to refine the model's prediction of RNA abundance under inflammatory conditions. Fine-tuning was performed for 20 epochs using the Adam optimizer (learning rate = 1×10^{-6}, mean-squared-error loss). To evaluate generalization, chromosomes

10 and 11 were withheld for validation and testing, respectively. Model loss steadily decreased over training for both sets, and the correlation between predicted and observed expression improved from Pearson $r = 0.51$ ($p = 4.44\times10^{-16}$) for the pretrained model to $r = 0.61$ ($p = 4.84\times10^{-24}$) after fine-tuning (Fig. 5A–B).

Feature attribution and motif discovery

To interpret the fine-tuned model's predictions, we applied DeepLIFT to compute nucleotide–resolution importance scores, quantifying each base's contribution to the predicted RNA signal within a 524 kb genomic window centered on each intron of interest. Positive importance scores indicate sequence positions contributing to higher predicted intronic RNA signal (i.e., delayed splicing), whereas negative scores correspond to features predictive of splice completion.

Importance profiles were analyzed using modisco-lite, which clusters recurrent high-importance patterns across examples into position weight matrices (PWMs) representing putative regulatory motifs. Enrichment of discovered motifs was assessed using FIMO, scanning the 50 (or 150) slowest-splicing introns—ranked by area under the CoSI–time curve—against all annotated murine introns (GENCODE M23). For targeted motif discovery, attributions input to modisco-lite were masked under different configurations to restrict pattern identification to either intronic or exonic regions adjacent to slow-splicing introns.

Motif enrichment and visualization

Relative motif frequencies were plotted as a scatter of percent representation in slow- versus all-intron backgrounds. Sequence logos of the top five enriched motifs were visualized alongside genomic examples where GA-rich and A-rich motifs coincided with regions of high attribution in DeepLIFT profiles (Fig. 6A–D).

Acknowledgements

The authors would like to thank Alex Shishkin and Mitchell Guttman (Dept. of Biology, Caltech) for assistance with hybrid capture strategy design; and Ann-Jay Tong, Stephen Smale, Doug Black, and Amy-Pandya Jones (Dept. of Biology, University of California, Los Angeles) for insights and advice; and Sergei Manakov, Evelyn Stuwe, Dubravka Pezic, Igor Antoshechkin, Sagar Damle, and Alok Joglekar (Dept. of Biology, California Institute of Technology) for experimental and computational assistance. This work was funded from a grant from NIH and from an endowment provided by the Raymond and Beverly Sackler Foundation. Research reported in this publication was supported by the National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health under award number P20GM125498.

Author Contributions

J.S. Dearborn, Conceptualization, Investigation, Data curation, Software, Formal analysis, Visualization, Methodology, Writing – review & editing. L. Frankiw, Conceptualization, Investigation, Resources, Methodology, Validation, Writing – review & editing. D.W. Limoge, Data curation, Formal analysis, Visualization, Writing – review & editing. C.H. Burns, L. Vlach, P. Turpin, T. Kirch, Z.D. Miller, W. Dowell, S. Languon, Y. Garcia-Flores, Investigation,

Data curation, Validation, Writing – review & editing. R.C. Cockrell, Resources, Supervision, Project administration, Writing – review & editing. D. Baltimore, Conceptualization, Supervision, Funding acquisition, Project administration, Writing – review & editing. D. Majumdar, Conceptualization, Supervision, Funding acquisition, Project administration, Methodology, Writing – review & editing.

References

Ameur, A., Zaghlool, A., Halvardson, J., Wetterbom, A., Gyllensten, U., Cavelier, L., and Feuk, L. (2011). Total RNA sequencing reveals nascent transcription and widespread co-transcriptional splicing in the human brain. *Nature Structural & Molecular Biology*, 18(12):1435–1440.

Avsec, Ž., Agarwal, V., Visentin, D., Ledsam, J. R., Grabska-Barwinska, A., Taylor, K. R., Assael, Y., Jumper, J., Kohli, P., and Kelley, D. R. (2021). Effective gene expression prediction from sequence by integrating long-range interactions. *Nature Methods*, 18(10):1196–1203.

Bentley, D. L. (2014). Coupling mRNA processing with transcription in time and space. *Nature Reviews. Genetics*, 15(3):163–175.

Bergkessel, M., Whitworth, G. B., and Guthrie, C. (2011). Diverse environmental stresses elicit distinct responses at the level of pre-mRNA processing in yeast. *RNA (New York, N.Y.)*, 17(8):1461–1478.

Bhatt, D. M., Pandya-Jones, A., Tong, A.-J., Barozzi, I., Lissner, M. M., Natoli, G., Black, D. L., and Smale, S. T. (2012). Transcript dynamics of proinflammatory genes revealed by sequence analysis of subcellular RNA fractions. *Cell*, 150(2):279–290.

Braunschweig, U., Barbosa-Morais, N. L., Pan, Q., Nachman, E. N., Alipanahi, B., Gonatopoulos-Pournatzis, T., Frey, B., Irimia, M., and Blencowe, B. J. (2014). Widespread intron retention in mammals functionally tunes transcriptomes. *Genome Research*, 24(11):1774–1786.

Brugiolo, M., Herzel, L., and Neugebauer, K. M. (2013). Counting on co-transcriptional splicing. *F1000prime Reports*, 5:9.

Carrillo Oesterreich, F., Preibisch, S., and Neugebauer, K. M. (2010). Global analysis of nascent RNA reveals transcriptional pausing in terminal exons. *Molecular Cell*, 40(4):571–581.

Chen, J. and Chen, Z. J. (2013). Regulation of NF-κB by ubiquitination. *Current Opinion in Immunology*, 25(1):4–12.

Cho, V., Mei, Y., Sanny, A., Chan, S., Enders, A., Bertram, E. M., Tan, A., Goodnow, C. C., and Andrews, T. D. (2014). The RNA-binding protein hnRNPLL induces a T cell alternative splicing program delineated by differential intron retention in polyadenylated RNA. *Genome Biology*, 15(1):R26.

Davis-Turak, J. C., Allison, K., Shokhirev, M. N., Ponomarenko, P., Tsimring, L. S., Glass, C. K., Johnson, T. L., and Hoffmann, A. (2015). Considering the kinetics of mRNA synthesis in the analysis of the genome and epigenome reveals determinants of co-transcriptional splicing. *Nucleic Acids Research*, 43(2):699–707.

Eissa, N. T., Strauss, A. J., Haggerty, C. M., Choo, E. K., Chu, S. C., and Moss, J. (1996). Alternative splicing of human inducible nitric-oxide synthase mRNA. tissue-specific regulation and induction by cytokines. *The Journal of Biological Chemistry*, 271(43):27184–27187.

Engreitz, J. M., Pandya-Jones, A., McDonel, P., Shishkin, A., Sirokman, K., Surka, C., Kadri, S., Xing, J., Goren, A., Lander, E. S., Plath, K., and Guttman, M. (2013). The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. *Science (New York, N.Y.)*, 341(6147):1237973.

Freund, M., Hicks, M. J., Konermann, C., Otte, M., Hertel, K. J., and Schaal, H. (2005). Extended base pair complementarity between U1 snRNA and the 5′ splice site does not inhibit splicing in higher eukaryotes, but rather increases 5′ splice site recognition. *Nucleic Acids Research*, 33(16):5112–5119.

Gautier, E. L., Shay, T., Miller, J., Greter, M., Jakubzick, C., Ivanov, S., Helft, J., Chow, A., Elpek, K. G., Gordonov, S., Mazloom, A. R., Ma'ayan, A., Chua, W.-J., Hansen, T. H., Turley, S. J., Merad, M., Randolph, G. J., and Immunological Genome Consortium (2012). Gene-expression profiles and transcriptional regulatory pathways that underlie the identity and diversity of mouse tissue macrophages. *Nature Immunology*, 13(11):1118–1128.

Girard, C., Will, C. L., Peng, J., Makarov, E. M., Kastner, B., Lemm, I., Urlaub, H., Hartmuth, K., and Lührmann, R. (2012). Post-transcriptional spliceosomes are retained in nuclear speckles until splicing completion. *Nature Communications*, 3:994.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., and Regev, A. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, 29(7):644–652.

Grant, C. E., Bailey, T. L., and Noble, W. S. (2011). FIMO: Scanning for occurrences of a given motif. *Bioinformatics (Oxford, England)*, 27(7):1017–1018.

Guilgur, L. G., Prudêncio, P., Sobral, D., Liszekova, D., Rosa, A., and Martinho, R. G. (2014). Requirement for highly efficient pre-mRNA splicing during Drosophila early embryonic development. *eLife*, 3:e02181.

Han, J., Brown, T., and Beutler, B. (1990). Endotoxin-responsive sequences control cachectin/tumor necrosis factor biosynthesis at the translational level. *The Journal of Experimental Medicine*, 171(2):465–475.

Hao, S. and Baltimore, D. (2009). The stability of mRNA influences the temporal order of the induction of genes encoding inflammatory molecules. *Nature Immunology*, 10(3):281–288.

Hao, S. and Baltimore, D. (2013). RNA splicing regulates the temporal order of TNF-induced gene expression. *Proceedings of the National Academy of Sciences of the United States of America*, 110(29):11934–11939.

Hargreaves, D. C., Horng, T., and Medzhitov, R. (2009). Control of inducible gene expression by signal-dependent transcriptional elongation. *Cell*, 138(1):129–145.

Honda, K., Yanai, H., Negishi, H., Asagiri, M., Sato, M., Mizutani, T., Shimada, N., Ohba, Y., Takaoka, A., Yoshida, N., and Taniguchi, T. (2005). IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature*, 434(7034):772–777.

Jaganathan, K., Kyriazopoulou Panagiotopoulou, S., McRae, J. F., Darbandi, S. F., Knowles, D., Li, Y. I., Kosmicki, J. A., Arbelaez, J., Cui, W., Schwartz, G. B., Chow, E. D., Kanterakis, E., Gao, H., Kia, A., Batzoglou, S., Sanders, S. J., and Farh, K. K.-H. (2019). Predicting Splicing from Primary Sequence with Deep Learning. *Cell*, 176(3):535–548.e24.

Khodor, Y. L., Rodriguez, J., Abruzzi, K. C., Tang, C.-H. A., Marr, M. T., and Rosbash, M. (2011). Nascent-seq indicates widespread cotranscriptional pre-mRNA splicing in Drosophila. *Genes & Development*, 25(23):2502–2512.

Koh, C. M., Bezzi, M., Low, D. H. P., Ang, W. X., Teo, S. X., Gay, F. P. H., Al-Haddawi, M., Tan, S. Y., Osato, M., Sabò, A., Amati, B., Wee, K. B., and Guccione, E. (2015). MYC regulates the core pre-mRNA splicing machinery as an essential step in lymphomagenesis. *Nature*, 523(7558):96–100.

Kontoyiannis, D., Pasparakis, M., Pizarro, T. T., Cominelli, F., and Kollias, G. (1999). Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: Implications for joint and gut-associated immunopathologies. *Immunity*, 10(3):387–398.

Koodathingal, P., Novak, T., Piccirilli, J. A., and Staley, J. P. (2010). The DEAH box ATPases Prp16 and Prp43 cooperate to proofread 5′ splice site cleavage during pre-mRNA splicing. *Molecular Cell*, 39(3):385–395.

Lei, X. F., Ohkawara, Y., Stämpfli, M. R., Mastruzzo, C., Marr, R. A., Snider, D., Xing, Z., and Jordana, M. (1998). Disruption of antigen-induced inflammatory responses in CD40 ligand knockout mice. *The Journal of Clinical Investigation*, 101(6):1342–1353.

Leppek, K., Schott, J., Reitter, S., Poetz, F., Hammond, M. C., and Stoecklin, G. (2013). Roquin promotes constitutive mRNA decay via a conserved class of stem-loop recognition motifs. *Cell*, 153(4):869–881.

Linder, J., Srivastava, D., Yuan, H., Agarwal, V., and Kelley, D. R. (2025). Predicting RNA-seq coverage from DNA sequence as a unifying model of gene regulation. *Nature Genetics*, 57(4):949–961.

Martin, R. M., Rino, J., Carvalho, C., Kirchhausen, T., and Carmo-Fonseca, M. (2013). Live-cell visualization of pre-mRNA splicing with single-molecule sensitivity. *Cell Reports*, 4(6):1144–1155.

Michaelson, J. S., Bader, D., Kuo, F., Kozak, C., and Leder, P. (1999). Loss of Daxx, a promiscuously interacting protein, results in extensive apoptosis in early mouse development. *Genes & Development*, 13(15):1918–1923.

Mukherji, S., Ebert, M. S., Zheng, G. X. Y., Tsang, J. S., Sharp, P. A., and van Oudenaarden, A. (2011). MicroRNAs can generate thresholds in target gene expression. *Nature Genetics*, 43(9):854–859.

O'Connell, R. M., Rao, D. S., and Baltimore, D. (2012). microRNA regulation of inflammatory responses. *Annual Review of Immunology*, 30:295–312.

Pandya-Jones, A., Bhatt, D. M., Lin, C.-H., Tong, A.-J., Smale, S. T., and Black, D. L. (2013). Splicing kinetics and transcript release from the chromatin compartment limit the rate of Lipid A-induced gene expression. *RNA (New York, N.Y.)*, 19(6):811–827.

Patel, A. A., McCarthy, M., and Steitz, J. A. (2002). The splicing of U12-type introns can be a rate-limiting step in gene expression. *The EMBO journal*, 21(14):3804–3815.

Pessa, H. K. J., Ruokolainen, A., and Frilander, M. J. (2006). The abundance of the spliceosomal snRNPs is not limiting the splicing of U12-type introns. *RNA (New York, N.Y.)*, 12(10):1883–1892.

Rabani, M., Levin, J. Z., Fan, L., Adiconis, X., Raychowdhury, R., Garber, M., Gnirke, A., Nusbaum, C., Hacohen, N., Friedman, N., Amit, I., and Regev, A. (2011). Metabolic labeling of RNA uncovers principles of RNA production and degradation dynamics in mammalian cells. *Nature Biotechnology*, 29(5):436–442.

Rabani, M., Raychowdhury, R., Jovanovic, M., Rooney, M., Stumpo, D. J., Pauli, A., Hacohen, N., Schier, A. F., Blackshear, P. J., Friedman, N., Amit, I., and Regev, A. (2014). High-resolution sequencing and modeling identifies distinct dynamic RNA regulatory strategies. *Cell*, 159(7):1698–1710.

Ramirez-Carrozzi, V. R., Braas, D., Bhatt, D. M., Cheng, C. S., Hong, C., Doty, K. R., Black, J. C., Hoffmann, A., Carey, M., and Smale, S. T. (2009). A unifying model for the selective regulation of inducible transcription by CpG islands and nucleosome remodeling. *Cell*, 138(1):114–128.

Shrikumar, A., Greenside, P., and Kundaje, A. (2019). Learning Important Features Through Propagating Activation Differences.

Shrikumar, A., Tian, K., Avsec, Ž., Shcherbina, A., Banerjee, A., Sharmin, M., Nair, S., and Kundaje, A. (2020). Technical Note on Transcription Factor Motif Discovery from Importance Scores (TF-MoDISco) version 0.5.6.5.

Smale, S. T., Tarakhovsky, A., and Natoli, G. (2014). Chromatin contributions to the regulation of innate immunity. *Annual Review of Immunology*, 32:489–511.

Takashima, Y., Ohtsuka, T., González, A., Miyachi, H., and Kageyama, R. (2011). Intronic delay is essential for oscillatory expression in the segmentation clock. *Proceedings of the National Academy of Sciences of the United States of America*, 108(8):3300–3305.

Tilgner, H., Knowles, D. G., Johnson, R., Davis, C. A., Chakrabortty, S., Djebali, S., Curado, J., Snyder, M., Gingeras, T. R., and Guigó, R. (2012). Deep sequencing of subcellular RNA fractions shows splicing to be predominantly co-transcriptional in the human genome but inefficient for lncRNAs. *Genome Research*, 22(9):1616–1625.

Wan, F., Anderson, D. E., Barnitz, R. A., Snow, A., Bidere, N., Zheng, L., Hegde, V., Lam, L. T., Staudt, L. M., Levens, D., Deutsch, W. A., and Lenardo, M. J. (2007). Ribosomal protein S3: A KH domain subunit in NF-kappaB complexes that mediates selective gene regulation. *Cell*, 131(5):927–939.

Yeo, G. and Burge, C. B. (2004). Maximum entropy modeling of short sequence motifs with applications to RNA splicing signals. *Journal of Computational Biology: A Journal of Computational Molecular Cell Biology*, 11(2-3):377–394.

Yuan, H., Linder, J., and Kelley, D. R. (2025). Parameter-Efficient Fine-Tuning of a Supervised Regulatory Sequence Model.

Zhou, J. and Troyanskaya, O. G. (2015). Predicting effects of noncoding variants with deep learning-based sequence model. *Nature Methods*, 12(10):931–934.

Figure Legends

Figure 1. Sequencing of complete, chromatin-associated pre-mRNA during inflammatory stimulus reveals differential splicing dynamics among introns of IKBα. (A) Histogram of reads corresponding to the TNF-induced expression and splicing of IKBα pre-mRNA of BMDMs. RNA-seq was performed on chromatin-associated RNA, enriched for NFκB genes as a function of a TNF stimulation time course, time shown in minutes after stimulation. Reads are histogrammed in $\log_{10}$ scale and normalized to each time point's maximum value. (B) The Coefficient of Splicing (CoSI) metric quantifies extent of splicing as a function of time, expressed as a ratio of reads from each splice junction to total junctional reads. Dynamics of IKBα splicing as a function of each intron's CoSI is shown (C), where 1 = spliced and 0 = unspliced, with corresponding introns highlighted in a sample time point. (D) Differential dynamics of splicing for each *Nfkbia* intron are further demonstrated in the coverage plot for the transcript.

Figure 2. Heterogeneity of splicing at each intron reveals splicing "bottlenecks". The CoSI of each intron per time point is shown as a function of the entire inflammatory mRNA dataset as a box–whisker plot (A). Each point represents an intron of one of 230 genes, revealing high rates of splicing (median CoSI indicated by bar near 1.0 for each time point) for most genes with significant outliers. As an example, *Cxcl10* intron 2 (red arrowhead) is represented by the datapoint with arrowhead, and a histogram of reads is shown to demonstrate the relatively unspliced nature of this intron, which is not involved in alternative splicing. (B) Several similar introns that are relatively unspliced are found throughout the inflammatory transcriptome; shown are bottleneck introns within *Cd40*, *Daxx*, and *Irf7* as examples in the context of their neighboring introns.

Figure 3. Splicing kinetics of inflammatory introns are heterogeneous, ranging from seconds to minutes. CoSI of introns representing various splicing rates are measured and fit to half-lives. Cells were treated with Actinomycin D, from which hybrid capture of genes of interest and sequencing was performed on total (unfractionated) RNA. Shown are four representative samples of splicing kinetics.

Figure 4. Bottleneck introns can be repaired and account for significant alterations to gene expression. (A) Intron–GFP splicing reporters for each wild-type intron (red) and modified intron (green) are shown as BFP:GFP ratio. (B) Ratio of WT:fixed slopes is shown; whereas *Tfec* expression is not altered by improved 5′ sequence, *Malt1* intron sequence is significantly impaired owing to its 5′ donor sequence, exhibiting a roughly five-fold impairment in gene expression due to the 5′ splice donor.

Figure 5. Fine-tuning improves gene expression prediction in macrophages. (A) Training and test loss curves over 20 epochs of model fine-tuning. (B–C) Predicted versus measured RNA expression values for 230 transcripts induced by TNFα in macrophages, shown for the pretrained model (B) and the fine-tuned model (C).

Figure 6. Putative regulatory sequences are enriched in slow-splicing introns. (A) Scatterplot showing percent representation of position weight matrices (PWMs) scanned across slow-splicing introns versus all introns genome-wide using FIMO. Sequence logos of the top five enriched PWMs are shown to the right. (B–D) Attribution plots highlighting GA-rich sequences (source seqlets for enriched PWMs) and their locations mapped to gene schematics for GA-rich motif (B) and A-rich motif (C). Corresponding RNA-seq tracks are shown for each gene.

Supplementary Figure Legends

Supplementary Figure 1. Hybrid capture strategy for isolating chromatin-associated inflammatory transcripts. RNA was purified from chromatin-associated bone marrow–derived macrophages (BMDMs), and cDNA was generated using oligo(dT) priming to enrich for polyadenylated transcripts. Biotinylated RNA oligonucleotides complementary to the terminal exons of inflammatory genes were hybridized to the cDNAs, allowing for selective enrichment of these transcripts via streptavidin bead capture.

Supplementary Figure 2. Hybrid capture enriches NF-κB–responsive transcripts and yields robust intron coverage. (A) Fraction of sequencing reads corresponding to NF-κB–responsive genes (blue) in hybrid-captured versus poly(A)-selected RNA (orange), showing substantial enrichment after hybrid capture. (B) Histogram of read counts per intron at the 6-min TNF induction time point. Reads were detected for 1,024 introns out of 1,508 targeted introns, with undetected introns largely corresponding to transcripts induced at later time points (>60 min). (C) Distribution of intron fold induction across time points reveals that many NF-κB target genes begin to show induction as early as 4 min post-TNF stimulation, with both the number of induced introns and magnitude of induction increasing markedly by 14 min. (D) Scatterplot comparing induction at 4 min versus 14 min for individual introns demonstrates that most early-induced genes are further upregulated at later time points.

Supplementary Figure 3. Gene track visualization of *Nfkbia* induction dynamics. Genome browser tracks show *Nfkbia* (IKBα) transcript induction over time in chromatin-associated, hybrid-captured RNA. Signal intensity is displayed on a linear scale (left) normalized to the maximum height at 20 min, and on a log scale (right) normalized to the maximum at each time point.

Supplementary Figure 4. Splice site strength and expression kinetics among NF-κB–induced genes. (A) Representative introns from selected genes were scored for 5′ splice donor strength based on similarity to the canonical "GTAAG" motif; introns with weaker matches (e.g., *Irf7* intron 3) received lower scores. (B) NF-κB–induced genes display well-characterized variability in expression kinetics (RNA-seq data from Reference 25). Heatmap shows temporal expression profiles of NF-κB target genes following lipid A stimulation, categorized into immediate-early, early, and later expression groups. (C) All introns in the NF-κB transcriptome were scored using MaxEntScan for both 5′ splice donor (top) and 3′ splice acceptor (bottom) sequences, stratified by their gene expression group.

Supplementary Figure 5. Reporter assay design for assessing intron splicing efficiency. (A) Schematic of the bidirectional reporter assay. Individual introns were cloned into a bidirectional promoter context together with their flanking exons, positioned upstream of a self-cleaving 2A peptide and GFP reporter. In the opposite transcriptional direction, a BFP reporter served as a transcriptional control. (B) Evolutionarily conserved weak 5′ splice donors were "repaired" to the canonical GTAAG sequence within this reporter construct, and splicing efficiency was quantified by flow cytometry (FACS).

Supplementary Figure 6. Splicing completion across kinetic gene expression groups. Box-and-whisker plots depict time-course CoSI values for introns from inflammatory gene cohorts. Each point represents the CoSI value of an individual intron at a given time point. Plots are grouped by expression category: Immediate Early (top left), Early (top right), Intermediate (bottom left), and Late (bottom right).

Supplementary Figure 7. Relationships between intron length, GC content, transcript

position, and splicing kinetics. (A) Scatterplot of intron length (y-axis) versus GC content (%) (x-axis) for the 150 slowest-splicing introns. Pearson $r = -0.44$ ($p = 2.76\times10^{-45}$) and Spearman $r = -0.44$ ($p = 1.34\times10^{-46}$). (B) Histogram of intron length distributions (100-nt bins) for all 1,098 introns (red) and the 150 slowest-splicing introns (blue). Length distributions are largely similar between cohorts. (C) Splice completion (CoSI) trajectories across the time course for the 50 fastest (left, blue) and 50 slowest (right, orange) introns, ranked by area under the CoSI curve (AUC). (D) Same analysis as in (C) but extended to the top 150 fastest and bottom 150 slowest introns. (E) Minimum CoSI values for each intron plotted against transcript abundance (TPM, left), intron length (middle), and position along the transcript (5′ → 3′ %, right) for the bottom 150 and bottom 50 introns. Correlations: TPM—Bottom 150: Pearson $r = 0.204$ ($p = 1.24\times10^{-2}$), Spearman $r = 0.203$ ($p = 1.27\times10^{-2}$); Bottom 50: Pearson $r = 0.276$ ($p = 5.25\times10^{-2}$), Spearman $r = 0.316$ ($p = 2.53\times10^{-2}$). Length—Bottom 150: Pearson $r = -0.242$ ($p = 2.88\times10^{-3}$), Spearman $r = -0.266$ ($p = 9.92\times10^{-4}$); Bottom 50: Pearson $r = -0.160$ ($p = 0.268$), Spearman $r = -0.171$ ($p = 0.235$). Transcript position—Bottom 150: Pearson $r = 0.260$ ($p = 1.33\times10^{-3}$), Spearman $r = 0.231$ ($p = 4.44\times10^{-3}$); Bottom 50: Pearson $r = 0.259$ ($p = 6.90\times10^{-2}$), Spearman $r = 0.318$ ($p = 2.43\times10^{-2}$).

Supplementary Figure 8. Enrichment of sequence motifs near slow-splicing introns. Scatterplots show the percent representation of position weight matrices (PWMs) scanned across slow-splicing introns compared to all introns genome-wide using FIMO. Sequence logos of the top five enriched PWMs are displayed to the right of each scatterplot. (A) Scatterplot and enriched motifs identified in exons adjacent to the 50 slowest-splicing introns. (B) Scatterplot and enriched motifs identified within the 150 slowest-splicing introns. (C) Scatterplot and enriched motifs identified in exons adjacent to the 150 slowest-splicing introns.

Supplementary Table Legends

Supplementary Table 1. Oligonucleotide sequences used
Supplementary Table 2. Exons of interest for hybrid capture

Figures

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Supplementary Figures

Isolation of Chromatin-Associated RNA

cDNA generation (dT-primed)

Macrophage

cDNA Enrichment for 230 *full length* Inflammatory Genes

| Exon 1 | Exon 2 | Exon 3 | Last Exon |

Biotinylated probes tiled to last exon

Enrichment with Streptavidin Immobilization

Supplementary Figure 1.

Supplementary Figure 2.

Supplementary Figure 3.

Supplementary Figure 4.

Supplementary Figure 5.

Supplementary Figure 6.

Supplementary Figure 7.

Supplementary Figure 8.

Supplementary Tables

Supplementary Table 1. Oligonucleotides used in this study.

Oligo	Sequence (5′–3′)	Use
d14-952	AAGCAGTGGTATCAACGCAGAGTACACArGrGrG	Template-switch oligo
d14-954	AAGCAGTGGTATCAACGCAGAGTACT(30)	RT dT oligo
d14-955	AAGCAGTGGTATCAACGCAGAGTAC	Post-pulldown amplification oligo
L32-For	AAGCGAAACTGGCGGAAAC	qPCR primer
L32-Rev	TAACCGATGTTGGGCATCAG	qPCR primer
NF-KBIA Ex5–6-For	ACGGAGTCAGAATTCACAGAGG	qPCR primer
NF-KBIA Ex5–6-Rev	CACAAAGACAACAGCCGAATC	qPCR primer
Tn5MErev	[phos]CTGTCTCTTATACACATCT	Tn5 adaptor oligo
Tn5ME-A	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG	Tn5 adaptor oligo

Table 2. Supplementary Table 2. Exons targeted for hybrid capture. Exons are grouped by probe pool (A–I). Coordinates are given relative to the mm10 genome assembly.

Pool	Gene	Chr	Start	End	Strand
A	*Tnf*	chr17	35336336	35337508	–
A	*Ccl2*	chr11	81850428	81850954	+
A	*Cxcl10*	chr5	92775665	92776411	–
A	*Irg1*	chr14	103453729	103455790	+
A	*Cxcl2*	chr5	91334215	91334964	+
A	*Rsad2*	chr12	27127608	27130435	–
B	*Nfkbia*	chr12	56590396	56590956	–
B	*Ccl4*	chr11	83477796	83478185	+
B	*Tnfaip2*	chr12	112691632	112693229	+
B	*Rasgef1b*	chr5	99646439	99647817	–
B	*Ccl7*	chr11	81860487	81861025	+
B	*Tnfaip3*	chr10	18720717	18722821	–
B	*Clec4e*	chr6	123231807	123233714	–
B	*Mx1*	chr16	97668644	97669979	–
C	*Plek*	chr11	16871209	16874156	–
C	*Rgs1*	chr1	146091799	146092617	–
C	*Icam1*	chr9	20832215	20833240	+
C	*Tnfsf9*	chr17	57246464	57247180	+
C	*Sdc4*	chr2	164249747	164251737	–
C	*Atf3*	chr1	192994176	192995558	–
C	*Mapkapk2*	chr1	132950281	132951812	–
C	*Ccl5*	chr11	83339281	83339566	–

Pool	Gene	Chr	Start	End	Strand
C	*Nfkbiz*	chr16	55811490	55812896	–
C	*Cmpk2*	chr12	27162877	27164702	+
D	*Irf1*	chr11	53590874	53591876	+
D	*Dusp1*	chr17	26642536	26643595	–
D	*Zfp36*	chr7	29161803	29163512	–
D	*Il1b*	chr2	129190316	129190974	–
D	*Cd83*	chr13	43896926	43898501	+
D	*Plk2*	chr13	111190241	111191051	+
D	*Nlrp3*	chr11	59379666	59380458	+
D	*Cd44*	chr2	102651300	102654494	–
D	*Ccrl2*	chr9	110957339	110958943	–
D	*Helz2*	chr2	180962320	180963574	–
D	*Tnip3*	chr6	65581071	65584034	+
D	*Parp14*	chr16	35832964	35835021	–
D	*Tlr2*	chr3	83640194	83642711	–
D	*Gbp5*	chr3	142184004	142185308	+
D	*Kctd12*	chr14	103375798	103381854	–
D	*Ifrd1*	chr12	40929716	40930038	–
D	*Ppp1r15a*	chr7	52778288	52778692	–
D	*Cxcl1*	chr5	91321534	91322141	+
D	*Fam20c*	chr5	139285062	139286017	+
D	*Clic4*	chr4	134769885	134773180	–
D	*Mx2*	chr16	97781946	97782506	+
E	*Igsf6*	chr7	128207581	128209031	–

Pool	Gene	Chr	Start	End	Strand
E	*Ifih1*	chr2	62433850	62436145	–
E	*Ehd1*	chr19	6298074	6300096	+
E	*Ifit1*	chr19	34721961	34724499	+
E	*Socs3*	chr11	117827401	117829632	–
E	*Peli1*	chr11	21047960	21050330	+
E	*Gdf15*	chr8	73153293	73154073	–
E	*Junb*	chr8	87500808	87502647	–
E	*Icosl*	chr10	77540638	77542270	+
E	*Bcl2l11*	chr2	127983967	127988283	+
E	*Cd274*	chr19	29459880	29462584	+
E	*Cd40*	chr2	164896222	164897154	+
E	*Ccrn4l*	chr3	51053619	51055576	+
E	*Il27*	chr7	133732809	133733063	–
E	*Irf7*	chr7	148449082	148449279	–
E	*Insig1*	chr5	28403331	28405202	+
E	*Malt1*	chr18	65635642	65638446	+
E	*Btg2*	chr1	135971442	135974280	–
E	*Cd69*	chr6	129217343	129218387	–
E	*Sod2*	chr17	13200705	13200934	+
E	*Nfkbid*	chr7	31213153	31213765	+
E	*Rnf19b*	chr4	128760972	128761770	+
E	*Gbp2*	chr3	142300347	142300972	+
E	*Ier3*	chr17	35958983	35959856	+
E	*Skil*	chr3	31017377	31021845	+
E	*Zc3h12c*	chr9	51976136	51976216	–

Pool	Gene	Chr	Start	End	Strand
E	*Egr1*	chr18	35022119	35024610	+
E	*Tfec*	chr6	16783381	16784330	−
E	*Cers6*	chr2	68946499	68949347	+
E	*Nfkbie*	chr17	45698951	45700117	+
E	*Ccl12*	chr11	81916612	81916901	+
E	*Daxx*	chr17	34052288	34052535	+
E	*Plau*	chr14	21661496	21662610	+
E	*Srgn*	chr10	61957176	61957854	−
E	*Il12b*	chr11	44226726	44227519	+
F	*Arl5c*	chr11	97850894	97851653	−
F	*Itgav*	chr2	83643397	83647073	+
F	*Ptgs2*	chr1	151952503	151955142	+
F	*Pde4b*	chr4	102277579	102279867	+
F	*Casp4*	chr9	5336535	5336791	+
F	*Csf1*	chr3	107543966	107546063	−
F	*Ifit2*	chr19	34647557	34651024	+
F	*Sgms1*	chr19	32197217	32199014	−
F	*Gbp3*	chr3	142235438	142236176	+
F	*Ifi47*	chr11	48908891	48910489	+
F	*C3*	chr17	57343396	57343577	−
F	*Il1rn*	chr2	24204935	24207011	+
F	*Dusp4*	chr8	35881452	35882948	+
F	*Mmp13*	chr9	7282018	7283333	+
F	*Dusp2*	chr2	127163317	127164113	+
F	*Ell2*	chr13	75908128	75909806	+

Pool	Gene	Chr	Start	End	Strand
F	Jdp2	chr12	86979723	86980828	+
F	Fnbp1l	chr3	122241637	122245201	−
F	Ripk2	chr4	16050522	16051038	−
F	Marcksl1	chr4	129191961	129193225	+
F	Slc16a10	chr10	39753341	39757210	−
F	Traf1	chr2	34798778	34799611	−
F	Fabp4	chr3	10204343	10204570	−
F	Trim21	chr7	109706436	109708177	−
F	Jak2	chr19	29386231	29387570	+
F	Errfi1	chr4	150240425	150242989	+
F	Csrnp1	chr9	119880284	119882329	−
F	Orai2	chr5	136623332	136626822	−
F	Rnf19a	chr15	36169689	36171820	−
F	Fos	chr12	86816767	86818219	+
F	Oasl1	chr5	115386927	115387920	+
F	Icam4	chr9	20834826	20834975	+
F	Gm6377	chrX	106393115	106395784	+
F	Rab20	chr8	11453977	11454526	−
F	Isg15	chr4	155573533	155574175	−
F	Plaur	chr7	25260258	25260892	+
F	Klf7	chr1	64082247	64082338	−
F	Ifi204	chr1	175677425	175677751	−
F	Id3	chr4	135700795	135701307	+
F	AW112010	chr19	11122102	11122583	−
G	Ets2	chr16	95940530	95942656	+

Pool	Gene	Chr	Start	End	Strand
G	Nos2	chr11	78773156	78773626	+
G	Ier2	chr8	87185230	87186751	–
G	Gpr85	chr6	13785074	13787073	–
G	Tgm2	chr2	157942141	157943687	–
G	Igtp	chr11	58019546	58021094	+
G	Slc44a1	chr4	53562861	53563036	+
G	Snx18	chr13	114382387	114385070	–
G	Stx11	chr10	12659787	12661787	–
G	Nod2	chr8	91211459	91212373	+
G	Saa3	chr7	53967368	53967634	–
G	Herc6	chr6	57612694	57615130	+
G	Usp18	chr6	121220490	121220935	+
G	Egr2	chr10	67002493	67004936	+
G	Il6	chr5	30345921	30346508	+
G	Fchsd2	chr7	108430964	108432919	+
G	Med21	chr6	146598633	146599122	+
G	Gm12250	chr11	58001241	58003700	+
G	Pyhin1	chr1	175576122	175578059	+
G	Fam26f	chr10	33845873	33846372	–
G	Fam108c	chr7	91257866	91259311	–
G	Fgl2	chr5	20881072	20884204	+
G	Tagap	chr17	8125748	8127762	+
G	Tagap	chr17_random	39677	41691	–
G	Nfkbib	chr7	29551263	29551543	–
G	Il1a	chr2	129125346	129126635	–

Pool	Gene	Chr	Start	End	Strand
G	Ifit3	chr19	34661551	34663472	+
G	Slc28a2	chr2	122286156	122286866	+
G	G530011O06Rik	chrX	166412975	166414722	–
G	Maff	chr15	79187851	79189506	+
G	Il4i1	chr7	52094953	52096179	+
G	Iigp1	chr18	60549446	60552283	+
G	Gbp7	chr3	142209284	142213047	+
G	Ifnb1	chr4	88167929	88168698	–
G	Slc7a11	chr3	50168858	50176288	–
G	Samsn1	chr16	75859039	75859884	–
G	Slco3a1	chr7	81420304	81420974	–
G	Kdm6b	chr11	69212020	69213351	–
G	Hivep2	chr10	13868718	13871184	+
G	Plscr1	chr9	92166367	92167399	+
G	Nod1	chr6	54873936	54875003	–
G	Serpine1	chr5	137537376	137539081	–
G	Fpr2	chr17	18029697	18030916	+
G	Prdm1	chr10	44156981	44160048	–
G	Rab11fip1	chr8	28249245	28253866	–
G	Phldb1	chr9	44494391	44495781	–
G	Gpr84	chr15	103138666	103140087	–
G	Niacr1	chr5	124313579	124315525	–
G	Zswim4	chr8	86734841	86736704	–
G	Cxcl9	chr5	92750357	92752932	–
G	Tma16	chr8	69000245	69000808	–

Pool	Gene	Chr	Start	End	Strand
G	*Myc*	chr15	61820901	61821916	+
G	*Tnfsf15*	chr4	63385637	63391077	–
G	*Dusp5*	chr19	53615406	53615812	+
G	*Zbp1*	chr2	173032114	173032727	–
G	*Osgin2*	chr4	15924268	15926147	–
G	*Gem*	chr4	11640623	11642140	+
G	*Cxcl11*	chr5	92788571	92789832	–
G	*Trib1*	chr15	59485791	59488654	+
G	*Olfr56*	chr11	48947790	48948889	+
G	*Ch25h*	chr19	34548274	34549625	–
G	*Batf2*	chr19	6171306	6172475	+
G	*Il15ra*	chr2	11654777	11655612	+
G	*Agpat4*	chr17	12411851	12412506	+
G	*Gm4951*	chr18	60405029	60407474	+
G	*Ifi205*	chr1	175942129	175942419	–
G	*Rcan1*	chr16	92392198	92393873	–
G	*Vcan*	chr13	89794915	89796963	–
G	*Agrn*	chr4	155539399	155540762	–
H	*Cfb*	chr17	34993319	34993527	–
H	*Foxp4*	chr17	48004082	48005752	–
H	*Cish*	chr9	107202699	107204292	+
H	*Tnfsf10*	chr3	27234144	27238587	+
H	*Flnb*	chr14	8782819	8784101	+
H	*Phf11d*	chr14	59966244	59967888	–
H	*Ifi44*	chr3	151393887	151395517	–

Continued on next page

Pool	Gene	Chr	Start	End	Strand
H	*Gbp9*	chr5	105507413	105509283	–
H	*Edn1*	chr13	42402146	42403358	+
H	*Frmd6*	chr12	72000371	72003221	+
H	*Isg20*	chr7	86064932	86065282	+
H	*AA467197*	chr2	122466437	122466812	+
H	*Klhl25*	chr7	83017584	83019016	+
H	*Dhx40*	chr11	86582351	86583763	–
H	*Pmaip1*	chr18	66622924	66625212	+
H	*Zhx2*	chr15	57670146	57671387	+
H	*Gm14446*	chr19	34667378	34669513	–
H	*Irs2*	chr8	10986964	10987014	–
H	*Dpep3*	chr8	108497420	108497751	–
H	*Phlda1*	chr10	110945116	110945705	+
H	*Fmnl2*	chr2	52990874	52993236	+
H	*Slfn1*	chr11	82934523	82936161	+
H	*Ms4a4c*	chr19	11500686	11501736	+
H	*Gbp4*	chr5	105544786	105547518	–
H	*Cp*	chr3	19892509	19892998	+
H	*Slfn4*	chr11	83002165	83003718	+
H	*Phf11c*	chr14	59999670	59999984	–
H	*Snn*	chr16	11072325	11075078	+
H	*Gnb4*	chr3	32482450	32484148	–
H	*Ccnd2*	chr6	127075727	127080505	–
H	*Egr3*	chr14	70478703	70479964	+
I	*Extl2*	chr3	115729931	115731934	+

Pool	Gene	Chr	Start	End	Strand
I	*Nr4a1*	chr15	101104427	101105225	+
I	*Gm11435*	chr11	83325238	83325629	–
I	*Phf11b*	chr14	59939801	59940120	–
I	*Odc1*	chr12	17557413	17558308	+
I	*Dusp8*	chr7	149265392	149268935	–
I	*Gm10941*	chr10	76720518	76720731	–
I	*Adamts1*	chr16	85794073	85796345	–
I	*Gm889*	chr12	40902973	40903815	–