Programmed Delayed Splicing: A Mechanism for Timed Inflammatory Gene Expression

Jacob S. Dearborn[1], Luke Frankiw[2], Damas W. Limoge[3], Christian H. Burns[2], Logan Vlach[2], Patricia Turpin[2], Tylar Kirch[1], Zachary D. Miller[1], William Dowell[1], Sylvester Languon[1], Yvette Garcia-Flores[1], Robert C. Cockrell[1], David Baltimore[1], Devdoot Majumdar*[1,2]

[1]Dept of Surgical Research, Larner College of Medicine, University of Vermont; [2]Division of Biology, California Institute of Technology; [3]Nanotronics (Brooklyn, New York),

*Correspondence to dev.majumdar@uvm.edu

SUMMARY:

Inflammation involves timed gene expression, suggesting that the fine-tuned onset, amplitude, and termination of expression of hundreds of genes is of critical importance to organismal homeostasis. Recent study of post-transcriptional regulation of inflammatory gene expression led to the suggestion of a regulatory role for pre-mRNA splicing. Here, using a hybrid capture approach to purify incompletely spliced, chromatin-associated pre-mRNAs, we use deep sequencing to study pre-mRNA splicing of the NF-kB transcriptome. By freezing transcription and examining subsequent splicing of complete transcripts, we find many introns splice tens to hundreds of times slower than average. Investigating the basis of these delays, we focused on evolutionarily conserved introns with suboptimal splice donor sequences and found that strengthening these donor sites by as few as two nucleotides in minigene reporter assays markedly increased gene expression for several targets. This suggests that such sites can act as timing elements that both delay mRNA production and limit expression amplitude. To broaden this mechanistic view, we applied deep learning sequence-to-function models with feature attribution to identify additional regulatory sequences—both intronic and exonic—that may contribute to delayed splicing through mechanisms independent of donor site strength. This integrated approach revealed non-canonical motifs enriched in slow-splicing introns, pointing to a broader repertoire of cis-elements that can fine-tune transcript maturation during inflammation. Together, these findings support a model in which the temporal regulation of pre-mRNA splicing serves as a layer of control in inflammatory gene expression, and raise the possibility that similar timing mechanisms operate in other rapid-response transcriptional programs.

KEYWORDS: INFLAMMATION, INNATE IMMUNITY, MACROPHAGE, GENE EXPRESSION, SPLICING

INTRODUCTION

Gene expression in response to an inflammatory stimulus begins rapidly and is tightly controlled by conventional means (transcription and protein turnover (1–3)) and by an expanding list of modalities that have gained in appreciation as being general regulatory strategies (RNA stabilization, RNA deadenylation, ribosomal regulation, microRNA regulation, as examples (4–7)). We and others have recently investigated the role of RNA splicing kinetics – independently of alternative splicing – in gene expression (8–11). In macrophages, an inflammatory stimulus leads to upregulation of expression of pre-mRNAs from hundreds of genes, providing an experimentally favorable system to investigate whether differential kinetics of pre-mRNA splicing may control the timing of gene expression following an inducing stimulus.

Pre-mRNA conversion to mRNA has been implicated in regulation of gene expression in diverse systems. As part of the cellular response to various environmental stressors, mRNAs for ribosomal proteins were shown to be downregulated due to decreased splicing efficiency in yeast (12). Global changes in efficiency of pre-mRNA splicing have been shown to be a developmental prerequisite for *Drosophila* early embryonic development (13). The developing vertebrate embryo obeys a 'segmentation clock' determining body segment length whose very timing relies on delays attributable to control of the splicing rate of the *Hes7* transcriptional repressor (14).

In certain well-studied cases, as with the cytokine TNFα, regulatory mechanisms modulating RNA levels exert significant physiological effects (15–22). The insight that TNFα contains AU-rich elements in its 3' untranslated region that act as mRNA degradation signals (23), and subsequent observations that a mouse in which these AU-rich elements were removed resulted in a robust autoimmune phenotype (17), was an early indication of the importance of precisely tuned mRNA levels in the regulation of inflammation to avoid autoimmunity. Given the role of pre-mRNA splicing in biogenesis of mature mRNA, we and others (8, 24–27) suggested that regulation of splicing kinetics may influence the gene expression kinetics that define the inflammatory cascade. Consistent with this idea, Braunschweig et al. (28) demonstrated that intron retention is a widespread and regulated feature of mammalian transcriptomes, often associated with reduced mRNA abundance and subject to tissue-specific control.

Deep learning models have rapidly advanced our ability to interpret noncoding regions of the genome by learning complex regulatory logic directly from DNA sequence. Early models such as DeepSEA demonstrated that convolutional neural networks could predict the effects of noncoding variants on transcription factor binding and chromatin state, laying a foundation for sequence-to-function inference (29). Building on this, SpliceAI adapted deep convolutional architectures to model splicing dynamics directly from primary sequence, achieving high accuracy in predicting splice site

usage and the effects of sequence variants on splicing (30). More recent models such as Enformer and Borzoi have expanded this paradigm further by predicting RNA-based outputs, including gene expression levels, transcript isoform usage, and transcription start site precision (31, 32). These models also support feature attribution though interpretable machine learning approaches such as saliency mapping and deepSHAP, allowing inference of sequence elements—such as transcription factor binding sites and splicing regulators—that most influence RNA output. Together, these approaches provide an unprecedented modeling context—spanning hundreds of kilobases—that enables the systematic dissection of sequence-level determinants of intron retention and the broader landscape of RNA processing.

To examine the timing of intron removal from 230 different transcripts induced by TNFα in macrophages, we have developed a method for highly enriching transcript populations for mRNAs of interest, which is followed by deep sequencing of the largely pre-mRNA populations we purify. The induction of transcripts and removal of introns can be quantified with precision, with lifetimes of introns determined by blocking transcription early after induction. Among genes whose mRNAs appear more slowly after induction, we identify ones containing introns with poor binding sites for splicing factor U1, usually finding one per transcript. We call these "bottleneck" introns. Among the most rapidly induced genes we find no such introns. We show that in these pre-mRNAs, the sequence of the U1 binding site is critical for the speed of intron removal by building mini-genes with "repaired" introns and showing that these splice at the canonical rate (identified as about 20 seconds after polymerase has passed that point). To complement these findings and explore regulatory determinants at larger scale, we apply a deep learning model trained on genomic sequence to predict splicing dynamics. Feature attribution methods identified sequence features associated with the splicing delays. We observe and model fine-tuning improved detection of patterns associated with intron retention. We propose that bottleneck introns are important for determining either the rate of degradation of pre-mRNAs or the rate of appearance of mature mRNA or both.

RESULTS

Hybrid Capture of Chromatin-Associated Transcripts

It has been established that pre-mRNA is highly enriched in the chromatin-associated, polyadenylated RNA fraction (33). To examine splicing events in pre-mRNAs, we performed time-course experiments with TNF-stimulated bone marrow-derived macrophages (BMDMs), isolating RNA after biochemical separation of chromatin-associated material (24). Using a hybrid-capture approach, we targeted sequencing toward transcripts of 230 genes previously identified as TNF-induced inflammatory mRNAs (34). Purification of cDNA corresponding to these inflammatory

transcripts involved reverse transcription of chromatin-associated RNA using oligo(dT), and capture of desired cDNAs using biotinylated probes complementary to the last exon of each gene of interest. Oligo(dT) priming and the choice of the last exon as a capture target enabled us to sequence complete transcripts from the standpoint of the splicing machinery, because all introns will have been transcribed in such transcripts. The hybrid capture strategy, based on a published approach (35), involved: (1) microarray printing of 12,000 150-bp ssDNAs designed from tiled fragments of the last exon of each gene of interest; (2) conversion to a pool of biotinylated ssRNA probes using PCR followed by in vitro transcription with biotinylated ribonucleotides; (3) hybridization of ssRNA pools to cDNA from each biological experiment, and (4) and streptavidin-coated bead-mediated capture of the transcripts of interest (Fig. S1). This approach resulted in a ~30-fold enrichment of genes of interest, with 70% of the sequenced reads corresponding to the genes of interest; RNA submitted only to poly(A) selection contained only 2% of such reads (Fig. S2A). In this way we could analyze 1098 introns (after TPM filter >100) from genes induced by an inflammatory stimulus (Fig S2B).

The selected chromatin-associated transcripts collected across time points after induction are shown as read-density histograms, displaying sequencing read abundance along each gene to reveal exon–intron structure. In Fig. 1A, Nfκbia (encoding IκB-α) is shown over 1 hour after TNF induction, with log_{10}-scaled densities normalized within each time point to highlight transcriptional kinetics. New transcription becomes evident by 6 minutes, especially on an unnormalized linear scale (Fig. S3).The corresponding log-scaled histograms permit visualization of the intronic signal as a function of time after induction. Furthermore, from Fig. 1A it is evident that at all time points following induction, the 5'-proximal introns have been totally removed from the sequenced transcripts, indicating that the selection against partial transcripts is quite complete. Whereas excision of the first intron is always observed, the middle three introns are seen at intermediate states of excision in all time points such that intron definition for these introns is readily observable from read density histograms. We attribute this first exon excision largely to co-transcriptional splicing, consistent with other genome-wide splicing studies (36). Strikingly, the final intron deviates significantly in its kinetic trajectory, as its read density does not obey a similar relative reduction. This might be due to a lag in terminal intron splicing (37) or a feature of splicing that accompanies transcript release from chromatin.

Quantifying Splice Completion Across the Transcriptome

To better quantify the observed dynamics, we adapted the Coefficient of Splicing (CoSI) (Figure 1B), which quantifies the extent of splicing as a ratio of spliced to total (spliced and unspliced) junction reads such that CoSI values of ~1 and ~0 imply near-complete splicing and virtually unspliced states, respectively (33). Though we observed a decrease in read density as a function of distance from the 3' of the gene (Fig. S3), presumably as a consequence of premature termination of

the reverse transcriptase during copying of the pre-mRNA, the use of CoSI allows for an intron-specific splicing score regardless of read densities at neighboring introns. Using the CoSI metric, a time course plotting of the extent of splicing showed a dip in CoSI at ~6 minutes (Fig. 1C) corresponding to the aforementioned accumulation of new, unspliced transcripts. The splicing dynamics of each *Nfkbia* intron can be inferred from the CoSI dynamics, and the notable difference in splicing between the 5' and 3' introns (Fig 1D) is demonstrated by the amplitude of the 'dip' at 6 minutes and the time required for each intron to return to Co-SI of 1.

A surprising heterogeneity in CoSI was observed among all inflammatory introns (Fig 2A), implying diversity in their propensity to be spliced (Fig. 2). When considering all 1,024 introns in the chromatin-associated TNF time course we find most introns very near to CoSI ~1 relatively soon after induction, indicating that most introns do not remain unspliced long after induction begins between 4-8minutes (Fig. 2A). The observation that many TNF-responsive transcripts initiate within 4–8 minutes is consistent with that of other related studies (24) Unexpectedly, although the median CoSI value remains high, we identified considerable heterogeneity among introns, some at and remaining near CoSI values <0.5, indicating relatively poor splicing, found very late into the time course.

As an example of a poorly spliced intron, chemokine *Cxcl10* intron 2 (Fig. 2A) is notable as it remains poorly spliced despite clear excision of neighboring introns, remaining quite unspliced even ~30 minutes after induction. It is possible that this intron undergoes splicing after its nascent chromatin-associated state, as is likely the case for the 3'-terminal intron of *Nfkbia*. It is also possible that this intron targets *Cxcl10* transcripts for degradation and the relatively fixed nature of intron 2's splicing status throughout the time course is a function of a constant rate of degradation. Introns with low CoSI at late time points post-TNF induction were considered putative 'bottleneck introns'– borrowing from the language that accompanied the discovery and characterization of slowly splicing U12-type introns (38). These introns were so slow to splice that they may intrinsically delay gene expression. Notably, the distribution of CoSI values of the entire dataset (Fig 2A) was very broad, and though most introns spliced immediately, many introns showed evidence of splicing bottlenecks, noticeable by their significant deviation from mean CoSI. At 10 and 60 minutes post-induction, 14% and 11% of introns, respectively, had CoSI values below one standard deviation of the mean (0.86±0.25 and 0.91±0.19, respectively). Shown as examples are *Cd40*, *Cxcl10, Daxx*, and *Irf7* (Fig. 2A-B), genes whose immunological and inflammatory importance is well-established in studies with knockout mice (39–41).

Measurements of Intron Splicing Half-Lives

To quantify splicing kinetics, we used Actinomycin D (actD) to freeze transcription and followed the loss of intron and accumulation of splice junctions. In these experiments, splicing was analyzed at

many time points immediately following actD treatment on the same 230 transcripts of interest and selected using hybrid capture from the total pool of cellular RNA rather than chromatin-associated RNA. Fitting the accumulation of spliced transcripts (as measured by CoSI) with an exponential distribution, we were able to extrapolate intron excision half-lives (Fig. 3). Because total cellular RNA was used, observed rates were independent of chromatin localization. We find intron half-lives that range from 20-40s (56% of introns splicing in this timing window) to several minutes, reflecting the considerable heterogeneity that is observed from CoSI differences.

A Multifaceted Approach for Identification of Determinants of Slow Splicing

Building on our observation that certain introns exhibit markedly delayed splicing by both CoSI dynamics and direct half-life measurements, we sought to uncover the underlying sequence features responsible for these delays. To this end, we employed a multifaceted strategy: (1) computational analysis of 5′ splice donor strength using MaxEntScan, (2) experimental testing of splicing efficiency with a minigene reporter assay, and (3) interpretable deep learning to identify additional cis-regulatory elements—beyond canonical splice sites—that may modulate splicing kinetics.

<u>Computational Analysis of Splice Donor Strength</u>

A delay in splicing at certain sites could simply confer a delay in gene expression (by ~5 minutes in the slow case shown in Figure 3, IκBε), or, as is seen in yeast studies, it could result in both gene expression delay and gene expression diminution due to degradation of slowly splicing pre-mRNA (42). Prior studies placed IκBε in a delayed splicing category (8), suggesting a pronounced splicing delay relative to rapidly induced genes. To understand a potential mechanistic basis for these differences in splicing time, each intron within our dataset was assessed computationally for the concurrence of its 5' splice donor sequence to a consensus sequence (43). The 5' splice donor is a highly conserved sequence that directly base pairs with splicing factor U1 (44); deviation from consensus sequence confers a significantly reduced ability to engage the splicing machinery.

A maximum entropy model (45) was used to calculate an intron quality score measuring extent of deviation from consensus splice sequence (Fig. S4). Among the inflammatory transcripts studied, many examples of introns with poor 5' donor scores were identified such as *Irf7* and *Il12b*, where lower scores indicate significant deviation from consensus. We suggest that having non-consensus splice sites may be a regulatory mechanism affecting gene expression. We considered that splicing might show profound differences in the previously defined categories of induction (immediate/early/intermediate/late) characteristic of the inflammatory gene expression kinetics (24). We found that the 'immediate' genes showed consistently fast splicing (highest CoSI values) at all of

their introns but the most 3', but that the other three groups shared similar CoSI distributions (Fig. S6). Using the bioinformatics "intron quality score" we also found that the introns of the later gene classes have significantly lower scoring 5' splice donor sequences (Fig. S4). Therefore, from experimental measurement of splicing and sequence-based prediction, genes expressed immediately following inflammatory stimulus are spliced faster, whereas all other inflammatory genes have a complex and heterogeneous distribution of splicing efficiency that does not stratify cleanly into the later kinetic categories (early/intermediate/late). Slowly splicing introns are found throughout these later kinetic categories in similar abundance, perhaps playing very gene-specific roles in diverse kinetic categories.

Experimental Validation with Minigene Fluorescent Reporter Assay

To test whether delays in splicing result in changes to gene expression, we identified a set of introns with the following criteria: (1) introns that splice poorly as defined by RNA-Seq, (2) introns that contain a low-scoring (non-consensus) 5' splice donor, and (3) introns whose weak 5' splice donor is evolutionarily conserved across many mammalian species. These introns were tested in the context of a splicing reporter expressed on a bidirectional promoter (46). For each intron of interest, the reporter construct consists of a single transcript containing: (1) the 5' neighboring exon from the gene of interest, the intron of interest, and the 3' neighboring exon; (2) a 2A 'self-cleaving' peptide; and (3) the GFP gene. In the opposite orientation but from the same promoter, a blue fluorescent protein (BFP) mRNA is made in equal amounts to the intron-GFP construct ,GFP fluorescence of cells transfected with this reporter is a readout of splicing efficiency of the intron-GFP construct, when normalized to BFP fluorescence levels. Transfected into HEK293T cells and expressed for 24 hours, this bidirectional reporter enabled us to understand, at steady state, whether gene expression is affected by slow splicing introns. Measured using flow cytometry, the slope of the line corresponding to BFP:GFP ratios provides a relative metric of splicing efficiency, where slopes ~.9 and ~0.1 imply efficient and inefficient splicing, respectively.

To test the effect of a poor splice donor, we 'rescued' some poorly splicing introns in the context of the reporter by mutation to consensus splice donor 'GTAAG.' For instance, for *IL12* intron 3, the splice donor sequence of "GTAAT" that is conserved among many mammalian species, was altered to "GTAAG" (Fig. S5). Expression levels of the reporter construct with the wild-type *IL12* intron were found to be about half (57%) of the levels of the same construct with a single base pair alteration to make the stronger splice donor. *IRF7* intron 5 was tested against a 'fixed' intron as well as a wild-type intron from an actin gene, both resulting in two-fold improvements of gene expression.

In one case, expression of *TFEC* (transcription factor EC) was not altered by splice site repair, suggesting that other mechanisms beyond 5'-splice site deficiencies may be involved in mediating slow splicing. Generally, when the BFP:GFP slopes of wild-type and mutated introns were compared by taking a ratio of their slopes, a change of ~2 nucleotides dramatically altered the slope of the line (Fig. 4).

Deep Learning–Based Discovery of Non-Canonical Regulatory Motifs

To extend our understanding of regulatory features beyond canonical splice donor sequences, we applied a supervised regulatory sequence model (Borzoi) to investigate sequence determinants associated with slow splicing. Borzoi is a model originally trained on steady-state RNA profiles across diverse cell types and tissues, which may limit its ability to capture stimulus-specific regulatory dynamics at play during TNFα-induced intron retention. Fine-tuning of this model offers a way to adapt a large, pre-trained sequence model to a specialized context by continuing training on domain-specific datasets, leveraging the general regulatory knowledge already encoded in the model weights while learning new patterns relevant to the task at hand. Fine-tuning has been shown to enhance Borzoi's performance in other specialized contexts, including tissue-specific expression, transcription factor knockdowns, and cellular aging (47). We fine-tuned it using RNA-seq coverage data from BMDMs at 18, 20, and 30 minutes post-TNFα induction—timepoints that capture dynamic intron retention. Fine-tuning was performed over 20 epochs using the Adam optimizer (learning rate = 1e-6, MSE loss). To evaluate generalization, chromosomes 10 and 11 were held out for training and used for validation and testing, respectively. Model loss steadily decreased over the training period for both datasets (Fig. 5A), and Pearson correlation (reflecting the strength of their linear relationship) between predicted and observed expression improved from $r = 0.51$ ($p = 4.44 \times 10^{-16}$) for the pretrained model to $r = 0.61$ ($p = 4.84 \times 10^{-24}$) after fine-tuning (Fig. 5B).

To interpret the model's predictions and identify candidate regulatory elements, we applied a widely adopted backpropagation based technique, DeepLIFT (48), to compute nucleotide-resolution importance scores, capturing the contribution of each base to splicing of the retained intron. Attribution scores were computed for each intron along with sequence within the model input window (524kb). In these scores, larger positive values indicate nucleotides that the model predicts as contributing to intron retention (i.e., delaying splicing), whereas negative values indicate nucleotides predicted to be associated with splice completion. We analyzed these scores using modisco-lite, which clusters recurring high-importance patterns into position weight matrices (PWMs) representing putative regulatory motifs (49). To assess motif enrichment among delayed introns, we focused on the 50 slowest-splicing introns as ranked by area under the CoSI-time curve (Fig. S7C). Motif occurrences were identified using a conventional motif scanning approach, Finding Individual Motif

Occurrences (FIMO) (50), and enrichment was calculated relative to either the remaining introns in our dataset or all introns in the transcriptome.

We compared the frequency of putative regulatory motifs in the 50 slowest-splicing introns to their frequency across all introns in the murine transcriptome, highlighting both the most enriched motifs overall and specific examples of their genomic locations within delayed introns. Among these, a GA-repeat–like motif showed enrichment in the slow-splicing set (FIMO, 80.0% vs 55.2%) (Fig. 6A). Representative instances of this motif were also recovered from the original source seqlets used to generate the PWM by modisco-lite, providing further support for its regulatory relevance (Fig. 6B-D). These findings illustrate how sequence-to-expression modeling, combined with attribution-based motif discovery, can uncover non-canonical cis-regulatory elements that may influence splicing efficiency during inflammatory gene activation.

DISCUSSION

In this study we sought to understand splicing kinetics of the large number of genes that comprise the inflammatory response and to assess whether splicing itself might play a regulatory role in inflammation. We developed a targeted sequencing strategy, purifying transcripts containing each gene's terminal exon. This approach allowed us to sequence the 1,024 introns within inflammatory genes and permitted direct assessment of the structures of nearly-completed transcripts. We found considerable heterogeneity in splicing efficiency among these introns. In studying evolutionarily conserved weak 5' splice donors, we have isolated one cause of slow appearance of mRNA following a pulse of stimulus; many other slowly spliced introns without such sequences were also identified in this study and suggest other regulatory mechanisms may be responsible.

Crucially, the hybrid capture approach averts a common ambiguity in analyzing splicing kinetics of not being able to differentiate completed pre-mRNA from nascent transcripts during an induction pulse – this often leads to an overestimation of the unspliced status of early introns and complicates quantification of splicing kinetics. To the contrary, we rarely found genes containing unspliced first introns. This was true of chromatin-associated RNA and of whole cell RNA after inhibition of transcription with actD. These effects are consistent with the emerging model of co-transcriptional pre-mRNA splicing, where the splicing machinery has been suggested to lag 3-5kb behind the polymerase. Indeed, several recent global studies of RNA splicing bolster the claim that much pre-mRNA is spliced co-transcriptionally: 74% (yeast) or 75-84% (human) of introns are found to be at least 50% spliced by the time of transcription termination in several other studies (33, 37, 51–54). Surprisingly, this ~80% figure remains constant whether total RNA or chromatin-associated RNA

is measured, implying that our choice to analyze chromatin-associated RNA does not significantly overrepresent splicing intermediates.

We found that most introns are spliced very efficiently, appearing and disappearing as a rapid dip of CoSI immediately following induction, returning to a CoSI of ~0.95 within minutes after induction. Notably, the distribution of CoSI values of the entire dataset (Figure 2) was very broad. Though most introns spliced immediately, there were several 'bottleneck introns'. In order to determine more specifically the rates of slowly splicing introns, studies employing actD to stall transcription and examine intron splicing half-lives corroborated the idea that there is tremendous intron-to-intron heterogeneity. Most delayed introns ultimately reached higher CoSI values over the time course, consistent with completion of splicing rather than stable, long-term retention (see Figs. 2–3). Whereas most introns spliced within 20-40s, some were delayed significantly (upwards of 5 minutes). Of note, however, is that our 20-40s rate of splicing is somewhat at odds with other figures in the literature of 8-10 minutes for intron excision (9) after a washout of the drug D-ribofuranosylbenzimidazole. There is some debate as to the perturbative role of actD in splicing, with one report observing that splicing intermediates in the context of the MS2 reporter system are prematurely liberated from chromatin upon actD treatment. Even in this case, the rapid actD-based rates, are likely underestimating even faster kinetics if one considers that the co-transcriptional splicing machinery targets chromatin mRNA faster than released mRNA (55). However, even in the absence of actD, stimulation revealed that most of IKBα's introns are spliced in less than two minutes when one takes into account the 'dip' in CoSI due to induction and the time to reach steady CoSI levels (Figure 2). While the terminal intron of IKBα appears to have a longer half-life, this unique feature of terminal introns is consistent with prior studies (37).

In testing gene expression differences in bottleneck introns among introns with poor splice sites that are also evolutionarily conserved, we found that steady state levels of reporter proteins were upregulated when the 5' splice donor sequence was mutated to the consensus sequence 'GTAAG' in all cases but one. Attenuated U1 binding provides a mechanistic insight for bottleneck introns that were chosen for their weak 5' splice donors. This implies that at the level of splicing, either due to delays in expression or perhaps degradation due to delayed expression, significant differences in gene expression arise from small differences in nucleotide sequence. These reporter assays measure steady-state expression influenced by intron sequence; while consistent with changes in splicing efficiency, other post-transcriptional mechanisms may be at work to preclude these introns from efficient splicing. We also find many slowly spliced introns not explained by weak 5' splice donors. In some cases, we find multiple bottlenecks introns per gene, as is the case of IRF7, where one bottleneck (intron 5) was attributable to an evolutionarily conserved weak 5' splice donor

while another (intron 7) was observed experimentally but of unknown cause. This may be due to any of a number of potential mechanisms that may also serve in tuning the speed of splicing: cis-regulatory protein recruitment, 3' splice acceptor sequence or other sequence elements, or alterations of RNA polymerase speed or chromatin marks or three-dimensional gene structure.

Central to our inquiry is the enigmatic nature of these bottlenecks remaining in physiologically critical genes, often evolutionarily conserved, and yet intrinsically mediating an inefficiency in gene expression. Importantly, the conservation of these weak donor sites suggests they confer a regulatory advantage—such as fine-tuning of expression timing or transcript abundance—rather than representing neutral or deleterious features merely tolerated by selection. We posit that the gene expression changes that are shown in bone marrow-derived macrophages offer a regulatory strategy to slow up and maybe restrict expression of genes in a manner dependent on the composition of mRNA processing factors in the cell ('the splicing landscape'), the cell type, or the stimulus type in question. Recent studies have demonstrated global changes in intron retention preferences in B cell lymphomagenesis and granulocyte differentiation (56). In a similar manner, we suggest that selection of splicing and kinetics of splicing might allow a previously unappreciated level of specificity to gene expression decisions in cells presented with an inflammatory stimulus (57–63).

Induction with TNF is a particularly favorable situation because many of the genes we examined were up-regulated in their transcription within 4-6 minutes of adding inducer (Supplemental Fig. 2B), allowing examination of large numbers of pre-mRNA transcripts. This, in concert with the hybrid capture approach that provides a large number of junctional sequencing reads, has permitted unique insight into the kinetics of splicing of mature transcripts and revealed surprising heterogeneity. We suggest that this methodology and analysis could have wider applicability for other gene induction situations.

Despite the promise of deep learning models, challenges remain in translating their predictions into biological insight. While performance on benchmark tasks continues to improve—often surpassing traditional motif-finding tools—generalization across conditions and cell types is still constrained by the scope of training data. Interpretability is another major hurdle, particularly for genomic sequences. Interpretable machine learning methods can highlight sequence elements that influence model predictions, but connecting these regions to underlying biological mechanisms often demands extensive experimental validation. In the case of RNA splicing, complexity of overlapping regulatory layers—such as RNA secondary structure, co-transcriptional dynamics, and nuclear export—can make it difficult to disentangle causal sequence features from correlated signals. Nonetheless, sequence-to-expression modeling provides a powerful framework for identifying

sequence features correlated with delayed splicing and for guiding experimental discovery of novel regulators through fine-tuning and perturbation-informed learning.

Our deep learning-based analysis further supports the notion that splicing efficiency is shaped by a broad array of cis-regulatory features, many of which fall outside canonical splice site motifs. By leveraging, a sequence-to-expression transformer trained on multimodal regulatory data, we identified non-canonical sequence motifs enriched in the slowest-splicing introns—motifs that may act as silencers or delay elements involved in splicing. These results offer a complementary, agnostic perspective to our mutational analyses, and suggest that intronic bottlenecks can arise from diverse sequence architectures. The enrichment of these motifs in bottleneck introns—independent of donor site strength—points to additional layers of splicing control that may be particularly relevant in rapid-response transcriptional programs like inflammation. We propose that such regulatory elements may encode a form of temporal tuning, modulating transcript availability through fine control of intron excision. More broadly, our work demonstrates interpretable machine learning can uncover latent regulatory features that elude traditional sequence analysis, advancing efforts to decode the logic of splicing regulation in a cell-type or stimulus-specific context.

EXPERIMENTAL PROCEDURES:

Cells
C56BL6/J mice were sacrificed via CO_2 euthanasia and sterilized with 70% ethanol. Femur and tibia bones harvested and stripped of muscle tissue. Bone marrow cells were resuspended in 20mL of fresh DMEM. 2.5e6 bone-marrow cells plated in a 15-cm dish in 20mL of BMDM Media (DMEM, 20% FBS, 30% L929 condition media, and 1% Pen/Strep) and grown at 5% CO_2 and 37°C. BMDM media completely replaced on day 3 as well as a supplemental addition of 5mL L929 condition media on day 5.

RNA fractionation.
RNA was fractionated into cytoplasmic, nucleoplasmic, and chromatin-associated pools as previously described (24) with modifications. Confluent 15 cm dishes of mature BMDMs were scraped into 400 µL cold NP-40 lysis buffer (10 mM Tris-HCl pH 7.5, 0.08% NP-40, 150 mM NaCl) and layered onto a 1 mL sucrose cushion (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 24% w/v sucrose). Samples were centrifuged at 13,000 rpm for 10 min at 4 °C. The supernatant (cytoplasmic fraction) was mixed with 3 volumes of 100% ethanol and 2 volumes of buffer RLT (4 M GuSCN, 0.1 M β-mercaptoethanol, 0.5% N-lauroylsarcosine, 25 mM Na-citrate, pH 7.2) and stored at -80 °C.

The nuclear pellet was resuspended in 200 µL cold glycerol buffer (20 mM Tris-HCl pH 7.5, 75 mM NaCl, 0.5 mM EDTA pH 8.0, 50% glycerol, 0.85 mM DTT) and lysed with an equal volume of nuclear lysis buffer (20 mM HEPES pH 7.5, 7.5 mM $MgCl_2$, 0.2 mM EDTA pH 8.0, 1 M urea, 1% NP-40). After vortexing and centrifugation (14,000 rpm, 5 min, 4 °C), the supernatant (nucleoplasmic fraction) was processed as above.

The remaining chromatin pellet was hydrated in 1× PBS and dissolved in 500 μL TRIzol reagent at 50 °C with intermittent vortexing. After phase separation with 100 μL chloroform, the aqueous phase was recovered and processed as above. All RNA fractions were purified using the Qiagen RNeasy protocol and eluted in nuclease-free water. Typical yields were 300–500 ng/μL for cytoplasmic RNA, 100–250 ng/μL for nucleoplasmic RNA, and 300–500 ng/μL for chromatin-associated RNA. RNA was DNase-treated (TURBO DNase, Thermo Fisher) and stored at -80 °C.

Template-switch reverse transcription

One microgram of RNA was mixed with 1 μM oligo(dT)$_{30}$ (5′-AAGCAGTGGTATCAACGCAGAGTACT$_{30}$-3′), heated to 80 °C for 2.5 min, and snap-cooled on ice. A 10 μL reverse-transcription mix containing 10 μM template-switch oligo (5′-AAGCAGTGGTATCAACGCAGAGTACACArGrGrG-3′), 20 mM DTT, 2× First-Strand Buffer (Invitrogen), 1 mM dNTPs, 40 U Murine RNase Inhibitor (NEB), and 200 U SuperScript II (Invitrogen) was added. Reactions were incubated sequentially at 42 °C (30 min), 45 °C (30 min), 50 °C (10 min), then heat-inactivated at 80 °C (10 min).

RNA templates were degraded by adding NaOH (final 0.1 M) and EDTA (5 mM) and heating to 70 °C for 10 min, followed by neutralization with HCl. cDNA was purified using 2× Sera-Mag carboxylate-modified magnetic beads (GE Healthcare) with standard PEG/ethanol washes and eluted in nuclease-free water.

Hybrid capture probe design and synthesis

Biotinylated RNA probes were designed against the terminal exons of inflammatory genes (see Supplemental Table 2). For each gene, 100-bp overlapping oligonucleotides were synthesized (CustomArray Inc.) and pooled into nine expression-matched subgroups. Each subpool was PCR-amplified to append a T7 RNA polymerase promoter and transcribed in vitro using the AmpliScribe T7 Biotin Flash Kit (Epicentre). Purified RNA probe subpools were combined in weighted ratios (A–I) to normalize capture representation across expression levels as below:

Pool	Pool composition
A	0.01113
B	0.0211
C	0.03235
D	0.05514
E	0.10542
F	0.21908
G	0.4674
H	1
I	1

cDNA hybrid capture and elution.

Biotinylated probes were hybridized to cDNA at 74 °C for 4.5 min, followed by addition of 2× hybridization buffer (1 M LiCl, 40 mM Tris-HCl pH 7.5, 20 mM EDTA pH 8.0, 4 M urea, 0.5% Triton X-100, 1% SDS, 0.2% Na-deoxycholate). Hybridization continued 30 min at 70 °C. Streptavidin BioMag

beads (0.3 mg) were washed and incubated 20 min at 70 °C to capture cDNA–probe complexes. Beads were washed twice with 1× HYB, once each with Wash 4 and Wash 5 buffers, and eluted in 35 µL base elution buffer (125 mM NaOH, 10 mM EDTA pH 8.0, 10 mM Tris-HCl pH 7.5) at 74 °C for 5 min. The eluate was neutralized and purified with 1× Sera-Mag beads and eluted in 45µL and stored at -80°C.

Determining the efficiency of cDNA pulldown
Pulldown efficiency was evaluated by qPCR to quantify enrichment of target transcripts and depletion of background RNA. qPCR reactions (KAPA SYBR 2× Master Mix) compared pre- and post-pulldown cDNA at a 2:1 ratio. Primers:
• L32 (background control): F 5′-AAGCGAAACTGGCGGAAAC-3′; R 5′-TAACCGATGTTGGGCATCAG-3′
• NF-κBIA (spliced exon 5–6 junction): F 5′-ACGGAGTCAGAATTCACAGAGG-3′; R 5′-CACAAAGACAACAGCCGAATC-3′

Cycle-threshold (Ct) values were used to calculate ΔCt between L32 and NF-κBIA. Successful pulldowns typically showed ΔL32 = –7 to –9 cycles and ΔNF-κBIA = –2 to –4, corresponding to ΔL32/ΔNF-κBIA > 2.0.

Post-pulldown cDNA amplification
Pulldown cDNA was amplified prior to tagmentation. PCR reactions contained Q5 High-Fidelity 2× Master Mix (NEB), 1 µM primer (5′-AAGCAGTGGTATCAACGCAGAGTACT-3′), and ~5% of the pulldown reaction. Cycling: 95 °C (2 min) → 20–25 cycles of 95 °C (30 s), 62.5 °C (30 s), 72 °C (150 s) → 72 °C (5 min). PCR products were purified (0.9× Sera-Mag) and eluted in 25 µL H_2O. Concentrations were determined using a Qubit HS dsDNA Assay.

Tagmentation of cDNA libraries with Tn5 transposase
Tn5 transposase was purified as in Picelli et al., Genome Res. (2014) and pre-loaded with hybridized adapter oligos (Tn5MErev 5′-[phos]CTGTCTCTTATACACATCT-3′; Tn5ME-A 5′-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-3′; Tn5ME-B 5′-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-3′). Tagmentation reactions contained ~40 ng amplified cDNA, 0.2 µL Tn5, 5% PEG8000, 10 mM TAPS (pH 8.5), 5 mM $MgCl_2$, incubated 10 min at 55 °C. SDS (0.02%) was added and incubation continued 10 min to inactivate Tn5. Reactions were 1.4× Sera-Mag purified and eluted in 20 µL H_2O.

Library barcoding and sequencing
Tagmented libraries were PCR-amplified with paired barcode oligos using Q5 High-Fidelity 2× Master Mix (NEB). Cycling: 72 °C (3 min), 98 °C (2 min), then 25 cycles of 98 °C (10 s), 63 °C (30 s), 72 °C (30 s), and 72 °C (5 min). Libraries were double-purified (1.0×/1.4× Sera-Mag), quantified by Qubit HS dsDNA assay, pooled equimolarly, repurified, and analyzed with an Agilent Bioanalyzer 2100. Sequencing was performed on an Illumina HiSeq 2500 (50 bp, single-end mode).

RNA-seq alignment and analysis
Single-end 50 bp reads were aligned to the mm10 genome using STAR. Junctions were retained only if both exonic and intronic segments were ≥ 3 bp. Using pysam, splice and intron junctions were classified as types a, b, or c, and Completion of Splicing Index (CoSI) values were computed using custom Python scripts. All analyses were performed in Python and R. Sequencing data are being deposited to NCBI GEO, and all analysis code will be made available on GitHub.

Deep Learning Analysis of Intron Regulatory Features

Model architecture and initialization.

We used Borzoi, a transformer-based sequence-to-function model trained to predict steady-state RNA abundance from genomic DNA sequence. The pretrained model captures general patterns of cis-regulatory architecture by integrating multi-omic training targets (CAGE, ATAC-seq, and ChIP-seq tracks). However, because Borzoi is optimized for baseline expression rather than stimulus-responsive contexts, we adapted it to our macrophage TNFα time course through targeted fine-tuning.

Fine-tuning on TNFα-stimulated macrophage data.

We generated genome-wide RNA-seq coverage tracks from bone marrow–derived macrophages (BMDMs) collected at 18, 20, and 30 minutes post-TNFα induction—timepoints representative of dynamic intron retention. These coverage values served as supervised training targets to refine the model's prediction of RNA abundance under inflammatory conditions. Fine-tuning was performed for 20 epochs using the Adam optimizer (learning rate = 1×10^{-6}, mean-squared-error loss). To evaluate generalization, chromosomes 10 and 11 were withheld for validation and testing, respectively. Model loss steadily decreased over training for both sets, and the correlation between predicted and observed expression improved from Pearson r = 0.51 (p = 4.44×10^{-16}) for the pretrained model to r = 0.61 (p = 4.84×10^{-24}) after fine-tuning (Fig. 5A–B).

Feature attribution and motif discovery.

To interpret the fine-tuned model's predictions, we applied DeepLIFT to compute nucleotide-resolution importance scores, quantifying each base's contribution to the predicted RNA signal within a 524 kb genomic window centered on each intron of interest. Positive importance scores indicate sequence positions contributing to higher predicted intronic RNA signal (i.e., delayed splicing), whereas negative scores correspond to features predictive of splice completion.

Importance profiles were analyzed using modisco-lite, which clusters recurrent high-importance patterns across examples into position weight matrices (PWMs) representing putative regulatory motifs. Enrichment of discovered motifs was assessed using FIMO, scanning the 50 (or 150) slowest-splicing introns—ranked by area under the CoSI–time curve—against all annotated murine introns (GENCODE M23). For targeted motif discovery, attributions input to modisco-lite were masked under different configurations to restrict pattern identification to either intronic or exonic regions adjacent to slow-splicing introns.

Motif enrichment and visualization.

Relative motif frequencies were plotted as a scatter of percent representation in slow- versus all-intron backgrounds. Sequence logos of the top five enriched motifs were visualized alongside genomic examples where GA-rich and A-rich motifs coincided with regions of high attribution in DeepLIFT profiles (Fig. 6A–D).

ACKNOWLEDGEMENTS:

The authors would like to thank Alex Shishkin and Mitchell Guttman (Dept. of Biology, Caltech) for assistance with hybrid capture strategy design; and Ann-Jay Tong Stephen Smale, Doug Black and Amy-Pandya Jones (Dept. of Biology, University of California, Los Angeles) for insights and advice; and Sergei Manakov, Evelyn Stuwe, Dubravka Pezic, Igor Antoshechkin, Sagar Damle, and Alok Joglekar (Dept. of Biology, California Institute of Technology) for experimental and computational assistance. This work was funded from a grant from NIH and from an endowment provided by the Raymond and Beverly Sackler Foundation. Research reported in this publication was supported by the National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health under award number P20GM125498

AUTHOR CONTRIBUTIONS:

J.S. Dearborn, Conceptualization, Investigation, Data curation, Software, Formal analysis, Visualization, Methodology, Writing – review & editing. L. Frankiw, Conceptualization, Investigation, Resources, Methodology, Validation, Writing – review & editing. D.W. Limoge, Data curation, Formal analysis, Visualization, Writing – review & editing. C.H. Burns, L. Vlach, P. Turpin, T. Kirch, Z.D. Miller, W. Dowell, S. Languon, Y. Garcia-Flores, Investigation, Data curation, Validation, Writing – review & editing. R.C. Cockrell, Resources, Supervision, Project administration, Writing – review & editing. D. Baltimore, Conceptualization, Supervision, Funding acquisition, Project administration, Writing – review & editing. D. Majumdar, Conceptualization, Supervision, Funding acquisition, Project administration, Methodology, Writing – review & editing.

REFERENCES

1. Chen, J., and Chen, Z. J. (2013) Regulation of NF-κB by ubiquitination. *Curr. Opin. Immunol.* **25**, 4–12
2. Gautier, E. L., Shay, T., Miller, J., Greter, M., Jakubzick, C., Ivanov, S., Helft, J., Chow, A., Elpek, K. G., Gordonov, S., Mazloom, A. R., Ma'ayan, A., Chua, W.-J., Hansen, T. H., Turley, S. J., Merad, M., Randolph, G. J., and Immunological Genome Consortium (2012) Gene-expression profiles and transcriptional regulatory pathways that underlie the identity and diversity of mouse tissue macrophages. *Nat. Immunol.* **13**, 1118–1128
3. Smale, S. T., Tarakhovsky, A., and Natoli, G. (2014) Chromatin contributions to the regulation of innate immunity. *Annu. Rev. Immunol.* **32**, 489–511
4. Hao, S., and Baltimore, D. (2009) The stability of mRNA influences the temporal order of the induction of genes encoding inflammatory molecules. *Nat. Immunol.* **10**, 281–288
5. Leppek, K., Schott, J., Reitter, S., Poetz, F., Hammond, M. C., and Stoecklin, G. (2013) Roquin promotes constitutive mRNA decay via a conserved class of stem-loop recognition motifs. *Cell*. **153**, 869–881
6. O'Connell, R. M., Rao, D. S., and Baltimore, D. (2012) microRNA regulation of inflammatory responses. *Annu. Rev. Immunol.* **30**, 295–312
7. Wan, F., Anderson, D. E., Barnitz, R. A., Snow, A., Bidere, N., Zheng, L., Hegde, V., Lam, L. T., Staudt, L. M., Levens, D., Deutsch, W. A., and Lenardo, M. J. (2007) Ribosomal protein S3: a KH domain subunit in NF-kappaB complexes that mediates selective gene regulation. *Cell*. **131**, 927–939
8. Hao, S., and Baltimore, D. (2013) RNA splicing regulates the temporal order of TNF-induced gene expression. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 11934–11939
9. Pandya-Jones, A., Bhatt, D. M., Lin, C.-H., Tong, A.-J., Smale, S. T., and Black, D. L. (2013) Splicing kinetics and transcript release from the chromatin compartment limit the rate of Lipid A-induced gene expression. *RNA N. Y. N.* **19**, 811–827
10. Rabani, M., Levin, J. Z., Fan, L., Adiconis, X., Raychowdhury, R., Garber, M., Gnirke, A., Nusbaum, C., Hacohen, N., Friedman, N., Amit, I., and Regev, A. (2011) Metabolic labeling of RNA uncovers principles of RNA production and degradation dynamics in mammalian cells. *Nat. Biotechnol.* **29**, 436–442
11. Rabani, M., Raychowdhury, R., Jovanovic, M., Rooney, M., Stumpo, D. J., Pauli, A., Hacohen, N., Schier, A. F., Blackshear, P. J., Friedman, N., Amit, I., and Regev, A. (2014) High-resolution sequencing and modeling identifies distinct dynamic RNA regulatory strategies. *Cell*. **159**, 1698–1710
12. Bergkessel, M., Whitworth, G. B., and Guthrie, C. (2011) Diverse environmental stresses elicit distinct responses at the level of pre-mRNA processing in yeast. *RNA N. Y. N.* **17**, 1461–1478
13. Guilgur, L. G., Prudêncio, P., Sobral, D., Liszekova, D., Rosa, A., and Martinho, R. G. (2014) Requirement for highly efficient pre-mRNA splicing during Drosophila early embryonic development. *eLife*. **3**, e02181
14. Takashima, Y., Ohtsuka, T., González, A., Miyachi, H., and Kageyama, R. (2011) Intronic delay is essential for oscillatory expression in the segmentation clock. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 3300–3305
15. Eissa, N. T., Strauss, A. J., Haggerty, C. M., Choo, E. K., Chu, S. C., and Moss, J. (1996) Alternative splicing of human inducible nitric-oxide synthase mRNA. tissue-specific regulation and induction by cytokines. *J. Biol. Chem.* **271**, 27184–27187
16. Hargreaves, D. C., Horng, T., and Medzhitov, R. (2009) Control of inducible gene expression by signal-dependent transcriptional elongation. *Cell*. **138**, 129–145

17. Kontoyiannis, D., Pasparakis, M., Pizarro, T. T., Cominelli, F., and Kollias, G. (1999) Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: implications for joint and gut-associated immunopathologies. *Immunity*. **10**, 387–398

18. Mahtani, K. R., Brook, M., Dean, J. L., Sully, G., Saklatvala, J., and Clark, A. R. (2001) Mitogen-activated protein kinase p38 controls the expression and posttranslational modification of tristetraprolin, a regulator of tumor necrosis factor alpha mRNA stability. *Mol. Cell. Biol.* **21**, 6461–6469

19. Mino, T., Murakawa, Y., Fukao, A., Vandenbon, A., Wessels, H.-H., Ori, D., Uehata, T., Tartey, S., Akira, S., Suzuki, Y., Vinuesa, C. G., Ohler, U., Standley, D. M., Landthaler, M., Fujiwara, T., and Takeuchi, O. (2015) Regnase-1 and Roquin Regulate a Common Element in Inflammatory mRNAs by Spatiotemporally Distinct Mechanisms. *Cell*. **161**, 1058–1073

20. Rao, N., Nguyen, S., Ngo, K., and Fung-Leung, W.-P. (2005) A novel splice variant of interleukin-1 receptor (IL-1R)-associated kinase 1 plays a negative regulatory role in Toll/IL-1R-induced inflammatory signaling. *Mol. Cell. Biol.* **25**, 6521–6532

21. Ruggiero, T., Trabucchi, M., De Santa, F., Zupo, S., Harfe, B. D., McManus, M. T., Rosenfeld, M. G., Briata, P., and Gherzi, R. (2009) LPS induces KH-type splicing regulatory protein-dependent processing of microRNA-155 precursors in macrophages. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* **23**, 2898–2908

22. Stoecklin, G., Lu, M., Rattenbacher, B., and Moroni, C. (2003) A constitutive decay element promotes tumor necrosis factor alpha mRNA degradation via an AU-rich element-independent pathway. *Mol. Cell. Biol.* **23**, 3506–3515

23. Han, J., Brown, T., and Beutler, B. (1990) Endotoxin-responsive sequences control cachectin/tumor necrosis factor biosynthesis at the translational level. *J. Exp. Med.* **171**, 465–475

24. Bhatt, D. M., Pandya-Jones, A., Tong, A.-J., Barozzi, I., Lissner, M. M., Natoli, G., Black, D. L., and Smale, S. T. (2012) Transcript dynamics of proinflammatory genes revealed by sequence analysis of subcellular RNA fractions. *Cell*. **150**, 279–290

25. Cho, V., Mei, Y., Sanny, A., Chan, S., Enders, A., Bertram, E. M., Tan, A., Goodnow, C. C., and Andrews, T. D. (2014) The RNA-binding protein hnRNPLL induces a T cell alternative splicing program delineated by differential intron retention in polyadenylated RNA. *Genome Biol.* **15**, R26

26. Davis-Turak, J. C., Allison, K., Shokhirev, M. N., Ponomarenko, P., Tsimring, L. S., Glass, C. K., Johnson, T. L., and Hoffmann, A. (2015) Considering the kinetics of mRNA synthesis in the analysis of the genome and epigenome reveals determinants of co-transcriptional splicing. *Nucleic Acids Res.* **43**, 699–707

27. Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., and Regev, A. (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* **29**, 644–652

28. Braunschweig, U., Barbosa-Morais, N. L., Pan, Q., Nachman, E. N., Alipanahi, B., Gonatopoulos-Pournatzis, T., Frey, B., Irimia, M., and Blencowe, B. J. (2014) Widespread intron retention in mammals functionally tunes transcriptomes. *Genome Res.* **24**, 1774–1786

29. Zhou, J., and Troyanskaya, O. G. (2015) Predicting effects of noncoding variants with deep learning–based sequence model. *Nat. Methods*. **12**, 931–934

30. Jaganathan, K., Kyriazopoulou Panagiotopoulou, S., McRae, J. F., Darbandi, S. F., Knowles, D., Li, Y. I., Kosmicki, J. A., Arbelaez, J., Cui, W., Schwartz, G. B., Chow, E. D., Kanterakis, E., Gao, H., Kia, A., Batzoglou, S., Sanders, S. J., and Farh, K. K.-H. (2019) Predicting Splicing from Primary Sequence with Deep Learning. *Cell*. **176**, 535-548.e24

31. Avsec, Ž., Agarwal, V., Visentin, D., Ledsam, J. R., Grabska-Barwinska, A., Taylor, K. R., Assael, Y., Jumper, J., Kohli, P., and Kelley, D. R. (2021) Effective gene expression prediction from sequence by integrating long-range interactions. *Nat. Methods*. **18**, 1196–1203

32. Linder, J., Srivastava, D., Yuan, H., Agarwal, V., and Kelley, D. R. (2025) Predicting RNA-seq coverage from DNA sequence as a unifying model of gene regulation. *Nat. Genet.* **57**, 949–961

33. Tilgner, H., Knowles, D. G., Johnson, R., Davis, C. A., Chakrabortty, S., Djebali, S., Curado, J., Snyder, M., Gingeras, T. R., and Guigó, R. (2012) Deep sequencing of subcellular RNA fractions shows splicing to be predominantly co-transcriptional in the human genome but inefficient for lncRNAs. *Genome Res.* **22**, 1616–1625

34. Ramirez-Carrozzi, V. R., Braas, D., Bhatt, D. M., Cheng, C. S., Hong, C., Doty, K. R., Black, J. C., Hoffmann, A., Carey, M., and Smale, S. T. (2009) A unifying model for the selective regulation of inducible transcription by CpG islands and nucleosome remodeling. *Cell*. **138**, 114–128

35. Engreitz, J. M., Pandya-Jones, A., McDonel, P., Shishkin, A., Sirokman, K., Surka, C., Kadri, S., Xing, J., Goren, A., Lander, E. S., Plath, K., and Guttman, M. (2013) The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. *Science*. **341**, 1237973

36. Bentley, D. L. (2014) Coupling mRNA processing with transcription in time and space. *Nat. Rev. Genet.* **15**, 163–175

37. Carrillo Oesterreich, F., Preibisch, S., and Neugebauer, K. M. (2010) Global analysis of nascent RNA reveals transcriptional pausing in terminal exons. *Mol. Cell*. **40**, 571–581

38. Patel, A. A., McCarthy, M., and Steitz, J. A. (2002) The splicing of U12-type introns can be a rate-limiting step in gene expression. *EMBO J.* **21**, 3804–3815

39. Honda, K., Yanai, H., Negishi, H., Asagiri, M., Sato, M., Mizutani, T., Shimada, N., Ohba, Y., Takaoka, A., Yoshida, N., and Taniguchi, T. (2005) IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature*. **434**, 772–777

40. Lei, X. F., Ohkawara, Y., Stämpfli, M. R., Mastruzzo, C., Marr, R. A., Snider, D., Xing, Z., and Jordana, M. (1998) Disruption of antigen-induced inflammatory responses in CD40 ligand knockout mice. *J. Clin. Invest.* **101**, 1342–1353

41. Michaelson, J. S., Bader, D., Kuo, F., Kozak, C., and Leder, P. (1999) Loss of Daxx, a promiscuously interacting protein, results in extensive apoptosis in early mouse development. *Genes Dev.* **13**, 1918–1923

42. Koodathingal, P., Novak, T., Piccirilli, J. A., and Staley, J. P. (2010) The DEAH box ATPases Prp16 and Prp43 cooperate to proofread 5' splice site cleavage during pre-mRNA splicing. *Mol. Cell*. **39**, 385–395

43. Pessa, H. K. J., Ruokolainen, A., and Frilander, M. J. (2006) The abundance of the spliceosomal snRNPs is not limiting the splicing of U12-type introns. *RNA N. Y. N.* **12**, 1883–1892

44. Freund, M., Hicks, M. J., Konermann, C., Otte, M., Hertel, K. J., and Schaal, H. (2005) Extended base pair complementarity between U1 snRNA and the 5' splice site does not inhibit splicing in higher eukaryotes, but rather increases 5' splice site recognition. *Nucleic Acids Res.* **33**, 5112–5119

45. Yeo, G., and Burge, C. B. (2004) Maximum entropy modeling of short sequence motifs with applications to RNA splicing signals. *J. Comput. Biol. J. Comput. Mol. Cell Biol.* **11**, 377–394

46. Mukherji, S., Ebert, M. S., Zheng, G. X. Y., Tsang, J. S., Sharp, P. A., and van Oudenaarden, A. (2011) MicroRNAs can generate thresholds in target gene expression. *Nat. Genet.* **43**, 854–859

47. Yuan, H., Linder, J., and Kelley, D. R. (2025) Parameter-Efficient Fine-Tuning of a Supervised Regulatory Sequence Model. 10.1101/2025.05.26.656171

48. Shrikumar, A., Greenside, P., and Kundaje, A. (2019) Learning Important Features Through Propagating Activation Differences. 10.48550/arXiv.1704.02685

49. Shrikumar, A., Tian, K., Avsec, Ž., Shcherbina, A., Banerjee, A., Sharmin, M., Nair, S., and Kundaje, A. (2020) Technical Note on Transcription Factor Motif Discovery from Importance Scores (TF-MoDISco) version 0.5.6.5. 10.48550/arXiv.1811.00416

50. Grant, C. E., Bailey, T. L., and Noble, W. S. (2011) FIMO: scanning for occurrences of a given motif. *Bioinforma. Oxf. Engl.* **27**, 1017–1018

51. Ameur, A., Zaghlool, A., Halvardson, J., Wetterbom, A., Gyllensten, U., Cavelier, L., and Feuk, L. (2011) Total RNA sequencing reveals nascent transcription and widespread co-transcriptional splicing in the human brain. *Nat. Struct. Mol. Biol.* **18**, 1435–1440

52. Brugiolo, M., Herzel, L., and Neugebauer, K. M. (2013) Counting on co-transcriptional splicing. *F1000prime Rep.* **5**, 9

53. Girard, C., Will, C. L., Peng, J., Makarov, E. M., Kastner, B., Lemm, I., Urlaub, H., Hartmuth, K., and Lührmann, R. (2012) Post-transcriptional spliceosomes are retained in nuclear speckles until splicing completion. *Nat. Commun.* **3**, 994

54. Khodor, Y. L., Rodriguez, J., Abruzzi, K. C., Tang, C.-H. A., Marr, M. T., and Rosbash, M. (2011) Nascent-seq indicates widespread cotranscriptional pre-mRNA splicing in Drosophila. *Genes Dev.* **25**, 2502–2512

55. Martin, R. M., Rino, J., Carvalho, C., Kirchhausen, T., and Carmo-Fonseca, M. (2013) Live-cell visualization of pre-mRNA splicing with single-molecule sensitivity. *Cell Rep.* **4**, 1144–1155

56. Koh, C. M., Bezzi, M., Low, D. H. P., Ang, W. X., Teo, S. X., Gay, F. P. H., Al-Haddawi, M., Tan, S. Y., Osato, M., Sabò, A., Amati, B., Wee, K. B., and Guccione, E. (2015) MYC regulates the core pre-mRNA splicing machinery as an essential step in lymphomagenesis. *Nature*. **523**, 96–100

57. Brinster, R. L., Allen, J. M., Behringer, R. R., Gelinas, R. E., and Palmiter, R. D. (1988) Introns increase transcriptional efficiency in transgenic mice. *Proc. Natl. Acad. Sci. U. S. A.* **85**, 836–840

58. Damgaard, C. K., Kahns, S., Lykke-Andersen, S., Nielsen, A. L., Jensen, T. H., and Kjems, J. (2008) A 5' splice site enhances the recruitment of basal transcription initiation factors in vivo. *Mol. Cell*. **29**, 271–278

59. Furger, A., O'Sullivan, J. M., Binnie, A., Lee, B. A., and Proudfoot, N. J. (2002) Promoter proximal splice sites enhance transcription. *Genes Dev.* **16**, 2792–2799

60. Juneau, K., Miranda, M., Hillenmeyer, M. E., Nislow, C., and Davis, R. W. (2006) Introns regulate RNA and protein abundance in yeast. *Genetics*. **174**, 511–518

61. Kornblihtt, A. R., de la Mata, M., Fededa, J. P., Munoz, M. J., and Nogues, G. (2004) Multiple links between transcription and splicing. *RNA N. Y. N*. **10**, 1489–1498

62. Parenteau, J., Durand, M., Véronneau, S., Lacombe, A.-A., Morin, G., Guérin, V., Cecez, B., Gervais-Bird, J., Koh, C.-S., Brunelle, D., Wellinger, R. J., Chabot, B., and Abou Elela, S. (2008) Deletion of many yeast introns reveals a minority of genes that require splicing for function. *Mol. Biol. Cell*. **19**, 1932–1941

63. Shabalina, S. A., Ogurtsov, A. Y., Spiridonov, A. N., Novichkov, P. S., Spiridonov, N. A., and Koonin, E. V. (2010) Distinct patterns of expression and evolution of intronless and intron-containing mammalian genes. *Mol. Biol. Evol.* **27**, 1745–1749

FIGURES

Figure 1

Fig. 1. Sequencing of complete, chromatin-associated pre-mRNA during inflammatory stimulus reveals differential splicing dynamics among introns of IKBα. (A) Histogram of reads corresponding to the TNF-induced expression and splicing of IKBα pre-mRNA of BMDMs. RNA-Seq was performed on chromatin associated RNA, enriched for NFkB genes as a function of a TNF stimulation timecourse, time shown in minutes after stimulation. Reads are histogrammed in $\log_{10}$ scale and normalized to each time point's maximum value. (B) The Coefficient of Splicing (CoSI) metric quantifies extent of splicing as a function of time, expressed as a ratio of reads from each splice junction to total junctional reads. Dynamics of IKBα splicing as a function of each intron's CoSI is shown (C), where 1=spliced and 0=unspliced, with corresponding introns highlighted in sample timepoint. (D) Differential dynamics of splicing for each *Nfikbia* intron are further demonstrated in the coverage plot for the transcript.

Figure 2

Fig. 2. Heterogeneity of splicing at each intron reveals splicing 'bottlenecks'. The Co-SI of each intron per timepoint is shown as a function of the entire inflammatory mRNA dataset as box-whisker plot (A). Each point represents an intron of one of 230 genes, revealing high rates of splicing (median CoSI indicated by bar near 1.0 for each timepoint) for most genes with significant outliers. As an example, Cxcl10 intron 2 (red arrowhead) is represented by the datapoint with arrowhead, and a histogram of reads is shown to demonstrate relative unspliced nature of this intron, which is not involved in alternative splicing. (B) Several similar introns that are relatively unspliced are found throughout the inflammatory transcriptome; shown are bottleneck introns within Cd40, Daxx, and Irf7 as examples in the context of their neighboring introns.

Figure 3

Fig. 3. Splicing kinetics of inflammatory introns are heterogeneous, ranging from seconds to minutes. CoSI of introns representing various splicing rates are measured and fit to half-lives. Cells were treated with Actinomycin D-treated, from which hybrid capture of genes of interest and sequencing was performed on total (unfractionated) RNA. Shown are four representative samples of splicing kinetics.

Figure 4

Fig. 4. Bottleneck introns can be repaired, and account for significant alterations to gene expression. (A) Intron-GFP splicing reporters for each wild-type intron (red) and modified intron (green) are shown as BFP:GFP ratio. (B) Ratio of WT:Fixed slopes is shown; whereas Tfec expression is not altered by improved 5' sequence, Malt1 intron sequence is significantly impaired owing to its 5' donor sequence, exhibiting a roughly 5-fold impairment in gene expression due to the 5' splice donor.

Figure 5

Fig. 5. Fine-tuning improves gene expression prediction in macrophages. (A) Training and test loss curves over 20 epochs of model fine-tuning. (B-C) Predicted versus measured RNA expression values for 230 transcripts induced by TNFα in macrophages, shown for the pretrained model (B) and the fine-tuned model (C).

Fig. 6. Putative regulatory sequences are enriched in slow-splicing introns (A) Scatterplot showing percent representation of position weight matrices (PWMs) scanned across slow-splicing introns versus all introns genome-wide using FIMO. Sequence logos of the top 5 enriched PWMs are shown to the right. (B–D) Attribution plots highlighting GA-rich sequences (source seqlets for enriched PWMs) and their locations mapped to gene schematics for GA-rich motif (B), and A-rich motif (C). Corresponding RNA-seq tracks are shown for each gene.

SUPPLEMENTAL MATERIALS

Table 1. Oligos used

Oligo	Sequence	Use
d14-952	5'-AAGCAGTGGTATCAACGCAGAGTACACArGrGrG	Template-Switch Oligo
d14-954	5'-AAGCAGTGGTATCAACGCAGAGTACT(30)	RT dT Oligo
d14-955	5'-AAGCAGTGGTATCAACGCAGAGTAC	Post Pulldown Amplification Oligo
L32-For	5'-AAGCGAAACTGGCGGAAAC	qPCR Primer
L32-Rev	5'-TAACCGATGTTGGGCATCAG	qPCR Primer
NF-KBIA Ex5-6-For	5'-ACGGAGTCAGAATTCACAGAGG	qPCR Primer
NF-KBIA Ex5-6-Rev	5'-CACAAAGACAACAGCCGAATC	qPCR Primer
Tn5MErev	5'-[phos]CTGTCTCTTATACACATCT	Tn5 Adaptor Oligo
Tn5ME-A	5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG	Tn5 Adaptor Oligo

Table 2. Exons of interest for hybrid capture

```
A

chr17  exon    35336336      35337508      .    -    .        gene_id "Tnf"; gene_name "Tnf"; p_id "P2280"; transcript_id "NM_013693"; tss_id "TSS22666";
chr11  exon    81850428      81850954      .    +    .        gene_id "Ccl2"; gene_name "Ccl2"; p_id "P3671"; transcript_id "NM_011333"; tss_id "TSS2627";
chr5   exon    92775665      92776411      .    -    .        gene_id "Cxcl10"; gene_name "Cxcl10"; p_id "P12069"; transcript_id "NM_021274"; tss_id "TSS22149";
chr14  exon    103453729     103455790     .    +    .        gene_id "Irg1"; gene_name "Irg1"; p_id "P21444"; transcript_id "NM_008392"; tss_id "TSS15107";
chr5   exon    91334215      91334964      .    +    .        gene_id "Cxcl2"; gene_name "Cxcl2"; p_id "P16237"; transcript_id "NM_009140"; tss_id "TSS25761";
chr12  exon    27127608      27130435      .    -    .        gene_id "Rsad2"; gene_name "Rsad2"; p_id "P15143"; transcript_id "NM_021384"; tss_id "TSS6404";

B

chr12  exon    56590396      56590956      .    -    .        gene_id "Nfkbia"; gene_name "Nfkbia"; p_id "P9152"; transcript_id "NM_010907"; tss_id "TSS10273";
chr11  exon    83477796      83478185      .    +    .        gene_id "Ccl4"; gene_name "Ccl4"; p_id "P10062"; transcript_id "NM_013652"; tss_id "TSS13177";
chr12  exon    112691632     112693229     .    +    .        gene_id "Tnfaip2"; gene_name "Tnfaip2"; p_id "P9235"; transcript_id "NM_009396"; tss_id "TSS24384";
chr5   exon    99646439      99647817      .    -    .        gene_id "Rasgef1b"; gene_name "Rasgef1b"; p_id "P13047"; transcript_id "NM_145839"; tss_id "TSS17562";
chr11  exon    81860487      81861025      .    +    .        gene_id "Ccl7"; gene_name "Ccl7"; p_id "P16317"; transcript_id "NM_013654"; tss_id "TSS4230";
chr10  exon    18720717      18722821      .    -    .        gene_id "Tnfaip3"; gene_name "Tnfaip3"; p_id "P20822"; transcript_id "NM_009397"; tss_id "TSS6673";
chr6   exon    123231807     123233714     .    -    .        gene_id "Clec4e"; gene_name "Clec4e"; p_id "P24237"; transcript_id "NM_019948"; tss_id "TSS20858";
chr16  exon    97668644      97669979      .    -    .        gene_id "Mx1"; gene_name "Mx1"; p_id "P12190"; transcript_id "NM_010846"; tss_id "TSS16276";

C

chr11  exon    16871209      16874156      .    -    .        gene_id "Plek"; gene_name "Plek"; p_id "P24391"; transcript_id "NM_019549"; tss_id "TSS3895";
chr1   exon    146091799     146092617     .    -    .        gene_id "Rgs1"; gene_name "Rgs1"; p_id "P5146"; transcript_id "NM_013642"; tss_id "TSS11168";
chr9   exon    20832215      20833240      .    +    .        gene_id "Icam1"; gene_name "Icam1"; p_id "P10266"; transcript_id "NM_010493"; tss_id "TSS13118";
chr17  exon    57246464      57247180      .    +    .        gene_id "Tnfsf9"; gene_name "Tnfsf9"; p_id "P625"; transcript_id "NM_009404"; tss_id "TSS8246";
chr2   exon    164249747     164251737     .    -    .        gene_id "Sdc4"; gene_name "Sdc4"; p_id "P1421"; transcript_id "NM_011521"; tss_id "TSS724";
chr1   exon    192994176     192995558     .    -    .        gene_id "Atf3"; gene_name "Atf3"; p_id "P15464"; transcript_id "NM_007498"; tss_id "TSS6664";
chr1   exon    132950281     132951812     .    -    .        gene_id "Mapkapk2"; gene_name "Mapkapk2"; p_id "P16800"; transcript_id "NM_008551"; tss_id "TSS23282";
chr11  exon    83339281      83339566      .    -    .        gene_id "Ccl5"; gene_name "Ccl5"; p_id "P8842"; transcript_id "NM_013653"; tss_id "TSS5049";
chr16  exon    55811490      55812896      .    -    .        gene_id "Nfkbiz"; gene_name "Nfkbiz"; p_id "P11871"; transcript_id "NM_001159395"; tss_id "TSS16396";
chr12  exon    27162877      27164702      .    +    .        gene_id "Cmpk2"; gene_name "Cmpk2"; p_id "P8441"; transcript_id "NM_020557"; tss_id "TSS7390";

D

chr11  exon    53590874      53591876      .    +    .        gene_id "Irf1"; gene_name "Irf1"; p_id "P1653"; transcript_id "NM_001159393"; tss_id "TSS23488";
chr17  exon    26642536      26643595      .    -    .        gene_id "Dusp1"; gene_name "Dusp1"; p_id "P21593"; transcript_id "NM_013642"; tss_id "TSS23195";
chr7   exon    29161803      29163512      .    -    .        gene_id "Zfp36"; gene_name "Zfp36"; p_id "P20599"; transcript_id "NM_011756"; tss_id "TSS22952";
chr2   exon    129190316     129190974     .    -    .        gene_id "Il1b"; gene_name "Il1b"; p_id "P19892"; transcript_id "NM_008361"; tss_id "TSS14421";
chr13  exon    43896926      43898501      .    +    .        gene_id "Cd83"; gene_name "Cd83"; p_id "P20497"; transcript_id "NM_009856"; tss_id "TSS21514";
chr13  exon    111190241     111191051     .    +    .        gene_id "Plk2"; gene_name "Plk2"; p_id "P20272"; transcript_id "NM_152804"; tss_id "TSS25357";
chr11  exon    59379666      59380458      .    +    .        gene_id "Nlrp3"; gene_name "Nlrp3"; p_id "P12031"; transcript_id "NM_145827"; tss_id "TSS2239";
chr2   exon    102651300     102654494     .    -    .        gene_id "Cd44"; gene_name "Cd44"; p_id "P1048"; transcript_id "NM_009851"; tss_id "TSS25791";
chr9   exon    110957339     110958943     .    -    .        gene_id "Ccrl2"; gene_name "Ccrl2"; p_id "P24054"; transcript_id "NM_017466"; tss_id "TSS5201";
chr2   exon    180962320     180963574     .    -    .        gene_id "Helz2"; gene_name "Helz2"; p_id "P12385"; transcript_id "NM_183162"; tss_id "TSS196";
chr6   exon    65581071      65584034      .    +    .        gene_id "Tnip3"; gene_name "Tnip3"; p_id "P15078"; transcript_id "NM_001001495"; tss_id "TSS23603";
chr16  exon    35832964      35835021      .    +    .        gene_id "Parp14"; gene_name "Parp14"; p_id "P13379"; transcript_id "NM_001039530"; tss_id "TSS5442";
chr3   exon    83640194      83642711      .    -    .        gene_id "Tlr2"; gene_name "Tlr2"; p_id "P3963"; transcript_id "NM_011905"; tss_id "TSS4833";
chr3   exon    142184004     142185308     .    +    .        gene_id "Gbp5"; gene_name "Gbp5"; p_id "P21165"; transcript_id "NM_153564"; tss_id "TSS9418";
chr14  exon    103375798     103381854     .    -    .        gene_id "Kctd12"; gene_name "Kctd12"; p_id "P13935"; transcript_id "NM_177715"; tss_id "TSS7235";
chr12  exon    40929716      40930038      .    -    .        gene_id "Ifrd1"; gene_name "Ifrd1"; p_id "P14611"; transcript_id "NM_013562"; tss_id "TSS25090";
chr7   exon    52778288      52778692      .    -    .        gene_id "Ppp1r15a"; gene_name "Ppp1r15a"; p_id "P4705"; transcript_id "NM_008654"; tss_id "TSS10655";
chr5   exon    91321534      91322141      .    +    .        gene_id "Cxcl1"; gene_name "Cxcl1"; p_id "P4705"; transcript_id "NM_008176"; tss_id "TSS18478";
chr5   exon    139285062     139286017     .    +    .        gene_id "Fam20c"; gene_name "Fam20c"; p_id "P8478"; transcript_id "NM_030565"; tss_id "TSS2047";
chr4   exon    134769885     134773180     .    -    .        gene_id "Clic4"; gene_name "Clic4"; p_id "P11568"; transcript_id "NM_013885"; tss_id "TSS10071";
chr16  exon    97781946      97782506      .    +    .        gene_id "Mx2"; gene_name "Mx2"; p_id "P20533"; transcript_id "NM_013606"; tss_id "TSS15654";

E
```

```
chr7    exon    128207581   128209031   .   .   -   .   gene_id "Igsf6"; gene_name "Igsf6"; p_id "P11261"; transcript_id "NM_030691"; tss_id "TSS16286";
chr2    exon    62433850    62436145    .   .   -   .   gene_id "Ifih1"; gene_name "Ifih1"; p_id "P4870"; transcript_id "NM_027835"; tss_id "TSS19275";
chr19   exon    6298074     6300096     .   .   +   .   gene_id "Ehd1"; gene_name "Ehd1"; p_id "P19125"; transcript_id "NM_010119"; tss_id "TSS24002";
chr19   exon    34721961    34724499    .   .   +   .   gene_id "Tfit1"; gene_name "Tfit1"; p_id "P22750"; transcript_id "NM_008331"; tss_id "TSS13500";
chr11   exon    117827401   117829632   .   .   .   .   gene_id "Socs3"; gene_name "Socs3"; p_id "P3923"; transcript_id "NM_007707"; tss_id "TSS26657";
chr11   exon    21047960    21050330    .   .   +   .   gene_id "Peli1"; gene_name "Peli1"; p_id "P23395"; transcript_id "NM_023324"; tss_id "TSS17987";
chr8    exon    73153293    73154073    .   .   -   .   gene_id "Gdf15"; gene_name "Gdf15"; p_id "P5185"; transcript_id "NM_011819"; tss_id "TSS7076";
chr8    exon    87500808    87502647    .   .   -   .   gene_id "Junb"; gene_name "Junb"; p_id "P14294"; transcript_id "NM_008416"; tss_id "TSS14734";
chr10   exon    77540638    77542270    .   .   +   .   gene_id "Icosl"; gene_name "Icosl"; p_id "P12741"; transcript_id "NM_015790"; tss_id "TSS25520";
chr2    exon    127983967   127988283   .   .   +   .   gene_id "Bcl2l11"; gene_name "Bcl2l11"; p_id "P21609"; transcript_id "NM_009754"; tss_id "TSS12620";
chr19   exon    29459880    29462584    .   .   +   .   gene_id "Cd274"; gene_name "Cd274"; p_id "P24521"; transcript_id "NM_021893"; tss_id "TSS25796";
chr2    exon    164896222   164897154   .   .   +   .   gene_id "Cd40"; gene_name "Cd40"; p_id "P289"; transcript_id "NM_011611"; tss_id "TSS22026";
chr3    exon    51053619    51055576    .   .   +   .   gene_id "Ccrn4l"; gene_name "Ccrn4l"; p_id "P42"; transcript_id "NM_009834"; tss_id "TSS17093";
chr7    exon    133732809   133733063   .   .   -   .   gene_id "Il27"; gene_name "Il27"; p_id "P17495"; transcript_id "NM_145636"; tss_id "TSS15214";
chr7    exon    148449082   148449279   .   .   -   .   gene_id "Irf7"; gene_name "Irf7"; p_id "P23208"; transcript_id "NM_016850"; tss_id "TSS20549";
chr5    exon    28403331    28405202    .   .   +   .   gene_id "Insig1"; gene_name "Insig1"; p_id "P6141"; transcript_id "NM_153526"; tss_id "TSS14384";
chr18   exon    65635642    65638446    .   .   +   .   gene_id "Malt1"; gene_name "Malt1"; p_id "P17757"; transcript_id "NM_172833"; tss_id "TSS23592";
chr1    exon    135971442   135974280   .   .   -   .   gene_id "Btg2"; gene_name "Btg2"; p_id "P9815"; transcript_id "NM_007570"; tss_id "TSS364";
chr6    exon    129217343   129218387   .   .   +   .   gene_id "Cd69"; gene_name "Cd69"; p_id "P9137"; transcript_id "NM_001033122"; tss_id "TSS434";
chr17   exon    13200705    13200934    .   .   +   .   gene_id "Sod2"; gene_name "Sod2"; p_id "P6002"; transcript_id "NM_013671"; tss_id "TSS18083";
chr7    exon    31213153    31213765    .   .   +   .   gene_id "Nfkbid"; gene_name "Nfkbid"; p_id "P22378"; transcript_id "NM_172142"; tss_id "TSS20875";
chr4    exon    128761072   128761770   .   .   -   .   gene_id "Rnf19b"; gene_name "Rnf19b"; p_id "P5648"; transcript_id "NM_029219"; tss_id "TSS5924";
chr3    exon    142300347   142300972   .   .   +   .   gene_id "Gbp2"; gene_name "Gbp2"; p_id "P5816"; transcript_id "NM_010260"; tss_id "TSS1441";
chr17   exon    35958983    35959856    .   .   +   .   gene_id "Ier3"; gene_name "Ier3"; p_id "P16019"; transcript_id "NM_133662"; tss_id "TSS23149";
chr3    exon    31017377    31021845    .   .   +   .   gene_id "Skil"; gene_name "Skil"; p_id "P16978"; transcript_id "NM_001039090"; tss_id "TSS20990";
chr9    exon    51976136    51976216    .   .   -   .   gene_id "Zc3h12c"; gene_name "Zc3h12c"; p_id "P19129"; transcript_id "NM_001162921"; tss_id "TSS15697";
chr18   exon    35022119    35024610    .   .   +   .   gene_id "Egr1"; gene_name "Egr1"; p_id "P3179"; transcript_id "NM_007913"; tss_id "TSS5272";
chr6    exon    16783381    16784330    .   .   -   .   gene_id "Tfec"; gene_name "Tfec"; p_id "P3938"; transcript_id "NM_031198"; tss_id "TSS12610";
chr2    exon    68946499    68949347    .   .   +   .   gene_id "Cers6"; gene_name "Cers6"; p_id "P23142"; transcript_id "NM_172856"; tss_id "TSS24901";
chr17   exon    45698951    45700117    .   .   +   .   gene_id "Nfkbie"; gene_name "Nfkbie"; p_id "P16819"; transcript_id "NM_008690"; tss_id "TSS2517";
chr11   exon    81916612    81916901    .   .   +   .   gene_id "Ccl12"; gene_name "Ccl12"; p_id "P21246"; transcript_id "NM_011331"; tss_id "TSS4090";
chr17   exon    34052288    34052535    .   .   +   .   gene_id "Daxx"; gene_name "Daxx"; p_id "P7384"; transcript_id "NM_007829"; tss_id "TSS7741";
chr14   exon    21661496    21662610    .   .   +   .   gene_id "Plau"; gene_name "Plau"; p_id "P5851"; transcript_id "NM_008873"; tss_id "TSS14484";
chr10   exon    61957176    61957854    .   .   -   .   gene_id "Srgn"; gene_name "Srgn"; p_id "P15966"; transcript_id "NM_011157"; tss_id "TSS17012";
chr11   exon    44226726    44227519    .   .   +   .   gene_id "Il12b"; gene_name "Il12b"; p_id "P19893"; transcript_id "NM_008352"; tss_id "TSS584";

F

chr11   exon    97850894    97851653    .   .   -   .   gene_id "Arl5c"; gene_name "Arl5c"; p_id "P14239"; transcript_id "NM_207231"; tss_id "TSS9964";
chr2    exon    83643397    83647073    .   .   +   .   gene_id "Itgav"; gene_name "Itgav"; p_id "P22324"; transcript_id "NM_008402"; tss_id "TSS21510";
chr1    exon    151952503   151955142   .   .   +   .   gene_id "Ptgs2"; gene_name "Ptgs2"; p_id "P8580"; transcript_id "NM_011198"; tss_id "TSS16611";
chr4    exon    102277579   102279867   .   .   +   .   gene_id "Pde4b"; gene_name "Pde4b"; p_id "P3538"; transcript_id "NM_001177981"; tss_id "TSS995";
chr9    exon    5336535     5336791     .   .   +   .   gene_id "Casp4"; gene_name "Casp4"; p_id "P4979"; transcript_id "NM_007609"; tss_id "TSS769";
chr3    exon    107543946   107546063   .   .   -   .   gene_id "Csf1"; gene_name "Csf1"; p_id "P4042"; transcript_id "NM_007778"; tss_id "TSS22705";
chr19   exon    34647557    34651024    .   .   +   .   gene_id "Ifit2"; gene_name "Ifit2"; p_id "P8949"; transcript_id "NM_008332"; tss_id "TSS2685";
chr19   exon    32197217    32199014    .   .   +   .   gene_id "Sgms1"; gene_name "Sgms1"; p_id "P20224"; transcript_id "NM_001168526"; tss_id "TSS7412";
chr3    exon    142235438   142236176   .   .   +   .   gene_id "Gbp3"; gene_name "Gbp3"; p_id "P13206"; transcript_id "NM_018734"; tss_id "TSS15170";
chr11   exon    48908891    48910489    .   .   +   .   gene_id "Ifi47"; gene_name "Ifi47"; p_id "P1015"; transcript_id "NM_008330"; tss_id "TSS10281";
chr17   exon    57343396    57343577    .   .   -   .   gene_id "C3"; gene_name "C3"; p_id "P2319"; transcript_id "NM_009778"; tss_id "TSS1016";
chr2    exon    24204935    24207011    .   .   +   .   gene_id "Il1rn"; gene_name "Il1rn"; p_id "P526"; transcript_id "NM_001039701"; tss_id "TSS6957";
chr8    exon    35881452    35882948    .   .   +   .   gene_id "Dusp4"; gene_name "Dusp4"; p_id "P7460"; transcript_id "NM_176933"; tss_id "TSS1452";
chr9    exon    7282018     7283333     .   .   +   .   gene_id "Mmp13"; gene_name "Mmp13"; p_id "P13295"; transcript_id "NM_008607"; tss_id "TSS15376";
chr2    exon    127163317   127164113   .   .   +   .   gene_id "Dusp2"; gene_name "Dusp2"; p_id "P9444"; transcript_id "NM_010090"; tss_id "TSS2906";
chr13   exon    75908128    75909806    .   .   +   .   gene_id "Ell2"; gene_name "Ell2"; p_id "P9427"; transcript_id "NM_001205052"; tss_id "TSS19027";
chr12   exon    86979723    86980828    .   .   +   .   gene_id "Jdp2"; gene_name "Jdp2"; p_id "P1564"; transcript_id "NM_001205052"; tss_id "TSS2002";
chr11   exon    122241637   122243201   .   .   -   .   gene_id "Fnbp1l"; gene_name "Fnbp1l"; p_id "P23393"; transcript_id "NM_001114665"; tss_id "TSS1974";
chr4    exon    16050522    16051038    .   .   +   .   gene_id "Ripk2"; gene_name "Ripk2"; p_id "P9162"; transcript_id "NM_138952"; tss_id "TSS16489";
chr2    exon    129191961   129193225   .   .   +   .   gene_id "Marcksl1"; gene_name "Marcksl1"; p_id "P12626"; transcript_id "NM_010807"; tss_id "TSS2577";
chr10   exon    39753341    39757210    .   .   -   .   gene_id "Slc16a10"; gene_name "Slc16a10"; p_id "P791"; transcript_id "NM_001114332"; tss_id "TSS14295";
chr2    exon    34798778    34799611    .   .   +   .   gene_id "Traf1"; gene_name "Traf1"; p_id "P23707"; transcript_id "NM_009421"; tss_id "TSS6544";
chr3    exon    10204343    10204570    .   .   -   .   gene_id "Fabp4"; gene_name "Fabp4"; p_id "P2902"; transcript_id "NM_024406"; tss_id "TSS6338";
chr7    exon    109706436   109708177   .   .   -   .   gene_id "Trim21"; gene_name "Trim21"; p_id "P23556"; transcript_id "NM_001082552"; tss_id "TSS11938";
chr19   exon    29386231    29387570    .   .   +   .   gene_id "Jak2"; gene_name "Jak2"; p_id "P5298"; transcript_id "NM_001048177"; tss_id "TSS16849";
chr4    exon    150240425   150242989   .   .   +   .   gene_id "Errfi1"; gene_name "Errfi1"; p_id "P9372"; transcript_id "NM_133753"; tss_id "TSS4716";
chr9    exon    119880284   119882329   .   .   +   .   gene_id "Csrnp1"; gene_name "Csrnp1"; p_id "P1038"; transcript_id "NM_153287"; tss_id "TSS10788";
chr5    exon    136623332   136626822   .   .   -   .   gene_id "Orai2"; gene_name "Orai2"; p_id "P8692"; transcript_id "NM_178751"; tss_id "TSS11015";
chr15   exon    36169689    36171820    .   .   -   .   gene_id "Rnf19a"; gene_name "Rnf19a"; p_id "P15766"; transcript_id "NM_001033923"; tss_id "TSS3635";
chr12   exon    86816767    86818219    .   .   +   .   gene_id "Fos"; gene_name "Fos"; p_id "P6877"; transcript_id "NM_010234"; tss_id "TSS15173";
chr5    exon    115386927   115387920   .   .   +   .   gene_id "Oas1l"; gene_name "Oas1l"; p_id "P22147"; transcript_id "NM_145209"; tss_id "TSS4493";
chr9    exon    20834826    20834975    .   .   +   .   gene_id "Icam4"; gene_name "Icam4"; p_id "P16107"; transcript_id "NM_023892"; tss_id "TSS5040";
chrX    exon    106393115   106395784   .   .   +   .   gene_id "Gm6377"; gene_name "Gm6377"; p_id "P22506"; transcript_id "NM_001037917"; tss_id "TSS20699";
chr8    exon    11453977    11454526    .   .   +   .   gene_id "Rab20"; gene_name "Rab20"; p_id "P9178"; transcript_id "NM_011227"; tss_id "TSS9231";
chr4    exon    155573533   155574175   .   .   +   .   gene_id "Isg15"; gene_name "Isg15"; p_id "P6589"; transcript_id "NM_015783"; tss_id "TSS19779";
chr7    exon    25260258    25260892    .   .   +   .   gene_id "Plaur"; gene_name "Plaur"; p_id "P9755"; transcript_id "NM_011113"; tss_id "TSS24418";
chr1    exon    64082247    64082338    .   .   -   .   gene_id "Klf7"; gene_name "Klf7"; p_id "P9777"; transcript_id "NM_033563"; tss_id "TSS7500";
chr1    exon    175677425   175677751   .   .   -   .   gene_id "Ifi204"; gene_name "Ifi204"; p_id "P15649"; transcript_id "NM_008329"; tss_id "TSS17138";
chr4    exon    135700795   135701307   .   .   +   .   gene_id "Id3"; gene_name "Id3"; p_id "P21755"; transcript_id "NM_008321"; tss_id "TSS23418";
chr19   exon    11122102    11122583    .   .   -   .   gene_id "AW112010"; gene_name "AW112010"; p_id "P21084"; transcript_id "NM_001177351"; tss_id "TSS12924";

G

chr16   exon    95940530    95942656    .   .   +   .   gene_id "Ets2"; gene_name "2015-03-02Ets2"; p_id "P20783"; transcript_id "NM_011809"; tss_id "TSS12733";
chr11   exon    78773156    78773626    .   .   +   .   gene_id "Nos2"; gene_name "Nos2"; p_id "P23999"; transcript_id "NM_010927"; tss_id "TSS5656";
chr8    exon    87185230    87186751    .   .   +   .   gene_id "Ier2"; gene_name "Ier2"; p_id "P3914"; transcript_id "NM_010499"; tss_id "TSS11043";
chr6    exon    13785074    13787073    .   .   +   .   gene_id "Gpr85"; gene_name "Gpr85"; p_id "P9006"; transcript_id "NM_145066"; tss_id "TSS13111";
chr2    exon    157942141   157943687   .   .   -   .   gene_id "Tgm2"; gene_name "Tgm2"; p_id "P20583"; transcript_id "NM_009373"; tss_id "TSS2703";
chr11   exon    58019546    58021094    .   .   +   .   gene_id "Igtp"; gene_name "Igtp"; p_id "P10081"; transcript_id "NM_018738"; tss_id "TSS3154";
chr4    exon    53562861    53563036    .   .   +   .   gene_id "Slc44a1"; gene_name "Slc44a1"; p_id "P9505"; transcript_id "NM_001159633"; tss_id "TSS19315";
chr13   exon    114382387   114385070   .   .   -   .   gene_id "Snx18"; gene_name "Snx18"; p_id "P18333"; transcript_id "NM_130796"; tss_id "TSS9534";
chr10   exon    12659787    12661787    .   .   -   .   gene_id "Stx11"; gene_name "Stx11"; p_id "P1435"; transcript_id "NM_001163590"; tss_id "TSS20471";
chr8    exon    91211459    91212373    .   .   +   .   gene_id "Nod2"; gene_name "Nod2"; p_id "P13132"; transcript_id "NM_145857"; tss_id "TSS7189";
chr7    exon    53967368    53967634    .   .   +   .   gene_id "Saa3"; gene_name "Saa3"; p_id "P5870"; transcript_id "NM_011315"; tss_id "TSS9385";
chr6    exon    57612694    57615130    .   .   +   .   gene_id "Herc6"; gene_name "Herc6"; p_id "P9678"; transcript_id "NM_025992"; tss_id "TSS14171";
chr6    exon    121220490   121220935   .   .   +   .   gene_id "Usp18"; gene_name "Usp18"; p_id "P19317"; transcript_id "NM_011909"; tss_id "TSS3918";
chr10   exon    67002493    67004936    .   .   +   .   gene_id "Egr2"; gene_name "Egr2"; p_id "P18543"; transcript_id "NM_010118"; tss_id "TSS12595";
chr5    exon    30345921    30346508    .   .   +   .   gene_id "Il6"; gene_name "Il6"; p_id "P9970"; transcript_id "NM_031168"; tss_id "TSS6882";
chr7    exon    108430964   108432919   .   .   +   .   gene_id "Fchsd2"; gene_name "Fchsd2"; p_id "P13022"; transcript_id "NM_001146010"; tss_id "TSS1726";
chr6    exon    146598633   146599122   .   .   +   .   gene_id "Med21"; gene_name "Med21"; p_id "P199"; transcript_id "NM_025315"; tss_id "TSS1643";
chr11   exon    58001241    58003700    .   .   +   .   gene_id "Gm12250"; gene_name "Gm12250"; p_id "P6220"; transcript_id "NM_001135115"; tss_id "TSS1143";
chr1    exon    175576122   175578059   .   .   +   .   gene_id "Pyhin1"; gene_name "Pyhin1"; p_id "P19336"; transcript_id "NM_175026"; tss_id "TSS14108";
chr10   exon    33845873    33846372    .   .   -   .   gene_id "Fam26f"; gene_name "Fam26f"; p_id "P10436"; transcript_id "NM_175449"; tss_id "TSS7315";
chr7    exon    91257866    91259311    .   .   -   .   gene_id "Fam108c"; gene_name "Fam108c"; p_id "P22064"; transcript_id "NM_133722"; tss_id "TSS1034";
chr5    exon    20881072    20884204    .   .   +   .   gene_id "Fgl2"; gene_name "Fgl2"; p_id "P19695"; transcript_id "NM_008013"; tss_id "TSS20933";
chr17   exon    8125748     8127762     .   .   +   .   gene_id "Tagap"; gene_name "Tagap"; p_id "P8999"; transcript_id "NM_145968_1"; tss_id "TSS4248";
chr17_random    exon    39677   41691   .   .   -   .   gene_id "Tagap"; gene_name "Tagap"; p_id "P23244"; transcript_id "NM_145968"; tss_id "TSS6373";
chr7    exon    29551263    29551543    .   .   -   .   gene_id "Nfkbib"; gene_name "Nfkbib"; p_id "P10826"; transcript_id "NM_010908"; tss_id "TSS22699";
chr2    exon    129125346   129126635   .   .   +   .   gene_id "Il1a"; gene_name "Il1a"; p_id "P9941"; transcript_id "NM_010554"; tss_id "TSS14538";
chr19   exon    34661151    34663472    .   .   +   .   gene_id "Ifit3"; gene_name "Ifit3"; p_id "P3803"; transcript_id "NM_010501"; tss_id "TSS22430";
chr2    exon    122286156   122286866   .   .   +   .   gene_id "Slc28a2"; gene_name "Slc28a2"; p_id "P8445"; transcript_id "NM_172980"; tss_id "TSS16659";
chrX    exon    166412975   166414722   .   .   -   .   gene_id "G530011006Rik"; gene_name "G530011006Rik"; p_id "P24135"; transcript_id "NR_029457"; tss_id "TSS23870";
chr15   exon    79187851    79189506    .   .   +   .   gene_id "Maff"; gene_name "Maff"; p_id "P21898"; transcript_id "NM_010755"; tss_id "TSS7572";
chr7    exon    52094953    52096179    .   .   +   .   gene_id "Il4i1"; gene_name "Il4i1"; p_id "P14475"; transcript_id "NM_010215"; tss_id "TSS21834";
chr18   exon    60549446    60552283    .   .   +   .   gene_id "Iigp1"; gene_name "Iigp1"; p_id "P24220"; transcript_id "NM_001146275"; tss_id "TSS22447";
chr3    exon    142209284   142213047   .   .   +   .   gene_id "Gbp7"; gene_name "Gbp7"; p_id "P10586"; transcript_id "NM_001083312"; tss_id "TSS23222";
chr4    exon    88167929    88168698    .   .   -   .   gene_id "Ifnb1"; gene_name "Ifnb1"; p_id "P21845"; transcript_id "NM_010510"; tss_id "TSS25442";
chr3    exon    50168858    50176288    .   .   +   .   gene_id "Slc7a11"; gene_name "Slc7a11"; p_id "P4123"; transcript_id "NM_011990"; tss_id "TSS24352";
chr16   exon    75859039    75859884    .   .   -   .   gene_id "Samsn1"; gene_name "Samsn1"; p_id "P14457"; transcript_id "NM_019578"; tss_id "TSS20090";
chr7    exon    81420304    81420974    .   .   +   .   gene_id "Slco3a1"; gene_name "Slco3a1"; p_id "P121"; transcript_id "NM_001038643"; tss_id "TSS8114";
chr11   exon    69212020    69213351    .   .   -   .   gene_id "Kdm6b"; gene_name "Kdm6b"; p_id "P9055"; transcript_id "NM_001017426"; tss_id "TSS18373";
chr10   exon    13868718    13871184    .   .   -   .   gene_id "Hivep2"; gene_name "Hivep2"; p_id "P22703"; transcript_id "NM_010437"; tss_id "TSS6381";
chr9    exon    92166367    92167399    .   .   +   .   gene_id "Plscr1"; gene_name "Plscr1"; p_id "P17054"; transcript_id "NM_011636"; tss_id "TSS11480";
chr6    exon    54873936    54875003    .   .   +   .   gene_id "Nod1"; gene_name "Nod1"; p_id "P17761"; transcript_id "NM_172729"; tss_id "TSS13431";
chr5    exon    137537376   137539081   .   .   -   .   gene_id "Serpine1"; gene_name "Serpine1"; p_id "P16074"; transcript_id "NM_008871"; tss_id "TSS17025";
chr17   exon    18029697    18030916    .   .   +   .   gene_id "Fpr2"; gene_name "Fpr2"; p_id "P200"; transcript_id "NM_008039"; tss_id "TSS20689";
chr10   exon    44156981    44160048    .   .   -   .   gene_id "Prdm1"; gene_name "Prdm1"; p_id "P5182"; transcript_id "NM_007548"; tss_id "TSS16926";
chr8    exon    28249245    28253866    .   .   -   .   gene_id "Rab11fip1"; gene_name "Rab11fip1"; p_id "P220"; transcript_id "NM_001080813"; tss_id "TSS19574";
chr9    exon    44494391    44495781    .   .   +   .   gene_id "Phldb1"; gene_name "Phldb1"; p_id "P8634"; transcript_id "NM_153537"; tss_id "TSS9456";
chr15   exon    103138666   103140087   .   .   -   .   gene_id "Gpr84"; gene_name "Gpr84"; p_id "P18218"; transcript_id "NM_030720"; tss_id "TSS17218";
chr5    exon    124313579   124315525   .   .   +   .   gene_id "Niacr1"; gene_name "Niacr1"; p_id "P4625"; transcript_id "NM_030701"; tss_id "TSS707";
chr8    exon    86734841    86736704    .   .   +   .   gene_id "Zswim4"; gene_name "Zswim4"; p_id "P24135"; transcript_id "NM_172503"; tss_id "TSS887";
chr8    exon    92750357    92752932    .   .   +   .   gene_id "Cxcl9"; gene_name "Cxcl9"; p_id "P19130"; transcript_id "NM_008599"; tss_id "TSS23167";
chr8    exon    69000245    69000808    .   .   +   .   gene_id "Tma16"; gene_name "Tma16"; p_id "P22272"; transcript_id "NM_025465"; tss_id "TSS4033";
chr15   exon    61820901    61821916    .   .   +   .   gene_id "Myc"; gene_name "Myc"; p_id "P18425"; transcript_id "NM_010849"; tss_id "TSS15915";
chr4    exon    63385637    63391077    .   .   +   .   gene_id "Tnfsf15"; gene_name "Tnfsf15"; p_id "P12252"; transcript_id "NM_177371"; tss_id "TSS3724";
chr19   exon    53615406    53615812    .   .   +   .   gene_id "Dusp5"; gene_name "Dusp5"; p_id "P6646"; transcript_id "NM_001085390"; tss_id "TSS19363";
chr2    exon    173032114   173032727   .   .   +   .   gene_id "Zbp1"; gene_name "Zbp1"; p_id "P4281"; transcript_id "NM_021394"; tss_id "TSS13610";
chr4    exon    15924268    15926147    .   .   +   .   gene_id "Osgin2"; gene_name "Osgin2"; p_id "P24566"; transcript_id "NM_145950"; tss_id "TSS5672";
chr4    exon    11640623    11642140    .   .   +   .   gene_id "Gem"; gene_name "Gem"; p_id "P845"; transcript_id "NM_010276"; tss_id "TSS16095";
chr5    exon    92788571    92789832    .   .   +   .   gene_id "Cxcl11"; gene_name "Cxcl11"; p_id "P2702"; transcript_id "NM_019494"; tss_id "TSS8759";
chr15   exon    59485791    59488654    .   .   +   .   gene_id "Trib1"; gene_name "Trib1"; p_id "P21350"; transcript_id "NM_144549"; tss_id "TSS14254";
chr11   exon    48947790    48948889    .   .   +   .   gene_id "Olfr56"; gene_name "Olfr56"; p_id "P9227"; transcript_id "NM_010999"; tss_id "TSS4614";
chr19   exon    34548274    34549625    .   .   +   .   gene_id "Ch25h"; gene_name "Ch25h"; p_id "P882"; transcript_id "NM_009890"; tss_id "TSS17189";
chr19   exon    6171306     6172475     .   .   +   .   gene_id "Batf2"; gene_name "Batf2"; p_id "P13629"; transcript_id "NM_028967"; tss_id "TSS15611";
chr2    exon    11654777    11655612    .   .   +   .   gene_id "Il15ra"; gene_name "Il15ra"; p_id "P10569"; transcript_id "NM_133836"; tss_id "TSS13507";
chr17   exon    12411851    12412506    .   .   +   .   gene_id "Agpat4"; gene_name "Agpat4"; p_id "P20641"; transcript_id "NM_026644"; tss_id "TSS3138";
chr18   exon    60405029    60407474    .   .   +   .   gene_id "Gm4951"; gene_name "Gm4951"; p_id "P3089"; transcript_id "NM_001033767"; tss_id "TSS19296";
chr1    exon    175942129   175942419   .   .   -   .   gene_id "Ifi205"; gene_name "Ifi205"; p_id "P4164"; transcript_id "NM_172648"; tss_id "TSS16317";
chr16   exon    92392198    92393873    .   .   -   .   gene_id "Rcan1"; gene_name "Rcan1"; p_id "P16767"; transcript_id "NM_019466"; tss_id "TSS831";
```

```
chr13   exon    89794915        89796963        .       -       .       gene_id "Vcan"; gene_name "Vcan"; p_id "P15421"; transcript_id "NM_001081249"; tss_id "TSS20677";
chr4    exon    155539399       155540762       .       -       .        gene_id "Agrn"; gene_name "Agrn"; p_id "P16425"; transcript_id "NM_021604"; tss_id "TSS15328";

H

chr17   exon    34993319        34993527        .       -       .       gene_id "Cfb"; gene_name "Cfb"; p_id "P13056"; transcript_id "NM_001142706"; tss_id "TSS23526";
chr17   exon    48004082        48005752        .       -       .       gene_id "Foxp4"; gene_name "Foxp4"; p_id "P16134"; transcript_id "NM_001110824"; tss_id "TSS19822";
chr9    exon    107202699       107204292       .       +       .        gene_id "Cish"; gene_name "Cish"; p_id "P8321"; transcript_id "NM_009895"; tss_id "TSS23598";
chr3    exon    27234144        27238587        .       +       .        gene_id "Tnfsf10"; gene_name "Tnfsf10"; p_id "P6001"; transcript_id "NM_009425"; tss_id "TSS7370";
chr14   exon    8782819         8784101         .       +       .      gene_id "Flnb"; gene_name "Flnb"; p_id "P6881"; transcript_id "NM_134080"; tss_id "TSS13225";
chr14   exon    59966244        59967888        .       -       .       gene_id "Phf11d"; gene_name "Phf11d"; p_id "P19480"; transcript_id "NM_199015"; tss_id "TSS13050";
chr3    exon    151393887       151395517       .       -       .        gene_id "Ifi44"; gene_name "Ifi44"; p_id "P4710"; transcript_id "NM_133871"; tss_id "TSS11790";
chr5    exon    105507413       105509283       .       -       .        gene_id "Gbp9"; gene_name "Gbp9"; p_id "P23637"; transcript_id "NM_172777"; tss_id "TSS22932";
chr13   exon    42402146        42403358        .       +       .       gene_id "Edn1"; gene_name "Edn1"; p_id "P15001"; transcript_id "NM_010104"; tss_id "TSS16855";
chr12   exon    72000371        72003221        .       +       .       gene_id "Frmd6"; gene_name "Frmd6"; p_id "P2929"; transcript_id "NM_028127"; tss_id "TSS4836";
chr7    exon    86064932        86065282        .       +       .      gene_id "Isg20"; gene_name "Isg20"; p_id "P23651"; transcript_id "NM_020583"; tss_id "TSS17804";
chr2    exon    122466437       122466812       .       +       .       gene_id "AA467197"; gene_name "AA467197"; p_id "P11166"; transcript_id "NM_001004174"; tss_id "TSS5413";
chr7    exon    83017584        83019016        .       +       .      gene_id "Klhl25"; gene_name "Klhl25"; p_id "P22799"; transcript_id "NM_001122780"; tss_id "TSS12992";
chr11   exon    86582351        86583763        .       -       .       gene_id "Dhx40"; gene_name "Dhx40"; p_id "P23840"; transcript_id "NM_026191"; tss_id "TSS24769";
chr18   exon    66622924        66625212        .       +       .       gene_id "Pmaip1"; gene_name "Pmaip1"; p_id "P7859"; transcript_id "NM_021451"; tss_id "TSS22625";
chr15   exon    57670146        57671387        .       +       .       gene_id "Zhx2"; gene_name "Zhx2"; p_id "P21154"; transcript_id "NM_199449"; tss_id "TSS3741";
chr19   exon    34667378        34669513        .       -       .       gene_id "Gm14446"; gene_name "Gm14446"; p_id "P16440"; transcript_id "NM_001101605"; tss_id "TSS23646";
chr8    exon    10986964        10987014        .       -       .       gene_id "Irs2"; gene_name "Irs2"; p_id "P3108"; transcript_id "NM_001081212"; tss_id "TSS25904";
chr8    exon    108497420       108497751       .       -       .       gene_id "Dpep3"; gene_name "Dpep3"; p_id "P1255"; transcript_id "NM_027960"; tss_id "TSS20475";
chr10   exon    110945116       110945705       .       +       .        gene_id "Phlda1"; gene_name "Phlda1"; p_id "P19275"; transcript_id "NM_009344"; tss_id "TSS19408";
chr2    exon    52990874        52993236        .       +       .      gene_id "Fmnl2"; gene_name "Fmnl2"; p_id "P4156"; transcript_id "NM_172409"; tss_id "TSS2704";
chr11   exon    82934523        82936161        .       +       .       gene_id "Slfn1"; gene_name "Slfn1"; p_id "P22608"; transcript_id "NM_011407"; tss_id "TSS25201";
chr19   exon    11500686        11501736        .       +       .       gene_id "Ms4a4c"; gene_name "Ms4a4c"; p_id "P2294"; transcript_id "NM_029909"; tss_id "TSS21383";
chr5    exon    105544786       105547518       .       -       .        gene_id "Gbp4"; gene_name "Gbp4"; p_id "P24799"; transcript_id "NM_008620"; tss_id "TSS16713";
chr3    exon    19892509        19892998        .       +       .      gene_id "Cp"; gene_name "Cp"; p_id "P6496"; transcript_id "NM_007752"; tss_id "TSS19796";
chr11   exon    83002165        83003718        .       +       .       gene_id "Slfn4"; gene_name "Slfn4"; p_id "P16897"; transcript_id "NM_011410"; tss_id "TSS23253";
chr14   exon    59999670        59999984        .       +       .       gene_id "Phf11c"; gene_name "Phf11c"; p_id "P2484"; transcript_id "NM_001164289"; tss_id "TSS18997";
chr16   exon    11072325        11075078        .       +       .       gene_id "Snn"; gene_name "Snn"; p_id "P22666"; transcript_id "NM_009223"; tss_id "TSS15465";
chr3    exon    32482450        32484148        .       -       .       gene_id "Gnb4"; gene_name "Gnb4"; p_id "P5107"; transcript_id "NM_013531"; tss_id "TSS18140";
chr6    exon    127075727       127080505       .       -       .        gene_id "Ccnd2"; gene_name "Ccnd2"; p_id "P4356"; transcript_id "NM_009829"; tss_id "TSS24028";
chr14   exon    70478703        70479964        .       +       .       gene_id "Egr3"; gene_name "Egr3"; p_id "P10547"; transcript_id "NM_018781"; tss_id "TSS18772";

I

chr3    exon    115729931       115731934       .       +       .       gene_id "Extl2"; gene_name "Extl2"; p_id "P23186"; transcript_id "NM_001163515"; tss_id "TSS13032";
chr15   exon    101104427       101105225       .       +       .        gene_id "Nr4a1"; gene_name "Nr4a1"; p_id "P11564"; transcript_id "NM_010444"; tss_id "TSS1779";
chr11   exon    83325238        83325629        .       -       .       gene_id "Gm11435"; gene_name "Gm11435"; p_id "P3561"; transcript_id "NM_001045543"; tss_id "TSS21732";
chr14   exon    59939801        59940120        .       -       .       gene_id "Phf11b"; gene_name "Phf11b"; p_id "P21005"; transcript_id "NM_001164327"; tss_id "TSS6272";
chr12   exon    17557413        17558308        .       +       .       gene_id "Odc1"; gene_name "Odc1"; p_id "P13979"; transcript_id "NM_013614"; tss_id "TSS312";
chr7    exon    149265392       149268935       .       -       .        gene_id "Gm10941"; gene_name "Gm10941"; transcript_id "NR_026944"; tss_id "TSS17723";
chr10   exon    76720518        76720731        .       -       .       gene_id "Adamts1"; gene_name "Adamts1"; p_id "P4336"; transcript_id "NM_009621"; tss_id "TSS24958";
chr12   exon    40902973        40903815        .       -       .       gene_id "Gm889"; gene_name "Gm889"; p_id "P23562"; transcript_id "NM_001033437"; tss_id "TSS7723";
```

Supplemental Figure 1. Hybrid capture strategy for isolating chromatin-associated inflammatory transcripts.

RNA was purified from chromatin-associated bone marrow–derived macrophages (BMDMs), and cDNA was generated using oligo(dT) priming to enrich for polyadenylated transcripts. Biotinylated RNA oligonucleotides complementary to the terminal exons of inflammatory genes were hybridized to the cDNAs, allowing for selective enrichment of these transcripts via streptavidin bead capture.

Supplemental Figure 2. Hybrid capture enriches NF-κB–responsive transcripts and yields robust intron coverage.

(A) Fraction of sequencing reads corresponding to NF-κB–responsive genes (blue) in hybrid-captured versus poly(A)-selected RNA (orange), showing substantial enrichment after hybrid capture. (B) Histogram of read counts per intron at the 6-min TNF induction time point. Reads were detected for 1,024 introns out of 1,508 targeted introns, with undetected introns largely corresponding to transcripts induced at later time points (>60 min). (C) Distribution of intron fold induction across time points reveals that many NF-κB target genes begin to show induction as early as 4 min post-TNF stimulation, with both the number of induced introns and magnitude of induction increasing markedly by 14 min. (D) Scatterplot comparing induction at 4 min versus 14 min for individual introns demonstrates that most early-induced genes are further upregulated at later time points.

Supplemental Figure 3. Gene track visualization of Nfkbia induction dynamics.

Genome browser tracks show Nfkbia (IκBα) transcript induction over time in chromatin-associated, hybrid-captured RNA. Signal intensity is displayed on a linear scale (left) normalized to the maximum height at 20 min, and on a log scale (right) normalized to the maximum at each time point.

Supplemental Figure 4. Splice site strength and expression kinetics among NF-κB–induced genes.

(A) Representative introns from selected genes were scored for 5′ splice donor strength based on similarity to the canonical 'GTAAG' motif; introns with weaker matches (e.g., Irf7 intron 3) received lower scores. (B) NF-κB–induced genes display well-characterized variability in expression kinetics (RNA-seq data from Reference 25). Heatmap shows temporal expression profiles of NF-κB target genes following lipid A stimulation, categorized into immediate-early, early, and later expression groups. (C) All introns in the NF-κB transcriptome were scored using MaxEntScan for both 5′ splice donor (top) and 3′ splice acceptor (bottom) sequences, stratified by their gene expression group.

Supplemental Figure 5. Reporter assay design for assessing intron splicing efficiency.

(A) Schematic of the bidirectional reporter assay. Individual introns were cloned into a bidirectional promoter context together with their flanking exons, positioned upstream of a self-cleaving 2A peptide and GFP reporter. In the opposite transcriptional direction, a BFP reporter served as a transcriptional control. (B) Evolutionarily conserved weak 5′ splice donors were "repaired" to the canonical GTAAG sequence within this reporter construct, and splicing efficiency was quantified by flow cytometry (FACS).

Supplemental Figure 6. Splicing completion across kinetic gene expression groups.

Box-and-whisker plots depict time-course CoSI values for introns from inflammatory gene cohorts. Each point represents the CoSI value of an individual intron at a given time point. Plots are grouped by expression category: Immediate Early (top left), Early (top right), Intermediate (bottom left), and Late (bottom right).

Supplemental Figure 7. Relationships between intron length, GC content, transcript position, and splicing kinetics.

(A) Scatterplot of intron length (y-axis) versus GC content (%) (x-axis) for the 150 slowest-splicing introns. Pearson r = −0.44 (p = 2.76 × 10⁻⁴⁵) and Spearman r = −0.44 (p = 1.34 × 10⁻⁴⁶). (B) Histogram of intron length distributions (100-nt bins) for all 1,098 introns (red) and the 150 slowest-splicing introns (blue). Length distributions are largely similar between cohorts. (C) Splice completion (CoSI) trajectories across the time course for the 50 fastest (left, blue) and 50 slowest (right, orange) introns, ranked by area under the CoSI curve (AUC). (D) Same analysis as in (C) but extended to the top 150 fastest and bottom 150 slowest introns. (E) Minimum CoSI values for each intron plotted against transcript abundance (TPM, left), intron length (middle), and position along the transcript (5′ → 3′ %, right) for the bottom 150 and bottom 50 introns. Correlations: TPM — Bottom 150: Pearson r = 0.204 (p = 1.24 × 10⁻²), Spearman r = 0.203 (p = 1.27 × 10⁻²); Bottom 50: Pearson r = 0.276 (p = 5.25 × 10⁻²), Spearman r = 0.316 (p = 2.53 × 10⁻²). Length — Bottom 150: Pearson r = −0.242 (p = 2.88 × 10⁻³), Spearman r = −0.266 (p = 9.92 × 10⁻⁴); Bottom 50: Pearson r = −0.160 (p = 0.268), Spearman r = −0.171 (p = 0.235). Transcript position — Bottom 150: Pearson r = 0.260 (p = 1.33 × 10⁻³), Spearman r = 0.231 (p = 4.44 × 10⁻³); Bottom 50: Pearson r = 0.259 (p = 6.90 × 10⁻²), Spearman r = 0.318 (p = 2.43 × 10⁻²).

Supplemental Figure 8. Enrichment of sequence motifs near slow-splicing introns.

Scatterplots show the percent representation of position weight matrices (PWMs) scanned across slow-splicing introns compared to all introns genome-wide using FIMO. Sequence logos of the top five enriched PWMs are displayed to the right of each scatterplot. (A) Scatterplot and enriched motifs identified in exons adjacent to the 50 slowest-splicing introns. (B) Scatterplot and enriched motifs identified within the 150 slowest-splicing introns. (C) Scatterplot and enriched motifs identified in exons adjacent to the 150 slowest-splicing introns.