Bud13 Promotes a Type I Interferon Response by Countering Intron Retention in Irf7

Luke S. Frankiw[1,2], Devdoot Majumdar[1,2], Christian Burns[1], Annie Moradian[1], Michael J. Sweredoski[1], David Baltimore[1,3]*

[1]*Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125, USA*
[2]*These authors contributed equally*
[3]*Lead Contact*
Correspondence: baltimo@caltech.edu

SUMMARY

Intron retention (IR) has emerged as an important mechanism of gene expression control. Despite this, the factors that control IR events remain poorly understood. We observed consistent IR in one intron of the Irf7 gene and identified Bud13 as an RNA-binding protein that acts at this intron to increase the amount of successful splicing. Deficiency in Bud13 led to increased IR, decreased mature Irf7 transcript and protein levels, and consequently to a dampened type I interferon response. This impairment of Irf7 production in Bud13-deficient cells compromised their ability to withstand VSV infection. Global analysis of Bud13 knockdown and BUD13 cross-linking to RNA revealed a subset of introns that share many characteristics with the one found in Irf7 and are spliced in a Bud13-dependent manner. Deficiency of Bud13 led to decreased mature transcript from genes containing such introns. Thus, by acting as an antagonist to IR, Bud13 facilitates the expression of genes at which IR occurs.

INTRODUCTION

Three forms of alternative processing of a pre-mRNA have been described: differential inclusion of an exon, alternative splice site selection, and intron retention (IR). The latter, IR, has emerged as a previously underappreciated mechanism of post-transcriptional gene regulation. Unlike the two alternative splicing events, IR rarely contributes to proteomic diversity.[1] However, IR events have the ability to act as negative regulators of gene expression by: (1) delaying onset of gene expression by slowing down splicing kinetics, (2) increasing potential nuclear degradation by nuclear exosomes, (3) increasing potential cytoplasmic degradation by nonsense mediated decay.[2]

Recent genomic studies suggest IR plays an important role in the regulation of gene expression in a wide range of processes including cellular differentiation[3,4] and tumorigenesis.[5] Further, widespread IR throughout mouse and human cell and tissue types has led to the idea that IR events act to functionally "tune" the transcriptome of a cell.[6] However, with few exceptions, the factors that control IR events and thus potentially shape gene expression programs of cells, remain poorly understood.

Irf7 is an interferon-inducible master regulator of the type-I interferon-dependent immune response and is crucial to the production of interferon-α and β.[7] Aberrant Irf7 production is linked to a wide range of pathologies, from life-threatening influenza[8] to autoimmunity[9], because precise regulation of Irf7 ensures a proper immune response. Notably, intron 4 of Irf7 is short, GC-rich, and has a poor splice donor sequence, characteristics shared by many poorly spliced introns. We and others have previously shown that intron 4 of Irf7 splices inefficiently[10], affecting gene expression and opening a new line of inquiry as to the mechanism of IR regulation in *Irf7*.

Using RNA antisense purification-mass spectrometry (RAP-MS)[11], we identified the protein Bud13 as one that regulates IR in Irf7. Bud13 was found to aid splicing efficiency and expression of the Irf7 mature transcript and protein, thus increasing the downstream type-I interferon-dependent immune response. We show that Irf7 is able to trigger a robust interferon response in the presence but not in the absence of Bud13. Further, Bud13 was found to increase the splicing efficiency of a multitude of other junctions with similar characteristics to the one found in Irf7. By aiding in splicing

136   efficiency, Bud13 limits intron retention and increases gene expression levels of transcripts

137   containing Bud13 dependent junctions.

139   **RESULTS**

140
141   **Irf7 contains an intron that splices poorly following stimulation.**

142
To study the role of mRNA splicing during an innate immune response, we sequenced the

RNA from mouse bone marrow-derived macrophages (BMDMs) stimulated with either TNFα, IFNα,

or Poly(I:C). From this sequencing, we identified an increased number of intronic reads in the fourth

intron of the most abundant transcript of Irf7 independent of time or stimulant (Fig. 1A, I, J). A variety

of features of this intron make it a likely candidate for retention.[6] It is extremely small at 69

nucleotides and has a high G/C content in both the flanking exons and within the intron itself.

Furthermore, the intron contains a 'weak' 5' splice site, one that deviates significantly from a

consensus splice site sequence. This is quantified using a maximum entropy model to calculate the

splice site quality score (Fig. 1F).[12]

To quantify the extent of retention across RNA-seq data-sets, we use a metric we designate

the "splicing ratio" (SR) (Fig. 1G; see methods), which is a length normalized ratio of intronic reads

to total reads at each junction. Low SR values indicate a junction is primarily spliced whereas high

SR values indicate a junction is primarily unspliced. Using this metric, we quantified the extent of

retention for all junctions in the most abundant Irf7 transcript. We observed that for all types of

stimulation, the retention of the fourth intron of the transcript is much greater than that seen for any

of the other introns (Fig. 1H and S1A, B). This intron remains poorly spliced despite the fact that

there is clear excision of neighboring introns. It is worth noting that quantitation of the IFNα

stimulation shows increased intronic signal throughout the Irf7 transcript. This increased intronic

signal is due to faster and stronger induction of Irf7 via stimulation with IFNα and as such, an

increase in the amount of pre-mRNA at a given stimulation time-point. Despite this increase in

intronic signal throughout the transcript, we observed a corresponding increase in the level of

retention for the poorly spliced fourth intron (Fig. 1J, S1B). Thus, we conclude this intron of Irf7

splices poorly following many forms of stimulation.

RAP-MS identifies Bud13 as an RNA binding protein that interacts with IRF7 mRNA.

To understand how cells handle a retained intron, we sought to identify trans-acting proteins

that might affect the process using RNA Antisense Purification followed by Mass Spectrometry

(RAP-MS) (Fig. 2A)[11]. RAP-MS employs antisense biotin-containing ssDNAs complementary to *Irf7*

exons to purify the proteins associated with the total pool of *Irf7* transcripts, containing both nascent

pre-mRNAs and mature mRNA. Using this proteomic approach, we identified the RNA-binding

protein Bud13 to be highly enriched (~6-fold) on Irf7 transcripts as compared to β-*actin* transcripts,

which were used as a control (Fig. 2B). Bud13 has been characterized in yeast as a member of the

Retention and Splicing complex (RES),[13] forming a trimeric complex with Pml1p and Snu17p, and

aids in the splicing and nuclear retention of a subset of transcripts. It is not well characterized in

mammalian systems. We captured a variety of other known RNA-binding proteins (Pum2, Prpf40a,

Son); however, no other protein was enriched greater than two fold on Irf7 transcripts. We observed

specificity in the RNA antisense purification for the intended transcripts (Fig. 2C).

Following RAP-MS, we confirmed Bud13 enrichment on Irf7 transcripts by performing RNA

Immunoprecipitation (RIP) followed by qPCR. Using formaldehyde cross-linked, BMDMs stimulated

with TNFα for 30 minutes or Poly(I:C) for 12 hours, we observed >7-fold enrichment of *Irf7*

transcripts associated with Bud13 immunoprecipitates as compared to Rabbit IgG control

immunoprecipitates (Fig. 2D, E). In contrast, no differential enrichment of Rpl32 was observed.

Thus, isolating the proteins associated with Irf7 mRNA transcripts led to the identification of Bud13,

and immunoprecipitation of Bud13 protein confirmed enrichment of Irf7 mRNA.

Bud13 knockdown leads to increased retention in the weak Irf7 intron.

To determine whether the enrichment of Bud13 had an effect on Irf7 mRNA processing, we

used an shRNA approach to knockdown Bud13 protein levels in BMDMs (Fig. S1A, B). To quantify

differences in splicing between the shBud13 sample and the scrambled control sample, we

calculated the difference in the previously mentioned splicing ratio (SR) metric between shBud13

and control for each junction at each time point. This resulting value was designated the ΔSR. A

positive ΔSR indicates a junction is more unspliced in the shBud13 sample while a negative ΔSR

indicates a junction is more unspliced in the control sample.

RNA-seq was performed on RNA from unstimulated BMDMs, as well as macrophages

stimulated with TNFα for 0, 30, 60, and 120 minutes. Bud13 knockdown led to a further increased

retention of the fourth intron in *Irf7* (Fig. 3A – highlighted intron). Further, the sequencing coverage

plots showed little variation in splicing for the other seven introns in the transcript. This was

confirmed when splicing was quantified using the ΔSR metric (Fig. 3B). At all stimulation time-

points, the ΔSR value for the fourth intron was significantly greater than 0, indicating an increase in

retention when Bud13 levels were reduced. In contrast, the ΔSR values for the other introns in the

Irf7 transcript are negligible. It appears that Bud13 plays a specific role of aiding in the excision of

the poorly spliced junction but is not required for total splicing of other introns in the transcript, at

least as indicated by the partial knockdown with an shRNA. We next looked at how this retention

affected the induction kinetics of Irf7. We observed that the increased retention seen upon

knockdown of Bud13 led to decreased induction of Irf7 mRNA in response to TNFα stimulation (Fig.

3C, S2 C, D, E), consistent with the idea that intron retention leads to transcript degradation.[14]

Bud13 knockdown alters the type I interferon response.

213
Because Irf7 is known as a 'master regulator' for robust type I interferon production[7], we next

investigated the effect of Bud13 knockdown on a type I interferon response. To do so, we stimulated

BMDMs with the TLR3 agonist Poly(I:C) for up to 24 hours. Activation of TLR3 leads to the

production of type I interferons followed by the downstream induction and activation of Irf7, which

serves to amplify the type I interferon response via positive feedback[8]. We again observed

differential splicing between the shBud13 time-points and the control time-points in intron 4 of Irf7

(Fig. 4A). Quantified using the ΔSR metric, five out of the six poly(I:C) time-points have high intron 4

ΔSR values (Fig. 4B). The ΔSR values for the other introns in the Irf7 transcript again fluctuate

around 0, indicating that the fourth intron of Irf7 has a specific dependence on Bud13 for efficient

splicing. As is the case with TNFα, this decrease in splicing efficiency alters Irf7 induction kinetics.

Less Irf7 mRNA is induced at 240, 720, and 1440mins of poly(I:C) stimulation (Fig. 4C). This

reduction in Irf7 mRNA leads to a decrease in nuclear Irf7 protein levels (Fig. 4D).

Next we looked at how this reduction in Irf7 would alter the production of RNA from interferon

signature genes (ISGs). Expression of 119 ISGs (selected based on upregulation in response to

IFNα [15]; see methods) was examined. In unstimulated BMDMs, used as a baseline, the median $\log_2$

expression fold change (FPKM shBud13/ FPKM control) is 0.1655 (Fig. 4E). In contrast, at 720 mins

of stimulation, the median $\log_2$ expression fold change shifts to -0.1007 (Fig. 4F), indicating a

significant decrease in ISG expression in the shBud13 sample compared to the control sample at

this time-point compared to the baseline (Wilcoxon rank-sum, $P< .001$). This significant decrease in

ISG expression remained true when comparing any of the 'early' time points (0, 15, 60 mins) to any

of the 'late' timepoints (240, 720, 1440 mins) (Wilcoxon rank-sum, $P<0.001$). Dampened levels of

Irf7 led to defects in the amplification of the type I interferon response, and a decrease in ISG

expression later in the time-course (Fig. 4G). Trim21, Stat1, Stat2, Ifih1, and Dhx58 are among the

notable ISGs with reduced expression at 720 mins in the shBud13 sample (Fig. 4H). Expression of

both IFNα and IFNβ were significantly reduced at 720 and 1440 mins (Fig. 4I, J). To ensure

differential expression of ISGs was not due to splicing defects from Bud13 knockdown, we quantified

the ΔSR for every ISG junction at 720 mins. The fourth intron of Irf7 has the greatest ΔSR at 0.227.

Only four other junctions of the 375 that were examined have ΔSRs greater than 0.1, and the

majority of junctions have ΔSRs close to 0 (Fig. S3; mean = 0.002, median = 0). Similar results

were obtained when BMDMs were stimulated with the TLR9 agonist CpG (Figure S4). Taken

together, we conclude that Bud13 deficiency results in a highly compromised type I interferon

response.

Finally, we examined whether Irf7 pre-mRNA with a retained fourth intron was able to exit the

nucleus and enter the cytoplasm. BMDMs were stimulated with poly(I:C) and fractionated into a

nuclear fraction (containing the nucleoplasm and chromatin) and a cytoplasmic fraction. RNA-seq

was performed on RNA from the cytoplasmic fraction and the Irf7 mRNA was found to be almost

completely spliced (Fig. 4K, L). In the nucleus, we again observed retention of Irf7 intron 4 (Fig. 4K,

highlighted intron, ΔSR. = 0.35 at 720 mins). Taking the intronic reads across stimulation time-

points, we notice both that the shBud13 nuclear sample is more unspliced and that the cytoplasmic

samples in both the shBud13 and control samples are completely spliced. Thus, unspliced Irf7 is

either being degraded in the nucleus, or it makes it to the cytoplasm and is degraded extremely

quickly, such that virtually no signal can be detected via RNA-seq.

Global analysis of the role of Bud13 in BMDMs.

We next investigated global splicing differences caused by Bud13 knockdown. Using the

TNFα stimulated data-set, ΔSR was calculated for every junction in every expressed gene.   To

determine whether splicing differences caused by Bud13 knockdown led to altered gene expression,

we compared the effect of the Bud13 knockdown on genes that contained a Bud13 dependent

junction to those that did not. (see methods). The median $\log_2$ expression fold change (FPKM

shBud13/ FPKM control) for genes containing a Bud13 dependent junction was -0.5084. In contrast,

the median $\log_2$ expression fold change (FPKM shBud13/ FPKM control) for genes without any

junctions affected by Bud13 knockdown was -0.2170. Thus, we conclude there is an inverse

relationship between IR due to Bud13 knockdown and gene expression (Wilcoxon rank-sum, *P*< .01)

(Fig. 5A).

Next, it was of interest to us to identify sequence elements that led Bud13 to have its specific

splicing effect. The most evident element to explore was the effect of splice site strength on Bud13-

dependent splicing. Previous work has shown that the yeast orthologue of Bud13 plays a role in

efficient splicing for a junction with a weak 5' splice site[13]. Further, the junction affected in Irf7 has a

non-consensus 5' splice site. To investigate this issue, we first quantified every 5' and 3' splice site

using a maximum entropy model.[12] Then, we took progressively weaker splice site thresholds, and

compared the mean ΔSR for every junction below that threshold to the mean ΔSR of every junction

in the data-set (Fig. 5B). We saw that as the splice site threshold for the 5' splice site became

progressively weaker, the mean ΔSR for junctions weaker than that threshold increased and thus

there was a greater Bud13 splicing effect. This result was not seen when the same analysis was

applied to the 3' splice site. We then analyzed the Bud13 splicing effect with respect to other

features known to correlate with IR.[6] Across all time-points for both TNFα (Fig. 5E, F, H, I) and

Poly(I:C) (Fig. S7 A-C), Bud13 dependent introns were dramatically smaller and had increased G/C

content in both the intron and in the flanking exons. We also noticed that the distance from the

branch point to the 3' splice site was smaller in the Bud13 dependent introns than in the total data-

set (Fig. 5I, Fig. S7 D). This could be a byproduct of the smaller intron length; however, it is of

interest because Bud13 has been shown in yeast to bind just downstream of the branch point.[16] A

significant difference was not seen in branch point strength and Bud13 splicing effect (Fig. S7 E, F).

Of note, the fourth intron of Irf7 is among the most Bud13 dependent junctions in both the TNFα and

Poly(I:C) data-sets (Fig. 5D., Fig. S7H, see methods for analysis details). Further, almost all

transcripts contain only a single Bud13 dependent junction, even when low thresholds are used to

quantify dependency (Fig. 5G.). This is again similar to what we observed in Irf7.

Finally, using enhanced crosslinking and immunopreciptation (eCLIP)-seq data from the

ENCODE Project Consortium,[17] we investigated Bud13 binding specificity across the genome. We

found that in K562 and Hep G2 cells, the majority of Bud13 eCLIP-sequencing reads were located

downstream of the branchpoint near the 3' splice site (Fig. 6A, B), consistent with what is seen in

yeast.[16] There is some read density near the 5' splice site, which we hypothesize is due to Bud13's

association with the spliceosome. Although it may not bind near the 5' splice site, factors in the

spliceosome that interact with Bud13 may immunoprecipitate with Bud13, leading to a 5' signal.

Data for Sf3b4 and Prpf8, known RBPs that interact with the 3' and 5' splice site respectively, is also

shown (Fig. 6A, B). After peak calling, we counted the location of significant peaks. The majority of

peaks are in intronic regions or intron-exon junctions. Further, of the peaks that lie in intron-exon

junction, most are located at the 3' junction (Fig. 6C, D). Lastly, when comparing introns that have

an overlapping eCLIP peak to all introns from expressed transcripts, we see both a length and G/C%

bias (Fig. 6E, F). Bud13 peaks tend to fall in smaller introns that are GC rich, a finding consistent

with the ΔSR data.

Bud13 knockdown alters the BMDM response to VSV.

305   Vesicular stomatitis virus (VSV) is a (-)ssRNA virus known to induce type I IFN through

TLR7[18]. To test whether impairment of Irf7 due to Bud13 knockdown was present in VSV stimulated

BMDMs, we infected both shBud13 and control BMDMs at an MOI of 5 and 10. At all time-points

throughout infection in both MOIs, there was dampened Irf7 induction (Fig. 7A, B) as quantified by

Taqman qPCR. Next, in order to determine the consequences of impaired Irf7 induction, we

determined the yield of virus from BMDMs following a period of infection with a given input MOI.

shBud13 BMDMs produce >10 times the amount of VSV as compared to control BMDMs (Fig. 7C).

This difference in viral production is presumably due to the dampened type I interferon response, as

well as the corresponding deficiency in ISG production associated with Bud13 knockdown. Finally,

we notice increased infectivity across a 24-hour time-course, a finding again consistent with our

previous data showing that Bud13 alters Irf7 splicing and thus hinders its ability to amplify the type I

interferon response (Fig. S8).

317
318   **DISCUSSION**
319
320   In this study, we sought proteins that might relate to the poor splicing of an intron in Irf7

transcripts.   Using RAP-MS, we identified Bud13 as a protein that has the ability to increase splicing

of the Irf7 intron. In the absence of *Bud13*, in response to inflammatory stimulus, macrophages

produced Irf7 with increased intron retention (IR) and notably less mature Irf7 transcript and protein

(Fig. 3C, 4C, 4D, S4C). Irf7 is the interferon-inducible master regulator of the type-I interferon-

dependent immune response.[7] Correspondingly, depletion of Bud13 led to a general reduction in

ISG and cytokine production, implying a compromised type I interferon response (Fig. 4E-J, S4D-G).

This splicing and corresponding expression defect upon Bud13 depletion was observed under

various stimulation regimens and times. We found that macrophages deficient for Bud13 were

strikingly more susceptible to infection by VSV, presumably owing to the reduction in Irf7 transcript

levels (Fig. 7).

We observed the Bud13 splicing dependence in other introns of other genes. A number of

short, GC-rich introns with non-consensus splice donor sites were excised inefficiently when Bud13

levels were depleted (Fig. 5). As was the case with Irf7, this increased IR reduced mature transcript

levels (Fig. 5A). Transcripts containing retained introns have been shown in the literature to be

degraded by two mechanisms: (1) nuclear degradation via the RNA exosome, (2) cytoplasmic

degradation upon detection of a pre-termination codon (PTC) via the NMD decay machinery.

Although the majority of these introns contain a PTC, it remains to be determined whether

degradation is occurring in the nucleus or cytoplasm.[14] [19]

Bud13 was originally identified as a part of a "Retention and Splicing" (RES) complex[13] in

yeast. However, yeast Bud13 (ScBud13) and mammalian Bud13 are significantly different lengths

(266 vs. 637 amino acids) [20], with only the mammalian protein containing a large, disordered

arginine-rich N-terminal domain. ScBud13 counteracts IR in introns within the mediator complex,

mating genes, and tRNA modifying genes[21–23], which in turn impair yeast budding. In connection

with the RES complex, ScBud13 is thought to safeguard formation of the 'B^{act} complex' of the

spliceosome.[24] In the B^{act} stage, the 5' splice donor and branch point are recognized by the

spliceosome. However, progression to catalysis of the first step of the splicing reaction requires

remodeling of several spliceosome components.[25] Lack of the RES complex has been shown to lead

to premature binding of Prp2, a quality control factor that is responsible for spliceosome remodeling

as well as the disassembly of suboptimal substrates. It has been hypothesized that ScBud13 and

the RES complex temporally regulate Prp2 binding.[24] In the mammalian context, short, GC-rich

introns with weak donor sites may be particularly susceptible to Prp2-mediated disassembly, which

may explain the specificity of IR events upon Bud13 depletion.

In yeast, differential studies using mass spectrometry[26] and cross-linking have established

that some ScBud13 is detectable in preparations of stalled B, B[act], and B* complexes. One cryo-EM

structure of the yeast spliceosome found density corresponding to ScBud13 in a stalled B[act] pre-

catalytic complex, although a structure of the stalled B complex found only weak density for

ScBud13.[22,27] In mammals, structural evidence of Bud13 is limited. Given the partial sequence

homology between all members of the yeast RES complex and their mammalian counterparts, it is

not surprising that Bud13 (and other RES complex members) are often undetectable in preparations

of stalled spliceosomes using cross-linking and mass-spectrometry. Furthermore, Bud13 was not

detected in a recent human cryo-EM structure of a stalled B complex.[28] Taken together, it is not yet

possible to determine if the sub-stoichiometric nature of Bud13 in mammalian spliceosome

complexes is because it is constitutively associated but highly transient or because it serves as a

non-essential accessory to spliceosome function. Cryo-EM studies, as well as single molecule

studies, would seem to suggest compositional heterogeneity of the spliceosome, and that the

Bud13-endowed spliceosome may catalyze the splicing reaction in a fundamentally different way

than is used in its absence.[24,29,30]

Recently, Bud13 in zebrafish was shown to regulate levels of IR in short, GC-rich introns in

knockout studies.[31] Indeed, both in Zebrafish[31] and C. Elegans,[32] Bud13 deficiency has been

reported to lead to embryonic lethality. Our results show that mammalian Bud13 shares this splicing

fidelity function, and deficiency may prevent proper development. Despite this, knockdown and

knock-out cell lines have displayed no overt growth defects, suggesting a developmental but not

immune-cell intrinsic dependence on Bud13 for survival.

Bud13 might simply represent a mechanism that evolved to counter intron retention in a

subset of introns that require splicing but happen to be inherently weak. Furthermore, as previously

mentioned, IR has emerged as a post-transcriptional mechanism used by the cell to fine-tune gene

expression. It is possible that the presence of these weak introns, coupled with regulation of Bud13,

aid in the fine-tuning of the transcriptome. Mammalian Bud13 undergoes intensive phosphorylation

and acetylation.[20] Additionally, it is intrinsically disordered and co-localizes with nuclear speckles.[33]

Alterations in the levels of Bud13 or regulation of post-translation modifications that could affect

Bud13 activity or lead to speckle sequestration might allow a cell the ability to quickly adjust the

expression of a subset of transcripts.

In summary, we found that Bud13 modulates gene expression through its ability to alter IR,

often in notably small, GC-rich introns with weak splice sites. Deficiency of Bud13 results in IR and

concomitant decreased gene expression in transcripts such as Irf7, dampening the type I interferon

response and increasing viral susceptibility. Therefore Bud13, in mediating Irf7 gene expression,

presents a potential therapeutic target for the treatment of infections or autoimmune conditions.

Future studies should seek to understand why Bud13 is vital for the efficient splicing of only a subset

of junctions and whether or not this junction specificity plays an active role in regulating gene

expression. If modulated, this strategy by which components associated with the spliceosome

rescue transcripts from intron retention and degradation may represent a previously

underappreciated layer of regulation in many gene expression programs.

ACKNOWLEDGEMENTS
395

The authors would like to thank Mario Blanco and Mitchell Guttman (Dept. of Biology, Caltech) for
assistance with eCLIP analysis; and Logan Vlach, Patricia Turpin, Mati Mann and Alok Joglekar
(Dept. of Biology, California Institute of Technology) for experimental and computational assistance.
Additionally, the authors would like to thank Genhong Cheng (Department of Microbiology,
Immunology & Molecular Genetics, UCLA) for VSV and Jae Jung (Department of Molecular
Microbiology & Immunology, USC) for VSV-GFP. This work was funded from a grant from NIH
(5R21AI126344) and from an endowment provided by the Raymond and Beverly Sackler
Foundation.

404
405 **AUTHOR CONTRIBUTIONS**
406

407 L.S.F, D.M., and D.B., conceived and designed experiments. L.S.F. conducted experiments. C.B.
408 helped develop RAP-MS and knockdown experiments. L.S.F. and D.M. analyzed sequencing data.
409 A.M. oversaw mass spectrometry and M.J.S. performed mass spectrometry analysis. L.S.F, D.M.,
410 and D.B wrote the manuscript with input from all authors.

415 **References:**
416
1. Schmitz, U. *et al.* Intron retention enhances gene regulatory complexity in vertebrates.

Genome Biol. **18,** 216 (2017).

2. Wong, J. J.-L., Au, A. Y., Ritchie, W. & Rasko, J. E. Intron retention in mRNA: No longer

nonsense. *Bioessays* **38,** 41–49 (2016).

3. Wong, J. J.-L. *et al.* Orchestrated intron retention regulates normal granulocyte

differentiation. *Cell* **154,** 583–595 (2013).

4. Yap, K., Lim, Z. Q., Khandelia, P., Friedman, B. & Makeyev, E. V. Coordinated regulation

of neuronal mRNA steady-state levels through developmentally controlled intron retention.

Genes Dev. **26,** 1209–1223 (2012).

5. Dvinge, H. & Bradley, R. K. Widespread intron retention diversifies most cancer

transcriptomes. *Genome Med.* **7,** 45 (2015).

6. Braunschweig, U. *et al.* Widespread intron retention in mammals functionally tunes

transcriptomes. *Genome Res.* **24,** 1774–1786 (2014).

7. Honda, K. *et al.* IRF-7 is the master regulator of type-I interferon-dependent immune

responses. *Nature* **434,** 772–777 (2005).

8. Ciancanelli, M. J. *et al.* Life-threatening influenza and impaired interferon amplification in

human IRF7 deficiency. *Science* **348,** 448–453 (2015).

9. Harley, J. B. *et al.* Genome-wide association scan in women with systemic lupus

erythematosus identifies susceptibility variants in ITGAM, PXK, KIAA1542 and other loci.

Nat. Genet. **40,** 204 (2008).

10. Shalek, A. K. *et al.* Single-cell transcriptomics reveals bimodality in expression and splicing

in immune cells. *Nature* **498,** 236 (2013).

11. McHugh, C. A. *et al.* The Xist lncRNA interacts directly with SHARP to silence

transcription through HDAC3. *Nature* **521,** 232–236 (2015).

12. Yeo, G. & Burge, C. B. Maximum entropy modeling of short sequence motifs with

applications to RNA splicing signals. *J. Comput. Biol.* **11,** 377–394 (2004).

13. Dziembowski, A. *et al.* Proteomic analysis identifies a new complex required for nuclear

pre-mRNA retention and splicing. *EMBO J.* **23,** 4847–4856 (2004).

14. Jacob, A. G. & Smith, C. W. Intron retention as a component of regulated gene expression

programs. *Hum. Genet.* 1–15 (2017).

15. Mostafavi, S. *et al.* Parsing the interferon transcriptional network and its disease

associations. *Cell* **164,** 564–578 (2016).

16. Schneider, C. *et al.* Dynamic Contacts of U2, RES, Cwc25, Prp8 and Prp45 Proteins with the

Pre-mRNA Branch-Site and 3' Splice Site during Catalytic Activation and Step 1 Catalysis

in Yeast Spliceosomes. *PLOS Genet.* **11,** e1005539 (2015).

17. Consortium, E. P. An integrated encyclopedia of DNA elements in the human genome.

Nature **489,** 57–74 (2012).

18. Lund, J. M. *et al.* Recognition of single-stranded RNA viruses by Toll-like receptor 7. *Proc.*

Natl. Acad. Sci. U. S. A. **101,** 5598–5603 (2004).

19. Sayani, S. & Chanfreau, G. F. Sequential RNA degradation pathways provide a fail-safe

mechanism to limit the accumulation of unspliced transcripts in Saccharomyces cerevisiae.

Rna **18,** 1563–1572 (2012).

20. Na, I., Meng, F., Kurgan, L. & Uversky, V. N. Autophagy-related intrinsically disordered

proteins in intra-nuclear compartments. *Mol. Biosyst.* **12,** 2798–2817 (2016).

21. Zhou, Y., Chen, C. & Johansson, M. J. The pre-mRNA retention and splicing complex

controls tRNA maturation by promoting TAN1 expression. *Nucleic Acids Res.* **41,** 5669–

5678 (2013).

22. Zhou, Y. & Johansson, M. J. The pre-mRNA retention and splicing complex controls

expression of the Mediator subunit Med20. *RNA Biol.* 1–7 (2017).

23. Ni, L. & Snyder, M. A Genomic Study of the Bipolar Bud Site Selection Pattern

inSaccharomyces cerevisiae. *Mol. Biol. Cell* **12,** 2147–2170 (2001).

24. Bao, P., Will, C. L., Urlaub, H., Boon, K.-L. & Lührmann, R. The RES complex is required

for efficient transformation of the precatalytic B spliceosome into an activated Bact complex.

Genes Dev. (2018).

25. Ohrt, T. *et al.* Prp2-mediated protein rearrangements at the catalytic core of the spliceosome

as revealed by dcFCCS. *RNA* **18,** 1244–1256 (2012).

26. Fabrizio, P. *et al.* The evolutionarily conserved core design of the catalytic activation step of

the yeast spliceosome. *Mol. Cell* **36,** 593–608 (2009).

27. Plaschka, C., Lin, P.-C. & Nagai, K. Structure of a pre-catalytic spliceosome. *Nature* **546,**

617 (2017).

28. Bertram, K. *et al.* Cryo-EM structure of a pre-catalytic human spliceosome primed for

activation. *Cell* **170,** 701–713 (2017).

29. Blanco, M. R. *et al.* Single Molecule Cluster Analysis dissects splicing pathway

conformational dynamics. *Nat. Methods* **12,** 1077 (2015).

30. Hoskins, A. A. & Moore, M. J. The spliceosome: a flexible, reversible macromolecular

machine. *Trends Biochem. Sci.* **37,** 179–188 (2012).

31. Fernandez, J. *et al.* RES complex is associated with intron definition and required for

zebrafish early embryogenesis. *bioRxiv* 197939 (2017). doi:10.1101/197939

32. Jiang, M. *et al.* Genome-wide analysis of developmental and sex-regulated gene expression

profiles in Caenorhabditis elegans. *Proc. Natl. Acad. Sci.* **98,** 218–223 (2001).

33. Sundararaman, B. *et al.* Resources for the comprehensive discovery of functional RNA

elements. *Mol. Cell* **61,** 903–913 (2016).

34. Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of insertions,

deletions and gene fusions. *Genome Biol.* **14,** R36 (2013).

35. Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals unannotated

transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **28,** 511–515

(2010).

36. Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31,** 166–169 (2015).

37. Corvelo, A., Hallegger, M., Smith, C. W. & Eyras, E. Genome-wide association between

branch point properties and alternative splicing. *PLoS Comput. Biol.* **6,** e1001016 (2010).

38. Liu, J. M., Sweredoski, M. J. & Hess, S. Improved 6-Plex Tandem Mass Tags Quantification

Throughput Using a Linear Ion Trap–High-Energy Collision Induced Dissociation MS3

Scan. *Anal. Chem.* **88,** 7471–7475 (2016).

39. Van Nostrand, E. L. *et al.* Robust transcriptome-wide discovery of RNA-binding protein

binding sites with enhanced CLIP (eCLIP). *Nat. Methods* **13,** 508–514 (2016).

40. Shen, L., Shao, N., Liu, X. & Nestler, E. ngs. plot: Quick mining and visualization of next-

generation sequencing data by integrating genomic databases. *BMC Genomics* **15,** 284

(2014).

41. Lovci, M. T. *et al.* Rbfox proteins regulate alternative mRNA splicing through evolutionarily

conserved RNA bridges. *Nat. Struct. Mol. Biol.* **20,** 1434–1442 (2013).

42. Budak, G., Srivastava, R. & Janga, S. C. Seten: a tool for systematic identification and

comparison of processes, phenotypes, and diseases associated with RNA-binding proteins

from condition-specific CLIP-seq profiles. *RNA* **23,** 836–846 (2017).

     FIGURE LEGENDS

Figure 1: *Irf7* contains a weak intron that is following many forms of stimulation. (A) Histogram of
mapped reads corresponding to the TNFα-induced expression of Irf7. The poorly spliced fourth intron is
highlighted. For all read density plots, reads are histogrammed in $\log_{10}$ scale and normalized to the
maximum value across the stimulation. **(B)** Comparison of Irf7 splice donor and acceptor sites in mice,
rats, and humans. **(C)** Comparison of median intron length across all annotated transcripts to the length
of Irf7 intron 4. **(D)** Comparison of mean exon G/C% across all annotated exons compared to the G/C%
of the exons directly upstream and downstream of Irf7 intron 4. **(E)** Mean intron G/C% across all
annotated introns as compared to Irf7 intron 4. **(F)** Comparison of donor splice site strength (calculated
using a maximum entropy model; see methods) for the consensus donor splice site as compared to the
Irf7 donor splice site. **(G)** Outline of Splicing Ratio (SR) metric. **(H)** Splicing ratio for all introns in Irf7
plotted against time stimulated with TNFα. **(I, J)** Histogram of mapped reads corresponding to the IFNα
(I) and poly(I:C) **(J)** induced expression of Irf7 focused on the slow splicing fourth intron.
545
**Figure 2: RAP-MS and RIP identifies Bud13 as an RNA binding protein that interacts with IRF7
mRNA. (A)** Outline of the RAP-MS procedure used to identify RNA-binding proteins on transcritps of
interest. **(B)** TMT ratio (Irf7/Actb) for proteins identified as enriched on either Irf7 (TMT ratio >1) or Actb
(TMT ratio <1) transcripts. **(C)** qRT-PCR analysis of transcripts captured via RAP for Irf7 (blue) and Actb
(gold) probes. **(D)** RIP followed by qRT-PCR for Irf7 and Rpl32 in TNFα stimulated BMDMs. Shown is
the relative enrichment of transcripts captured in Bud13 RIP as compared to Rabbit IgG RIP. **(E)** Same as
(d) except stimulation with poly(I:C). Data are representative of two independent experiments (**(C-E),**
mean + SEM). *P < 0.05, **P < 0.01 and ***P < 0.001 (t-test).

Figure 3: Bud13 knockdown leads to increased retention in the poorly splicing intron of Irf7. (A)
Description of ΔSR metric used to calculate differences in splicing at individual time-points for given
junctions. **(B)** Histogram of mapped reads corresponding to the TNFα-induced expression of Irf7. The
poorly spliced fourth intron is highlighted. shBud13 samples are shown in green. Control samples are
shown in grey. **(C)** ΔSRs calculated for each junction in the Irf7 transcript. Shade represents columns
different stimulation time-point. **(D)** Irf7 FPKM fold change with respect to time stimulated. shBud13 is
shown in green, control is shown in grey. **(E)** Splicing gel from RNA extracted from BMDMS stimulated for
30 mins. TNFα.

Figure 4: Bud13 knockdown alters the type I interferon response. (A) Histogram of mapped reads
corresponding to the TNFα-induced expression of Irf7. The poorly spliced fourth intron is highlighted.
shBud13 samples are shown in blue. Control samples are shown in grey. **(B)** ΔSRs calculated for each
junction in the Irf7 transcript. Shade of columns represents different stimulation time-point. **(C)** Irf7 FPKM
fold change with respect to time stimulated. **(D)** Immunoblot analysis of Irf7 protein after nuclear
fractionation from BMDMs left untreated (UT) or treated with poly(I:C) (PIC) or CpG for 12h. Lamin B1
serves as loading control. **(E)** $\log_2$ expression fold change (shBud13/control) for 119 ISGs in
unstimulated BMDMs (median = 0.1655). **(F)** As in **(E)** for stimulated BMDMs (720 mins poly(I:C) (median
= -0.1007). Wilcoxon rank-sum between **(E)** and **(F)**, *P*< .001. **(G)** Median $\log_2$ expression fold change
(shBud13/control) for selected ISGs in unstimulated BMDMs, and BMDMs stimulated for 15, 60, 240, 720,
and 1440 mins. (Wilcoxon rank-sum, *P*< .001, for any of the 'early' time-points (0, 15, 60 mins) compared
to any of the 'late' time-points (240, 720, 1440 mins). **(H)** Normalized FPKM expression levels in shBud13
and control samples at 720 mins poly(I:C) stimulation for select ISGs. **(I)** RT-qPCR analysis of IFNα
mRNA levels in unstimulated BMDMs and BMDMs stimulated with poly(I:C) for 720 mins and 1440 mins.
(J) Same as **(I)** for IFNβ. **(K)** Nuclear fraction (top) and cytoplasmic fraction (bottom) histograms of
mapped reads corresponding to the poly(I:C)-induced expression of Irf7 (720 mins). The poorly spliced
fourth intron is highlighted. shBud13 samples are shown in blue. Control samples are shown in grey.
Nuclear ΔSR = 0.35. **(L)** Nuclear and cytoplasmic FPKM for Irf7 intron 4 from fractionated BMDMs
stimulated with poly(I:C). Data is representative of two independent experiments (i,j) and is represented
as mean + SEM. * denotes p < 0.05, ** denotes p < 0.01, and *** denotes p < 0.001 using a Student's t
test. Results are presented relative to those of Rpl32 **(I,J).**

Figure 5: Global analysis of the role of Bud13. (A) Transcripts were classified as 'Bud13 dependent' if
they had a junction with a ΔSR. >0.15. The $\log_2$ expression fold change (FPKM shBud13/ FPKM control)
for each gene represented by the transcripts in the 'Bud13 dependent' category as well as all other genes
is shown. Median 'increased IR' = -0.5084. Median 'decreased IR' = -0.2170. (Wilcoxon rank-sum, *P*<
.01). **(B)** Mean ΔSR. for junctions below the indicated threshold (x-axis) vs. mean ΔSR. for all junctions.
Threshold applied for the 5' splice site (blue) and the 3' splice site (green). **(C)** 5'SS motif for all
expressed junctions as compared to junctions that show retention upon Bud13 knockdown (ΔSR. > 0.15).
(**D**) Ranked bar chart showing genes with a junction most affected by Bud13 knock-down in all samples
during TNFα stimulation. See S7 for histograms relating to most affected junctions. **(E)** Size of intron for
introns retained upon Bud13 knockdown (ΔSR. > 0.15) (blue), in introns located in the same transcript as
those affected by Bud13 (green), and in introns from all expressed transcripts (orange). **(F)** Same as **(E)**
for GC content. (G) Grouped bar chart depicting the number of genes that have a single Bud13
affected junction vs. multiple Bud13 affected junctions using three different ΔSR thresholds. **(H)**
Flanking exon GC content for exons that flank introns retained upon Bud13 knockdown (ΔSR. > 0.15)
(dark green) as compared to exons that flank introns from all expressed transcripts (light green). **(I)**
Distance from the branch point to the 3' splice site for introns retained upon Bud13 knockdown (ΔSR. >
0.15) (dark blue) as compared to introns from all expressed transcripts (light blue). **(E, F, H, I)** data from
BMDM TNFα stimulation. Box plots show median (center line), interquartile range (box) and tenth and
ninetieth percentiles. *P < 0.05, **P < 0.01 and ***P < 0.001 (Mann-Whitney *U*-test).

Figure 6: Bud13 interacts primarily near the 3' splice site of small, GC rich introns. (A) eCLIP-seq
read density plots in K562 cells. Bud13 eCLIP-seq is shown in blue (top), Sf3b4 is shown in maroon
(middle), and Prpf8 is shown in green (bottom). **(B)** Same as in **(A)** but for Hep G2 cells. **(C)** Bud13
eCLIP-seq peak distribution. Peaks fell within either intronic regions, intron-exon junctions, or exonic
regions. Peaks that fell within intron-exon junction were further classified as 5' junction peaks or 3'
junction peaks (bottom). **(D)** Same as **(C)** but for Hep G2. **(E)** Size of all introns in expressed transcripts
for the given cell line (dark blue) vs size of introns with overlapping eCLIP peak (maroon). Shown in
K562 (left) and Hep G2 (right) cells. Box plots show median (center line), interquartile range (box) and
tenth and ninetieth percentiles (whiskers). *P < 0.05, **P < 0.01 and ***P < 0.001 (Mann-Whitney *U*-
test). **(F)** Same as **(E)** for GC content. **(G)** GO terms (biological process) enriched among Bud13
eCLIP peaks in K562(dark blue) and Hep G2 (maroon) cells.

Figure 7: Bud13 knockdown alters the BMDM response to VSV. (A) RT-qPCR analysis of Irf7 mRNA
levels in infected control or shBud13 BMDMs stimulated with VSV (MOI 5) across 24 hours. **(B)** Same as
in **(A)** except stimulated at an MOI of 10. Results are presented relative to those of Rpl32. **(C)** PFU/mL for
viral supernatant from infected control or shBud13 BMDMs. Data is representative of two **(A, B)** or three
independent experiments **(C)** and is shown as mean + SEM. * denotes p < 0.05, ** denotes p < 0.01, and
*** denotes p < 0.001 using a Student's t test

636   **STAR METHODS**
637
638   **Contact for Reagent and Resource Sharing**
639
640   Further information and requests for resources and reagents should be directed to and will be
641   fulfilled by the Lead Contact, David Baltimore (baltimo@caltech.edu).
642
643   **Experimental Model and Subject Detail**
644
Animals
The California Institute of Technology Institutional Animal Care and Use Committee approved all
experiments. C57BL/6 WT mice were bred and housed in the Caltech Office of Laboratory Animal
Resources (OLAR) facility. C56BL6/J mice were sacrificed via CO2 euthanasia and sterilized with
70% ethanol. Femur and tibia bones harvested and stripped of muscle tissue. Bone marrow cells
were resuspended in 20mL of fresh DMEM. $2.5 *10^6$ bone-marrow cells plated in a 150mm dish in
20mL of BMDM Media (DMEM, 20% FBS, 30% L929 condition media, and 1% Pen/Strep) and
grown at 5% CO_2 and 37°C. BMDM media was completely replaced on day 3 as well as a
supplemental addition of 5mL L929 condition media on day 5.
654
655   **Cell Culture**
664
665   Human embryonic kindey cells (HEK293T) from ATCC were cultured in DMEM supplemented with
666   10% FBS and 1% Pen/Strep. Cell line was maintened at 37°C in 5% CO_2.
667
668   **Method Detail**
669
Knockdown Experiments
BMDMs for knockdown experiments were grown as described above with a few additions. On days 3
and 4, retrovirus encoding shRNAs were added to cells. On day 5, cells were selected with
puromycin (5ug/mL). On day 8, following ~72 hours of puromycin treatment, media was removed
and 10mL of PBS w/ 2mM EDTA was added. Cells were lightly scraped and replated in either 6 well
plates or 10-cm dishes depending on the experiment. Cells were left in BMDM media overnight.
The following day, cells were stimulated with either 20ng/mL of TNFα, 5ug/mL Poly(I:C) (Sigma), or
ODN 1585 (1-5 µM) (InvivoGen).
678
RNA Isolation:
Total RNA was purified from BMDMs using TRIzol reagent (Ambion) as per the manufacturer's
instructions. Genomic DNA in RNA purifications was eliminated through treatment with Turbo
DNase (Thermo Fisher Scientific) for 30 min at 37°C. 0.1-1µg RNA and 1µM dT(30) oligo (d14-954:
5'-AAGCAGTGGTATCAACGCAGAGTACT(30)) was heated at 80°C for 2.5min followed by snap
cooling on ice. 10µL template-switch RT mix added (10µM template-switch oligo (TSO: 5'-
AAGCAGTGGTATCAACGCAGAGTACACArGrGrG), 20mM DTT, 2X ProtoScriipt II Reverse
Transcriptase Reaction Buffer (NEB), 1mM dNTPs, 40U Murine RNAse Inhibitor (NEB), and 200U
ProtoScript II (NEB) Reverse Transcriptase. Reaction incubated in thermocycler with the following
program: 1. 42°C for 30min, 2. 45°C for 30min, 3. 50°C for 10min, followed by deactivation of RT for
10min at 80°C.
690
RNA Fractionation:
Confluent 10-cm dish of mature BMDMs were scraped into 400µL cold NP-40 lysis buffer, APJ1
(10mM Tris-HCl (pH 7.5), 0.08% NP-40, 150mM NaCl). Lysed cells layered onto 1mL cold sucrose
322 cushion, APJ2 (10mM Tris-HCl (pH 7.5), 150mM NaCl, 24% w/v sucrose) and centrifuged for
10min at 4°C and 13000 rpm. The supernatant from this spin represents the cytoplasmic RNA
fraction, which is immediately added to 3 volumes of 100% ethanol and 2 volumes of buffer RLT (4M

GuSCN, 325 0.1M β-mercaptoethanol, 0.5% N-lauroyl sarcosine, 25mM Na-citrate, pH7.2) and
stored at -80°C until ready to purify RNA. Pellet, containing intact nuclei, is resuspended in 500µL
TRIzol reagent. If the pellet was difficult to dissolve, it was heated at 50°C with occasional vortexing.
100µL chloroform added and shaken vigorously for 15-20s; allowed to phase separate at room
temperature for 5min. Tube centrifuged at 4°C and 12000 x g for 15min. Clear upper aqueous phase
removed to a new tube, ensuring white DNA mid-phase is not removed, and is immediately added to
3 volumes of 100% ethanol and 2 volumes of buffer RLT and stored at -80°C until ready to purify
RNA. RNA is purified according to Qiagen RNeasy column protocol and eluted in 30µL nuclease-
free H2O. RNA samples are DNAse treated with Turbo-DNAse and stored at -80°C.
706
Library preparation and RNA-Seq Analysis
Limited PCR amplifications was performed prior to library preparation. PCR reaction done with
KAPA HiFi HotStart 2x ReadyMix, 5% cDNA, and 1µM primer (d14-955: 5'-
AAGCAGTGGTATCAACGCAGAGTACT). Thermal cycler programmed for 120 seconds at 95°C as
initial denaturation, followed by 14 cycles of 30sec at 95°C for denaturation, 30sec at 62.5°C as
annealing, 150sec at 72°C for extension, and final extension at 72°C for 5 min. PCR reactions 0.9X
SeraMag and eluted in 25µL. Concentrations of purified library determined using Qubit High
Sensitivity dsDNA kit (Invitrogen) as described. Full length cDNA libraries were barcoded using the
Nextera XT Tagmentation protocol (Illumina).
716
RNA-Antisense Purification
RNA antisense purification-mass spectrometry (RAP-MS) was performed as described in McHugh et
al. with a few alterations. Briefly, we designed three 90-mer DNA oligonucleotide probes that were
antisense to the complementary target RNA sequence in both *Irf7* and *Actb* transcripts. Each probe
was targeted to a different location on the transcript and modified with a biotin in order to enable
capture of DNA:RNA hybrids on streptavidin coated magnetic beads.
RNA Prep and Lysis: ~250million cells, or 25 150mm plates of BMDMs were used for each capture.
Following stimulation with TNFα (20ng/ml) for 30 minutes, ~5-10 mL of PBS w/ 2mM EDTA was
added to each plate and cells were removed by lightly scraping. Cells were pelleted, resuspended in
PBS, and poured into a new 150mm plate. The cells were then crosslinked in Spectrolinker at 254
nm wavelength with 0.8 J/cm2 (instrument setting: 8000 x 100 uJ/cm^2). Following crosslinking, cells
were again pelleted, at which point the pellet could be frozen and stored at -80°C. Cells were lysed
in 2mL of lysis buffer per capture (10 mM Tris pH 7.5, 500 mM LiCl, 0.5% Triton X-100, 0.2% sodium
dodecyl sulphate, 0.1% sodium deoxycholate) supplemented with Protease Inhibitor Cocktail (EMD
Millipore) and 1000 U of Murine RNase Inhibitor (New England Biolabs). We found the smaller the
volume used per sample, the more efficient the capture was downstream and thus the minimum
volume needed to lyse cells should be optimized. Samples were incubated for 10 min on ice to
allow lysis. Following lysis, sample was passed through 20-gauge needle once and then 26-gauge
needle 3-5 times to disrupt the pellet and shear genomic DNA. In between passing the sample
through the 26-gauge needle, the sample was sonicated on ice with a microtip set at 5W power for a
total of 30 s in intermittent pulses (0.7 s on, 1.3 s off). Samples were then mixed with twice the
lysate volume of 1.5x LiCl/Urea Buffer (the final buffer contains 10 mM Tris pH 7.5, 500 mM LiCl,
0.5% Triton X-100, 0.2% SDS, 0.1% deoxycholate, 4 M urea). Lysates were incubated on ice for 10
min then cleared by centrifugation for 10 min at 4,000g.
Pre-clearing lysate: BioMag streptavidin beads (Bang Laboratories Inc.) were first washed 3x in
0.25-0.5ml of 500mM LiCl/4M Urea buffer (10 mM Tris pH 7.5, 500 mM LiCl, 0.5% Triton X-100,
0.2% SDS, 0.1% deoxycholate, 4 M urea). 50ul of beads were added to each sample and the
samples were incubated at 37°C for 30 min with shaking. Streptavidin beads were then magnetically
separated from lysate samples using a magnet. The beads used for preclearing lysate were
discarded and the lysate sample was transferred to fresh tubes twice to remove all traces of
magnetic beads. Input for quality control experiments can be removed at this point.
Hybridization, Capture of Probes and Elution of Associated Protein: Following pre-clearing, the
biotinylated 90-mer DNA oligonucleotide probes specific for the RNA target of interest (vary per

sample but ~5ul of 25uM per probe) were heat-denatured at 85°C for 3 min and then snap-cooled on
ice. Probes and pre-cleared lysate were mixed and incubated at 55°C with shaking for 2 h to
hybridize probes to the capture target RNA. 500mL of washed streptavidin beads (Bang
Laboratories Inc.) were then added to each sample at 55°C with shaking for 30 mins. Beads with
captured hybrids were washed 6 times with LiCl/Urea Hybridization Buffer. If needed, 1% of the
beads can be removed for qPCR quality control experiment. TRIzol reagent can be added directly to
beads to elute RNA. Beads were then resuspended in Benzonase Elution Buffer (20 mM Tris pH
8.0, 2 mM MgCl2, 0.05% NLS, 0.5 mM TCEP) and 125 U of Benzonase nonspecific RNA/DNA
nuclease was added. Incubation occurred for 1-2 h at 37°C. Beads were then separated from the
sample using a magnet. Supernatant was collected. Contaminant beads were removed by 5 rounds
of magnetic separation on supernatant. Protein was precipitated overnight at 4°C with 10%
trichloroacetic acid (TCA). TCA treated protein elution samples were pelleted by centrifugation for 30
min at 20,000g, then washed with 1 ml cold acetone and recentrifuged. Final protein elution pellets
were air dried to remove acetone, resuspended in fresh 8 M urea dissolved in 40 ml of 100 mM Tris-
HCl pH 8.5, and stored at -20°C.
Mass Spec Prep. and Analysis Performed as in McHugh et al. with few exceptions. Instead of
SILAC we label proteins at the mass spec prep step using TMT (Thermo). After desalting on a
Microm Bioresources C8 peptide MicroTrap column and lyophilization of peptide fraction, lyophilized
protein pellets were resuspended in 100mM TEAB at a concentration of 1ug/ul. We then added
1.64ul of TMT labelling reagent to each ug of sample. The reaction was incubated for one hour at
room temperature. The reaction was quenched with 0.32ul of 5% hydroxylamine per ug of protein
used and incubated for 15 mins at room temperature. Following quenching, the samples were
mixed, desalted as before, lyophilized, and mass spec was performed on Orbitrap Fusion mass
spectrometer using a TMT instrument method as described in Liu et al.[38]
Raw files were searched using MaxQuant (v. 1.5.3.30) against the UniProt mouse database (59550
sequences) and a contaminant database (248 sequences). TMT 6plex was selected as the
quantitation method with a reporter mass tolerance of 0.3. Oxidation of methionine and protein N-
terminal acetylation were variable modifications and carbamidomethylation of cysteine was fixed
modification. A 1% protein and peptide false discovery rate as estimated by the target-decoy
approach was used for identification.
780
RNA Immunoprecipitation
RNA immunoprecipitations were performed as previously described. Between 5-10 confluent 15 cm^2
dishes of BMDMs per sample were stimulated with either 20ng/mL of TNFα for 30 minutes or
5ug/mL Poly(I:C) for 12 hours. Following stimulation, proteins were cross-linked to DNA by adding
formaldehyde directly to the media to a final concentration of 0.75%, with light shaking at room
temperature for 10 mins. To quench the crosslinking reaction, glycine to a final concentration of 125
mM was added to the media and incubated with shaking for 5 mins at room temp. Media was then
aspirated and cells were rinsed twice with 10 mL of cold PBS. Following the second wash, cells
were scraped into 10mL of PBS and spun down gently (5 min, 4°C, 1,000xg). Final cell pellet was
resuspended in 0.1-1mL of polysome lysis buffer (100 mM KCl, 5 mM MgCl$_2$, 10 mM HEPES (pH
7.0), 0.5% NP40, 1 mM DTT, 100 U.ml RNase Inhibitor (NEB)) supplemented with Protease Inhibitor
Cocktail (EMD Millipore). At this point the mRNP lysate was frozen. If needed, passing the lysate
through a small gauge needle can help with lysate. Protein-G beads were pre-treated at 4°C with
NT2 (50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1 mM MgCl$_2$, 0.05% NP40) supplemented with 5%
BSA to a final ratio of 1:5 for at least 1h before use. Appropriate amount of antibody per sample
(optimized based on antibodyused but typically ~1-10ug) was added ot 250-500ul of bead/BSA
slurry and incubated at 4°C. Following incubation, beads were spun down and washed with 1 ml of
ice-cold NT2 buffer 4–5 times. Following final wash, beads were resuspended in 850ul of NT2 and
supplemented with 200U of RNase inhibitor, 10 µl of 100 mM DTT and EDTA to 20 mM. Frozen
lysate was thawed and centrifuged at 15,000*g for 15 mins. The cleared supernatant was removed
and 100ul was added to the prepared beads. Input removed at this step. Beads and lysate were
incubated for 4h at 4°C with mixing. The beads were washed 4-5 times with ice-cold NT2 and then

resuspended in 100ul of NT2 buffer. 4ul of 5M NaCl was added incubated with shaking at 65°C for 2
hours. NT2 buffer can also be supplemented with 30 μg of proteinase K to release the RNP
component. RNA was isolated by adding TRIzol reagent (Ambion) as per the manufacturer's
instructions. RNA was reverse transcribed and quantification was performed using TaqMan qPCR.
807
Immunoblot
BMDM samples were prepared as described previously. On day 8, they were plated at similar
density. Following adherence, BMDMs were stimulated with Poly(I:C) or CpG for the indicated time.
Cells were scraped into subcellular fractionation buffer (20mM HEPES (pH 7.4), 10 mM KCl, 2 mM
$MgCl_2$, 1mM EDTA, 1 mM EGTA). The cells were then passed through a 27 gauge needle 10 times,
incubated on ice for 10 mins, and spun down at 720xg for 5 min. The pellet contained the nuclei,
which was washed with fractionation buffer, passed through a 25 gauge needle 10 times, and
centrifuged again at 720xg for 10 mins. The resulting pellet was resuspended in RIPA lysis buffer.
Queal amounts of proteins were analyzed by immunoblot using the following reagents: anti-IRF7
(Millipore, ABF130), anti-Lamin B1 HRP conjugate (Cell Signalling, D9V6H), and anti-rabbit IgG
HRP conjugate (Cell Signalling).
819
Viral Plaque Assays
Plaque assays were done one Vero cells. $2.5*10^5$ vero cells were plated in a 12 well plate the night
before infection. Prior to infection, cells were checked to ensure confluence. VSV was serially
diluted and infected in 12 well plate for 1 h. VSV was then removed and cells were layered carefully
with DMEM supplemented with 2% FBS and 0.4% agarose. Plate was incubated for 2 days, and
then fixed with 10% formaldehyde, for 1 h to overnight. Finally, agarose plugs were removed
carefully and cells were stained with crystal violet.
827
VSV-GFP Infection Experiment
BMDMs were grown as described above in 150mm dishes. On day 8, following ~72 hours of
puromycin treatment, media was removed and 10mL of PBS w/ 2mM EDTA was added. Cells were
lightly scraped and 250,000 cells/well were replated in 12 well plates in BMDM media. Cells were
left for 12 hours to adhere. Following adherence, VSV-GFP was added at the specified MOI for the
specified amount of time. Following the time-course, cells were lightly scraped, washed and spun
down, and resuspended in PBS. Samples were analyzed on a MACSQuant10 Flow Cytometry
machine (Miltenyi). Gating strategy depicted in Fig. 7.
836
VSV-GFP Viral Supernatant Experiment
BMDMs were grown as described above in 150mm dishes.   On day 8, following ~72 hours of
puromycin treatment, media was removed and 10mL of PBS w/ 2mM EDTA was added. Cells were
lightly scraped and 400,000 cells/well were replated in 12 well plates in BMDM media. Cells were
left to adhere for 12 hours, before being infected at an MOI of 25 for 8 hours. Following infection,
virus was removed and the cells were washed with PBS three times. Then, 500ul of BMDM media
(DMEM, 20% FBS, 30% L929 condition media, and 1% Pen/Strep) was added to each well. 18
hours later, media was collected and stored at -80°C. To titer viral supernatant, Vero cells were
plated in a 96-well plate at 30,000 cells per well in 90ul of D10 media. 12 hours after plating, 90ul
supernatant was added to the 90ul of D10 at different dilutions. PFU/mL was calculated from a
standard curve with a virus of known concentration.
848
849 **Quantification and Statistical Analysis**
850 All statistical analysis was performed in Python (version 2.7.9). Unless otherwise indicated in figure
851 legends, statistical significance measurements were marked as follows: * denotes $p < 0.05$, ** denotes
852 $p < 0.01$, *** denotes $p < 0.001$, and n.s. denotes not significant. RNA-Seq expression and splicing
853 analysis as well as eCLIP analysis is described in more detail below.
854
855

RNA-Sequencing Analysis

Sequencing was performed on a HiSeq 2500 High Throughput Sequencer (Illumina). Single-end 50-mer reads were aligned using Tophat v2.1.1.[34] Gene expression was determined using Cufflinks v2.2.1 and the FPKM (Fragments Per Kilobase Million) metric.[35]

Splicing Ratio and ΔSR Calculation

A custom script was written in Python using the HTSeq[36] library to calculate Splicing Ratio. First, reads that map to an intron or exon feature are summed. To map to a feature, reads must have >1 bp overlap with the feature. If a read maps to more than one feature, such as in the case of a splice junction read, the read is split between the features. SR is calculated by taking the length normalized number of reads that map to each intron, divided by the average length normalized number of exon reads plus the length normalized intron value. When SR is equal to 0, this indicates a junction is completely spliced. In contrast, large SR values indicate intron retention. We use the SR value to calculate ΔSR, which is equal to SR(shBud13) – SR(Ctl). Values greater than 0 indicate the junction is more unspliced in the shBud13 sample, whereas values less than 0 indicate the junction is more unspliced in the Ctl sample. For the global analysis, in order for the ΔSR of a junction to be considered, it must pass through a number of filters. To account for transcripts that are annotated in Ensembl version 67, but not expressed, we set an FPKM threshold of 15. Further, a local normalized read count threshold on the upstream/downstream exons was implemented to ensure a level of sequencing depth needed to get accurate splicing values. To pass this threshold, the sum of the reads that map to the the upstream/downstream exons divided by the length of these exons must be ≥ 0.25.

ISG and Genome-Wide Analysis

ISGs used in Fig. 4 E-H were selected based on induction 2 hours after *in vivo* IFNa injection.[15] We classified ISGs to be any gene with a fold change ≥ 3.5 following 2 hours of induction. Intron RPKM was calculated using a custom python script with the HTSeq library. In Fig. 5a, transcripts from the 30 min. TNFα data-set that had a junction with a ΔSR value above 0.15 were sorted into an 'increased IR' category (ΔSR >0.15), whereas all other transcripts were sorted into an unaffected category (ΔSR <-0.15). The selected data-set is representative of all time-points from the TNFα, Poly(I:C) and CpG datasets. A maximum entropy model was used to calculate 3' and 5' splice site strengths.[12] To determine differences in 5' splice site sequence for Bud13 dependent junctions, the nine base pair sequence near the 5' splice site junctions for junctions that had a ΔSR >0.15 was compared to all expressed junctions (FPKM>1). The top Bud13 dependent junctions were plotted based on the average ΔSR value across all time-points from the TNFα data-set (Fig. 5D) as well as the Poly (I:C) data-set (Fig. SH). Junctions that had a ΔSR value <0.15 in a time-point were filtered out in the TNFα data-set, while junctions that had a ΔSR value <0.15 in two time-points were filtered out in the Poly (I:C) data-set. SVMBPfinder was used to determine BP related features (BP strength and distance from BP to 3' splice site).[37]

eCLIP

Data for eCLIP experiments were downloaded from ENCODE Project Consortium.[17] Analysis of eCLIP data is the same as has been described previously.[39] Fold change of eCLIP read density compared to input read density along a normalized intron was calculated using ngs.plot.[40] Peaks were called using CLIPper.[41] Peaks were deemed significant if they were >3-fold enriched and had p-value<10^{-5}. Peak locations were determined using a custom python script with the HTSeq library. Enriched GO terms were determined using Seten.[42]

Data and Software Availability

The next-generation sequencing data reported in this study will be deposited to the Gene Expression Omnibus (GEO). Upon completion of deposit, the accession number for this data will be provided.

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit polyclonal anti-IRF7 antibody	Millipore	Cat #ABF130
Rabbit monoclonal anti-LaminB1 (D9V6H) (HRP-conjugate)	Cell Signaling	Cat #15068
Anti-rabbit IgG (HRP-conjugate)	Cell Signaling	Cat #7074
Rabbit polyclonal anti-Bud13 antibody	Thermo Fisher Scientific	Cat #PA5-58351
Bacterial and Virus Strains		
Vesicular Stomatitis Virus-GFP (Indiana Strain)	Jae U. Jung	N/A
Vesicular Stomatitis Virus (Indiana Strain)	Genhong Cheng	N/A
Chemicals, Peptides, and Recombinant Proteins		
Mouse-TNFa	Biolegend	Cat #575206
ODN 1585	Invivogen	tlrl-1585
Poly (I:C)	Sigma-Aldrich	CAS Number 42424-50-0
Dulbecco's Modification of Eagle's Medium	Corning	10-013-CV
Fetal Bovine Serum	Corning	35-015-CV
Penicillin Streptomycin Solution (100x)	Corning	30-002-CI
0.25% Trypsin	Corning	25-053-CI
TRIzol Reagent	Invitrogen	15596026
Protoscript II	New England Biolabs	M0368L
Antarctic phosphatase	New England Biolabs	M0289S
Puromycin	InvivoGen	Cat#ant-pr-1
Critical Commercial Assays		
BioT	Bioland	B01-01
Nextera XT DNA Library Preparation Kit	illumina	FC-131-1096
Deposited Data		
RNA-Seq RAW Data	This Paper	GEO: TBD
Experimental Models: Cell Lines		
HEK293T	ATCC	CRL-3216
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	Jackson Laboratory	000664
Oligonucleotides		
shRNA targeting Bud13	IDT	TRCN0000123412
shRNA control (5' - CCTAAGGTTAAGTCGCCCTCGCTCGAGCGAGGGCGACTTAACCTTAGG - 3'	IDT	N/A
IFNa Forward qPCR Primer (5' - GCCTTGACACTCCTGGTACAAATGAG - 3')	Honda et. al., 2005	N/A
IFNa Reverse qPCR Primer (5' - CAGCACATTGGCAGAGGAAGACAG - 3')	Honda et. al., 2005	N/A
IFNb Reverse qPCR Primer (5' - AAGAGTTACACTGCCTTTGCCATC - 3')	Honda et. al., 2005	N/A
IFNb Reverse qPCR Primer (5' - CACTGTCTGCTGGTGGAGTTCATC - 3')	Honda et. al., 2005	N/A
Rpl32 Forward qPCR Primer (5' - AAGCGAAACTGGCGGAAC - 3')	Hao et. al., 2013	N/A
Rpl32 Reverse qPCR Primer (5' - TAACCGATGTTGGGCATCAG - 3')	Hao et. al., 2013	N/A
Irf7 TaqMan	Thermo Fisher Scientific	Mm00516793_g1
Rpl32 TaqMan	Thermo Fisher Scientific	Mm02528467_g1

Bud13 TaqMan	Thermo Fisher Scientific	Mm00524501_m1
d14-954: (5'-AAGCAGTGGTATCAACGCAGAGTACT(30))	IDT	N/A
Template-Switch Oligo (5'-AAGCAGTGGTATCAACGCAGAGTACACArGrGrG)	IDT	N/A
Software and Algorithms		
Tophat	Kim et. al., 2013	v2.1.1
Cufflinks	Trapnell et. al., 2010	v2.2.1
HTSeq	Anders et. al., 2015	Release_0.10.0
Clipper	Lovci et. al., 2013	N/A
SVMBPfinder	Corvelo et. al., 2010	N/A
Samtools	http://samtools.sourceforge.net/	v1.4.1
Other		
eCLIP Raw Reads	ENCODE	https://www.encodeproject.org

Figure 1

Figure 1: *Irf7* contains a weak intron that is retained following many forms of stimulation.

Figure 2

Figure 2: RAP-MS and RIP identify Bud13 as an RNA binding protein that interacts with IRF7 mRNA

Figure 3: Bud13 knockdown leads to increased retention in the poorly splicing intron of Irf7

Figure 4

Figure 4: Bud13 knockdown alters the type I interferon response

Figure 5

Figure 5: Global analysis of the role of Bud13.

Figure 6

Figure 6: Bud13 interacts primarily near the 3' splice site of small, GC rich introns.

Figure 7

Figure 7: Bud13 knockdown alters the BMDM response to VSV.

SUPPLEMENTAL FIGURES

Figure S1: Splicing Ratios across all junctions in Irf7

Figure S2: shBud13 knocks down Bud13 protein and mRNA.

A.

Figure S3: Irf7 Intron 4 is the most Bud13 knockdown affected junction of all ISGs.

Figure S4: Bud13 knockdown alters the type I interferon response in response to CpG

Figure S5: BMDM fractionation and effect of Bud13 on nuclear Irf7 splicing

Figure S6: Supplemental global analysis of shBud13

Figure S7: Histograms of mapped reads corresponding to hits identified in TNFα and PIC data-sets.

Figure S8: Bud13 knockdown alters the BMDM infection via VSV.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1: Splicing Ratios across all junctions in Irf7. Splicing ratios calculated for all junctions in the most abundant transcript of Irf7. Color represents time-point indicated in legend. **(A)** Poly(I:C) **(B)** IFNα.

Figure S2: shBud13 knocks down Bud13 protein and mRNA. (A) Immunoblot analysis of Bud13 in BMDMs infected with control or shBud13. ActB serves as loading control. **(B)** qRT-PCR analysis of Bud13 mRNA in BMDMs infected with control or shBud13. **(C-E)** FPKM fold change with respect to time stimulated **(C)** Zfp36, **(D)** IκBε, and **(E)** CD83. shBud13 is shown in green, control is shown in grey. Data is representative of two individual experiments **(A, B)** and is shown as mean + SEM **(B)**. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$.

Figure S3: Irf7 Intron 4 is the most Bud13 knockdown affected junction of all ISGs. (A) ΔSR was calculated at 720 mins of poly(I:C) stimulation for each ISG junction that passed the transcript and local read count threshold (see methods). Mean ΔSR = 0.002, Median ΔSR = 0.

Figure S4: Bud13 knockdown alters the type I interferon response in response to CpG. (A) Histogram of mapped reads corresponding to the CpG-induced expression of Irf7. The poorly spliced fourth intron is highlighted. shBud13 samples are shown in pink. Control samples are shown in grey. **(B)** ΔSR calculated for the fourth intron of Irf7 at stimulated time-points (720, 1440 mins). **(C)** Irf7 FPKM fold change with respect to time stimulated. shBud13 is shown in pink, control is shown in grey. **(D)** Log_2 expression fold change (shBud13/Control) for 119 ISGs (selected based on upregulation in response to IFNα) in unstimulated BMDMs (median = -0.2442). **(E)** As in **(D)** for stimulated BMDMs (720 mins CpG (median = -0.4776). Wilcoxon rank-sum between **(D)** and **(E)**, $P < .001$. **(F)** Normalized FPKM expression levels in shBud13 and control samples at 720 mins CpG stimulation for select ISGs. **(G)** RT-qPCR analysis of IFNα mRNA levels in unstimulated BMDMs and BMDMs stimulated with CpG for 720 and 1440 mins. Data is represented as mean + SEM. * denotes p < 0.05, ** denotes p < 0.01, and *** denotes p < 0.001 using a Student's t test. Results are presented relative to those of Rpl32 **(G).**

Figure S5: BMDM fractionation and effect of Bud13 on nuclear Irf7 splicing. (A) Ratio of cytoplasmic FPKM levels to cytoplasmic and nuclear FPKM levels for transcripts that are primarily nuclear (Malat1, Neat1, Xist, U2; left), and primarily cytoplasmic (Rpl32, Rps5, Actb, Rpl5; right) (BMDMs – 720 mins poly(I:C) stimulation). **(B)** ΔSR for intron 4 of Irf7 from nuclear fraction of unstimulated and poly(I:C) stimulated (15, 60, 240, 720 mins.) BMDMs.

Figure S6: Supplemental global analysis of Bud13. (A) Size of intron for introns retained upon Bud13 knockdown (ΔSR. > 0.15) (blue), in introns located in the same transcript as those affected by Bud13 (green), and in introns from all expressed transcripts (orange). **(B)** Same as **(A)** for GC content. **(C)** Flanking exon GC content for exons that flank introns retained upon Bud13 knockdown (ΔSR. > 0.15) (dark green) as compared to exons that flank introns from all expressed transcripts (light green). **(D)** Distance from the branch point to the 3' splice site for introns retained upon Bud13 knockdown (ΔSR. > 0.15) (dark blue) as compared to introns from all expressed transcripts (light blue). **(A-D)** data from BMDM poly(I:C) stimulation. **(E)** Branch point score for introns retained upon Bud13 knockdown (ΔSR. > 0.15) (beige) as compared to introns from all expressed transcripts (dark brown) in TNFα stimulated BMDMs. **(F)** Same as **(E)** but for poly(I:C) stimulated BMDMs. Box plots show median (center line), interquartile range (box) and tenth and ninetieth percentiles. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$ (Mann-Whitney *U*-test).

Figure S7: Histograms of mapped reads corresponding to hits identified in TNFα and PIC data-sets. (A-G) Histogram of mapped reads corresponding to the TNFα-induced expression Imp4, Psme1, Dpp7, Flii, Emg1, Hnrnpc, and Psma3 respectively. The poorly spliced intron in each transcript is highlighted. For all read density plots, reads are histogrammed in log_{10} scale and normalized to the maximum value across the stimulation. **(G)** Ranked bar chart showing genes with a junction most affected by Bud13 knock-down in all samples during PIC stimulation. **(I-L)** Histogram of mapped reads corresponding to the PIC-induced expression Imp4, Psma3, Emg1, and Psme1 respectively.

Figure S8: Bud13 knockdown alters the BMDM infection via VSV. **(A)** FSC/SSC plot showing the gating of live BMDMs in an uninfected control sample and the subsequent threshold used to calculate infectivity. **(B)** Same as in **(A)** but for a control sample infected with VSV-GFP for 12 hours. **(C)** Percent of live cells infected with VSV-GFP (MOI 10) in both control and shBud13 BMDMs across a 24-hour time-course. **(D)** Percent of live cells infected in both control and shBud13 BMDMs at 12 hours across a range of VSV-GFP MOIs.