Individual differences in visual salience vary along semantic dimensions

[1,2]Benjamin de Haas*, [2]Alexios L. Iakovidis, [2,3]D. Samuel Schwarzkopf, and [1]Karl R. Gegenfurtner

[1]Department of Psychology, Justus Liebig Universität Giessen, Germany

[2]Experimental Psychology, University College London, UK

[3]School of Optometry & Vision Science, University of Auckland, New Zealand

*Corresponding Author: Benjamin de Haas
benjamindehaas@gmail.com
Department of Psychology, Justus Liebig Universität Giessen, Germany
Otto-Behagel-Str. 10F
35394 Giessen, Germany
Fon: +49-641-9926125

Abstract: 150 words; Main text: 2615 words (excl. Methods, fig. and table captions); Methods: 1526 words; 5 figures:1 table; 1 supplementary figure

What determines where we look? Theories of attentional guidance hold that image

features and task demands govern fixation behaviour, while differences between observers

are 'noise'. Here, we investigated the fixations of > 100 human adults freely viewing a large

set of complex scenes. We found systematic individual differences in fixation frequencies

along six semantic stimulus dimensions. These differences were large (> twofold) and

highly stable across images and time. Surprisingly, they also held for *first* fixations directed

towards each image, commonly interpreted as 'bottom-up' visual salience. Their

perceptual relevance was documented by a correlation between individual face salience and

recognition skills. The dimensions of individual salience and their covariance pattern

replicated across samples from three different countries, suggesting they reflect

fundamental biological mechanisms of attention. Our findings show stable individual

salience differences along semantic dimensions, with meaningful perceptual implications.

Salience reflects features of the observer as well as the image.

Introduction

Humans constantly move their eyes[1]. The foveated nature of the human visual system

balances detailed representations with a large field of view. On the retina[2] and in visual cortex[3],

resources are heavily concentrated towards the central visual field, resulting in the inability to

resolve peripheral clutter[4] and the need to fixate visual objects of interest. Where we move our

eyes determines which objects and details we make out in a scene[5,6].

Models of attentional guidance aim to predict which parts of an image will attract

fixations based on image features[7–10] and task demands[11,12]. Classic salience models compute

image discontinuities of low level attributes, such as luminance, colour and orientation[13]. These

low-level models are inspired by 'early' visual neurons and their output correlates with neural

responses in subcortical[14] and cortical[15] areas thought to represent neural 'salience maps'.

However, while these models work relatively well for impoverished stimuli, human gaze

behaviour towards richer scenes can be predicted at least as well by the locations of objects[16] and

perceived meaning[9]. When sematic object properties are taken into account, their weight for gaze

prediction far exceeds that of low-level attributes[8,17]. A common thread of low- and high-level

salience models is that they interpret salience as a property of the image and treat inter-individual

differences as unpredictable[7,18], often using them as a 'noise ceiling' for model evaluations[18].

However, even the earliest studies of fixation behaviour noted considerable individual

differences[19,20] and basic occulomotor traits, like mean saccadic amplitude and velocity, reliably

vary between observers[21–27]. Moreover, recent twin-studies revealed that social attention and

gaze traces across complex scenes are highly heritable[28,29]. This suggests individual differences

in fixation behaviour are not random, but systematic. However, it is largely unclear, *how*

individuals differ in their fixation behaviour and what may explain these differences. Can

individual fixation behaviour be captured along a limited set of dimensions?

Here, we tested the hypothesis that individual gaze reflects individual *salience differences*

along a limited number of semantic dimensions. We investigated the fixation behaviour of > 100

human adults freely viewing 700 complex scenes, containing thousands of semantically

annotated objects[8]. We quantified salience differences as the *individual* proportion of cumulative

fixation time or *first* fixations landing on objects with a given semantic attribute. In free viewing,

the *first* fixations after image onset are thought to reflect 'automatic' or 'bottom-up' salience[30–32],

especially for short saccadic latencies[33,34]. They may therefore reveal individual differences with

a deep biological root. We tested the reliability of such differences across random subsets of

images and re-tests after several weeks. To test the generalizability of salience differences, we

replicated their set and covariance pattern across independent samples from three different

countries. Finally, we explored whether individual salience differences are related to personality

and perception, focussing on the example of face salience and face recognition skills for the

latter.

Results

Reliable Salience Differences along Semantic Dimensions

We tracked the gaze of healthy human adults freely viewing a broad range of images

depicting complex everyday scenes[8]. A first sample was tested at University College London,

UK (*Lon*; $n = 51$), and a replication sample at University of Giessen, Germany (*Gi_1*; n = 51).

The replication sample was also invited for a re-test after two weeks (*Gi_2*; $n = 48$). Additionally

we re-analysed a public dataset from Singapore (*Xu et al.*[8]; $n = 15$).

First, we probed the individual tendency to fixate objects with a given semantic attribute,

measuring duration-weighted fixations across a free-viewing period of 3s. We considered a total

of twelve semantic properties, which have previously been shown to carry more weight for

predicting gaze behaviour (on an aggregate group level) than geometric or pixel-level attributes[8].

To test the consistency of individual salience differences across independent sets of images, we

probed their reliability across 1000 random (half-)splits of 700 images. We found consistent

individual salience differences ($r > .6$) for six of the 12 semantic attributes: Neutral *Faces*,

*Emotion*al Faces, *Text*, objects being *Touched*, objects with a characteristic *Taste* (i.e. food and

beverages) and objects with implied *Motion* (Figure 1, grey scatter plots).

Observers showed up to two-fold differences in the cumulative fixation time attracted by

a given semantic attribute and the median consistency of individual differences across image

splits for these six dimensions ranged from $r = .64$ $P < .001$ (*Motion*) to $r = .94$, $P < .001$ (*Faces*;

Table 1, left hand side; *P*-Values Bonferroni corrected for 12 consistency correlations).

Previous studies have argued that extended viewing behaviour is governed by cognitive

factors, while first fixations towards a free-viewed image are governed by 'bottom-up'

salience[30–32], especially for short saccadic latencies[33,34]. Others have found that perceived

meaning[9,35] and semantic stimulus properties[8,36] are important predictors of gaze behaviour from

the first fixation. We found consistent individual differences also in the proportion of *first*

fixations directed towards each attribute. The range of individual differences in the proportion of

first fixations directed to each of the six attributes was even larger than that for all fixations (up

to threefold). Importantly, these inter-observer differences were consistent for all dimensions

found for cumulative fixation time except *Motion* ($r = .34$, *n.s.*), ranging from $r = .57$, $P < .001$

(*Taste*) to $r = .88$, $P < .001$ (*Faces*; Table 1, right hand side and green scatter plots in Figure 1; *P*-

Values Bonferroni corrected for 12 consistency correlations).

These salience differences proved robust for different splits of images and replicated

across datasets from three different countries (Figure 2a; Table 1). For the confirmatory *Gi_1*

dataset we tested the same number of observers as in the original *Lon* set. A power analysis

confirmed that this sample size yields > 95% power to detect consistencies with a population

effect size of $r > .5$. For cumulative fixation time (grey histograms in Figure 2a), the six

dimensions identified in the *Lon* sample (top row), closely replicated in the *Gi_1* and *Gi_2*

samples (middle rows), as well as in a re-analysis of the public *Xu et al.* dataset (bottom row),

with consistency correlations ranging from .65 (*Motion* in the *Gi_1* set) to .95 (*Faces* in the *Xu et*

al. set; left column of Table 1). Similar was true for *first* fixations, although the consistency

correlation for *Emotion* missed statistical significance in the small *Xu et al.* dataset (green

histograms in Figure 2a; right column of Table 1).

The individual salience differences we found were consistent across subsets of diverse,

complex images. To test whether they reflected stable observer traits, we additionally tested their

re-test reliability for the full image set across a period of 6-43 days (average 16 days; the *Gi_1*

and *Gi_2* datasets). Salience differences along all six semantic dimensions were highly

consistent over time (Figure 2b). This was true for both, cumulative fixation time (re-test

reliabilities ranging from $r = .68$, $P < .001$ (*Motion*) to $r = .85$, $P < .001$ (*Faces*); left column of

Table 1; grey bars in Figure 2b) and *first* fixations (re-test reliabilities ranging from $r = .62$, $P <$

$.001$ (*Taste*) to $r = .89$, $P < .001$ (*Text*); right column of Table 1; green bars in Figure 2b).

Finally, we tested the relationship between individual salience and visual field biases.

Different types of objects tend to appear at different parts of the visual field and recent findings

suggest that observers are attuned to these contingencies[37–39]. The most prominent spatial bias in

the scene stimuli we used was a strong tendency for faces to appear in the upper visual field

(~75%). Individual face salience (% *first* fixations) was indeed correlated with a general upper

visual field bias (as indexed by the median elevation of *first* fixations towards images not

containing a face; $r = .49-.78$ across samples, all $P < .05$). Crucially, however, individual

differences in face salience persisted independent of this spatial bias. Individual face salience in

the *lower* and upper visual field were highly correlated with each other ($r = .61-.87$ across

samples, all $P < .05$; see Supplementary Methods and Supplementary Fig. 1 for details).

Covariance Structure of Individual Differences in Semantic Salience

Having established reliable individual differences in fixation behaviour along semantic

dimensions, we further explored the space of these difference by quantifying the covariance

between them. For this analysis we collapsed neutral and emotional faces into a single *Faces*

label, because they are semantically related and corresponding differences were strongly

correlated with each other ($r = .74$, $P < .001$; $r = .81$, $P < .001$ for cumulative fixation times and

first fixations, respectively). Note that we decided to keep these two dimensions separated for the

analyses above because the residuals of fixation times for emotional faces still varied

consistently when controlling for neutral faces ($r = .73$, $P < .001$), indicating an independent

component (however the same was not true for *first* fixations, $r = .24$, *n.s.*).

The resulting five dimensions showed a pattern of pairwise correlations that allowed the

identification of two clusters (Fig. 3b, left hand side). This was illustrated by the projection of

the pairwise (dis-)similarities onto a two-dimensional space, using metric dimensional scaling

(Fig. 3b, right hand side). *Faces* and *Motion* were positively correlated with each other, but

negatively with the remaining three attributes *Text*, *Touched* and *Taste*. Interestingly, *Faces,* the

most prominent dimension of individual fixation behaviour, was strongly anti-correlated with

Text and *Touched*, the second and third most prominent dimensions (*Text*: $r = -.62$, $P < .001$ and

$r = -.47$, $P < .001$ for cumulative fixation times and first fixations, respectively, Fig. 3a, left hand

side; *Touched*: $r = -.58$, $P < .001$ and $r = -.80$, $P < .001$, Fig. 3a, right hand side). These findings

closely replicated across all four datasets (Fig. 4). Pair-wise correlations between (z-converted)

correlation matrices from different samples ranged from .68 to .95 for cumulative fixation times

and from .91 to .98 for first fixations.

Perceptual and Personality Correlates of Salience Differences

If salience differences are indeed deeply rooted in the visual cortices of our observers,

then this might have an effect on their perception of the world. We aimed to test this hypothesis

by focussing on the most prominent dimension of salience differences: *Faces*, as indexed by the

individual proportion of *first* fixations landing on faces (which is thought to be an indicator of

bottom-up salience). 46 observers from the *Gi* sample took the Cambridge Face Memory Test

(CFMT) and we tested the correlation between individual face salience and face recognition

skills. CFMT scores and the individual proportion of *first* fixations landing on faces correlated

with $r = .41$, $P < .005$. Interestingly, this correlation did not hold for the individual proportion of

total cumulative fixation time landing on faces, which likely represents more voluntary

differences in viewing behaviour ($r = .21$, *n.s.*; Fig. 5a).

Having established a correlation of semantic salience differences with perception, we

then explored potential relationships with personality variables. Observers in the *Lon* sample

completed questionnaires for seven personality dimensions, including the Big Five[40], Sensation

Seeking[41] and High Sensitivity[42] (see Methods for details). We tested all 7x6 pairwise

correlations between personality traits and individual salience differences, applying family-wise

error correction for multiple comparisons[43]. Results showed no significant relationship between

personality variables and cumulative fixation times (all $r < .39$, *n.s.*; Fig. 5b, left hand side), or

first fixations (all $r < .23$, *n.s.*; Fig. 5b, right hand side).

Discussion

Individual differences in gaze traces have been documented since the earliest days of

eyetracking[19,20]. However, salience models have routinely ignored them and the nature of these

differences was unclear. Our findings show that what was thought to be noise can actually be

explained by a canonical set of semantic salience differences. These salience differences were

highly consistent across hundreds of complex scenes, proved reliable in a re-test after several

weeks and persisted independently of correlated visual field biases. This shows that visual

salience is not just a factor of the image; individual salience differences are a stable trait of the

observer.

Not only the set of these differences, but also their covariance structure replicated across

independent samples from three different countries. This may partly be driven by environmental

and image statistics (for instance, faces are more likely to move than food). But it probably also

reflects a deep-rooted neurobiological basis of these differences. This possibility is underscored

by earlier studies showing that the visual salience of social stimuli is reduced in individuals with

autism spectrum disorder[28,44,45]. Most importantly, recent twin studies in infants and children

show that individual differences in gaze traces are heritable[28,29]. The gaze trace dissimilarities

investigated in these twin studies might be a manifestation of the salience differenceswe found

here. This suggests a strong genetic component for individual salience differences.

Recent findings in macaque show that face deprived monkeys have a drastically reduced

fixation tendency towards faces and an increased fixation tendency for hands. These differences

in fixation behaviour were accompanied by an underdevelopment of domain specific face

patches in the temporal cortex and a relative overdevelopment of hand preferring patches[46]. In

our study, high individual salience for the *Touched* label implies strong attentional guidance by

hands (see Fig. 1 for examples). It is tempting to speculate that the strong anti-correlation we

found between individual salience for *Faces* and *Touched* reflects individual differences in

developmental exposure and cortical layout. The same could apply to the anti-correlation

between *Faces* and *Text*, two stimulus classes which have previously been proposed to compete

for cortical territory[47]. It is worth noting that most of the reliable dimensions of individual

salience differences we found correspond to domain specific patches of the ventral path (as is

true for *Faces*[48–50], *Text*[47,51], *Motion*[52], *Touched*[46,53,54] and maybe *Taste*[55]). Our findings open up

a whole new field of research, where the relationship between salience differences and the

individual functional architecture of the ventral stream is directly compared.

Our findings raise important questions about the individual nature of visual perception.

Two observers presented with the same image likely end up with a different percept[5,6] and

interpretation[56] of this image when executing systematically different eye movements. Vision

scientists may be chasing a phantom when 'averaging out' individual differences to study the

'typical observer'[57–59] and *vice versa* perception may be crucial to understand individual

differences in cognitive abilities[60,61], personality[62,63], social behaviour[28,44], clinical traits[64–66] and

development[45].

We took a major first step towards uncovering the perceptual implications of salience

differences here. Individual face salience was significantly correlated with face recognition

skills. Interestingly, this was only true when considering the proportion of *first* fixations attracted

by faces. Immediate saccades towards faces can have very short latencies and be under limited

voluntary control[67,68], likely reflecting bottom-up processing. This raises questions about the

ontological interplay between face salience and recognition. Small initial differences may grow

through mutual reinforcement of face fixations and superior perceptual processing.

To our surprise, we found no significant correlations between salience differences and

personality factors captured by standard psychometric questionnaires. However, given the large

number of tests and control for multiple testing, we cannot rule out the existence of small or

medium effects. Either way, the salience differences we found should inform and enrich the

science of individual differences. They have unusually high magnitudes (up to factor 3) and

reliability (up to and greater than .9) for objective psychological traits and appear to document

fundamental differences in visual attention and perception, which are main filters of incoming

information about our surroundings

In summary, we found a small set of semantic dimensions that span a space of individual

differences in fixation behaviour. These dimensions replicated across culturally diverse samples

and also applied to the first fixations directed towards an image. Visual salience is not just a

function of the image, but also of the individual observer.

Methods

Subjects

The study comprised three original datasets (the *Lon*, *Gi_1* and *Gi_2* samples) and the re-

analysis of a public dataset (the *Xu et al.* sample[8]).

54 healthy adults with normal or corrected-to normal vision participated in the

experiment conducted at University College London, UK. The local institutional review board

approved the study and participants provided written informed consent. Reimbursement was 25£.

Eyetracking failed for three participants due to technical problems, leaving 51 in the *Lon* sample

(25 males, 4 left handed, mean age 23 with a standard deviation of 4 years). For two of these

participants, data from one block was missing due to a recording failure; the remaining data from

these participants were entered into the analysis nonetheless. Control-analysis showed that the

results remained virtually unchanged when these data were excluded. 46 of the *Lon* participants

additionally completed three personality questionnaires administered online: The IPIP-NEO 120

(NEO)[40] to measure the Big Five[69]; the Highly Sensitive Person Scale (HSP)[42] and the Brief

Sensation Seeking Scale (BSSS)[41].

51 healthy adults with normal or corrected-to normal vision participated in the

experiment conducted at Justus-Liebig Universität Giessen, Germany (the *Gi_1* sample; 11

males, 5 left handed, mean age 24, with a standard deviation of 4 years). For one of these

participants, data from one block was missing due to a recording failure; the remaining data from

this participants were entered into the analysis nonetheless. Control-analysis showed that results

remained virtually unchanged when these data were excluded. The local institutional review

board approved the study and participants provided written informed consent. Reimbursement

was 30€ or course credits.

48 of the *Gi_1* participants returned to the lab for a re-test (the *Gi_2* sample). The period

between tests was 16 days on average (minimum 6 days; maximum: 43 days; standard deviation:

7 days). 46 of the *Gi* participants additionally completed an online version of the Cambridge

Face Memory Test (CMFT)[70].

The public *Xu et al*. dataset comprises 15 participants with a reported age range of 18-30

years[8].

Stimuli

Participants were presented with 700 natural images depicting a wide variety of complex

everyday scenes (http://www-users.cs.umn.edu/~qzhao/predicting.html). Semantic metadata[8]

consist of binary pixel maps for 5551 objects in these images and accompanying labels for 12

semantic dimensions. We modified the provided labels to minimize overlap between them in the

following way: The (neutral) *Faces* label was removed from all objects with the *Emotion* label

(i.e. emotional faces); The *Smell* label was removed from all objects with the *Taste* label; The

Operable and *Gazed* labels were removed from all objects with the *Touched* label; The

Watchable label was removed from all objects with the *Text* label. This step allowed to test

individual differences in fixation attraction for a given dimension largely independently from the

others. Without this step some of the attributes would have been perfectly confounded (for

instance, all text is watchable; all emotional faces are faces).

In each experiment, participants viewed seven blocks of 100 images on a screen.

Stimulus presentation and data collection was coded in MATLAB Version R2016b (MathWorks,

Natick, MA) using Psychtoolbox Version 3.0.12[71,72]. For the *Lon* and *Gi_1* samples the order of

images was fixed across all participants and each image was presented for 3s, with a self-paced

period of a central fixation dot in between. Participants were simply instructed to 'look at the

images in any way [they] want' and initiated the onset of each image with a press of the space

bar. The re-test of the Giessen sample (*Gi_2*) followed the same procedure as the first

appointment, with the exception of the order of images presented, which was shuffled relative to

the first appointment (but again constant across participants).

Participants in the *Lon* sample sat at a distance of ~65 cm from a screen and saw the

stimuli at a resolution of 800 x 600 pixels and a size of 19.2 x 14.4 degrees visual angle.

Participants in the *Gi* samples sat with their head in a chinrest, at a distance of 46 cm from the

screen and saw the stimuli at a resolution and size of 1000x750 pixels and a size of 37.2 x 27.9

degrees visual angle. The *Xu et al*. sample saw stimuli at a size of 33.7 x 25.3 degrees visual

angle. Further details on this sample can be found in the original publication by Xu et al.[8].

Data Collection

The gaze of participants in the *Lon* sample was sampled remotely and binocularly with a

Tobii EyeX (Tobii Technologies, Danderyd, Sweden) at a frequency of ~55 Hz[73]. Eyetracking

data from the *Gi* samples was collected from the left eye with a tower mounted Eyelink 1000

(SR Research, Ottawa, Canada) at a frequency of 2 kHz (the *Xu et al*. data were collected with a

remote Eyelink 1000[8]).

At the beginning of each block, participants completed a nine-point calibration and

validation procedure, which was repeated if necessary. For the *Gi* samples, fixation data was

collected online using the 'normal' setting of the Eyelink parser (saccade velocity and

acceleration thresholds of 30 d.v.a./s and 9500 d.v.a./s^2, respectively) and the default drift check

procedure in each inter-trial interval. Raw data from the *Lon* sample was converted to fixation

data offline (see below).

Data Processing and Analyses

Raw eyetracking data from the *Lon* sample were transformed into fixation data applying a saccade threshold of 30 d.v.a./s and taking the median x and y position of fixation samples as the respective fixation location. Fixations were drift-corrected in a block-wise fashion, centering on the median of fixation locations registered at image onsets during the respective block.

Onset fixations and fixations with a duration below 100ms were disregarded for all datasets (minimum fixation duration following standard recommendations by SR Research and Xu *et al*.[8]). Fixations that fell on or within a distance of ~0.5 d.v.a from a labelled object were assigned the corresponding label. Unlabelled fixations were disregarded for the calculation of cumulative fixation times and the individual proportion of *first* fixations (see below).

In order to quantify the individual tendency to fixate objects bearing a given attribute label, we first calculated the cumulative fixation time for all labelled fixations made by a given observer to a given image set. This allowed us to calculate the proportion of this time spent on a given attribute in a second step.

The *first* fixations analysis considered the proportion of labelled *first* fixations (after image onset) landing on objects with a given attribute for a given observer and image set. Individual proportions of cumulative fixation time and first fixations were expressed in %.

Consistency and re-test correlations

To estimate the consistency of individual differences in % cumulative fixation time or %

first fixations along a given attribute dimension, we calculated these measures independently for

two random halves of the image for each observer and then correlated individual differences for

one half of the images with the other using Pearson correlation coefficients. This procedure was

repeated 1000 times. We inspected the frequency histograms of all correlations (Fig. 2) and

considered the median correlation coefficient across random image splits as an indicator of

consistency. The accompanying two-sided *P*-values were Bonferroni adjusted for twelve

comparisons in the *Lon* sample,

To estimate the re-test reliability of individual differences in % cumulative fixation time

or % first fixations for a given attribute, we correlated the corresponding values for the first

session of the *Gi* sample with those of the second.

Covariance patterns between dimensions and correlations with other measures

To investigate the covariance pattern between the dimensions of consistent individual

differences in fixation behaviour, we inspected pair-wise Pearson correlation matrices (Fig. 3B

and 4, left hand side). We further used metric multidimensional scaling to project the pair-wise

dissimilarities between dimensions (defined as $(1-r)^2$) onto distances in a two-dimensional space

(Fig. 3B and 4, right hand side). To further test the stability of covariance patterns between

datasets we calculated pair-wise Pearson correlations between Fisher Z-transformed correlation

matrices from different samples.

To explore the relationship between personality and individual gaze behaviour, we

computed all pairwise correlations between reliable dimensions of salience differences and

personality. Specifically we correlated individual salience differences for *Faces*, *Emotion*, *Text*,

Taste, *Touched* and *Motion* with the Big Five as determined by the IPIP-NEO 120[40] *Neuroticism*

(N), *Extraversion* (E), *Openness* (O), *Assertiveness* (A), *Conscientiousness* (C), with high

sensitivity (HS) as determined by the Highly Sensitive Person scale[42], and with sensation seeking

(SS) as determined by the Brief Sensation Seeking Scale[41]. The resulting 6x7 correlation matrix

was computed separately for the proportion of cumulative fixation time and for that of *first*

fixations; accompanying *p*-values were adjusted for multiple testing using the Holm-Bonferroni

correction for family-wise errors[43].

To test the perceptual implications of salience differences, we concentrated of the

example of faces. Specifically, we probed a correlation between individual face salience as

indicated by the proportion of cumulative dwell time or *first* fixations landing on faces and total

scores in the Cambridge Face Memory Test[70]. This standard test of individual face recognition

skills was administered online to participants from the *Gi* sample, after the first eyetracking

session.

Availability of data and code

Anonymised fixation data and code to reproduce the results presented here are freely

available at https://osf.io/n5v7t/

References

1. Gegenfurtner, K. R. The Interaction Between Vision and Eye Movements. *Perception* 1–25 (2016). doi:10.1177/0301006616657097

2. Curcio, C. A. & Allen, K. A. Topography of ganglion cells in human retina. *J. Comp. Neurol.* **300,** 5–25 (1990).

3. Dougherty, R. F. *et al.* Visual field representations and locations of visual areas V1/2/3 in human visual cortex. *J. Vis.* **3,** 586–98 (2003).

4. Rosenholtz, R. Capabilities and Limitations of Peripheral Vision. *Annu. Rev. Vis. Sci.* 437–459 (2016). doi:10.1146/annurev-vision-082114-035733

5. Henderson, J. M., Williams, C. C., Castelhano, M. S. & Falk, R. J. Eye movements and picture processing during recognition. *Percept. {&} Psychophys.* **65,** 725–734 (2003).

6. Nelson, W. W. & Loftus, G. R. The functional visual field during picture viewing. *J. Exp. Psychol. Hum. Learn.* **6,** 391–399 (1980).

7. Harel, J., Koch, C. & Perona, P. Graph-Based Visual Saliency. *Proceedings of the 19th International Conference on Neural Information Processing Systems* 545–552 (2006).

8. Xu, J., Jiang, M., Wang, S., Kankanhalli, M. S. & Zhao, Q. Predicting human gaze beyond pixels. *J. Vis.* **14,** (2014).

9. Henderson, J. M. & Hayes, T. R. Meaning-based guidance of attention in scenes as revealed by meaning maps. *Nat. Hum. Behav.* **1,** 743–747 (2017).

10. Einhauser, W., Spain, M. & Perona, P. Objects predict fixations better than early saliency. *J. Vis.* **8,** 18–18 (2008).

11. Borji, A. & Itti, L. Defending Yarbus: eye movements reveal observers' task. *J. Vis.* **14,** 29 (2014).

12. Tatler, B. W., Hayhoe, M. M., Land, M. F. & Ballard, D. H. Eye guidance in natural vision: Reinterpreting salience. *J. Vis.* **11,** 5–5 (2011).

13. Itti, L., Koch, C. & Niebur, E. A model of saliency-based visual attention for rapid scene analysis. *IEEE Trans. Pattern Anal. Mach. Intell.* **20,** 1254–1259 (1998).

14. White, B. J. *et al.* Superior colliculus neurons encode a visual saliency map during free viewing of natural dynamic video. *Nat. Commun.* **8,** 14263 (2017).

15. Bogler, C., Bode, S. & Haynes, J.-D. Decoding successive computational stages of saliency processing. *Curr. Biol.* **21,** 1667–71 (2011).

16. Stoll, J., Thrun, M., Nuthmann, A. & Einhäuser, W. Overt attention in natural scenes: objects dominate features. *Vision Res.* **107,** 36–48 (2015).

17. Kummerer, M., Wallis, T. S. A., Gatys, L. A. & Bethge, M. Understanding Low- and High-Level Contributions to Fixation Prediction. in *2017 IEEE International Conference on Computer Vision (ICCV)* 4799–4808 (IEEE, 2017). doi:10.1109/ICCV.2017.513

18. Kümmerer, M., Wallis, T. S. A. & Bethge, M. Information-theoretic model comparison unifies saliency metrics. *Proc. Natl. Acad. Sci. U. S. A.* **112,** 16054–9 (2015).

408   19.   Buswell, G. T. *How people look at pictures: a study of the psychology and perception in art. How*
409   *people look at pictures: a study of the psychology and perception in art.* (Univ. Chicago Press,
410   1935).

411   20.   Yarbus, A. L. in *Eye Movements and Vision* 171–211 (Springer US, 1967). doi:10.1007/978-1-4899-
412   5379-7_8

413   21.   Andrews, T. J. & Coppola, D. M. Idiosyncratic characteristics of saccadic eye movements when
414   viewing different visual environments. *Vision Res.* **39,** 2947–53 (1999).

415   22.   Castelhano, M. S. & Henderson, J. M. Stable individual differences across images in human
416   saccadic eye movements. *Can. J. Exp. Psychol.* **62,** 1–14 (2008).

417   23.   Henderson, J. M. & Luke, S. G. Stable individual differences in saccadic eye movements during
418   reading, pseudoreading, scene viewing, and scene search. *J. Exp. Psychol. Hum. Percept. Perform.*
419   **40,** 1390–400 (2014).

420   24.   Meyhöfer, I., Bertsch, K., Esser, M. & Ettinger, U. Variance in saccadic eye movements reflects
421   stable traits. *Psychophysiology* **53,** 566–578 (2016).

422   25.   Rigas, I. & Komogortsev, O. V. Current research in eye movement biometrics: An analysis based
423   on BioEye 2015 competition. *Image Vis. Comput.* (2016). doi:10.1016/j.imavis.2016.03.014

424   26.   Bargary, G. *et al.* Individual differences in human eye movements: An oculomotor signature?
425   *Vision Res.* **141,** 157–169 (2017).

426   27.   Ettinger, U. *et al.* Reliability of smooth pursuit, fixation, and saccadic eye movements.
427   *Psychophysiology* **40,** 620–628 (2003).

428   28.   Constantino, J. N. *et al.* Infant viewing of social scenes is under genetic control and is atypical in
429   autism. *Nature* **547,** 340–344 (2017).

430   29.   Kennedy, D. P. *et al.* Genetic Influence on Eye Movements to Complex Scenes at Short
431   Timescales. *Curr. Biol.* **27,** 3554–3560.e3 (2017).

432   30.   Parkhurst, D., Law, K. & Niebur, E. Modeling the role of salience in the allocation of overt visual
433   attention. *Vision Res.* **42,** 107–23 (2002).

434   31.   Foulsham, T. & Underwood, G. What can saliency models predict about eye movements? Spatial
435   and sequential aspects of fixations during encoding and recognition. *J. Vis.* **8,** 6 (2008).

436   32.   Einhauser, W., Rutishauser, U. & Koch, C. Task-demands can immediately reverse the effects of
437   sensory-driven saliency in complex visual stimuli. *J. Vis.* **8,** 2 (2008).

438   33.   Anderson, N. C., Ort, E., Kruijne, W., Meeter, M. & Donk, M. It depends on *when* you look at it:
439   Salience influences eye movements in natural scene viewing and search early in time. *J. Vis.* **15,** 9
440   (2015).

441   34.   Mackay, M., Cerf, M. & Koch, C. Evidence for two distinct mechanisms directing gaze in natural
442   scenes. *J. Vis.* **12,** 9–9 (2012).

443   35.   Henderson, J. M. & Hayes, T. R. Meaning guides attention in real-world scene images: Evidence
444   from eye movements and meaning maps. *J. Vis.* **18,** 10 (2018).

445   36.   Nyström, M. & Holmqvist, K. Semantic Override of Low-level Features in Image Viewing – Both
446   Initially and Overall. *J. Eye Mov. Res.* **2,** (2008).

447   37.   Kaiser, D. & Cichy, R. M. Typical visual-field locations enhance processing in object-selective
448   channels of human occipital cortex. *J. Neurophysiol.* **120,** 848–853 (2018).

38. de Haas, B. *et al.* Perception and Processing of Faces in the Human Brain Is Tuned to Typical
Feature Locations. *J. Neurosci.* **36,** (2016).

39. de Haas, B. & Schwarzkopf, D. S. Feature-location effects in the Thatcher illusion. *J. Vis.* **18,**
(2018).

40. Maples, J. L., Guan, L., Carter, N. T. & Miller, J. D. A test of the International Personality Item Pool
representation of the Revised NEO Personality Inventory and development of a 120-item IPIP-
based measure of the five-factor model. *Psychol. Assess.* **26,** 1070–84 (2014).

41. Hoyle, R. H., Stephenson, M. T., Palmgreen, P., Lorch, E. P. & Donohew, R. L. Reliability and
validity of a brief measure of sensation seeking. *Pers. Individ. Dif.* **32,** 401–414 (2002).

42. Aron, E. N. & Aron, A. Sensory-processing sensitivity and its relation to introversion and
emotionality. *J. Pers. Soc. Psychol.* **73,** 345–368 (1997).

43. Holm, S. A simple sequentially rejective multiple test procedure. *Scand. J. Stat.* (1979).

44. Wang, S. *et al.* Atypical Visual Saliency in Autism Spectrum Disorder Quantified through Model-
Based Eye Tracking. *Neuron* **88,** 604–16 (2015).

45. Jones, W. & Klin, A. Attention to eyes is present but in decline in 2-6-month-old infants later
diagnosed with autism. *Nature* **504,** 427–31 (2013).

46. Arcaro, M. J., Schade, P. F., Vincent, J. L., Ponce, C. R. & Livingstone, M. S. Seeing faces is
necessary for face-domain formation. *Nat. Neurosci.* **20,** 1404–1412 (2017).

47. Dehaene, S. *et al.* How learning to read changes the cortical networks for vision and language.
Science **330,** 1359–64 (2010).

48. Kanwisher, N. & Yovel, G. The fusiform face area: a cortical region specialized for the perception
of faces. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **361,** 2109–28 (2006).

49. Tsao, D. Y., Moeller, S. & Freiwald, W. A. Comparing face patch systems in macaques and
humans. *Proc. Natl. Acad. Sci. U. S. A.* **105,** 19514–9 (2008).

50. Grill-Spector, K. & Weiner, K. S. The functional architecture of the ventral temporal cortex and its
role in categorization. *Nat. Rev. Neurosci.* **15,** 536–548 (2014).

51. McCandliss, B. D., Cohen, L. & Dehaene, S. The visual word form area: expertise for reading in the
fusiform gyrus. *Trends Cogn. Sci.* **7,** 293–299 (2003).

52. Kourtzi, Z. & Kanwisher, N. Activation in human MT/MST by static images with implied motion. *J.
Cogn. Neurosci.* **12,** 48–55 (2000).

53. Orlov, T., Makin, T. R. & Zohary, E. Topographic representation of the human body in the
occipitotemporal cortex. *Neuron* **68,** 586–600 (2010).

54. Weiner, K. S. & Grill-Spector, K. Neural representations of faces and limbs neighbor in human
high-level visual cortex: evidence for a new organization principle. *Psychol. Res.* **77,** 74–97 (2013).

55. Adamson, K. & Troiani, V. Distinct and overlapping fusiform activation to faces and food.
Neuroimage **174,** 393–406 (2018).

56. Bush, J. C., Pantelis, P. C., Morin Duchesne, X., Kagemann, S. A. & Kennedy, D. P. Viewing
Complex, Dynamic Scenes "Through the Eyes" of Another Person: The Gaze-Replay Paradigm.
PLoS One **10,** e0134347 (2015).

57. Wilmer, J. B. How to use individual differences to isolate functional organization, biology, and
utility of visual functions; with illustrative proposals for stereopsis. *Spat. Vis.* **21,** 561–79 (2008).

58. Charest, I. & Kriegeskorte, N. The brain of the beholder: honouring individual representational
idiosyncrasies. *Lang. Cogn. Neurosci.* **30,** 367–379 (2015).

59. Peterzell, D. Discovering Sensory Processes Using Individual Differences: A Review and Factor
Analytic Manifesto. *Electron. Imaging* 1–11 (2016).

60. Haldemann, J., Stauffer, C., Troche, S. & Rammsayer, T. Processing Visual Temporal Information
and Its Relationship to Psychometric Intelligence. *J. Individ. Differ.* **32,** 181–188 (2011).

61. Hayes, T. R. & Henderson, J. M. Scan patterns during real-world scene viewing predict individual
differences in cognitive capacity. *J. Vis.* **17,** 23 (2017).

62. Troche, S. J. & Rammsayer, T. H. Attentional blink and impulsiveness: evidence for higher
functional impulsivity in non-blinkers compared to blinkers. *Cogn. Process.* **14,** 273–81 (2013).

63. Wu, D. W.-L., Bischof, W. F., Anderson, N. C., Jakobsen, T. & Kingstone, A. The influence of
personality on social attention. *Pers. Individ. Dif.* **60,** 25–29 (2014).

64. Hayes, T. R. & Henderson, J. M. Scan patterns during scene viewing predict individual differences
in clinical traits in a normative sample. *PLoS One* **13,** e0196654 (2018).

65. Armstrong, T. & Olatunji, B. O. Eye tracking of attention in the affective disorders: a meta-analytic
review and synthesis. *Clin. Psychol. Rev.* **32,** 704–23 (2012).

66. Molitor, R. J., Ko, P. C. & Ally, B. A. Eye Movements in Alzheimer's Disease. *J. Alzheimers. Dis.* **44,**
1 (2015).

67. Crouzet, S. M., Kirchner, H. & Thorpe, S. J. Fast saccades toward faces: Face detection in just 100
ms. *J. Vis.* **10,** 1–17 (2010).

68. Rösler, L., End, A. & Gamer, M. Orienting towards social features in naturalistic scenes is
reflexive. *PLoS One* **12,** e0182037 (2017).

69. Costa, P. T. & McCrae, R. R. Four ways five factors are basic. *Pers. Individ. Dif.* **13,** 653–665 (1992).

70. Duchaine, B. & Nakayama, K. The Cambridge Face Memory Test: results for neurologically intact
individuals and an investigation of its validity using inverted face stimuli and prosopagnosic
participants. *Neuropsychologia* **44,** 576–85 (2006).

71. Kleiner, M., Brainard, D. & Pelli, D. What's new in Psychtoolbox-3? *Perception* **36,** 14 (2007).

72. Pelli, D. G. The VideoToolbox software for visual psychophysics: transforming numbers into
movies. *Spat. Vis.* **10,** 437–42 (1997).

73. Gibaldi, A., Vanegas, M., Bex, P. J. & Maiello, G. Evaluation of the Tobii EyeX Eye tracking
controller and Matlab toolkit for research. *Behav. Res. Methods* **49,** 923–946 (2017).

Acknowledgements

This work was supported by a JUST'US fellowship from the University of Giessen (B.d.H), as

well as a research fellowship by the Deutsche Forschungsgemeinschaft (DFG HA 7574/1-1;

B.d.H.). KRG was supported by DFG Collaborative Research Center SFB/TRR 135. We would

like to thank Xu et al. for publishing their stimuli and dataset, Dr. Pete R. Jones for binding code

bridging between the Eye-X C library and MATLAB, as well as Ms Diana Weissleder for help

with collecting the *Gi* datasets.

Author Contributions

532   Conception and design: BdH, Data Collection: ALI, BdH, Data Analysis: BdH, in discussion

533   with KRG. Provision of lab space and resources: KRG, DSS. Manuscript writing: BdH wrote the

534   initial draft. All authors discussed the results and final manuscript.

Competing Interests

538   The authors declare no competing financial interests.

544 **Figures**

% cum. fix. **time** (odd / even images)
% **first** fix. (odd / even images)

Figure 1. Consistent individual differences in fixation behaviour along six semantic dimensions. For each semantic attribute, the grey scatter plot shows individual proportions of cumulative fixation time for the odd *versus* even numbered images in the *Lon* dataset. The green scatter plot shows the corresponding individual proportions of *first* fixations after image onset. For each dimension, two example images are given and overlaid with the fixations from one observer strongly attracted by the corresponding attribute (orange frames) and one observer weakly attracted by it (blue frames). The overlays show the first fixation after image onset as a green circle; any subsequent fixations are shown in purple. The two data points corresponding to the example observers are highlighted in the scatter plot, corresponding to the colour of the respective image frames.

Figure 2. Consistency of results across images, datasets and time. (a) Distribution plots of consistency correlations for each of the twelve semantic dimensions tested (as indicated by the labels on the x-axis in b). Results are shown separately for the four observer samples tested (as indicated by the row labels). Gi_1 and Gi_2 refer to the first and second appointment of the Gi sample. The grey left-hand leaf of each distribution plot shows a histogram of split-half

correlations for 1000 random splits of the image set, the green right hand leaf shows the
corresponding histogram for *first* fixations after image onset. Overlaid dots indicate the median
consistency correlation for each distribution. High split-half correlations indicate consistent
individual differences in fixation behaviour across images for a given dimension. The dashed red
line separates the six attributes found to be consistent dimensions of individual differences in the
Lon sample from the remaining dimensions. (**b**) Re-test reliability across Gi_1 and Gi_2. The
magnitude of re-test correlations for individual dwell time and proportion of first fixations is
indicated by grey and green bars, respectively. All correlation and *P*-values can be found in
Table 1.

Figure 3. Covariance of individual differences along semantic dimensions. (**a**) Grey scatter plots show the individual proportion of cumulative fixation time (in %) for *Faces versus Text* (left hand side) and *Faces versus* objects being *Touched* (right hand side). Green scatter plots show the corresponding data for the individual proportion of *first* fixations after image onset. (**b**) Correlation matrix for individual differences along five semantic dimensions (left hand side; note that the labels for emotional and neutral faces were collapsed for this analysis). Colour indicates pairwise Pearson correlation coefficients as indicated by the bar. *Motion* and *Face* are positively correlated with each other, but negatively correlated with the remaining dimensions, as captured by a two-cluster solution of metric dimensional scaling to two dimensions (right hand side). All data shown are based on individual proportions of fixation time in the *Lon* dataset. The corresponding supplemental Figure S1, shows the consistency of this pattern for the proportion of first fixations and across all four datasets.

Figure 4. Consistency of covariance pattern across datasets. (**a**) shows covariance patterns for dimensions of individual differences in cumulative fixation durations and (**b**) those for individual differences in the proportion of *first* fixations. Each row shows the results for one dataset as indicated (Lon, Gi_1, Gi_2 and Xu *et al*.). Colours of the left hand correlation matrices indicate pairwise Pearson correlation coefficients between dimensions as indicated by the bar. Please refer to Figure 3 for dimension labels (from top to bottom: taste, text, motion, face; from left to right: motion, text, taste, touched). Note the similarity of covariance patterns between datasets. All pairwise correlations between (Fisher z-transformed) correlation matrices were >.68 for cumulative fixation durations and >.90 for first fixations. The right hand side scatter plots show the results of metric multidimensional scaling onto two dimensions for cumulative fixation duration (**c**) and first fixations (**d**). This analysis yielded a distinct cluster for *Faces* and *Motion*, as well as a cluster for *Text*, *Touched Taste* for first fixations in all datasets. This structure was very similar for cumulative dwell time, with the exception of *Text* separating from *Touched* and *Taste* in the Gi_1 and Gi_2 data.

Figure 5. Correlations between fixation behaviour, face recognition skills and personality.
(**a**) Performance on the Cambridge Face Memory Test (CMFT) correlated significantly with the
individual proportion of first fixations landing on faces when freely viewing complex scenes
(green, right hand side). This correlation was not significant for cumulative fixation time across
the viewing time of three seconds (grey, left hand side). (**b**) Neither cumulative fixation time
(grey frame, left hand side), nor the proportion of first fixations (green frame, right hand side)
towards any of the six dimensions of individual gaze behaviour correlated significantly with any
of the tested dimensions of personality (N: Neuroticism, E: Extraversion, O: Openness, A:
Assertiveness, C: Conscientiousness, HS: High Sensitivity, SS: Sensation Seeking). All data
from the Gi_1 sample.

 Table

	Ind. % fixation time				Ind. % first fix			
	r	*P*	range [%]	max/min	*r*	*P*	range [%]	max/min
Faces - Lon	.94	<.001*	24-43	1.80	.88	<.001*	24-43	1.77
– Gi_1	.91	<.001	21-35	1.64	.88	<.001	15-42	2.82
– Gi_2	.91	<.001	17-34	1.99	.89	<.001	15-43	2.76
– Xu *et al.*	.95	<.001	23-35	1.50	.86	<.001	24-41	1.69
- Re-test (Gi_1-2)	.80	<.001	-	-	.86	<.001	-	-
Emotion - Lon	.82	<.001*	8-15	1.93	.69	<.001*	9-18	1.88
– Gi_1	.80	<.001	8-13	1.68	.76	<.001	8-17	2.20
– Gi_2	.82	<.001	7-13	1.79	.75	<.001	8-17	2.22
- Xu *et al.*	.89	<.001	8-12	1.56	.48	.07	13-17	1.31
- Re-test (Gi_1-2)	.85	<.001	-	-	.79	<.001	-	-
Text - Lon	.81	<.001*	9-16	1.76	.77	<.001*	6-17	2.96
– Gi_1	.90	<.001	9-19	1.97	.82	<.001	5-16	3.11
– Gi_2	.91	<.001	9-20	2.21	.82	<.001	4-17	3.91
- Xu *et al.*	.94	<.001	8-18	2.16	.83	<.001	4-13	3.25
- Re-test (Gi_1-2)	.84	<.001	-	-	.89	<.001	-	-
Touched - Lon	.73	<.001*	11-16	1.52	.73	<.001*	6-16	2.48

– Gi_1	.74	<.001	9-15	1.63	.71	<.001	6-15	2.65
– Gi_2	.75	<.001	9-16	1.67	.72	<.001	6-16	2.74
- Xu *et al.*	.91	<.001	9-16	1.85	.65	.008	7-13	1.81
- Re-test (Gi_1-2)	.70	<.001	-	-	.78	<.001	-	-
Taste - Lon	.70	<.001*	10-16	1.58	.57	<.001*	10-16	1.67
– Gi_1	.70	<.001	11-15	1.38	.29	.04	10-14	1.36
– Gi_2	.74	<.001	10-17	1.67	.50	<.001	9-16	1.75
- Xu *et al.*	.84	<.001	11-16	1.44	.69	.004	9-15	1.72
- Re-test (Gi_1-2)	.78	<.001	-	-	.62	<.001	-	-
Motion - Lon	.64	<.001*	28-36	1.31	.35	.15*	31-39	1.25
– Gi_1	.65	<.001	28-34	1.22	.52	<.001	30-38	1.29
– Gi_2	.73	<.001	26-33	1.29	.52	<.001	28-37	1.32
- Xu *et al.*	.85	<.001	28-33	1.19	.62	.01	33-40	1.22
- Re-test (Gi_1-2)	.68	<.001	-	-	.78	<.001	-	-

Table 1. Range and median consistency of individual differences in fixation behaviour towards six semantic dimensions. The left hand side of the table shows data for individual differences in the proportion of cumulative fixation time (across three seconds) spent on objects with the respective semantic attributes. The right hand side shows data on individual differences in the proportion of first fixations after image onset landing on objects with those attributes. For each semantic attribute, the data from different samples (*Lon, Gi_1, Gi_2* and *Xu et al.*) is presented in separate rows. The range across participants is given in % fixation time or % of first fixations and the max/min ratio indicates the relative difference between observers attracted the most or least by a given attribute. Pearson correlations (*r*) indicate the median split-half

correlation of individual differences across 1000 random image splits and the corresponding *P*-value (Bonferroni corrected for 12 semantic attributes in the *Lon* sample, as indicated by the *). Consistency correlations failing to reach statistical significance are highlighted in red. The final row for each attribute shows the corresponding re-test reliability across several weeks.

Supplementary Information

Supplementary Methods

To test the relationship between individual salience and visual field biases, we first explored the distribution of objects corresponding to dimensions of individual salience across image space (Supplementary Fig. 1a; neutral and emotional faces collapsed into *Faces*). The most prominent spatial bias in the scene stimuli was a strong tendency for faces to appear in the upper visual field (~75% of faces).

Therefore, we tested whether individual face salience (individual % of *first* fixations landing on faces) correlated with a general upper visual field bias. We defined the latter as the median elevation of *first* fixations towards images *not* containing a face (relative to central fixation and expressed in % image height). This was indeed the case (r = .49-.78 across samples, all P < .05; Supplementary Fig. 1b).

Given a correlation between individual face salience and a general bias towards the upper visual field, we wanted to test whether individual salience preferences *depended* on visual field biases. Therefore, we tested whether individual differences in face salience persisted for the lower visual field, correlating the individual percentage of first fixations landing on faces in the upper visual field (UVF) with that landing on faces in the lower visual field (LVF). For this analysis, we defined UVF and LVF faces as those for which the median height of the pixel mask was in the upper and lower image half, respectively. Importantly, individual face salience was stable across the UVF and LVF, showing that it persisted independently of the UVF bias (r = .61-.87 across samples, all P < .05; Supplementary Fig. 1c).

Supplementary Figure

Supplementary Figure 1. Individual salience and visual field biases. (a) Distribution of
different types of objects in image space across scenes. Notice the upwards bias of Faces. **(b)**
Individual face salience significantly correlated with a general upper visual field bias in all four
samples (i.e. with the elevation of first fixations towards images without a face; see
Suppelemtary Methods above for details). **(c)**. Nevertheless, individual face salience persisted
independent of this spatial bias. The individual proportion of first fixations landing in the upper
and lower visual field (UVF and LVF) were highly correlated with each other.

688
689