EEG signal diversity reflects capability for behavioral responsiveness during propofol sedation

Michał Bola[1], Paweł Orłowski[1,2], Martyna Płomecka[1,3], Artur Marchewka[1]

1: Laboratory of Brain Imaging, Nencki Institute of Experimental Biology of Polish Academy of Sciences,

Warsaw, Poland

2: Institute of Philosophy, University of Warsaw, Warsaw, Poland

3: Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, Warsaw, Poland

Corresponding author: Michał Bola, PhD; email: m.bola@nencki.gov.pl

Funding statement: This study was funded by Sonata grant from the National Science Centre Poland

(2015/17/D/HS6/00269) and Iuventus Plus grant from the Polish Ministry of Science and Higher

Education (082/IP3/2016/74). MB was additionally supported by a stipend from the Polish Ministry of

Science and Higher Education (555/STYP/11/2016).

Conflict of interest: The authors declare no competing interests.

Author Contributions: Conceived the study: MB; Analyzed data: PO, MB; Contributed analysis

methods: MP; Interpreted data: MB, AM; Wrote the paper: MB. Revised the paper: MB, AM.

Abstract

Background: Transitions between wakefulness and anesthesia are accompanied by profound changes in brain functioning. A key challenge is to disentangle neuronal mechanisms specific to loss and recovery of consciousness, from more general effects that are not directly related to the capacity for conscious experience. Measures of neuronal diversity have been recently proposed to constitute a robust correlate of the global states of consciousness. In the present study we investigated whether EEG signal diversity is indeed related to behavioral responsiveness during propofol sedation, or rather to the general drug-related effects.

Methods: We reanalyzed data collected from 20 subjects sedated with propofol. Based on the responsiveness to auditory stimuli all subjects were subdivided into two subgroups - *responsive* (n = 13), who remains awake throughout the experiment, and *drowsy* (n = 7), who becomes unresponsive during moderate sedation. Resting state EEG recorded during wakefulness and sedation was characterized by the Mean Information Gain - an information-theory measure estimating signal diversity.

Results: In line with our hypothesis, the *drowsy* group exhibited a decrease in diversity during sedation but, unexpectedly, the *responsive* group exhibited a robust increase in diversity (ANOVA *group* x *state* interaction: $F(3) = 7.81$, $p < 0.001$; $BF_{10} > 100$).

Conclusions: We revealed that propofol sedation is initially related to an increase in EEG signal diversity, and that only upon loss of responsiveness EEG diversity decreases. The qualitatively different pattern of changes in the *responsive* and *drowsy* groups makes EEG diversity a robust indirect index of responsiveness and, presumably, consciousness.

Introduction

What are the neuronal mechanisms of consciousness? One of the main hypotheses is that the capacity for conscious experience relies on the brain's ability to generate a large repertoire of functional states[1, 2]. However, the temporal diversity of brain states needs to be balanced between order and disorder, as both insufficient temporal diversity (i.e. dwelling in the same state for long time periods) and excessive diversity (i.e. chaotic transitions between states) will likely cause alterations in the global state of consciousness.

These hypotheses have been tested by recent studies, which established neuronal diversity as a robust marker of the global states of consciousness[3]. Specifically, it has been shown that brain signals are relatively complex and diverse during wakefulness, but they become simple and stereotypical during NREM sleep[4-7] and general anesthesia[8-10]. The diversity measures have been subsequently applied in the clinical context to successfully classify patients with disorders of consciousness[11, 12]. Interestingly, recent studies provided first evidence that not only decreases, but also increases in diversity of brain states are in principle possible, specifically during psychedelic states caused by LSD, ketamine, or psilocybin[13-15].

Based on these findings brain signal diversity has been interpreted as a correlate of consciousness. However, what limits this interpretation is that conscious and unconscious states differ not only with respect to the presence or absence of conscious experiences. In the context of anesthesia not only neuronal mechanisms specific to loss of consciousness can be observed (i.e. state-related), but also more general effects of the anesthetic that are not directly related to capacity for conscious experiences (i.e. drug-related). Thus, a key challenge is to disentangle these effects in order to distil the genuine neuronal mechanisms of consciousness. One possible approach is to take advantage of the between-subjects variability in susceptibility to anesthetics[16] and compare brain activity observed in subjects who are sedated and unconscious, with subjects who are sedated with similar anesthetic doses but remain conscious. Comparing such groups would allow testing whether a given measure is indeed related to

consciousness, as then it would change only in the unconscious group, or rather to the general effects of an

anesthetic, as then a change would be observed in both groups.

In the present study we adopted the outlined approach to investigate whether signal diversity is indeed

related to behavioral responsiveness and, presumably, consciousness. We reanalyzed data from an

experiment conducted by Chennu and colleagues[17], in which all subjects were sedated with similar doses

of propofol, but based on their responsiveness to auditory stimuli they could be subdivided into two

subgroups: sedated but responsive, and sedated and unresponsive. Resting-state EEG recorded during

wakefulness and sedation was characterized in terms of two information-theory measures: the Mean

Information Gain (MIG), which measures *diversity* understood as randomness or entropy of the data, and

Fluctuation Complexity (FC), which is a measure of *complexity* understood as the state intermediate

between order and disorder (**Fig. S1**). We hypothesized to observe a decrease of diversity and complexity

during sedation, but only in the unresponsive group.

Methods

Subjects and data collection

In the present study we reanalyzed data first published by Chennu et al.[17]. The procedures regarding data collection and preprocessing are briefly described below and the original paper should be consulted for additional details. Ethical approval for the study was provided by the local ethics committee and all subjects provided a written informed consent. The data are accessible from https://www.repository.cam.ac.uk/handle/1810/252736.

Data collected from 20 healthy subjects (11 females; mean age = 30.85, SD = 10.98) were used in the present analysis. Data were collected in four states: wakeful baseline, mild sedation, moderate sedation, and recovery. During each state subjects first performed an auditory discrimination task (for 3 to 5 minutes), and afterwards 7 minutes of resting-state EEG was recorded. States of mild and moderate sedation were obtained by performing a target-controlled infusion of propofol with a computerized syringe driver. The targeted blood plasma levels of propofol were 0.6 mcg/ml for mild sedation and 1.2 mcg/ml for moderate sedation. At each targeted level a period of 10 minutes was allowed to obtain a steady propofol concentration and only afterwards behavioral and EEG measurements were performed. Blood samples were taken in each state to evaluate the actual propofol concentration in blood.

To evaluate subjects' responsiveness in each state they were asked to perform a simple auditory discrimination task. Specifically, to indicate, by pressing on of two buttons, whether a presented binaurally stimulus was a broadband noise ("noise") or a harmonic complex with a 150 Hz fundamental frequency ("buzz"). 40 stimuli (20 of each kind) were presented in a random order with a mean inter-stimulus interval of 3 s. A *hit-rate* was defined as a proportion of correct responses. Binominal modelling of hit-rates obtained during baseline wakefulness and moderate sedation was used to classify subject into either drowsy or responsive subgroups (details of the procedure can be found in: [17]).

In each state approximately 7 minutes of eyes-closed resting-state EEG was recorded with high-density

128 electrodes caps and the Net Amps 300 amplifier (Electrical Geodesic Inc., Eugene, Oregon, USA).

Offline preprocessing was conducted with EEGlab[18]. First, 37 channels placed over neck, cheeks, and

forehead were excluded and only data from 91 channels covering the scalp area were retained and further

analyzed. EEG signals were filtered between 0.5 Hz and 45 Hz and segmented into 10-seconds long

epochs. Mean value was subtracted from signal from each channel in an epoch. Semi-automatic procedure

was used to remove noisy epochs and channels. The mean ± SD number of retained epochs was: baseline

wakefulness - 38 ± 5; mild sedation - 39 ± 4; moderate sedation - 38 ± 4; recovery - 40 ± 2 and did not

differ significantly across states. Missing channels were interpolated using the spherical spline

interpolation algorithm. As the last step, the data were re-referenced to the average of all channels.

The EEG data were published in the already preprocessed form and we did not apply any further data

transformations before calculating the measures described below.

Diversity and complexity analysis

a)   Mean information Gain (*MIG*)

The Mean Information Gain (*MIG*) is one of the information theory-based measures. *MIG* is a measure

of diversity, as it exhibits maximum values for random signals. The details and a formal definition of *MIG*

can be found in: [19, 20]. An application of *MIG* to EEG data was described by Wang et al.[15] and by Bola et

al.[21], and here we followed the same procedure.

Briefly, to calculate *MIG* a signal has to be binarized. Here, for each epoch and channel separately, a

median value was used as a threshold, and "1" was assigned to the time-points exceeding the median

value, while "0" was assigned to the time-points that were below or at the median value. Then binary

series were partitioned into "words" of L length, where L is the "window length' or 'information length',

and each "word" reflects the state of a system at a given time-point. Using $L = 4$, as in the main analysis

reported here, the possible states/words were 0000, 0001, 0010, 0100, 1000, 0011,0110, 1100, 1010, 0101,

1001, 0111, 1011, 1101, 1110, 1111. A transition to a next state occurs by a shifting a window forward by

one symbol.

Calculating *MIG* starts with the Shannon information content. We define the probability p_i that

the system is currently in a state '*i*' by dividing the number of occurrences of a state '*i*' by the total number

of occurrences[19]. Based on the probabilities of observing transition from state i to j, defined as p_{ij}, we

calculate the transition probability $p_{i \to j} = \frac{p_{ij}}{p_i}$. In the same way, the reverse conditional probability can be

obtained as $p_{i \leftarrow j} = \frac{p_{ij}}{p_j}$.

The net information gain can be written as $G_{ij} = \log_2(\frac{1}{p_{i \to j}})$, and by averaging the gains over all

transitions the *MIG* is estimated:

$$MIG = \sum_{i=1}^{N} p_{ij} \, G_{ij} = \sum_{i=1}^{N} p_{ij} \, log_2 \left(\frac{1}{p_{i \to j}} \right) = - \sum_{i=j}^{N} p_{ij} \, log_2 \, p_{i \to j}$$

b) Fluctuation Complexity (*FC*)

Another example of an information theory-based measure is Fluctuation Complexity (*FC*). The details

and a formal definition of *FC* can be found in: [19, 20]. While applying *FC* to EEG data we again followed

the procedure described in Wang et al.[15] and Bola et al.[21]. Unlike *MIG*, *FC* does not consider a random

signal to be complex. In the computation of *FC*, the EEG signals were binarized in the same way as

described for *MIG*.

While shifting from one state to another, the newly observed symbol contributes to a gain of

information, defined as $G_{ij} = \log_2(\frac{1}{p_{i \to j}})$. At the same time we forget the first symbol from the previous

word. The loss of information can be expressed as $L_{ij} = \log_2(\frac{1}{p_{i \leftarrow j}})$.

In average for the whole symbol sequence, information gain and loss eliminate each other. The

fluctuation complexity is the mean square deviation of the net information gain (i.e. the differences

between information gain and loss):

$$\text{FC} = \langle (\Gamma - \langle \Gamma \rangle)^2 \rangle = \sum_{I=1}^{N} p_{ij} \left(\log_2(\frac{p_i}{p_j}) \right)^2$$

The more this balance of information gain and loss is fluctuating in the given EEG time series the

more complex is the signal in the sense of the fluctuation complexity.

c) Lempel-Ziv (*LZs* and *LZc*)

MIG and *FC* have used in recent studies to evaluate diversity and complexity of EEG signals[15, 21]. But

majority of studies have employed the Lempel-Ziv algorithm to estimate EEG signal diversity[3, 10].

Therefore, for consistency with the previous studies in the present analysis we used two version of the

Lemepl-Ziv measures, single-channel Lempel-Ziv (*LZs*), which captures temporal variability, and multi-

channels Lempel-Ziv (*LZc*), which captures spatio-temporal variability. Similarly to *MIG*, Lempel-Ziv is a

measure of diversity, as it reaches maximum values for random signals. We used an implementation of the

method described by Schartner et al.[7]. Results of the *LZs* and *LZc* analysis are presented in the

supplementary materials.

To estimate *LZs* EEG signal from each channel and epoch was assessed independently by demeaning

(within 10 s long segment), dividing by standard deviation, and removing a linear trend. Hilbert transform

was used to estimate the envelope of a signal (an absolute value of the analytic signal) and the mean of the

envelope was used as a threshold to binarize the signal, i.e. "1" was assigned to time-points exceeding the

mean value, while "0" was assigned to measurements that were below or at the mean value. The binarized

signal was then segmented into blocks using the encoding step of the Lempel-Ziv compression algorithm.

The number of blocks is the raw *LZs* score for the segment. Next the raw *LZs* value was normalized by the

raw *LZs* value obtained from the same binary signal shuffled in time. Thus, the final (normalized) *LZs*

score varies between 0 and 1 indicating, respectively, minimally and maximally diverse signals.

To estimate *LZc* the 64 time series (one for each EEG channel) from a 10 s segment of data are

binarized as described above for *LZs*, then concatenated observation-by-observation into one binary string

such that the first 64 digits of that string are the observations of the 64 channels at time step 1, the next 64

are those at time step 2, etc. The diversity of this binary string is then assessed in the same way as

described for *LZs* above.

Statistical analysis

All values presented as Mean ± Standard Error of the Mean (SEM), unless stated otherwise. All

statistical analyses were conducted in JASP software[22]. The performed tests involved two-way repeated-

measures ANOVA, with within-subjects factor *state* (4 levels: wakeful baseline, mild sedation, moderate

sedation, recovery) and between-subjects factor *group* (2 levels: drowsy, responsive). Subsequent post-

hoc tests were computed, either paired-samples t-tests to compare between states, or independent-samples

t-tests to compare between groups. Post-hoc comparisons were corrected for multiple comparisons using

the Bonferroni-Holm procedure[23].

The traditional null-hypothesis significance testing approach was complemented with Bayesian

statistics in order to enable testing for the lack of differences between variables. The Bayes Factor (BF_{10})

is defined as the ratio of the probability of observing the data given the alternative hypothesis is true, to

the probability of observing the data given the null hypothesis is true. Thus BF_{10} provides a continuous

measure of evidential support and is typically interpreted according to Lee and Wagenmakers [24]: $BF_{10} <$

0.1 indicates strong evidence in favor of the null hypothesis (i.e. lack of an effect); $0.1 < BF_{10} < 0.33$

indicates moderate evidence in favor of the null hypothesis, $0.33 < BF_{10} < 3$ indicates inconclusive data, 3

$< BF_{10} < 10$ indicates moderate evidence in favor of the alternative hypothesis, $BF_{10} > 10$ indicates strong

evidence in favor of the alternative hypothesis,. Bayesian equivalents of a two-way ANOVA and t-test

were calculated using a medium prior scale (Cauchy scale: .707).

In an additional analysis we aimed to test whether between-group differences in MIG observed during

moderate sedation can be explained by differences in propofol concentration. We thus conducted a one-

way ANOVA with a *group* factor, which in this case is equivalent to an independent-samples t-test, and

included a propofol concentration as a covariate.

Results

In the present study we reanalyzed a data-set published by Chennu et al.[17]. In their study 20 subjects

were sedated with propofol and data were acquired in four states: baseline wakefulness, mild sedation,

moderate sedation, and recovery. In each state subjects performed an auditory discrimination task to

assess their responsiveness and next resting-state EEG was recorded. The profiles of hit rates, defined as a

proportion of correct responses, indicate that subjects can be subdivided into two subgroups - *responsive*

(n = 13), who remains awake throughout the experiment, and *drowsy* (n = 7), who becomes unresponsive

during moderate sedation (**Fig. 1**). Comparing both groups in terms of propofol concentration in blood we

found a significant *state* x *group* interaction ($F_{(2)}$ = 124.6, $p < 0.001$; $BF_{10} > 100$). Specifically, there was

no difference between groups during mild sedation (0.41 ± 0.04 vs. 0.50 ± 0.08 mcg/ml; $t_{(18)}$ = 1.05, p =

0.30; BF_{10} = 0.59) and recovery (0.27 ± 0.02 vs. 0.32 ± 0.03; $t_{(18)}$ = 1.37, p = 0.18; BF_{10} = 0.77), but

during moderate sedation the *drowsy* group had a higher in-blood propofol concentration (0.79 ± 0.05 vs.

1.09 ± 0.12; $t_{(18)}$ = 2.60, p = 0.018; BF_{10} = 3.37).

The recorded resting-state EEG signals were characterized in terms of diversity and complexity using

two information theory measures[19] (**Fig. S1**). Diversity, understood as randomness of the data, was

estimated by the Mean Information Gain (MIG). Statistical analysis of MIG indicates no main effect of

group ($F_{(1)}$ = 1.8, p = 0.19; BF_{10} = 0.80) and no effect of *state* ($F_{(3)}$ = 2.59, p = 0.062; BF_{10} = 1.27), but a

significant *group x state* interaction was found ($F_{(3)}$ = 7.81, $p < 0.001$; $BF_{10} > 100$). Subsequent post-hoc

tests indicate that, with respect to the wakeful baseline, MIG increases during mild sedation ($t_{(19)}$ = 3.34,

p = 0.020; BF_{10} = 12.7) and recovery ($t_{(19)}$ = 3.15, p = 0.026; BF_{10} = 8.83), but not during moderate

sedation ($t_{(19)}$ = 2.47, p = 0.093; BF_{10} = 2.56). Importantly, lack of the significant effect during moderate

sedation is due to a different pattern of changes in both subgroups, which is confirmed by a significant

difference between groups during moderate sedation ($t_{(18)}$ = 3.24, p = 0.005; BF_{10} = 9.20) but not during

other states (all $p > 0.10$; all $0.3 < BF_{10} < 1$). Specifically, the *drowsy* group exhibits a decrease in MIG

during moderate sedation (with respect to baseline) but the *responsive* group exhibits an increase.

Next we analyzed the Fluctuation Complexity (FC), which is a measures of complexity understood as a

state intermediate between an order and disorder. We found a significant effect of *state* ($F_{(3)} = 4.46$, p =

0.007; $BF_{10} = 9.56$), but no effect of *group* ($F_{(1)} = 0.36$, p = 0.55; $BF_{10} = 0.50$) and no *group x state*

interaction ($F_{(3)} = 0.74$, p = 0.53; $BF_{10} = 1.29$). Subsequent post-hoc tests indicate that, with respect to

baseline wakefulness, FC decreases during mild sedation ($t_{(19)} = 3.20$, p = 0.028; $BF_{10} = 9.65$).

Comparison of wake with moderate sedation is inconclusive ($t_{(19)} = 2.57$, p = 0.092; $BF_{10} = 3.08$), as *p*

value reaches a trend level only but BF_{10} indicates evidence for a difference between states. All other

comparisons between states were not significant (all p > 0.10; all $BF_{10} < 3$).

Further, we report results of three additional analyses. First, inspection of the hit-rates profiles suggests

that 2 subjects became unresponsive already during the mild sedation state (both had hit-rate < 25%; **Fig.**

1). Plotting hit-rates against changes in MIG demonstrates that exactly these two subjects exhibit a robust

decrease in MIG already during mild-sedation (**Fig. 2C**). While statistical analysis cannot be conducted on

such a small subsample of 2 subjects, this observation provides an additional, qualitative piece of evidence

that MIG is closely related to responsiveness. However, similar scatter-plots created for FC clearly show

that FC does not differentiate between responsive and drowsy groups in neither state.

Second, we interpret the difference between groups in MIG during moderate sedation as related to

behavioral responsiveness, but the groups differ also in terms of propofol concentration in blood. Thus, to

further support our interpretation we compared MIG between groups with propofol concentration included

as a covariate. We found a significant difference between groups in MIG ($F_{(1)} = 9.53$, p = 0.007) but the

effect of a propofol concentration was not significant ($F_{(1)} = 0.48$, p=0.49). Therefore, it is unlikely that

the between-groups difference in anesthetic concentration can account for the difference in diversity.

Third, for consistency with several previous studies[7, 10, 13] which used the Lemepl-Ziv algorithm to

estimate signal diversity we also applied this measure to our data. The results of the Lempel-Ziv analysis

can be found in the supplementary materials.

Discussion

Measures of neuronal diversity have been recently established as robust markers of the global states of consciousness[3-13]. However, it is not clear whether the diversity measures indeed reflect consciousness *per se*, or rather some general and unspecific effects related to a transition between conscious and unconscious states. Here we aimed to address this question in the context of anesthesia by investigating EEG signal diversity and complexity during propofol sedation. Importantly, despite receiving a similar dose of propofol all tested subjects could be divided into two subgroups: *responsive*, who remained awake throughout the experiment, and *drowsy*, who became unresponsive and unconscious. We hypothesized that the loss of responsiveness during sedation will be related to a decrease of EEG diversity. This was indeed confirmed, as 6/7 subjects from the *drowsy* group exhibited a drop in diversity upon loss of responsiveness. However, the unexpected finding is that EEG diversity increased in 12/13 subjects who were sedated but remained responsive. Such a qualitative difference - an increase in diversity in the responsive group and a decrease in the unresponsive group - makes it a robust index of responsiveness and, presumably, consciousness.

Several recent studies demonstrated that general anesthesia is accompanied by a marked decrease of neuronal diversity as estimated by the Lempel-Ziv algorithm[9, 10]. Further evidence for simplification and reduced repertoire of brain states during anesthesia was provided by studies characterizing brain signals in terms of functional interactions[25], long-range temporal correlations[26, 27], or using dynamic systems theory measures[28-30]. Our work adds to the literature by demonstrating this effect using "minimal" contrasts between states and groups. By minimal we mean that, first, we detected a robust decrease in diversity upon loss of responsiveness even though the doses of propofol were low (e.g. in comparison to Sarasso et al.[9] and Schartner et al.[10]). Second, we included also sedated but responsive subjects, who therefore constitute an important "control group" missing from previous experiments. Lack of a diversity decrease in this group further strengthens our conclusion that EEG diversity is closely related to the ability to respond to external stimuli.

Whereas the diversity decrease in the drowsy group during moderate sedation is in line with the

published results, an increase in the sedated but responsive group is a surprising finding. Further, during

mild sedation and recovery - states when all subjects were sedated but responsive - diversity increase was

indeed observed in all subjects (importantly, propofol was still detected in blood samples during recovery;

see: **Fig. 1**). To our knowledge, such an effect has not been reported so far, even though the state of mild

sedation has been included in some previous studies. For instance, Schartner et al.[10] analyzed EEG

recorded during mild propofol sedation, when subjects were partially conscious (Ramsay scale score = 3),

but found a decrease in EEG Lempel-Ziv diversity. However, in the study of Schartner and colleagues the

average concentration of propofol in blood during mild sedation was 1.9 ± 0.52 mcg/mL, whereas here it

was 0.44 ± 0.04 mcg/mL and 0.90 ± 0.06 mcg/mL during mild and moderate sedation, respectively.

Further, Huang et al.[31] investigated temporal dynamics of BOLD signals with the targeted propofol

concentration of 1.3 mcg/mL, whereas Liu et al.[32] analyzed entropy of BOLD signals with a targeted

propofol level of 0.98 mcg/mL. In neither study an effect, which could be interpreted as an increase in

diversity or variability of brain activity has been found which, most likely, is due to lower sensitivity of

fMRI (in comparison to EEG). However, increased signal diversity has been observed during psychedelic

states induced by LSD, ketamine, or psilocybin[13-15]. This provided first evidence that not only decreases,

but also increases of signal diversity with respect to the values observed during resting wakefulness, are in

principle possible. Our finding adds to this body of evidence by showing this effect is not specific to

psychedelic states only.

What is the neurophysiological mechanism behind changes in EEG diversity? While the decrease in

diversity in the drowsy group might be related to the occurrence of slow-waves, it is well established that

the slow-wave activity occurs mainly during deeper stages of anesthesia, which were not reached here[33-35].

Another hypothesis, which might explain both decreases and increases in diversity, is related to changes in

the higher frequency bands. Low anesthetic doses were reported to induce an increase in the beta/gamma

bands amplitude (EEG: [36], modeling: [37]). This effect is considered to represent an initial paradoxical

excitation caused by an anesthetic and might potentially result in a diversity increase. However, at higher

doses coherent alpha band oscillations occur over centro-frontal regions, and this might result in a more

regular and less diverse EEG signal (EEG: [34, 35, 38]; modeling: [39]). Further, during mild sedation

synchronization of fMRI BOLD signals within local brain regions increases, but at the same time

functional connectivity between distant brain regions decreases[31]. Such a greater independence of local

brain regions might also be reflected by greater diversity. Nevertheless, the precise neurophysiological

mechanism of diversity changes remains to be investigated.

In the present study we demonstrate that EEG diversity is closely related to the ability to recognize and

respond to auditory stimuli, which was used as a proxy for consciousness. Is there any other evidence that

signal diversity measures correlates with consciousness *per se*? First, different anesthetics have different

mechanisms of action but all suppress consciousness[40]. Therefore, one might compare changes in brain

activity under different anesthetics with a rationale that genuine correlates of loss and recovery of

consciousness should be common across anesthetics[41, 42]. Diversity measures have been indeed shown to

decrease irrespective of the anesthetic agent used[3, 9]. Second, unresponsive but potentially conscious states

– like ketamine sedation or REM sleep - provide an opportunity to identify correlates of consciousness

independent of responsiveness. In a study of Sarasso et al.[9] during ketamine anesthesia subjects

experienced vivid dreams and hallucinations and, indeed, a very subtle drop in diversity during ketamine

anesthesia was observed (in contrast to propofol and xenon anesthesia). Further, Schartner et al.[7] found

that only NREM sleep is related to decreases in signal diversity, whereas during REM sleep – a state when

dreams occur most frequently[43] – neuronal diversity is similar to wakefulness. Thus, these initial studies

indicate that signal diversity might be indeed a sensitive index of subjective experience, independent of

perception and responsiveness.

In the present analysis diversity of EEG signals was estimated by the Mean Information Gain (MIG),

which is an information theory measure[19]. However, multiple previous studies established the Lemepl-Ziv

algorithm as a robust correlate of the global conscious states[3, 5-10]. Therefore, we also calculated the

Lemepl-Ziv diversity to compare it with MIG (results reported in the supplementary materials). While

both measures behaved in a qualitatively similar way, MIG seems to distinguish responsive and drowsy

groups more robustly, and only MIG increased significantly during mild sedation and recovery. Greater

sensitivity of MIG might be explained by the fact that, for consistency with previous studies, MIG was

applied to the EEG signal[15, 21], whereas LZs was applied to the signal envelopes[7]. Further, Lempel-Ziv was

used in two versions, to estimate diversity of either temporal (LZs) or spatio-temporal (LZc) patterns. Our

results demonstrate that only LZs, but not LZc, was sensitive enough to distinguish between responsive

and drowsy groups. This suggests that temporal (rather than spatial) diversity of brain activity is the key

aspect and should be the focus of future studies.

Nevertheless, it is important to keep in mind that both MIG and Lempel-Ziv are measures of signal

diversity, defined as randomness or entropy of the data, and not *complexity,* understood as a state

intermediate between order and randomness. Therefore, in our study a measure of complexity, namely

Fluctuation Complexity (FC), was also included. FC has been proved capable of robustly differentiating

states of anesthesia and wakefulness[15]. However, FC decreased only during mild and not during moderate

sedation, and it did not distinguish responsive and drowsy subjects (**Fig. 2**). Lack of a reliable complexity

decrease during loss of responsiveness might be related to the fact, that moving from low diversity to

higher diversity causes an increase in complexity (i.e. moving towards the peak of the inverted "U" shape,

see: **Fig. S1**). Therefore, our analysis suggests that subjects' behavioral state during propofol sedation is

better represented by diversity than complexity of the EEG signal.

In summary, by comparing responsive and unresponsive subjects during propofol sedation we provide

further evidence that neuronal diversity is a promising correlate of responsiveness. It remains to be

investigated to what extent it is related to the subjective experience.

References

1. Tononi, G., & Edelman, G. M. (1998). Consciousness and complexity. *Science*, *282*(5395), 1846-
1851.

2. Seth, A. K., Izhikevich, E., Reeke, G. N., & Edelman, G. M. (2006). Theories and measures of
consciousness: an extended framework. *Proceedings of the National Academy of Sciences*, *103*(28),
10799-10804.

3. Casali, A. G., Gosseries, O., Rosanova, M., Boly, M., Sarasso, S., Casali, K. R., ... & Massimini, M.
(2013). A theoretically based index of consciousness independent of sensory processing and
behavior. *Science translational medicine*, *5*(198), 198ra105-198ra105.

4. Massimini, M., Ferrarelli, F., Huber, R., Esser, S. K., Singh, H., & Tononi, G. (2005). Breakdown of
cortical effective connectivity during sleep. *Science*, *309*(5744), 2228-2232.

5. Andrillon, T., Poulsen, A. T., Hansen, L. K., Léger, D., & Kouider, S. (2016). Neural markers of
responsiveness to the environment in human sleep. *Journal of Neuroscience*, *36*(24), 6583-6596.

6. Pigorini, A., Sarasso, S., Proserpio, P., Szymanski, C., Arnulfo, G., Casarotto, S., ... & Palva, J. M.
(2015). Bistability breaks-off deterministic responses to intracortical stimulation during non-REM
sleep. *Neuroimage*, *112*, 105-113.

7. Schartner, M. M., Pigorini, A., Gibbs, S. A., Arnulfo, G., Sarasso, S., Barnett, L., ... & Barrett, A. B.
(2017a). Global and local complexity of intracranial EEG decreases during NREM sleep.
Neuroscience of Consciousness, *3*(1).

8. Ferrarelli, F., Massimini, M., Sarasso, S., Casali, A., Riedner, B. A., Angelini, G., ... & Pearce, R. A.
(2010). Breakdown in cortical effective connectivity during midazolam-induced loss of
consciousness. *Proceedings of the National Academy of Sciences*, *107*(6), 2681-2686.

9. Sarasso, S., Boly, M., Napolitani, M., Gosseries, O., Charland-Verville, V., Casarotto, S., ... & Rex,
S. (2015). Consciousness and complexity during unresponsiveness induced by propofol, xenon, and
ketamine. *Current Biology*, *25*(23), 3099-3105.

10. Schartner, M., Seth, A., Noirhomme, Q., Boly, M., Bruno, M. A., Laureys, S., & Barrett, A. (2015).

Complexity of multi-dimensional spontaneous EEG decreases during propofol induced general

anaesthesia. *PloS one*, *10*(8), e0133532.

11. Rosanova, M., Gosseries, O., Casarotto, S., Boly, M., Casali, A. G., Bruno, M. A., ... & Massimini,

M. (2012). Recovery of cortical effective connectivity and recovery of consciousness in vegetative

patients. *Brain*, *135*(4), 1308-1320.

12. Casarotto, S., Comanducci, A., Rosanova, M., Sarasso, S., Fecchio, M., Napolitani, M., ... &

Gosseries, O. (2016). Stratification of unresponsive patients by an independently validated index of

brain complexity. *Annals of neurology*, *80*(5), 718-729.

13. Schartner, M. M., Carhart-Harris, R. L., Barrett, A. B., Seth, A. K., & Muthukumaraswamy, S. D.

(2017b). Increased spontaneous MEG signal diversity for psychoactive doses of ketamine, LSD and

psilocybin. *Scientific reports*, *7*, 46421.

14. Tagliazucchi, E., Carhart-Harris, R., Leech, R., Nutt, D., & Chialvo, D. R. (2014). Enhanced

repertoire of brain dynamical states during the psychedelic experience. *Human brain*

mapping, *35*(11), 5442-5456.

15. Wang, J., Noh, G. J., Choi, B. M., Ku, S. W., Joo, P., Jung, W. S., ... & Lee, H. (2017). Suppressed

neural complexity during ketamine-and propofol-induced unconsciousness. *Neuroscience letters*,

653, 320-325.

16. Palanca, B. J. A., Mashour, G. A., & Avidan, M. S. (2009). Processed electroencephalogram in

depth of anesthesia monitoring. *Current Opinion in Anesthesiology*, *22*(5), 553-559.

17. Chennu, S., O'Connor, S., Adapa, R., Menon, D. K., & Bekinschtein, T. A. (2016). Brain

connectivity dissociates responsiveness from drug exposure during propofol-induced transitions of

consciousness. *PLoS computational biology*, *12*(1), e1004669.

18. Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial

EEG dynamics including independent component analysis. *Journal of neuroscience*

methods, *134*(1), 9-21.

19. Bates, J. E., & Shepard, H. K. Measuring complexity using information fluctuation. *Physics Letters*
A, *172*(6), 416-425 (1993).

20. Wackerbauer, R., Witt, A., Atmanspacher, H., Kurths, J., & Scheingraber, H. A comparative
classification of complexity measures. *Chaos, Solitons & Fractals*, *4*(1), 133-173 (1994).

21. Bola, M., Orłowski, P., Płomecka, M., Koculak, M., Griskova-Bulanova, I. Subjective experience
measured by the Amsterdam Resting State Questionnaire is not related to the complexity of EEG
signal. (under review).

22. Wagenmakers, E. J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., Love, J., ... & Matzke, D. (2018).
Bayesian inference for psychology. Part I: Theoretical advantages and practical
ramifications. *Psychonomic bulletin & review*, *25*(1), 35-57.

23. Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian journal of*
statistics, 65-70.

24. Lee, M. D., and Wagenmakers, E.-J. (2013). Bayesian Modeling for Cognitive Science: A Practical
Course. London, UK: Cambridge University Press.

25. Bola, M., Barrett, A. B., Pigorini, A., Nobili, L., Seth, A. K., & Marchewka, A. (2018). Loss of
consciousness is related to hyper-correlated gamma-band activity in anesthetized macaques and
sleeping humans. *Neuroimage*, *167*, 130-142.

26. Tagliazucchi, E., Chialvo, D. R., Siniatchkin, M., Amico, E., Brichant, J. F., Bonhomme, V., ... &
Laureys, S. (2016). Large-scale signatures of unconsciousness are consistent with a departure from
critical dynamics. *Journal of the Royal Society Interface*, *13*(114), 20151027.

27. Krzemiński, D., Kamiński, M., Marchewka, A., & Bola, M. Breakdown of long-range temporal
correlations in brain oscillations during general anesthesia. *NeuroImage*, *159*, 146-158 (2017).

28. Alonso, L. M., Proekt, A., Schwartz, T. H., Pryor, K. O., Cecchi, G. A., & Magnasco, M. O. (2014).
Dynamical criticality during induction of anesthesia in human ECoG recordings. *Frontiers in neural*
circuits, *8*, 20.

29. Tajima, S., Yanagawa, T., Fujii, N., & Toyoizumi, T. Untangling brain-wide dynamics in consciousness by cross-embedding. *PLoS computational biology*, *11*(11), e1004537 (2015).

30. Solovey, G., Alonso, L. M., Yanagawa, T., Fujii, N., Magnasco, M. O., Cecchi, G. A., & Proekt, A. Loss of consciousness is associated with stabilization of cortical activity. *Journal of Neuroscience*, *35*(30), 10866-10877 (2015).

31. Huang, Z., Liu, X., Mashour, G. A., & Hudetz, A. G. (2018). Timescales of intrinsic BOLD signal dynamics and functional connectivity in pharmacologic and neuropathologic states of unconsciousness. *Journal of Neuroscience*, 2545-17.

32. Liu, X., Lauer, K. K., Ward, B. D., Roberts, C. J., Liu, S., Gollapudy, S., ... & Wang, L. (2018). Regional entropy of functional imaging signals varies differently in sensory and cognitive systems during propofol-modulated loss and return of behavioral responsiveness. *Brain imaging and behavior*, 1-12.

33. Breshears, J. D., Roland, J. L., Sharma, M., Gaona, C. M., Freudenburg, Z. V., Tempelhoff, R., ... & Leuthardt, E. C. (2010). Stable and dynamic cortical electrophysiology of induction and emergence with propofol anesthesia. *Proceedings of the National Academy of Sciences*, 201011949.

34. Murphy, M., Bruno, M. A., Riedner, B. A., Boveroux, P., Noirhomme, Q., Landsness, E. C., ... & Tononi, G. (2011). Propofol anesthesia and sleep: a high-density EEG study. *Sleep*, *34*(3), 283-291.

35. Purdon, P. L., Pierce, E. T., Mukamel, E. A., Prerau, M. J., Walsh, J. L., Wong, K. F. K., ... & Ching, S. (2013). Electroencephalogram signatures of loss and recovery of consciousness from propofol. *Proceedings of the National Academy of Sciences*, 201221180.

36. Gugino, L. D., Chabot, R. J., Prichep, L. S., John, E. R., Formanek, V., & Aglio, L. S. (2001). Quantitative EEG changes associated with loss and return of consciousness in healthy adult volunteers anaesthetized with propofol or sevoflurane. *British journal of anaesthesia*, *87*(3), 421-428.

37. McCarthy, M. M., Brown, E. N., & Kopell, N. (2008). Potential network mechanisms mediating
electroencephalographic beta rhythm changes during propofol-induced paradoxical
excitation. *Journal of Neuroscience*, *28*(50), 13488-13504.

38. Supp, G. G., Siegel, M., Hipp, J. F., & Engel, A. K. (2011). Cortical hypersynchrony predicts
breakdown of sensory processing during loss of consciousness. *Current biology*, *21*(23), 1988-1993.

39. Ching, S., Cimenser, A., Purdon, P. L., Brown, E. N., & Kopell, N. J. (2010). Thalamocortical
model for a propofol-induced α-rhythm associated with loss of consciousness. *Proceedings of the*
National Academy of Sciences, *107*(52), 22665-22670.

40. Mashour, G. A. (2014). Top-down mechanisms of anesthetic-induced unconsciousness. *Frontiers in*
systems neuroscience, *8*, 115.

41. Långsjö, J. W., Alkire, M. T., Kaskinoro, K., Hayama, H., Maksimow, A., Kaisti, K. K., ... &
Scheinin, H. (2012). Returning from oblivion: imaging the neural core of consciousness. *Journal of*
Neuroscience, *32*(14), 4935-4943.

42. Scheinin, A., Kallionpää, R. E., Li, D., Kallioinen, M., Kaisti, K., Långsjö, J., ... & Revonsuo, A.
(2018). Differentiating drug-related and state-related effects of dexmedetomidine and propofol on
the electroencephalogram. *Anesthesiology: The Journal of the American Society of*
Anesthesiologists.

43. Siclari, F., LaRocque, J. J., Postle, B. R., & Tononi, G. (2013). Assessing sleep consciousness within
subjects using a serial awakening paradigm. *Frontiers in psychology*, *4*, 542.

Figure 1. Hit-rates during an auditory detection task (left panels) and propofol concentration estimated

from blood samples taken during the experiment (right panels). Upper panels present data of individual

subjects from both groups, while bar-plots in lower panels present mean ± SEM of the same data.

Statistical analysis of hit-rates was not conducted in this study, and in case of propofol concentration only

comparisons between groups (but within states) were conducted. * - $p < 0.05$.

Figure 2. Diversity and complexity of the EEG signal across consciousness states. A) Data of individual subjects from both groups. Data of each were normalized by subtracting the baseline (wake) value to better depict profiles of changes. B) The same data presented as mean ± SEM without normalization. Results of between-states and between-groups comparisons depicted as: * - $p < 0.05$; ** - p

511 < 0.01. C) Individual hit-rates from the auditory detection task plotted against changes in

512 diversity/complexity measures (with respect to baseline, as in A). Data of two subjects who became

513 unresponsive already during mild sedation state are plotted as red diamonds (compare to Fig. 1).

Supplementary Materials

Supplementary Results

535  In the present study we focused on two information theory measures - MIG, which is a measure of

536 diversity, and FC, which is a measure of complexity. However, for consistency with multiple previous

537 studies which used the Lempel-Ziv algorithm we also calculated two version of Lempel-Ziv diversity on

538 our data. The implementation used here is the same as in[7].

539  First, we investigated temporal diversity, denoted here as LZs. Statistical analysis of LZs indicates no

540 main effect of *group* ($F(1) = 1.9$, $p = 0.18$; $BF_{10} = 0.75$) and no effect of *state* ($F(3) = 1.48$, $p = 0.22$; BF_{10}

541 $= 1.05$), but a significant *group x state* interaction was found ($F(3) = 3.27$, $p = 0.028$; $BF_{10} = 3.0$).

542 Subsequent post-hoc tests indicate no differences between states (all $p > 0.05$). However, comparisons

543 between groups indicate that the drowsy group exhibits lower LZs during moderate sedation ($t(18) = 3.32$,

544 $p = 0.004$; $BF_{10} = 10.4$) but not during other states (all $p > 0.10$; all $0.3 < BF_{10} < 1$).

545  Second, we investigated spatio-temporal diversity, denoted here as LZc. There was no main effect of

546 *group* ($F(1) = 0.35$, $p = 0.56$; $BF_{10} = 0.56$), a significant effect of *state* ($F(3) = 4.76$, $p = 0.005$; $BF_{10} =$

547 7.24), and no *group x state* interaction ($F(3) = 0.74$, $p = 0.53$; $BF_{10} = 1.0$). However, subsequent post-hoc

548 tests indicate a significant difference only between moderate sedation and recovery ($t(19) = 3.15$, $p =$

549 0.031; $BF_{10} = 8.82$), with all other comparisons not reaching significance threshold (all $p > 0.10$; all $0.3 <$

550 $BF_{10} < 1$).

Figure S1. A relation between complexity (FC) and diversity (MIG). Each dot represents a value

calculated from each state and subject (altogether 20 x 4 = 80 data-points). The data show an inverted "U"

shape, as complexity peaks for signals of intermediate diversity (between order and disorder).

Figure S2. Two versions of Lemepl-Ziv, measuring either temporal (LZs) or spatio-temporal (LZc) diversity of EEG data. A) Data of individual subjects from both groups. To depict profiles of changes data of each subject were normalized by subtracting the baseline (wake) value. B) The same data presented as mean ± SEM without normalization. Results of between-states and between-groups comparisons depicted as: * - p < 0.05; ** - p < 0.01.