Genetic networks underlying natural variation in basal and induced activity levels

in *Drosophila melanogaster*

Louis P. Watanabe, Cameron Gordon, and Nicole C. Riddle

The University of Alabama at Birmingham

Department of Biology

Birmingham, Alabama 35294

Running Title: Natural variation in activity levels

Key Words: Exercise, GWAS, DGRP2, Drosophila, REQS

Corresponding Author:

Nicole C. Riddle

1720 2nd Ave S

Birmingham, AL

35294

Phone: 205-975-4049

Email: riddlenc@uab.edu

ABSTRACT

Exercise is recommended by health professionals across the globe as part of a healthy

lifestyle to prevent and/or treat the consequences of obesity. While overall, the health

benefits of exercise and an active lifestyle are well understood, very little is known about

how genetics impacts an individual's inclination for and response to exercise. To

address this knowledge gap, we investigated the genetic architecture underlying natural

variation in activity levels in the model system *Drosophila melanogaster*. Activity levels

were assayed in the Drosophila Genetics Reference Panel 2 fly strains at baseline and

in response to a gentle exercise treatment using the Rotational Exercise Quantification

System. We found significant, sex-dependent variation in both activity measures and

identified over 100 genes that contribute to basal and induced exercise activity levels.

This gene set was enriched for genes with functions in the central nervous system and

in neuromuscular junctions and included several candidate genes with known activity

phenotypes such as flightlessness or uncoordinated movement. Interestingly, there

were also several chromatin proteins among the candidate genes, two of which were

validated here and shown to impact activity levels. Thus, the study described here

reveals the complex genetic architecture controlling basal and exercise-induced activity

levels in *D. melanogaster* and provides a resource for exercise biologists.

INTRODUCTION

Obesity is a disease associated with significantly higher all-cause mortality relative to

normal weight (CARMIENKE *et al.* 2013; FLEGAL *et al.* 2013; MASTERS *et al.* 2013). This

increase in mortality can be attributed largely to the elevated incidence of

cardiovascular disease (EBBERT *et al.* 2014; MOLICA *et al.* 2015; SISNOWSKI *et al.* 2015),

cancer (BERGER 2014; GARG *et al.* 2014; NAKAMURA *et al.* 2014; PARK *et al.* 2014), and

diabetes (NOMURA *et al.* 2010; RIOBO SERVAN 2013; POLSKY AND ELLIS 2015) observed in

obese individuals. The increasing prevalence of obesity is a serious international public

health issue that has warranted action from legislators worldwide (HARTEMINK *et al.*

2006). The upswing in rates of the disease throughout the United States caused the

national medical expenditure dedicated to treating obesity-related illnesses in adults to

increase by 29% between 2001 and 2015 (BIENER *et al.* 2018). Therefore, strategies to

counter the broadening obesity epidemic are needed to ensure the medical and

financial wellbeing of society at large.

Exercise is among the most common treatments for obesity, which also include surgical

procedures, medications, and other lifestyle modifications (BARETIC 2013; WYATT 2013;

KUSHNER 2014; MARTIN *et al.* 2015). Given the relatively risk-free nature of exercise as

a method of weight loss compared with many other treatment options, it is widely

considered to be an essential component of treatment regimes for obesity (MCQUEEN

2009; LASKOWSKI 2012; FONSECA-JUNIOR *et al.* 2013). In addition to treating obesity,

exercise imparts a number of health benefits including improved muscle function

(ANDERSEN *et al.* 2015; COEN *et al.* 2015; KIM *et al.* 2015) and cartilage integrity

(TONEVITSKY *et al.* 2013; BREIT *et al.* 2015; BLAZEK *et al.* 2016), increased insulin

sensitivity (MITROU *et al.* 2013; BROCKLEBANK *et al.* 2015), and prevention of many

chronic conditions (BOOTH *et al.* 2012). These benefits led exercise to be recognized as

an important facet of a healthy lifestyle, with government agencies such as the U.S.

Department of Health and Human Services issuing specific exercise recommendations

for adults, youths, and children (e.g. adults should do at least 150 minutes of moderate-

intensity, or 75 minutes of vigorous-intensity aerobic activity per week) (DHHS 2008).

Thus, exercise is an important component of many people's lives, whether to treat

obesity or to improve overall health.

Despite the growing relevance of exercise, there is a lack of clarity concerning a number

of factors that influence its physiological effects (KAROLY *et al.* 2012), most notably

genetic background. Although exercise has gained considerable popularity as both a

lifestyle choice and a treatment for obesity, there are large differences in how

individuals respond to exercise, and it is not universally effective (KEITH *et al.* 2006;

MCALLISTER *et al.* 2009). In fact, exercise provides no metabolic improvements to

certain individuals (BOUCHARD *et al.* 1999; BOUCHARD *et al.* 2011; STEPHENS AND SPARKS

2015), revealing an extreme disparity in exercise response that can likely be accounted

for, at least in part, by genetic variation. Moreover, existing data suggest a relationship

between exercise-induced improvements to muscle metabolism and exercise

performance in humans (LAREW *et al.* 2003). Thus, while some genes have been

identified as contributors to physical activity traits of an individual (STUBBE *et al.* 2006;

DE GEUS *et al.* 2014), the genetic architecture controlling exercise responses has yet to

be characterized.

As an emerging model organism for exercise studies, *Drosophila melanogaster*

possesses several characteristics advantageous for elucidating the relevant genetic

architecture (PIAZZA *et al.* 2009; TINKERHESS *et al.* 2012; SUJKOWSKI *et al.* 2015; MENDEZ

et al. 2016; WATANABE AND RIDDLE 2017). Traditional obstacles for exercise studies,

including difficulties in controlling for essential variables such as age, sex, fitness, and

diet, can be addressed using *D. melanogaster*. Furthermore, Drosophila is an

established model system with a well-characterized genome and ample tools for genetic

studies. These tools include the Drosophila Genetics Reference Panel 2 (DGRP2), a

fully sequenced population of 200 genetically diverse isogenic lines for quantitative

genetic studies (MACKAY *et al.* 2012; HUANG *et al.* 2014), in addition to large collections

of mutants and RNAi knockdown lines. Furthermore, a high degree of genetic

(SCHNEIDER 2000) and functional (HEWITT AND WHITWORTH 2017) conservation exists

between Drosophila and humans, particularly in areas such as energy-related pathways

(EDISON *et al.* 2016) and disease genes (PANDEY AND NICHOLS 2011; UGUR *et al.* 2016),

which allows findings from Drosophila to be tested to mammalian model systems.

Together, these features make Drosophila an excellent choice for studies of exercise

genetics.

Several innovative studies demonstrate that exercise treatments of Drosophila produce

significant physiological and behavioral responses, including increased lifespan and

improved climbing ability (PIAZZA *et al.* 2009; TINKERHESS *et al.* 2012; SUJKOWSKI *et al.*

2015; MENDEZ *et al.* 2016; WATANABE AND RIDDLE 2017). For example, treatments with

the Power Tower, which prompts exercise by exploiting the negative geotaxis of

Drosophila by repeatedly dropping their enclosures, causing the animals to fall to the

base and attempt another climb, improves mobility in aging animals (PIAZZA *et al.* 2009).

The TreadWheel exploits negative geotaxis through slow rotation of fly enclosures to

stimulate a response; responses to prolonged exercise on the TreadWheel include, for

example, changes in triglyceride and glycogen levels in the animals (MENDEZ *et al.*

2016). The Rotating Exercise Quantification System (REQS) is an offshoot of the

TreadWheel, which is able to record the activity levels of flies as they exercise,

facilitating the comparison of different exercise regimes and allowing for the

normalization of exercise levels (WATANABE AND RIDDLE 2017). The REQS validation

study also demonstrated that there is significant variability among different Drosophila

genotypes in how they respond to the rotational exercise stimulation (WATANABE AND

RIDDLE 2017), suggesting that genetic factors contribute to the difference in exercise

levels observed.

In this study, we investigate the genetic factors contributing to the level of exercise

induced through rotational stimulation. Using the REQS, we measured basal activity

levels (without rotation) as well as induced exercise levels (with rotation) in 161

genetically diverse strains from the DGRP2. Next, we used a genome-wide association

study (GWAS) to identify the genetic variants responsible for the approximately 10-fold

variation in activity levels observed within the DGRP2 lines. We identified over 100

annotated genes that contribute to basal and induced activity levels. The loci that

control activity levels are different for the untreated and exercise-treated conditions and

often also differ between males and females. Additional characterization of candidate

genes validate the results of our GWAS and confirm that genes with functions in the

central nervous system as well as some chromatin proteins impact Drosophila activity

levels. Together, our findings provide key insights into the number and types of genetic

factors that control basal and exercise-induced activity levels, provide an array of

candidate genes for follow-up studies, and identify chromatin modifiers as a new class

of proteins linked to exercise.

MATERIALS AND METHODS

Drosophila Lines and Husbandry

The DGRP2 fly lines used in this study were obtained from the Bloomington Drosophila

Stock Center or from our collaborator Dr. Laura Reed (University of Alabama). Fly lines

for the follow-up analysis of candidate genes (Supplemental Table S1) were obtained

from the Bloomington Drosophila Stock Center, with the exception of the *wde* mutant

stock, which was obtained from Dr. Wodarz (Georg-August-Universität Göttingen,

Germany). Drosophila were grown on media consisting of a cornmeal-molasses base

with the addition of Tegosept, propionic acid, agar, yeast, and Drosophila culture netting

(MENDEZ *et al.* 2016). All flies used for the assays were reared in an incubator at 25°C

and ~70% humidity with a 12-hour light-dark cycle for a minimum of three generations

prior to the start of the experiments. To minimize density effects, animals were grown in

vials established by mating seven male to ten female flies for each line. The resulting

progeny were collected as virgins, aged 3-7 days, separated by sex, and then used for

the exercise experiments.

Exercise Quantification Assay

Basal and induced exercise activity levels were determined for a total of 161 strains

(155 basal and 151 induced) from the DGRP2 using the REQS (MACKAY *et al.* 2012;

HUANG *et al.* 2014; WATANABE AND RIDDLE 2017; WATANABE AND RIDDLE 2018)

(Supplemental Table S2). For each DGRP2 line, 100 male and 100 female virgin flies

aged 3-7 days were used. The animals were anesthetized with CO_2, divided into

groups of 10 (n=10 groups per sex per line), and loaded into the vials of the REQS at

9AM [for additional details see (WATANABE AND RIDDLE 2018)]. The REQS was moved

into the incubator at 10AM, and the vials were rotated to a vertical orientation similar to

how flies are reared in a laboratory setting. The animals were allowed to recover from

anesthetization for one hour. The basal activity level of the animals was measured from

11AM to 12PM, keeping the REQS in a static position without rotation. At 12PM, the

REQS' rotational feature was turned on at 4 rotations per minute (rpm) to induce

exercise in the animals, and activity levels were monitored during this exercise until

2PM. Measurements were taken at five-minute intervals during both the basal and

induced activity phases.

Statistics

Basal and induced activity levels were calculated as the average activity level per five-

minute interval of each vial/genotype/sex combination. A GLM (general linear model

with gamma log-link) to investigate the impact of the factors vial, genotype, sex, and

treatment on activity levels was performed using SAS9.4 software (INC 2013). As the

initial analysis showed no effect of "vial", "vial" was removed from the final model.

Descriptive statistics were generated in SPSS25 (IBM 2017) and R (R DEVELOPMENT

CORE TEAM 2018). Custom Perl scripts were used for the SNP classification analysis in

addition to R (R DEVELOPMENT CORE TEAM 2018).

Genome Wide Association Study (GWAS)

Basal and induced activity levels were separately analyzed by calculating the average

activity level per five-minute interval of each genotype/sex combination. These

phenotypic values (Supplemental Table S3) were used for two separate GWASs using

the DGRP2 webtool (http://dgrp2.gnets.ncsu.edu/) developed by Dr. Trudy Mackay at

NC State University (MACKAY *et al.* 2012; HUANG *et al.* 2014). Genetic variants that met

a significance threshold of $p<10^{-5}$ in any of the analyses (mixed model, simple

regression model, female data, male data, combined data, and sex difference analysis)

were considered candidate loci (Supplemental Table S3). Candidate genes for follow-

up and validation were selected based on significance level, mutant availability from

Drosophila stock centers, and reports on FlyBase (GRAMATES *et al.* 2017) consistent

with phenotypes that might be linked to exercise/activity.

Gene Ontology (GO)

GO analysis was performed on the genes associated with the genetic variants that met

the significance threshold of $p<10^{-5}$ separately for both the basal and induced GWAS

results. Genetic variants lacking association with a specific gene were removed from

the list, and duplicate genes were removed as well. FlyBase gene IDs were retrieved

for the gene sets using The Database for Annotation, Visualization and Integrated

Discovery (DAVID) ID conversion function (HUANG DA *et al.* 2009b; HUANG DA *et al.*

2009a). GO analysis was carried out using PANTHER (Protein Analysis Through

Evolutionary Relationships) (MI *et al.* 2013; MI *et al.* 2017). The "biological process"

and "cellular component" enrichment terms were used.

Functional analysis of candidate genes

Candidate genes were selected for follow-up experiments based on their highly

significant *p*-values in the GWASs, as well as based on the annotation available on

FlyBase. Mutant lines for *eIF5B*, *Suz(2)*, *wde*, *Cirl*, *nwk*, *cpo*, *SmydA-9*, and *Prosap*

(see Supplemental Table S4 for genotypes) were characterized for basal and induced

activity levels using the same protocol as described above in the section entitled

"Exercise Quantification Assay." For UAS/Gal4 lines, the UAS transgene was brought

together with the nervous system GAL4 driver P{w[+mC]=GAL4-elav.L}2 through a

single generation cross.

Data availability

All data necessary for confirming the conclusions of this article are represented fully

within the article and its tables, figures, and supplemental files, with the exception of the

genotype data for the DGRP2 population and the GWAS model. This information can

be found at http://dgrp2.gnets.ncsu.edu/.

RESULTS

The DGRP2 shows extensive variation in basal and exercise-induced activity levels.

To investigate the genetic basis of variation in activity levels, both basal and exercise induced, we focused on the DGRP2 collection of Drosophila strains. The DGRP2 is a collection of 200 inbred lines of Drosophila derived from wild-caught females, representing genetic variation that is present in a natural population. To measure activity levels, we used the REQS, as it allowed us to record basal activity of the animals without rotation and induced activity levels during rotation. The rotation induces exercise (higher activity levels) through the animals' negative geotaxis response. Using 161 strains from the DGRP2, we measured basal activity of the animals as well as the activity during rotationally-induced exercise in a single experiment (Figure 1). The output from the REQS is the average activity level per 10 flies per five-minute interval, which was estimated based on a one-hour recording for the basal activity and a two-hour recording for the exercise activity.

Figure 2 shows the results from this set of experiments, with data from females (A+B) and males (C+D) shown separately. All activity measures described are given as crossings/five minutes/fly. We found a wide range of average activity levels for both basal and exercise-induced activity. The highest level of basal activity in males was found in line 57, which exhibited 144.5 +/- 9.4 activity units (mean +/- SEM), while for females line 595 had the highest basal activity with 58.55 activity units (+/- 3.9). The highest performing line was the same for exercise-induced activity in both sexes, line

808 with an average of 155.25 activity units (+/- 3.56) in males and an average of

133.94 activity units (+/- 3.60) in females. Interestingly, for males the lowest basal and

induced activity levels were measured both in line 383 with 0.275 activity units (+/- 0.1)

for basal activity and 1.63 activity units (+/- 0.79) for exercise-induced activity. The low

performer in the females was different between basal and induced activity: Line 390

females had the lowest basal activity with 0.53 activity units (+/- 0.14), while line 32 with

an average of 1.22 activity units (+/- 0.26) had the lowest exercise-induced activity level.

Looking across all factors, the variation in mean activity levels illustrated in Figure 2

range from a low of 0.275 +/- 0.1 activity units (line 383, basal males) to a high of

155.25 +/- 3.56 activity units (line 808, induced males). Thus, our highest activity

measurement showed an approximately 500-fold increase from the lowest

measurement, demonstrating that there is extensive variation in animal activity based

on genotype, sex, and treatment within the DGRP2 population.

Several general trends can be discerned from the bar graphs in Figure 2: 1) Across the

161 lines, female flies tend to have a lower basal activity level than males (compare A to

C); 2) rotational stimulation tends to increase activity levels (compare A to B and C to D),

and 3) while basal activity levels tend to be lower in females, exercise activity levels

tend to be more similar between the sexes (compare A and C to B and D). However,

there are exceptions to these general trends. For example when comparing male to

female activity, of the 155 basal lines measured, 27 lines displayed significantly higher

activity levels in males, but there is one genotype (line 83) which exhibited higher

activity levels in females (40.35 female activity, 4.87 male activity, p=0.044). Similarly,

for the induced activity levels, 12 lines showed significantly greater activity in males than

in females, while there was one line that showed the opposite trend, higher activity

levels in females than in males (line 796, 122.01 female activity, 75.6 male activity,

p<0.001). When observing the effects of exercise, as expected 41 genotypes

demonstrated significantly increased activity with only three lines showing decreased

activity. Thus, most genotypes showed increased activity when rotated, and males

exhibited higher activity levels than females.

Sex, genotype, and exercise treatment impact activity.

Next, we investigated the factors influencing the variation in activity levels we observed

in Figure 2 utilizing a general linear model (GLM) analysis. Specifically, the GLM

examined the impact of treatment (with or without rotation), sex, genotype, as well as

the interactions between these factors. Our results indicate that activity levels were

significantly impacted by treatment, illustrating that rotation indeed is able to increase

activity levels above baseline in this genetically diverse population of fly strains

(p<0.001, Table 1). In addition, sex and genotype significantly impacted activity levels,

as suggested by the descriptive data illustrated in Figures 2 and S1 (p<0.001).

Interaction effects between treatment, sex, and genotype also impacted the activity

levels measured by the REQS, indicating that to be able to predict activity phenotypes,

sex, treatment, and genotype must all be considered together (Table 1). Our

descriptive statistics combined with the GLM analysis thus indicate that for the activity

phenotype there is tremendous variation between the DGRP2 lines, some of which is

due to genetics (genotype effect). This finding suggested that individual genes

underlying the variation in activity levels might be identified by a GWAS.

GWAS identifies over 300 genetic variants impacting activity levels.

To identify the genetic factors impacting basal and exercise-induced activity levels in the

DGRP2 population, we carried out a GWAS. The analysis was run separately for the

basal and exercise-induced activity levels (155 and 151 lines respectively) using the

DGRP2 GWAS Webtool (http://dgrp2.gnets.ncsu.edu/). The webtool uses two different

models for the analysis, a simple regression model, as well as a more complex mixed

model, and the analysis is carried out for male data only, female data only, combined

data from both sexes, and for the difference between sexes. Together, this set of

GWASs identified over 400 genetic variants [single nucleotide polymorphisms (SNPs),

multiple nucleotide polymorphisms (MNPs), and deletions (DEL)] that impact basal and

exercise-induced activity levels ($p<0.005$; Figures 3 and S2, Tables 2 and S2),

illustrating the importance of genetic factors for exercise phenotypes.

The genomic variants identified by the GWAS are distributed throughout the genome,

and with the exception of the small 4[th] chromosome, all chromosome arms contain

genetic variants contributing to the basal and induced exercise activity phenotypes

(Table S2). Comparing the chromosomal distribution of the genetic variants identified

as significant in the basal activity or induced exercise activity analysis to that of the

overall distribution of variants using chi-square analysis, we found no significant

deviations from the expectation, indicating that the variants are not clustered in any

particular way in the genome. Overall, the results of this GWAS demonstrate that a

large number of loci contribute to the two activity phenotypes measured and that the

genetic architecture underlying the phenotypes is complex.

Basal activity variant distributions differ significantly from genome-wide set.

Next, we investigated what types of genetic variants were identified as significant in the

GWAS. In order to compare the classifications of the variants from the basal and

induced analyses to the entire genome, the variants from the induced and basal activity

were first characterized based on their genomic context (Introns, Exons, Upstream,

Downstream, UTR, and Unknown). We then compared the classification distributions

from the basal and induced analyses to a genome-wide set using a chi-square test. We

found that only the basal activity variant distributions were significantly different from the

genome-wide set ($p=5.413*10^{-7}$) with a greater number of exons, upstream, and

unknown elements (Figure 4). The classification for the variants contributing to induced

exercise activity were not significantly different from the genome-wide classification of

variants, possibly due to the smaller number of variants detected in this analysis and a

concomitant reduced power to detect differences. Thus, the GWASs carried out here

identify a diverse set of genetic variants as contributing to the activity phenotypes under

study, and the overrepresentation of exon variants in the basal activity analysis

suggests that these variants might indeed present genes important for animal activity

and exercise.

Genetic variants impacting activity levels differ between males and females.

The genetic variants identified as significantly associated with the two activity

phenotypes differed between the analyses, some linked to activity in the males, some

linked to activity in females, and some variants linked to the difference in activity levels

between males and females. While the separate analyses of both male and female

data lead to the identification of genetic variants, more genetic variants were identified

in males than in females, and the combined analysis recovered more variants identified

in males than in females. For example, the mixed model analysis of basal activity levels

identified 53 genetic variants in females, 93 genetic variants in the males, and in the

combined analysis, 46 variants are identified as significant, 22 of which overlap with the

variants identified using the male data, while only one variant also occurs in the female

analysis ($p < 5 * 10^{-5}$). These results illustrate the importance of collecting data in both

sexes, as the genetic factors contributing to both basal and induced exercise activity

levels differ between males and females. The results also suggest that the difference

between males and females is under genetic control.

GWAS discovers positive and negative effect variants impacting activity levels.

The genetic variants identified as significantly associated with basal and/or induced

exercise activity include loci with positive and negative effects on the phenotype. In the

analysis of the induced exercise activity data, only genetic variants with negative

impacts on activity levels were identified in either sex. Examining the data for basal

activity, in males, the majority of variants negatively impacted basal activity, and only

1.5% of variants (three out of 195; mixed model) increased activity levels. These rare

genetic variants leading to increased activity levels were associated with the genes *bdg*

and *slo.* In females, the results are similar, with the majority of variants (63; mixed

model) impacting basal activity levels negatively and positive effect variants being rare

(three variants associated with *CG32521* and *CG8420*). Overall we found that the vast

majority of our discovered variants resulted in a negative effect on activity.

Terms related to the central nervous system and muscle function are over-

represented in the GO analysis.

Next, we focused on the genetic variants identified in the GWAS as significant that were

associated with genes and asked what types of genes contributed to the basal and

induced activity phenotypes (146 genes for basal activity; 47 genes for induced exercise

activity). To do so, we explored the gene ontology (GO) terms associated with the gene

sets linked to basal and induced activity levels. In order to determine which gene

classes were over- or under-represented, GO analysis was carried out using the

PANTHER (Protein Analysis Through Evolutionary Relationships) tools (MI *et al.* 2013;

MI *et al.* 2017). Specifically, enrichment analysis was used to identify biological

processes, cellular components, or molecular functions over-represented within the

GWAS gene set relative to the genome as a whole.

For the GO term enrichment analysis for the genes contributing to basal activity levels,

30 GO terms were identified in the "biological process" category and 17 in the "cellular

component" category. Among the "biological process" and "cellular component" GO

terms, the largest fold enrichments were seen for two terms related to neurons, "axonal

growth cone" (26.24-fold enrichment; $p=3.3*10^{-4}$) and "neuron recognition" (7.51-fold

enrichment, p=5.79*10^{-5}) (Figure 5). Other significantly enriched GO terms include

"neuromuscular junction", "synaptic transmission", and "behavior" (Figure 5). The GO

term enrichment analysis of the gene set associated with exercise-induced activity

identified a 27-fold enrichment for genes involved in the Alzheimer disease-presenilin

pathway (p=2.01*10^{-4}). Thus, the GO term enrichment analysis identifies a variety of

terms associated with neuronal function and behavior as characterizing the gene set

involved in controlling basal and exercise-induced activity levels.

Candidate gene analyses support the GWAS results and suggest that the

chromatin proteins WDE and SU(Z)2 impact animal activity.

In order to assess the success of the GWAS analysis, we surveyed the information

available on FlyBase (GRAMATES *et al.* 2017) to determine if altered activity phenotypes

have been described associated with the candidate genes identified here. Mutants in

25 of the candidate genes have been described as either "flightless", or "flight defective"

on FlyBase (GRAMATES *et al.* 2017), five were described as "uncoordinated," and one

(*Cirl*) was described as "hyperactive" and "sleep defective" (VAN DER VOET *et al.* 2015).

Based on their molecular functions and previously described phenotypes, 10 genes

were selected for follow-up and mutant/RNAi lines were obtained: *eIF5B* (eukaryotic

translation initiation factor 5B; LASKO 2000; JOHNSTONE AND LASKO 2004), *Su(z)2* (a

member of the Polycomb Repressive Complex 1 [PRC1]; WU AND HOWE 1995; LO *et al.*

2009; NGUYEN *et al.* 2017), *wde* (a cofactor of the H3K9 histone methyltransferase

EGG; KOCH *et al.* 2009; YU *et al.* 2015), *Cirl* (a Calcium-independent receptor for α-

latrotoxin with hyperactivity phenotype; ANDERSEN *et al.* 2015; SCHOLZ *et al.* 2015;

SCHOLZ *et al.* 2017), *nwk* (an SH3 adaptor protein regulating synaptic growth; COYLE *et*

al. 2004b; O'CONNOR-GILES *et al.* 2008; RODAL *et al.* 2008a), *SmydA*-9 (an arthropod-

specific SET and MYND domain; THOMPSON AND TRAVERS 2008), *cpo* (controls various

neurological functions; BELLEN *et al.* 1992a; BELLEN *et al.* 1992b) (GLASSCOCK AND

TANOUYE 2005), *Prosap* (also an SH3 domain containing proteins involved in neuronal

development; DIAGANA *et al.* 2002; HARRIS *et al.* 2016; WU *et al.* 2017), *jus* (a gene with

links to seizures; ZHANG *et al.* 2002; MARLEY AND BAINES 2011; DEAN *et al.* 2018), and

Jarid2 (a Jumonji C domain-containing lysine demethylase; SASAI *et al.* 2007; HERZ *et al.*

2012). Two of the mutant lines (*jus* and *Jarid2*) obtained were very sickly, and we were

unable to conduct the exercise quantification assay we normally carry out. Thus, our

analysis focuses on *eIF5B*, *Su(z)2*, *wde*, *Cirl*, *nwk*, *cpo*, *SmydA-9*, and *Prosap.*

Because the mutant and RNAi lines we obtained for this analysis were in unknown

genetic backgrounds, we compared their activity measurements to our results from the

DGRP2 strains, with the understanding that this strain collection provided a good

overview of activity levels – basal and induced - within Drosophila in general (Figure 6

and Table S1). As with the initial analysis of the DGRP2, activity levels were measured

for each sex independently. For the eight candidate genes assayed in both sexes, the

basal activity levels for nine sex/genotype combinations are in the top or bottom 20% of

the phenotypic distribution. For the induced activity levels, seven of the sex/genotype

combinations are amongst the 20% most extreme phenotypes measured in the DGRP2

(Figure 6 and Table S2). One of these genes stands out, as in both sexes, mutants in

eIF5B showed some of the lowest basal and induced activity levels we recorded. These

animals generally were rather sickly, suggesting that the DGRP2 must harbor a weak

allele of this locus. For other mutants, the results were more mixed, with some affecting

one sex more strongly than the other, or affecting only basal and not induced exercise

or *vice versa*. For example, *Su(z)2* knockdown in the nervous system led to low basal

and induced activity levels in females, but did not consistently impact males. In contrast,

Cirl mutant animals showed relatively low induced activity in males, but relatively high

levels in females, and relatively normal basal activity levels. *wde* mutant animals show

the opposite: high basal activity in males, low basal activity levels in females, and

relatively normal activity levels after exercise induction. These results are not surprising,

as a significant number of the SNPs identified in the GWAS only impacted one sex, and

most candidate genes were specific to either basal or induced activity. Overall, we see

that approximately half of the sex/genotype combinations in our candidate mutants or

knockdown study exhibit phenotypes at the extreme ends of the DGRP2 phenotypic

distribution. This finding further validates the GWAS analysis presented here and

suggests that these candidate genes are worthy of further study.

DISCUSSION

As exercise is widely recommended as part of a healthy lifestyle and as a treatment for

obesity, we explored the contribution of genetic variation to exercise in Drosophila using

the DGRP2 strain collection. We found extensive variation in activity levels in this

population, both basal and exercise-induced, which was dependent on genotype and

sex. The GWASs identified over 300 genetic variants and more than 150 genes that

contributed to the basal and exercise-induced activity phenotypes. Some of these

genes had previously been associated with exercise performance or activity. This

group of genes includes *couch potato* [*cpo*], an RNA-binding, nuclear protein expressed

in the central nervous system that is essential for normal flight behavior (BELLEN *et al.*

1992b; GLASSCOCK AND TANOUYE 2005; SCHMIDT *et al.* 2008). *nervous wreck* [*nwk*]

encodes a very different kind of protein than *cpo*, but it also has been shown previously

to impact animal activity: mutants in this FCH and SH3 domain-containing adaptor

protein show paralysis due to problems at the synapse of neuromuscular junctions

(COYLE *et al.* 2004a; RODAL *et al.* 2008b). Another gene in this group is *bedraggled*

[*bdg*], which encodes a putative neurotransmitter transporter, and null mutants of which

are described as flightless and uncoordinated (RAWLS *et al.* 2007). The importance of

the factors identified in our study is further underscored by the GO terms enrichment

analysis, which revealed terms associated with the functions of the central nervous

system and its interaction with the musculature. Together, these findings illustrate that

the GWAS described here was successful in detecting variants associated with factors

involved in the control of animal activity levels and behavior.

In addition to the genes such as *cpo*, *nwk*, and *bdg*, that would have been expected to

impact basal activity or geotaxis-induced exercise activity, others candidate genes

represent pathways with no clear link to animal activity or behavior. For example,

variants in several chromatin proteins were identified as impacting animal activity levels,

including SmydA-9, a SET domain protein, Su(z)2, a Polycomb group protein related to

PSC, Jarid2, a Jumonji domain protein that interacts with PRC2, and Wde, an essential

co-factor of the H3K9 methyltransferase Egg. In addition, several sperm proteins were

identified as linked to activity (e. g. S-Lap8n, Sfp36F, Sfp51E) as were several members

of the immunoglobulin superfamily (e. g. Side-II, CG31814, CG13506), but how they

might contribute to an activity phenotype is unclear. As several of these genes are

among a total of 36 genes identified here that were identified also by Schnorrer and

colleagues as essential for normal muscle development (SCHNORRER *et al.* 2010), it is

likely they represent novel pathways linked to activity. Because it includes both

unexpected and expected gene classes, the gene set identified here as contributing to

both basal and exercise-induced activity levels provides a rich resource for researchers

interested in using Drosophila as a tool for the study of exercise.

Our study also revealed that there were significant differences in the genes contributing

to activity levels in males and in females, and that several genes could be identified that

were responsible for the difference between the sexes. This finding was surprising, as

we anticipated that the basic metabolic and sensory pathways involved in the control of

animal activity would be conserved between males and females. Mackay and

colleagues in 2012 discovered significant differences in the gene networks controlling

starvation response and chill coma recovery time, but not startle response, and for all

three traits, a large portion of the genetic variants that are identified in one sex show no

significant association with the trait in the other sex (OBER *et al.* 2012). Garlapow and

colleagues also found strong sex-specific differences in the factors controlling food

intake in the DGRP2 population (GARLAPOW *et al.* 2015), and Morozova and colleagues

detected sex differences in the networks controlling alcohol sensitivity (FOCHLER *et al.*

2017). These findings suggest that for many "basic" traits such as animal activity, sex

specific differences in the underlying genetic networks occur frequently, highlighting the

importance of studying both sexes.

When the list of candidate genes identified here is compared with those uncovered in

other studies of activity traits, substantial overlap can be seen. For example, Jordan and

colleagues investigated the genetic basis of the startle response and negative geotaxis

in the DGRP2 population using a GWAS (JORDAN *et al.* 2012). They identified

approximately 200 genes associated with each of these activity phenotypes, a number

of genes similar to that identified in our study. As our exercise system relies on

negative geotaxis, not surprisingly, 24 identified by Jordan and colleagues were also

identified in the basal activity analysis presented here, and an additional 17 genes from

the exercise induced activity analysis also overlapped with the gene set identified by

Jordan and colleagues. Five genes were identified as candidates in all three analyses

(*CG15630*, *CG33144*, *ed*, *nmo*, and *SKIP*) (JORDAN *et al.* 2012). The overlap seen

between the candidate genes identified in the two studies indicates that despite the

different activity traits measured shared pathways exist.

Interestingly, a second activity study utilizing the DGRP2 identified a completely

independent set of candidate genes, showing no overlap with the genes identified here.

The study by Rhode and colleagues used video-tracking to monitor the activity of male

Drosophila from the DGRP2 in a shallow petri dish by measuring distance traveled in a

5-minute interval (ROHDE *et al.* 2018). This "2D" activity study focused on specific

groups of genes linked to their phenotype, specifically genes involved in

transmembrane transport. While this study used a very different algorithm to identify

candidate genes, even when their phenotypic measures are analyzed with the standard

GWAS tool our study used, we find no overlap in the candidate gene sets identified.

This lack of overlap in candidate genes identified by two activity studies illustrates the

complexity in the pathways that impact basic animal behaviors such as activity. Given

current understanding, genes from basic energy metabolism pathways to genes

controlling the development of muscles and sensory organs to genes impacting the

processing of sensory information are all involved in controlling activity levels. Thus, it

is not surprising that studies using different activity types will identify distinct sets of

candidate genes; rather, these findings highlight that additional innovative studies are

needed to come to a comprehensive understanding of the genes involved in animal

activity, both basal and in response to stimulation.

The functional analysis of candidate genes presented here demonstrates that the REQS

can be used successfully to identify genes involved in controlling basal and exercise

induced activity levels in the DGRP2. Many of the candidate genes identified show

relatively small impacts in the DGRP2, likely due to the presence of weak, not null

alleles in this wild-derived population. Despite the fact that most of the candidate genes

identified are sex-specific and have weak effects, the functional analysis of candidate

genes showed that several clearly impact activity levels in the basal and exercise

assays. Likely, with additional analyses including several alleles per gene, RNAi

knockdown in tissues other than the central nervous system, and overexpression lines,

further candidate genes will be confirmed as impacting activity levels in Drosophila.

Although our analysis identified genes from many different categories of cell and

molecular functions, one category of note with regards to exercise response is

chromatin modifiers. Our assays on *wde* and *Suz(2)* mutants suggest that this group of

genes might indeed play a role in controlling exercise activity levels, and possibly

exercise response. Changes in an organism's activity levels have profound impacts,

both acute and long-term. Gene expression changes and alterations to the epigenome

have been identified as immediate consequences of exercise. For example, there are

several studies documenting DNA methylation changes as well as changes in histone

acetylation following exercise. These epigenetic changes are one possible mechanism

that might mediate the long-term consequences of exercise, many of which can persist

even in the absence of further exercise. Thus, it is of interest that two proteins linked to

different histone methylation marks, histone 3 lysine 9 methylation (Wde) and histone 3

lysine 27 methylation (Su(z)2), were identified as contributing to the variation in activity

levels between individuals in our study, and future studies in the role of these marks

with regard to exercise are needed.

In summary, our study has identified promising candidate genes contributing to the

variation in activity levels seen between genetically distinct individuals. Many of the

genes identified have not been linked to exercise previously, and they thus present

novel avenues for exploration to exercise biologists. Given the clear homology

relationships between many of the Drosophila genes identified and mammalian

genomes, the gene set presented here also provides new research directions for

exercise biology studies in rodents. Using the results from model systems such as

Drosophila as a guide, translational scientists will be able to accelerate biomarker

development to eventually allow medical professionals to prescribe individualized

exercise treatments for obesity related diseases and to guide athletes of all kinds.

ACKNOWLEDGEMENTS

The work described was supported partially by Award Number P30DK056336 from the

National Institute Of Diabetes And Digestive And Kidney Diseases through a pilot grant

from the UAB Nutrition and Obesity Research Center (NORC) to NCR. The content is

solely the responsibility of the authors and does not necessarily represent the official

views of the National Institute Of Diabetes And Digestive And Kidney Diseases or the

National Institutes of Health.

We would like to thank our undergraduate students Michael Azar, Makayla Nixon, Jamie

Hill, and Anthony Foster for their contributions to various aspects of these experiments.

The *wde* mutant stock *w;wde^{TD63}/Cyo[twi:GFP]* was obtained from Dr. A. Wodarz

(Georg-August-Universität Göttingen, Germany), and Dr. L. Reed (University of

Alabama) graciously shared many of the DGRP2 strains with us. Additional stocks

obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) were

used in this study.

 FIGURE LEGENDS

Figure 1. Experimental setup. Diagram illustrating the experimental setup, how

animals were processed, and data were collected.

Figure 2. Activity levels within the DGRP2 population vary significantly. This set

of bar graphs illustrate the variability in activity levels among the different lines of the

DGRP2 population. In each graph, the lines are ordered based on their activity levels

from smallest to largest (X-axis), with mean activity levels (average number of beam

crosses recorded by the activity monitor in a 5-minute interval) on the Y-axis. The error

bars are SEM.

A. Bar graph showing line means for basal activity – female.

B. Bar graph showing line means for induced activity – female.

C. Bar graph showing line means for basal activity – male.

D. Bar graph showing line means for induced activity – male.

Figure 3. GWAS identifies SNPs associated with basal and induced activity levels.

Chromosomal location of SNPs (X-axis) are plotted against the negative log of the p-

value testing for the likelihood of the SNP being associated with the measured

phenotype (mixed model; Y-axis). The blue line in each plot marks the $p=10^{-5}$

significance level, while the red line marks $p=10^{-7}$.

A. Manhattan plot for basal activity – female.

B. Manhattan plot for induced activity – female.

C. Manhattan plot for basal activity – male.

D. Manhattan plot of induced activity – male.

Figure 4. The genetic variants identified by the GWAS for basal activity are

biased towards exons and upstream genetic elements. The genetic variants

identified as significantly associated with either the basal (left) or exercise-induced

(middle) activity were classified based on their sequence context and compared to the

genome-wide set of variants forming the basis of this GWAS (right). The set of variants

identified in the basal activity GWAS is significantly different from the genome-wide set

($p=5.42*10^{-7}$; Chi-square test).

Figure 5. GO analysis highlights the importance of axonal growth cone and

neuron recognition for activity levels. Genetic variants identified in the GWAS for

basal analysis were subjected to GO enrichment analysis. Only high-level GO terms

meeting the significance threshold are shown (for complete results and individual p-

values, see Supplemental Table S5).

Figure 6. Mutations in candidate genes have varied impacts on activity levels.

Violin plot showing the average basal and induced activity levels of mutants in

candidate genes identified by GWAS overlaid over the distribution of activity levels

measured in the DGRP2 strains. The red square marks the median activity level

measured in the DGRP2 strain collection. For *Su(z)2*, two mutants were investigated: A

is a RNAi knockdown line, while B is a null allele.

Figure S1. Histograms illustrating the variability in activity levels among the lines of the DGRP2 population. All lines binned based on their mean activity levels (average number of beam crosses recorded by the activity monitor in a 5-minute interval; bin size: 10). The bins are indicated on the X-axis (average number of beam crosses recorded by the activity monitor in a 5 minute interval), and the number of lines in each bin is shown on the Y-axis.

A. Histogram for basal activity in females.

B. Histogram for induced activity in females.

C. Histogram for basal activity in males.

D. Histogram for induced activity in males.

Figure S2. Genetic variants associated with basal and induced activity levels in a GWAS using combined data from both sexes. Chromosomal location of genetic variants (X-axis) are plotted against the negative log of the p-value testing for the likelihood of the variant being associated with the measured phenotype (Y-axis). The blue line in each plot marks the $p=10^{-5}$ significance level, while the red line marks $p=10^{-7}$.

A. Manhattan plot for basal activity.

B. Manhattan plot for induced activity.

Figure S3. Small areas of linkage disequilibrium (LD) exist for the genetic variants associated with basal and exercise-induced activity.

Heat maps illustrating the extent of LD among the genetic variants identified as

significantly contributing to the basal (**A**) and exercise-induced (**B**) activity levels. The

level of LD is indicated by the color of each square with red indicating the highest level

of LD and blue indicating the absence of LD. The genetic variants are plotted based on

their chromosomal location (X-and Y-axis), with the diagonal showing areas of with the

strongest LD due to close physical proximity.

TABLES

Table 1. Treatment, sex, and genotype show strong effects on activity levels.
General linear model analysis of variance reveals the impact of treatment (+/- rotational stimulation), sex, and genotype on the activity phenotype measured.

Table 2. Summary of GWAS results. The table lists the number of genetic variants detected as significant by the different analysis types used in this study.

Supplemental Table S1. Mutant strains and candidate gene analysis. This table contains the genotypes and Bloomington stock numbers of the mutants analyzed, as well as the raw activity data collected from these strains.

Supplemental Table S2. List of DGRP lines included in this study.
Table is separated by treatment and sex for each line. Y = Yes, N = No.

Supplemental Table S3. Phenotypic measures used in the GWAS analysis.
Summary data as submitted to the DGRP2 GWAS webtool as well as raw data are provided.

Supplemental Table S4. Candidate genetic variants identified as significant by the GWAS. This table includes results from the basal activity GWAS and the exercise-induced activity GWAS.

689 **Supplemental Table S5. Detailed GO term enrichment analysis results for genetic**

690 **variants associated with basal activity.** This table includes the complete GO term

691 enrichment analysis results for the "Cellular component" and "Biological Process"

692 categories for genetic variants associated with basal activity.

LITERATURE CITED

Andersen, G., M. C. Orngreen, N. Preisler, T. D. Jeppesen, T. O. Krag *et al.*, 2015 Protein-

carbohydrate supplements improve muscle protein balance in muscular dystrophy

patients after endurance exercise: a placebo-controlled crossover study. Am J Physiol

Regul Integr Comp Physiol 308**:** R123-130.

Baretic, M., 2013 Obesity drug therapy. Minerva Endocrinol 38**:** 245-254.

Bellen, H. J., S. Kooyer, D. D'Evelyn and J. Pearlman, 1992a The Drosophila couch potato

protein is expressed in nuclei of peripheral neuronal precursors and shows homology to

RNA-binding proteins. Genes & Development 6**:** 2125-2136.

Bellen, H. J., H. Vaessin, E. Bier, A. Kolodkin, D. D'Evelyn *et al.*, 1992b The Drosophila *couch*

potato gene: an essential gene required for normal adult behavior. Genetics 131**:** 365-

375.

Berger, N. A., 2014 Obesity and cancer pathogenesis. Ann N Y Acad Sci 1311**:** 57-76.

Biener, A., J. Cawley and C. Meyerhoefer, 2018 The Impact of Obesity on Medical Care Costs

and Labor Market Outcomes in the US. Clinical Chemistry 64**:** 108-117.

Blazek, A. D., J. Nam, R. Gupta, M. Pradhan, P. Perera *et al.*, 2016 Exercise-Driven Metabolic

Pathways in Healthy Cartilage. Osteoarthritis Cartilage 24**:** 1210-1222.

Booth, F. W., C. K. Roberts and M. J. Laye, 2012 Lack of exercise is a major cause of chronic

diseases. Compr Physiol 2**:** 1143-1211.

Bouchard, C., P. An, T. Rice, J. S. Skinner, J. H. Wilmore *et al.*, 1999 Familial aggregation of

VO(2max) response to exercise training: results from the HERITAGE Family Study. J

Appl Physiol (1985) 87**:** 1003-1008.

Bouchard, C., T. Rankinen and J. A. Timmons, 2011 Genomics and Genetics in the Biology of

Adaptation to Exercise. Compr Physiol 1**:** 1603-1648.

Breit, M., M. Netzer, K. M. Weinberger and C. Baumgartner, 2015 Modeling and Classification

of Kinetic Patterns of Dynamic Metabolic Biomarkers in Physical Activity. PLOS

Computational Biology 11**:** e1004454.

Brocklebank, L. A., C. L. Falconer, A. S. Page, R. Perry and A. R. Cooper, 2015 Accelerometer-

measured sedentary time and cardiometabolic biomarkers: A systematic review. Prev

Med 76**:** 92-102.

Carmienke, S., M. H. Freitag, T. Pischon, P. Schlattmann, T. Fankhaenel *et al.*, 2013 General

and abdominal obesity parameters and their combination in relation to mortality: a

systematic review and meta-regression analysis. Eur J Clin Nutr 67**:** 573-585.

Coen, P. M., E. V. Menshikova, G. Distefano, D. Zheng, C. J. Tanner *et al.*, 2015 Exercise and

Weight Loss Improve Muscle Mitochondrial Respiration, Lipid Partitioning, and Insulin

Sensitivity After Gastric Bypass Surgery. Diabetes 64**:** 3737-3750.

Coyle, I. P., Y. H. Koh, W. C. Lee, J. Slind, T. Fergestad *et al.*, 2004a Nervous wreck, an SH3

adaptor protein that interacts with Wsp, regulates synaptic growth in Drosophila. Neuron

41**:** 521-534.

Coyle, I. P., Y. H. Koh, W. C. M. Lee, J. Slind, T. Fergestad *et al.*, 2004b Nervous wreck, an

SH3 adaptor protein that interacts with Wsp, regulates synaptic growth in Drosophila.

Neuron 41**:** 521-534.

de Geus, E. J., M. Bartels, J. Kaprio, J. T. Lightfoot and M. Thomis, 2014 Genetics of regular

exercise and sedentary behaviors. Twin Res Hum Genet 17**:** 262-271.

Dean, D., H. Weinstein, S. Amin, B. Karno, E. McAvoy *et al.*, 2018 Extending *julius seizure*, a

bang-sensitive gene, as a model for studying epileptogenesis: Cold shock, and a new

insertional mutation. Fly 12**:** 55-61.

DHHS, 2008 2008 physical activity guidelines for Americans.

Diagana, T. T., U. Thomas, S. N. Prokopenko, B. Xiao, P. F. Worley *et al.*, 2002 Mutation of

Drosophila *homer* disrupts control of locomotor activity and behavioral plasticity. Journal

of Neuroscience 22**:** 428-436.

Ebbert, J. O., M. Y. Elrashidi and M. D. Jensen, 2014 Managing overweight and obesity in

adults to reduce cardiovascular disease risk. Curr Atheroscler Rep 16**:** 445.

Edison, A. S., R. D. Hall, C. Junot, P. D. Karp, I. J. Kurland *et al.*, 2016 The Time Is Right to

Focus on Model Organism Metabolomes. Metabolites 6.

Flegal, K. M., B. K. Kit, H. Orpana and B. I. Graubard, 2013 Association of all-cause mortality

with overweight and obesity using standard body mass index categories: a systematic

review and meta-analysis. Jama 309**:** 71-82.

Fochler, S., T. V. Morozova, M. R. Davis, A. W. Gearhart, W. Huang *et al.*, 2017 Genetics of

alcohol consumption in *Drosophila melanogaster.* Genes Brain Behav 16**:** 675-685.

Fonseca-Junior, S. J., C. G. Sa, P. A. Rodrigues, A. J. Oliveira and J. Fernandes-Filho, 2013

Physical exercise and morbid obesity: a systematic review. Arq Bras Cir Dig 26 Suppl 1**:**

67-73.

Garg, S. K., H. Maurer, K. Reed and R. Selagamsetty, 2014 Diabetes and cancer: two diseases

with obesity as a common risk factor. Diabetes Obes Metab 16**:** 97-110.

Garlapow, M. E., W. Huang, M. T. Yarboro, K. R. Peterson and T. F. Mackay, 2015 Quantitative

Genetics of Food Intake in *Drosophila melanogaster*. PLoS One 10**:** e0138129.

Glasscock, E., and M. A. Tanouye, 2005 Drosophila *couch potato* mutants exhibit complex

neurological abnormalities including epilepsy phenotypes. Genetics 169**:** 2137-2149.

Gramates, L. S., S. J. Marygold, G. d. Santos, J.-M. Urbano, G. Antonazzo *et al.*, 2017 FlyBase

at 25: looking to the future. Nucleic Acids Research 45**:** D663-D671.

Harris, K. P., Y. Akbergenova, R. W. Cho, M. S. Baas-Thomas and J. T. Littleton, 2016 Shank

Modulates Postsynaptic Wnt Signaling to Regulate Synaptic Development. Journal of

Neuroscience 36**:** 5820-5832.

Hartemink, N., H. C. Boshuizen, N. J. Nagelkerke, M. A. Jacobs and H. C. van Houwelingen,

2006 Combining risk estimates from observational studies with different exposure

cutpoints: a meta-analysis on body mass index and diabetes type 2. Am J Epidemiol

163**:** 1042-1052.

Herz, H. M., M. Mohan, A. S. Garrett, C. Miller, D. Casto *et al.*, 2012 Polycomb repressive

complex 2-dependent and -independent functions of jarid2 in transcriptional regulation in

Drosophila. Molecular and Cellular Biology 32**:** 1683-1693.

Hewitt, V. L., and A. J. Whitworth, 2017 Mechanisms of Parkinson's Disease: Lessons from

Drosophila. Curr Top Dev Biol 121**:** 173-200.

Huang da, W., B. T. Sherman and R. A. Lempicki, 2009a Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res 37**:**

1-13.

Huang da, W., B. T. Sherman and R. A. Lempicki, 2009b Systematic and integrative analysis of

large gene lists using DAVID bioinformatics resources. Nat Protoc 4**:** 44-57.

Huang, W., A. Massouras, Y. Inoue, J. Peiffer, M. Ramia *et al.*, 2014 Natural variation in

genome architecture among 205 Drosophila melanogaster Genetic Reference Panel

lines. Genome Res 24**:** 1193-1208.

IBM, 2017 IBM SPSS Statistics 25.

Inc, S. I., 2013 SAS/ACCESS® 9.4 Interface to ADABAS, pp. SAS Institute Inc., Cary, NC.

Johnstone, O., and P. Lasko, 2004 Interaction with eIF5B is essential for Vasa function during

development. Development 131**:** 4167-4178.

Jordan, K. W., K. L. Craver, M. M. Magwire, C. E. Cubilla, T. F. Mackay *et al.*, 2012 Genome-

wide association for sensitivity to chronic oxidative stress in *Drosophila melanogaster*.

PLoS One 7**:** e38722.

Karoly, H. C., C. J. Stevens, R. E. Magnan, N. Harlaar, K. E. Hutchison *et al.*, 2012 Genetic

Influences on Physiological and Subjective Responses to an Aerobic Exercise Session

among Sedentary Adults. J Cancer Epidemiol 2012**:** 540563.

Keith, S. W., D. T. Redden, P. T. Katzmarzyk, M. M. Boggiano, E. C. Hanlon *et al.*, 2006

Putative contributors to the secular increase in obesity: exploring the roads less traveled.

Int J Obes (Lond) 30**:** 1585-1594.

Kim, H. J., B. So, M. Choi, D. Kang and W. Song, 2015 Resistance exercise training increases

the expression of irisin concomitant with improvement of muscle function in aging mice

and humans. Exp Gerontol 70**:** 11-17.

Koch, C. M., M. Honemann-Capito, D. Egger-Adam and A. Wodarz, 2009 Windei, the

Drosophila homolog of mAM/MCAF1, is an essential cofactor of the H3K9 methyl

transferase dSETDB1/Eggless in germ line development. PLoS Genetics 5**:** e1000644.

Kushner, R. F., 2014 Weight loss strategies for treatment of obesity. Prog Cardiovasc Dis 56**:**

465-472.

Larew, K., G. R. Hunter, D. E. Larson-Meyer, B. R. Newcomer, J. P. McCarthy *et al.*, 2003

Muscle metabolic function, exercise performance, and weight gain. Med Sci Sports

Exerc 35**:** 230-236.

Lasko, P., 2000 The Drosophila melanogaster genome: translation factors and RNA binding

proteins. Journal of Cell Biology 150**:** F51-F56.

Laskowski, E. R., 2012 The role of exercise in the treatment of obesity. Pm r 4**:** 840-844; quiz

844.

Lo, S. M., N. K. Ahuja and N. J. Francis, 2009 Polycomb group protein Suppressor 2 of zeste is

a functional homolog of Posterior Sex Combs. Molecular and Cellular Biology 29**:** 515-

525.

Mackay, T. F., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles *et al.*, 2012 The *Drosophila*

melanogaster Genetic Reference Panel. Nature 482**:** 173-178.

Marley, R., and R. A. Baines, 2011 Increased persistent Na+ current contributes to seizure in

the *slamdance* bang-sensitive Drosophila mutant. Journal of Neurophysiology 106**:** 18-

29.

Martin, K. A., M. V. Mani and A. Mani, 2015 New targets to treat obesity and the metabolic

syndrome. Eur J Pharmacol 763**:** 64-74.

Masters, R. K., D. A. Powers and B. G. Link, 2013 Obesity and US mortality risk over the adult

life course. Am J Epidemiol 177**:** 431-442.

McAllister, E. J., N. V. Dhurandhar, S. W. Keith, L. J. Aronne, J. Barger *et al.*, 2009 Ten Putative

Contributors to the Obesity Epidemic. Crit Rev Food Sci Nutr 49**:** 868-913.

McQueen, M. A., 2009 Exercise Aspects of Obesity Treatment. Ochsner J 9**:** 140-143.

Mendez, S., L. Watanabe, R. Hill, M. Owens, J. Moraczewski *et al.*, 2016 The TreadWheel: A

Novel Apparatus to Measure Genetic Variation in Response to Gently Induced Exercise

for Drosophila. PLOS ONE 11**:** e0164706.

Mi, H., X. Huang, A. Muruganujan, H. Tang, C. Mills *et al.*, 2017 PANTHER version 11:

expanded annotation data from Gene Ontology and Reactome pathways, and data

analysis tool enhancements. Nucleic Acids Research 45**:** D183-D189.

Mi, H., A. Muruganujan, J. T. Casagrande and P. D. Thomas, 2013 Large-scale gene function

analysis with the PANTHER classification system. Nature Protocols 8**:** 1551.

Mitrou, P., S. A. Raptis and G. Dimitriadis, 2013 Insulin action in morbid obesity: a focus on

muscle and adipose tissue. Hormones (Athens) 12**:** 201-213.

Molica, F., S. Morel, B. R. Kwak, F. Rohner-Jeanrenaud and S. Steffens, 2015 Adipokines at

the crossroad between obesity and cardiovascular disease. Thromb Haemost 113**:** 553-

566.

Nakamura, K., J. J. Fuster and K. Walsh, 2014 Adipokines: A link between obesity and

cardiovascular disease. J Cardiol 63**:** 250-259.

Nguyen, S. C., S. Yu, E. Oberlick and C. T. Wu, 2017 An Unexpected Regulatory Cascade

Governs a Core Function of the Drosophila PRC1 Chromatin Protein Su(z)2. Genetics

205**:** 551-558.

Nomura, D. K., J. Z. Long, S. Niessen, H. S. Hoover, S. W. Ng *et al.*, 2010 Monoacylglycerol

lipase regulates a fatty acid network that promotes cancer pathogenesis. Cell 140**:** 49-61.

O'Connor-Giles, K. M., L. L. Ho and B. Ganetzky, 2008 Nervous wreck interacts with thickveins

and the endocytic machinery to attenuate retrograde BMP signaling during synaptic

growth. Neuron 58**:** 507-518.

Ober, U., J. F. Ayroles, E. A. Stone, S. Richards, D. Zhu *et al.*, 2012 Using whole-genome

sequence data to predict quantitative trait phenotypes in *Drosophila melanogaster*. PLoS

Genet 8**:** e1002685.

Pandey, U. B., and C. D. Nichols, 2011 Human disease models in *Drosophila melanogaster* and

the role of the fly in therapeutic drug discovery. Pharmacol Rev 63**:** 411-436.

Park, J., T. S. Morley, M. Kim, D. J. Clegg and P. E. Scherer, 2014 Obesity and cancer--

mechanisms underlying tumour progression and recurrence. Nat Rev Endocrinol 10**:**

455-465.

Piazza, N., B. Gosangi, S. Devilla, R. Arking and R. Wessells, 2009 Exercise-training in young

Drosophila melanogaster reduces age-related decline in mobility and cardiac

performance. PLoS One 4**:** e5886.

Polsky, S., and S. L. Ellis, 2015 Obesity, insulin resistance, and type 1 diabetes mellitus. Curr

Opin Endocrinol Diabetes Obes 22**:** 277-282.

R Development Core Team, 2018

R: A language and environment for statstical computing, pp. R Foundation for Statistical

Computing, Vienna Austria.

Rawls, A. S., S. A. Schultz, R. D. Mitra and T. Wolff, 2007 Bedraggled, a putative transporter,

869   influences the tissue polarity complex during the R3/R4 fate decision in the Drosophila

870   eye. Genetics 177**:** 313-328.

871   Riobo Servan, P., 2013 Obesity and diabetes. Nutr Hosp 28 Suppl 5**:** 138-143.

872   Rodal, A. A., R. N. Motola-Barnes and J. T. Littleton, 2008a Nervous wreck and Cdc42

873   cooperate to regulate endocytic actin assembly during synaptic growth. Journal of

874   Neuroscience 28**:** 8316-8325.

875   Rodal, A. A., R. N. Motola-Barnes and J. T. Littleton, 2008b Nervous wreck and Cdc42

876   cooperate to regulate endocytic actin assembly during synaptic growth. J Neurosci 28**:**

877   8316-8325.

878   Rohde, P. D., S. Østergaard, T. N. Kristensen, P. Sørensen, V. Loeschcke *et al.*, 2018

879   Functional Validation of Candidate Genes Detected by Genomic Feature Models. G3:

880   Genes|Genomes|Genetics 8**:** 1659-1668.

881   Sasai, N., Y. Kato, G. Kimura, T. Takeuchi and M. Yamaguchi, 2007 The Drosophila *jumonji*

882   gene encodes a JmjC-containing nuclear protein that is required for metamorphosis.

883   FEBS Journal 274**:** 6139-6151.

884   Schmidt, P. S., C.-T. Zhu, J. Das, M. Batavia, L. Yang *et al.*, 2008 An amino acid polymorphism

885   in the *couch potato* gene forms the basis for climatic adaptation in Drosophila

886   melanogaster. Proceedings of the National Academy of Sciences of the United States of

887   America 105**:** 16207-16211.

888   Schneider, D., 2000 Using Drosophila as a model insect. Nat Rev Genet 1**:** 218-226.

889   Schnorrer, F., C. Schönbauer, C. C. H. Langer, G. Dietzl, M. Novatchkova *et al.*, 2010

890   Systematic genetic analysis of muscle morphogenesis and function in Drosophila.

891   Nature 464**:** 287.

892   Scholz, N., J. Gehring, C. Guan, D. Ljaschenko, R. Fischer *et al.*, 2015 The Adhesion GPCR

893   Latrophilin/CIRL Shapes Mechanosensation. Cell reports 11**:** 866-874.

Scholz, N., C. Guan, M. Nieberler, A. Grotemeyer, I. Maiellaro *et al.*, 2017 Mechano-dependent

signaling by Latrophilin/CIRL quenches cAMP in proprioceptive neurons. eLife 6**:**

e28360.

Sisnowski, J., E. Handsley and J. M. Street, 2015 Regulatory approaches to obesity prevention:

A systematic overview of current laws addressing diet-related risk factors in the

European Union and the United States. Health Policy 119**:** 720-731.

Stephens, N. A., and L. M. Sparks, 2015 Resistance to the beneficial effects of exercise in type

2 diabetes: are some individuals programmed to fail? J Clin Endocrinol Metab 100**:** 43-

52.

Stubbe, J. H., D. I. Boomsma, J. M. Vink, B. K. Cornes, N. G. Martin *et al.*, 2006 Genetic

Influences on Exercise Participation in 37.051 Twin Pairs from Seven Countries. PLOS

ONE 1**:** e22.

Sujkowski, A., B. Bazzell, K. Carpenter, R. Arking and R. J. Wessells, 2015 Endurance exercise

and selective breeding for longevity extend Drosophila healthspan by overlapping

mechanisms. Aging (Albany NY) 7**:** 535-552.

Thompson, E. C., and A. A. Travers, 2008 A Drosophila Smyd4 homologue is a muscle-specific

transcriptional modulator involved in development. PLoS ONE 3**:** e3008.

Tinkerhess, M. J., L. Healy, M. Morgan, A. Sujkowski, E. Matthys *et al.*, 2012 The Drosophila

PGC-1α Homolog *spargel* Modulates the Physiological Effects of Endurance Exercise.

PLOS ONE 7**:** e31633.

Tonevitsky, A. G., D. V. Maltseva, A. Abbasi, T. R. Samatov, D. A. Sakharov *et al.*, 2013

Dynamically regulated miRNA-mRNA networks revealed by exercise. BMC Physiology

13**:** 9.

Ugur, B., K. Chen and H. J. Bellen, 2016 Drosophila tools and assays for the study of human

diseases. Dis Model Mech 9**:** 235-244.

van der Voet, M., B. Harich, B. Franke and A. Schenck, 2015 ADHD-associated dopamine

920   transporter, latrophilin and neurofibromin share a dopamine-related locomotor signature

921   in Drosophila. Molecular Psychiatry 21**:** 565.

922   Watanabe, L. P., and N. C. Riddle, 2017 Characterization of the Rotating Exercise

923   Quantification System (REQS), a novel Drosophila exercise quantification apparatus.

924   PLOS ONE 12**:** e0185090.

925   Watanabe, L. P., and N. C. Riddle, 2018 Measuring Exercise Levels in *Drosophila melanogaster*

926   Using the Rotating Exercise Quantification System (REQS). JoVE**:** e57751.

927   Wu, C., and M. Howe, 1995 A genetic analysis of the suppressor 2 of zeste complex of

928   *Drosophila melanogaster*. Genetics 140**:** 139-181.

929   Wu, S., G. Gan, Z. Zhang, J. Sun, Q. Wang *et al.*, 2017 A Presynaptic Function of Shank

930   Protein in Drosophila. Journal of Neuroscience 37**:** 11592-11604.

931   Wyatt, H. R., 2013 Update on Treatment Strategies for Obesity. J Clin Endocrinol Metab 98**:**

932   1299-1306.

933   Yu, Y., J. Gu, Y. Jin, Y. Luo, J. B. Preall *et al.*, 2015 Panoramix enforces piRNA-dependent

934   cotranscriptional silencing. Science 350**:** 339-342.

935   Zhang, H., J. Tan, E. Reynolds, D. Kuebler, S. Faulhaber *et al.*, 2002 The Drosophila

936   *slamdance* gene: a mutation in an aminopeptidase can cause seizure, paralysis and

937   neuronal failure. Genetics 162**:** 1283-1299.

12hr light/dark
70% humidity
10 ♀ and 7 ♂

Collect virgins;
age 3-7 days.

10 vials of 10
animals per sex
per genotype

Recovery
period: 1hr

Assay basal
activity for 1h.

Assay induced
activity for 2h.