OSCA: a tool for omic-data-based complex trait analysis

Futao Zhang[1], Wenhan Chen[1], Zhihong Zhu[1], Qian Zhang[1], Ian J. Deary[2], Naomi R. Wray[1,3], Peter M. Visscher[1,3], Allan F. McRae[1], Jian Yang[1,3,4,*]

1. Institute for Molecular Bioscience, The University of Queensland, Brisbane, Queensland 4072, Australia
2. Centre for Cognitive Ageing and Cognitive Epidemiology, Department of Psychology, University of Edinburgh, 7 George Square, Edinburgh, EH8 9JZ, UK
3. Queensland Brain Institute, The University of Queensland, Brisbane, Queensland 4072, Australia
4. Institute for Advanced Research, Wenzhou Medical University, Wenzhou, Zhejiang 325027, China

* Correspondence: Jian Yang (jian.yang@uq.edu.au)

Abstract

The rapid increase of omic data in the past decades has greatly facilitated the investigation of associations between omic profiles such as DNA methylation (DNAm) and complex traits in large cohorts. Here, we proposed a mixed-linear-model-based method (called MOMENT) that tests for association between a DNAm probe and trait with all other distal probes fitted in multiple random-effect components to account for the effects of unobserved confounders as well as the correlations between distal probes induced by the confounders. We demonstrated by simulations that MOMENT showed a lower false positive rate and more robustness than existing methods. MOMENT has been implemented in a versatile software package (called OSCA) together with a number of other implementations for omic-data-based analysis including the estimation of variance in a trait captured by all measures of multiple omic profiles, omic-data-based quantitative trait locus (xQTL) analysis, and meta-analysis of xQTL data.

Introduction

The rapid proliferation of genetic and omic data in large cohort-based samples in the past decade have greatly advanced our understanding of the genetic architecture of omic profiles and the molecular mechanisms underpinning the genetic variation of human complex traits [1-3]. These advances include the identification of a large number of genetic variants associated with gene expression [4, 5], DNA methylation [6, 7], histone modification [8, 9], and protein abundance [10, 11]; the discovery of omic measures associated with complex traits [12, 13]; the improved accuracy in predicting a trait using omic data [14, 15]; and the prioritization of gene targets for complex traits by integrating genetic and omic data in large samples [3, 13, 16-18]. These advances have also led to the development of software tools, focusing on a range of different aspects of omic data analysis. Therefore, a software tool that implements reliable and robust statistical methods for comprehensive analysis of omic data with high-performance computing efficiency is required.

A well-recognised challenge in omic-data-based analysis is to control for false positive rate (FPR) in the presence of confounding factors, as failing to model the confounders may lead to spurious associations [19-21] and/or a loss of statistical power [22]. While some confounders (e.g., age and sex) are known and available in most data so that their effects can be accounted for by fitting them as covariates in linear models, others are either uncharacterised or difficult to measure. For example, in DNA methylation (DNAm) data from whole blood, cell type compositions (CTCs) are evident confounders in a methylome-wide association study (MWAS; also known as an epigenome-wide association study or EWAS) [21, 23, 24] although CTCs may be useful for the prediction of some phenotypes. CTCs tend to be correlated with the DNAm at CpG sites that are differentially methylated in different cell types (namely differentially methylated sites) and have been shown to be associated with age and multiple traits and diseases [19, 21, 25, 26]. MWAS analysis without accounting for CTCs could give rise to inflated test-statistics unless neither CTCs nor DNAm sites are associated with the trait in question. Although it is possible to measure CTCs directly or predict them by reference-based prediction methods [27, 28], reference-free methods that are able to correct for confounding effects without the need of characterizing all the confounders have broader applications [22, 29-32]. Moreover, the predicted CTCs often only explain a certain proportion of variation in CTCs resulting in inflated test-statistics due to the uncaptured variation in CTCs. Existing reference-free methods are mainly based on the strategy of fitting a number of covariates (estimated from factor analysis or similar approaches with or without reference [22, 29, 31, 32]) in a fixed-effect model or a set of selected DNAm probes in a mixed linear model (MLM) [30]. However, uncharacterized confounders

with small effects and numerous correlations between distal DNAm probes (e.g. those on different chromosomes) induced by the confounders may not be well captured by either a fixed number of principal features or a subset of selected probes.

In this study, we proposed a reference-free method (called MOA: <u>M</u>LM-based <u>o</u>mic <u>a</u>ssociation) that fits all probes as random effects in an MLM-based association analysis to account for the confounding effects, including the correlations among distal probes induced by the confounding. We then extended the method to stratify the probes into multiple random-effect components (called MOMENT: <u>m</u>ulti-component <u>M</u>LM-based omic association <u>e</u>xcludi<u>n</u>g the <u>t</u>arget) to model a scenario where some probes are much more strongly associated with the phenotype than others. We evaluated the performance of MOA and MOMENT by extensive simulations and demonstrated their reliability and robustness in comparison with existing methods. We have implemented MOA and MOMENT together with a comprehensive set of other methods for omic data analysis in an easy-to-use and computationally efficient software package, OSCA (<u>o</u>mic-data-ba<u>s</u>ed <u>c</u>omplex trait <u>a</u>nalysis).

Results
Overview of the OSCA software
OSCA comprises four main modules: 1) data management for which we designed a binary format to efficiently store and manage omic data; 2) linear-regression- and MLM-based methods (including the methods proposed in this study) to test for associations between omic measures and complex traits; 3) methods to estimate the proportion of variance in a complex trait captured by all the measures of one or multiple omic profiles (e.g., all SNPs and DNAm probes), and to predict the trait phenotype in a new sample based on the joint effects of all omic measures estimated in a discovery sample; and 4) an efficient implementation of the methods to identify genetic variants associated with an omic profile, e.g., DNA methylation quantitative trait loci (mQTL) analysis. We will describe the methods based on DNAm data but the methods and software tool are in principle applicable to other types of omic data including gene expression, DNAm, histone modification, and protein abundance. The computer code of OSCA is written in C++ programming language and supports multi-threading based on OpenMP for high-performance computing. The compiled binary files are freely available at http://cnsgenomics.com/software/osca/.

MLM-based omic association analysis methods

One of the primary applications of OSCA is to test for associations between omic measures (e.g. DNAm probes) and a complex trait (e.g. body mass index (BMI)) correcting for confounding effects. In an MWAS, the test-statistics can be inflated because of the associations of probes with confounders that are correlated with the phenotype. Note that, even if the confounders are not directly associated with the phenotype, the presence of confounders (e.g., CTCs or experimental batches) can cause correlations between the trait-associated probes and the null probes in distal genomic regions or even on different chromosomes, giving rise to inflated test-statistics of the null probes (see the simulation results below). Existing methods that fit a number of covariates computed from dimension reduction approaches in a fixed-effect model [22, 31, 32] or a set of selected DNAm probes in an MLM [30] may not be sufficient to correct for confounding effects widely spread among a large number of probes or correlations between distal probes induced by the confounding. We propose two MLM-based approaches (MOA and MOMENT) that include all the (distal) probes as random effects in the model to account for the effects of the confounders on the trait and the probes as well as the correlations among distal probes. We show by simulations (see below) that both MOA and MOMENT are more robust than existing methods in controlling for false positive rate (FPR) in MWAS.

Here we start with a general model that fits all probes as random effects, i.e.,

$$\mathbf{y} = \mathbf{C\beta} + \mathbf{Wu} + \mathbf{e} \qquad [1]$$

where $\mathbf{y}$ is an $n \times 1$ vector of phenotype values with n being the sample size, $\mathbf{C}$ is an $n \times p$ matrix for covariates (e.g., age and sex) with p being the number of covariates, $\mathbf{\beta}$ is a $p \times 1$ vector of the effects of covariates on the phenotype, $\mathbf{W}$ is an $n \times m$ matrix of standardised DNAm measures of all m probes, $\mathbf{u}$ is an $m \times 1$ vector of the joint effects of all probes on the phenotype, and $\mathbf{e}$ is an $n \times 1$ vector of residuals. In this model, $\mathbf{\beta}$ are fixed effects whereas $\mathbf{u}$ and $\mathbf{e}$ are random effects with $\mathbf{u} \sim N(\mathbf{0}, \mathbf{I}\sigma_u^2)$ and $\mathbf{e} \sim N(\mathbf{0}, \mathbf{I}\sigma_e^2)$. The variance-covariance matrix for $\mathbf{y}$ is $var(\mathbf{y}) = \mathbf{V} = \mathbf{WW}'\sigma_u^2 + \mathbf{I}\sigma_e^2$. This equation can be re-written as

$$\mathbf{V} = \mathbf{A}\sigma_o^2 + \mathbf{I}\sigma_e^2 \text{ with } \mathbf{A} = \mathbf{WW}'/m \text{ and } \sigma_o^2 = m\sigma_u^2 \qquad [2]$$

where $\mathbf{A}$ is defined as the omic-data-based relationship matrix (ORM) (**Methods**) and σ_o^2 is the amount of phenotypic variance captured by all probes. The variance components (σ_o^2 and σ_e^2) in such an MLM can be estimated by REML algorithms [33]. Analogous to the method for estimating SNP-based heritability [34, 35], the proportion of variance in the phenotype captured by all the probes can be defined as $\rho^2 = \sigma_o^2/(\sigma_o^2 + \sigma_e^2)$. We name this variance-estimation method OREML following the nomenclature of GREML [34].

Model [1] can be extended to test for association between a probe i and the trait, i.e.,

$$\mathbf{y} = \mathbf{w}_i b_i + \mathbf{C}\boldsymbol{\beta} + \mathbf{W}\mathbf{u} + \mathbf{e} \text{ with } \mathbf{V} = \mathbf{W}\mathbf{W}'\sigma_u^2 + \mathbf{I}\sigma_e^2 \qquad [3]$$

In comparison to model [1], model [3] has two additional terms, $\mathbf{w}_i$ (an $n \times 1$ vector of standardised DNAm measures of a probe i, i.e., the target probe) and b_i (the effect of probe i on the phenotype; fixed effect). The probe effect b_i (together with the covariates' effects) can be estimated by the generalized least squares (GLS) approach, i.e., $\left[\hat{b}_i\ \widehat{\boldsymbol{\beta}}\right]' = (\mathbf{X}'\mathbf{V}^{-1}\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ and $\text{var}\left[\hat{b}_i\ \widehat{\boldsymbol{\beta}}\right]^T = (\mathbf{X}'\mathbf{V}^{-1}\mathbf{X})^{-1}$ with $\mathbf{X} = [\mathbf{w}_i\ \mathbf{C}]$. The sampling variance (standard error (SE) squared) of $\hat{b}_i$ is the first diagonal element of $\text{var}\left[\hat{b}_i\ \widehat{\boldsymbol{\beta}}\right]^T$. The null hypothesis ($H_0: b_i = 0$) can be tested by a two-sided t-test (or approximately chi-squared test if sample size is large) given $\hat{b}_i$ and its SE. We call this method MOA. As with the approximate MLM-based GWAS methods [36, 37], to speed up the computation, we assume that the estimates of σ_u^2 and σ_e^2 under the null (i.e., $b_i = 0$) are approximately equal to those under the alternative (i.e., $b_i \neq 0$) so that we do not need to compute $\mathbf{V}^{-1}$ repeatedly for every target probe.

There are two properties of the MOA method worthy of consideration. First, the target probe is fitted twice in the MOA model, once as a fixed effect (b_i) and again as a random effect (the i-th element of $\mathbf{u}$), resulting in a loss of power to detect b_i (a recognised issue in MLM-based association analysis with SNP data [37, 38]). This problem can be solved by leaving out probes in close physical proximity of the target probe (including the target) from the random-effect term because DNAm status of CpG sites in close physical proximity are likely to be regulated by the same mechanism and therefore tend to be highly correlated. This strategy has been used previously in both GWAS (genome-wide association study) [37, 38] and MWAS [30]. In practice, we exclude the probes <50Kb from the target probe (note that the distance parameter may differ for other types of omic data). Second, MOA assumes a single distribution to all the probe effects in the random-effect term, which may not be well fitted to data if some probes have much stronger associations with the trait than other probes. For example, if CTCs are associated with the phenotype, then all the probes that are highly differentially methylated in different cell types [39-41] may present a very different distribution of effects from the other probes. One solution to this issue is to stratify the probes into multiple groups by the association test statistics (from linear regression) and fit them as separate random-effect terms in the model. We extended the MOA method with the two modifications mentioned above and named it as MOMENT (multi-component MLM-based omic association excluding the target). The MOMENT model can be written as

$$\mathbf{y} = \mathbf{w}_i b_i + \mathbf{C}\boldsymbol{\beta} + \sum_j \mathbf{W}_j \mathbf{u}_j + \mathbf{e} \text{ with } \mathbf{V} = \sum_j \mathbf{W}_j \mathbf{W}_j{}' \sigma_{u_j}^2 + \mathbf{I}\sigma_e^2 \quad [4]$$

where $\mathbf{W}_j$ is an $n \times m_j$ matrix of standardised DNAm measures of the probes in the j-th group with m_j being the number of probes in the group (excluding the probes <50Kb from the target probe). In practice, the probes are split into two groups by association p-values from a linear regression model (i.e., $\mathbf{y} = \mathbf{w}_i b_i + \mathbf{C\beta} + \mathbf{e}$) at a genome-wide significant threshold (or by the proportion of phenotypic variance captured per probe at a certain threshold). The GLS method described in model [3] can be used to estimate b_i and its SE for hypothesis testing with $\sigma_{u_j}^2$ and σ_e^2 (under the null) re-estimated every time when a different set of probes are excluded from the random-effect terms.

Simulation analysis

To quantify the FPR and power for MOMENT (implemented in OSCA), we performed simulations based on DNAm and CTCs measures from the Lothian Birth Cohorts (LBC) in three scenarios (**Supplementary Note 1**). That is, we simulated a phenotype: 1) with effects from a set of "causal probes" (randomly selected from all probes on the odd chromosomes) but no effects from the CTCs; 2) with small to large effects from CTCs but no direct effects from the probes; and 3) with effects from both the causal probes and CTCs (**Supplementary Note 1**). Note that we only sampled the causal probes from the odd chromosomes in scenarios 1 and 3, leaving the probes on the even chromosomes to quantify the FPR under the null. Results from our models were compared to 6 different methods including: 1) Unadj: linear regression without adjustment; 2) SVA: linear regression with the SVA surrogate variables fitted as covariates [22]; 3) LFMM2-ridge: a latent factor mixed model (LFMM) using ridge algorithm for confounder estimation [32]; 4) LFMM2-lasso: an LFMM using lasso algorithm for confounder estimation [32]; 5) ReFACTor: linear regression with the first 5 sparse principal components (PCs) from ReFACTor fitted as covariates [31]; 6) 5PCs: linear regression with the first 5 PCs, computed from a principal component analysis (PCA), fitted as covariates; and 7) FaST-LMM-EWASher: a set of selected probes fitted as random effect in an MLM [30]. For completeness of the analysis, we also included MOA (implemented in OSCA) in the comparison.

In simulation scenario 1, although there were no direct effects of the CTCs on the phenotype, the test-statistics from Unadj at the null probes were inflated (**Figure 1a** and **Supplementary Table 1**) because the null and causal probes, although on different sets of chromosomes, are correlated through their correlations with the CTCs (and/or other systematic biases such as experimental batches). The mean genomic inflation factor for Unadj from 100 simulation replicates was 7.67 (**Supplementary Table 1**), where the genomic inflation factor (λ) is defined as the median of χ^2 test-statistics of all probes divided by its expected value under the null. While all the other methods were

much less inflated compared to Unadj, MOMENT and MOA showed the least inflation with a mean λ value close to 1. It is important to note that the FPRs of all the methods except MOA and MOMENT are highly inflated (FPR > 0.6) (**Figure 1b** and **Supplementary Table 1**) despite the relatively small genomic inflation for most of the methods (**Figure 1a**). Here, FPR is defined as the number of simulation replicates with at least one genome-wide significant probe (i.e., MWAS p-values < 0.05 / m with m being the number of probes) divided by the number of simulation replicates, which can be interpreted as the probability of observing one or more false positives at a genome-wide significance level in a single experiment. There was no inflation in FPR for MOMENT, and a marginal inflation for MOA (**Figure 1b** and **Supplementary Table 1**), showing the effectiveness of using all (distal) probes to account for the probe correlations. We also compared the power among the methods. Since the test-statistics for many approaches were inflated, for a fair comparison, we recalibrated the power of each method by dividing the test-statistics of the causal probes (on the odd chromosomes) by the λ value of the null probes (on the even chromosomes). Apart from Unadj, the powers of all the methods were on a similar level with small differences reflecting the differences in the λ value of the null probes (**Figures 1c** and **1d**). The conclusions held in additional simulations varying the number of causal probes and the proportion of phenotypic variance captured by the causal probes (**Supplementary Figures 1** and **2**) despite that the inflation in FPR for the existing methods appeared to increase with the increase of the proportion of variance captured per causal probe.

In simulation scenario 2 where there is no direct probe-trait association, the χ^2 test-statistics of all the probes are expected to follow a χ^2 distribution with 1 degree of freedom if the effects of CTCs have been well accounted for. The results showed that the λ value was close to 1 for all the methods except Unadj and FaST-LMM-EWASher (**Figure 2a**). It seems that, for some of the methods (e.g. 5PCs and ReFACTor), the λ value slightly increased with the increase of the proportion of variance explained by the CTCs (R^2_{CTCs}) (**Figure 2a**). Nevertheless, a non-inflated median test-statistic does not necessarily mean that the FPR has been well controlled for. In fact, most methods showed inflated FPR in this simulation scenario, and the FPRs of all the methods increased with increasing R^2_{CTCs} (**Figure 2b**). The FPRs of 5PCs, ReFACTor, LFMM2-ridge, and LFMM2-lasso were close to the expected value (i.e., 0.05) when $R^2_{\text{CTCs}} = 0.005$ and increased to a level between 0.15 and 0.2 when $R^2_{\text{CTCs}} = 0.05$ (**Figure 2b**). The relationship between FPR and R^2_{CTCs} was relatively flat for SVA with its FPR varying from 0.05 to 0.1 when R^2_{CTCs} increased from 0.005 to 0.05. Although FaST-LMM-EWASher showed the most deflated test-statistics among all the methods (**Figure 2a**), its FPR was substantially higher than all the other methods except Unadj (**Figure 2b**), likely due to its feature selection strategy

(**Supplementary Note 2**). MOA and MOMENT performed similarly in this simulation scenario and showed the lowest inflation in FPR among all the methods with their FPRs being lower than 0.05 when R^2_{CTCs} = 0.005 and increased to about 0.1 when R^2_{CTCs} = 0.05 (**Figure 2b**). In addition, we performed a linear regression analysis with the known CTCs fitted as covariates; as expected, the FPR of was close to 0.05 irrespective of the level of R^2_{CTCs} (see below for the analysis with predicted CTCs).

We further compared the methods under the circumstance (simulation scenario 3) where there were associations between the phenotype and CTCs ($R^2_{\text{CTCs}} = 0.05$) and the null probes were correlated with distal causal probes because both of them were correlated with CTCs (**Supplementary Note 1**). The results were similar to those above (**Figure 1**). That is, the FPR of MOMENT was close to the expected value, demonstrating the reliability and robustness of the method. The FPR of MOA is slightly higher than that of MOMENT but much lower than those of the other methods that showed strong inflation in FPR due to the correlations between causal and null probes (**Supplementary Figures 3a** and **3b** and **Supplementary Table 2**). The mean test-statistic of the causal probes from MOMENT was significantly lower than that from MOA but slightly higher than the other methods after recalibration (dividing the test-statistics of the causal probes by the λ value of the null probes) (**Supplementary Figure 3c**). The statistical powers of MOA, MOMENT, LFMM2-ridge, LFMM2-lasso and FaST-LMM-EWASher were similar and slightly higher than those of 5PCs and SVA (**Supplementary Figure 3d**). The conclusions held with different sample sizes (**Supplementary Figures 4** and **5**) or different numbers of causal probes with smaller or larger variance explained per causal probe (**Supplementary Figures 6** and **7**). The conclusions also held if we simulated confounding effects on experimental batches in lieu of CTCs (**Supplementary Figures 8** and **9**). We further demonstrated the consistency between the result from MOA/MOMENT analysis of the whole sample and that from a meta-analysis of summary statistics from MOA/MOMENT analyses in two halves of the sample (**Supplementary Figure 10**).

An application of MOMENT to real data

We applied MOMENT and the other methods to four real quantitative traits in the LBC cohorts. These traits, including BMI, height, lung function (measured in the highest score of forced expiratory volume in one second), and walking speed (measured in the time taken to walk 6 meters), were standardised and corrected for age in each gender group within each sub-cohort (LBC1936 or LBC1921) (**Methods**). The standardised phenotypes were further processed by a rank-based inverse-normal

transformation. The DNAm probes were adjusted for age, sex, and experimental batches. We did not adjust the probes for CTCs or smoking status for the purpose of testing methods (see below).

Consistent with the results from simulations, the test-statistics from MOA and MOMENT were not inflated whereas all the other methods showed modest inflation for all the traits (**Figure 3, Table 1,** and **Supplementary Figures 11-14**). There were three associations that were identified by multiple methods, including one for BMI (cg11202345, detected by all methods), in line with a previous study [42], and two for lung function (cg05575921 and cg05951221, detected by all methods except MOMENT) (**Supplementary Table 3, Supplementary Figures 11** and **13**). It should be noted that cg05575921 was reported to be associated with smoking in a previous study [43], indicating that the association between cg05575921 and lung function might be confounded by smoking status. Moreover, MOA, LFMM2-ridge, LFMM2-lasso, and ReFACTor consistently identified 12 additional probes associated with lung function, and most of the probes have been linked to smoking in a previous study [44]. Almost all the associations were not significant when smoking status was fitted as a covariate in the models (6.5% of variance in lung function associated with smoking status; **Supplementary Table 4** and **Supplementary Figure 15**), suggesting that most (if not all) of the probe associations with lung function identified by the four methods mentioned above were owing to the confounding of smoking. None of the smoking-associated probes were genome-wide significant for lung function in the analysis using MOMENT (**Supplementary Figure 13**) and the result remained the same when smoking status was fitted as a covariate in MOMENT (**Supplementary Figure 15**), showing the capability of MOMENT in correcting for unobserved confounding factors as demonstrated by the simulation above.

It has been demonstrated in previous GWASs that MLM-based association analysis methods developed for quantitative traits are applicable to discrete traits including ascertained case-control status [36, 37, 45, 46]. To demonstrate the applicability of the proposed methods to discrete phenotypes, we analysed smoking status (non-smoker, former smoker or current smoker; coded as 0, 1 or 2) in LBC by MOA and MOMENT in comparison with the existing methods. All the methods detected a large number (at least 112) of probes at a genome-wide significance level ($P < 2.19e-7$) except for MOMENT and EWASher which only identified 4 and 2 genome-wide significant probes, respectively (**Supplementary Figure 16**). To validate the association signals other than those identified by MOMENT, we fitted the 4 MOMENT probes as fixed covariates in MOA. None of the additional associations remained genome-wide significant conditioning on the 4 MOMENT probes

(**Supplementary Figure 17**), suggesting that those additional associations detected by MOA were driven by their correlations with the 4 MOMENT signals. MOA failed in this scenario likely because the associations of the 4 MOMENT signals were too strong to be fitted in a signal normal distribution with the other probes. This conclusion is further supported by the result that the accuracy of predicting/classifying smoking status in a cross-validation setting using the large number of probes detected by linear regression or MOA was even lower than that using the small number of probes detected by MOMENT (**Supplementary Table 5**). In addition, we recoded the smoking status data to a binary phenotype (0 for non-smoker and 1 for former or current smoker) and applied all the methods to the recoded binary phenotype; the conclusions were similar as above but it seemed that the analyses with the binary phenotype were less powerful than those with the categorical phenotype above (**Supplementary Figure 18**). All these results show the applicability of MOMENT to discrete traits and again demonstrate the robustness and reliability of MOMENT in controlling for false positive associations.

Estimating variance in a phenotype captured by all probes by OREML

We have demonstrated the performance of the omic-data-based association analysis methods in OSCA by simulation and real data analysis. We then turned to evaluate the performance of OREML in estimating the proportion of variance in a complex trait captured by all probes (ρ^2) by simulation in two scenarios (**Supplementary Note 1**). The results showed that under either scenario, OREML reported an unbiased estimate of ρ^2 (**Supplementary Table 6**). Here the unbiasedness is defined as that the mean ρ^2 estimate from 500 independent simulations is not significantly different from the ρ^2 parameter used in the simulation. There are two methods implemented in OSCA to compute the ORM (**Methods**). Our simulation results showed that the estimates of ρ^2 based on the two methods were similar (**Supplementary Table 6**).

We also attempted to partition and estimate the proportions of phenotypic variation captured by all SNPs (i.e., h^2_{SNP}) and all the DNAm probes respectively when they were fitted jointly in one model. We first investigated the correlation between genomic relationship matrix (GRM) and methylomic relationship matrix (MRM) in the LBC dataset. We found that the off-diagonal elements of the GRM were almost independent of those of the MRM (r = 0.0045; **Supplementary Figure 19**). From an OREML analysis that fits the GRM and MRM jointly, we estimated that all the DNAm probes captured 6.5% (SE = 0.038) of the variance for BMI but the estimate for height was nearly zero ($\hat{\rho}^2$ = -0.005 and SE = 0.0086) (**Supplementary Table 7**). These results are in line with the finding from a previous

study that the accuracy of genetic risk prediction can be improved by incorporating DNAm data for BMI but not height [14].

Discussion

In this study, we developed a versatile software tool—OSCA—to manage omic data generated from high-throughput experiments in large cohorts and to facilitate the analyses of complex traits using omic data (**Supplementary Note 4**). The primary applications of OSCA are to identify omic measures associated with a complex trait accounting for unobserved confounding factors (MOMENT) and to estimate the proportion of phenotypic variation captured by all measures of one or multiple omic profiles (OREML). A by-product of the OREML application is to estimate the joint effects of all measures of one or multiple omic profiles (i.e., OBLUP analysis) to predict the phenotype in a new sample. This has been shown to be a powerful and robust approach in age prediction using gene expression or DNAm data [47, 48]. We have also provided computationally efficient implementations in OSCA to manage large-scale omic data, and to perform omic-data-based quantitative trait locus (xQTL) analysis and meta-analysis of xQTL summary data. OSCA is an ongoing software development project so that any further methods or functions related to omic-data-based analysis can be included in the software package in the future.

We showed, by simulation, a surprisingly high FPR for all the existing MWAS/EWAS methods, owing to the correlations between distal probes induced by CTCs (and/or other systematic biases) in DNAm data (**Figure 1**). These correlations are widespread at a large number of probes across the genome (as demonstrated by the proportion of null probes with $P_{MWAS} < 0.05$ in simulation scenario 1; **Supplementary Figure 20**) and thus are not adequately accounted for by a fixed number of principal features computed from the data (e.g. 5PCs, ReFACTor, LFMM2, and SVA) nor a set of selected probes (e.g. FaST-LMM-EWASher). This conclusion is likely to be applicable to other types of omic data if the measures in distal genomic regions are correlated due to unmeasured confounding factors such as systematic experimental biases or unwanted biological variation. We show by simulation that this confounding effect can be corrected for by fitting the target probe as a fixed effect and all the other (distal) probes as random effects (i.e., the MOA or MOMENT method).

Our simulation also showed that, if CTCs or batches explain a large proportion of variation in the phenotype, the FPRs of all the methods tended to be inflated (**Supplementary Figures 21** and **22**) despite that the genomic inflation factor is close to unity for most methods (**Figure 2**). We re-ran the

simulation under a more extreme setting with R^2_{CTCs} varying from 10% to 70%. In this case, the genomic inflation factors of the fixed-effect models (i.e., SVA, ReFACTor, LFMM2, and 5PCs) and the FPRs of all the methods increased as R^2_{CTCs} increased (to a lesser extent for FaST-LMM-EWASher), suggesting that there were a set of probes strongly associated with CTCs (**Supplementary Figure 23**). Note that even in this extreme case, MOMENT showed the lowest FPR on average amongst all the methods. It is also of note that the FPR of FaST-LMM-EWASher was relatively low in this scenario (**Supplementary Figure 21**), opposite to its performance when R^2_{CTCs} was low (**Figure 2**), possibly due to its variable selection strategy (**Supplementary Note 2**). The inflation in FPR was only slightly alleviated by fitting the predicted CTCs as covariates (**Supplementary Figures 24** and **25**). The results also suggest that it may be worth fitting measured CTCs as fixed-effect covariates in MLM-based association analyses such as MOA and MOMENT in practice although this approach is likely to be conservative as indicated by the deflated λ and FPR (**Supplementary Figure 26**). These conclusions are likely to be applicable to other confounding factors such as smoking status, as demonstrated in the real data analysis for lung function (**Supplementary Figure 15**). Our results also caution the interpretation of associations identified from MWAS for traits that are highly correlated with CTCs and/or other biological confounders.

In conclusion, we showed by simulation the inflation in test-statistics of the existing MWAS methods because of the ubiquitous correlations between distal probes caused by confounding factors, and developed two new MWAS methods (MOA and MOMENT) to correct for the inflation. We demonstrated the reliability and robustness of MOMENT by simulations in a number of scenarios and real data analysis. We recommend the use of MOMENT in practice because of its robustness in the presence of unobserved confounders despite that it is slightly less powerful than MOA. We implemented both MOA and MOMENT in a computationally efficient and easy-to-use software tool OSCA together with many other functions for omic-data-based analyses (**Supplementary Figure 27**).

Methods

Omic-data-based relationship matrix (ORM)

We have described in equations [1] and [2] the OREML model to estimate the proportion of variance in a phenotype captured by the DNAm probes all together. In equation [1], i.e., $\mathbf{y} = \mathbf{C\beta} + \mathbf{Wu} + \mathbf{e}$, we define $\mathbf{W}$ as a matrix of standardised measures of DNAm probes, and in equation [2] we define the ORM as $\mathbf{A} = \mathbf{WW'}/m$. Therefore, the omic relationship between individual j and k (the jk-th element of $\mathbf{A}$) can be computed as $A_{jk} = \frac{1}{m}\sum_i (x_{ij} - \mu_i)(x_{ik} - \mu_i)/\sigma_i^2$, where x_{ij} is the unstandardised DNAm level of probe i in individual j, μ_i and σ_i^2 are the mean and variance of the i-th probe over all the individuals respectively, and m is the number of probes. This model implicitly assumes that the probes of smaller variance in DNAm level (unstandardised) tend to have larger effects on the phenotype (strictly speaking, stronger associations with the phenotype), and that there is no relationship between the proportion of trait variance captured by a probe and the variance of the probe. We also provide in OSCA an alternative method to compute the ORM, i.e., $A_{jk} = \sum_i (x_{ij} - \mu_i)(x_{ik} - \mu_i) / \sum_i \sigma_i^2$. If we use this definition of ORM in the OREML analysis, we implicitly assume that there is no relationship between the probe effect on the trait and the variance of the probe but the proportion of trait variance associated with a probe increases as the variance of the probe increases. We showed by simulation and real data analysis that the difference between the OREML results using the two methods was very small (**Supplementary Tables 6** and **7**).

OREML: estimating the proportion of trait variance captured by all DNAm probes

We have shown in equations [1] and [2] an OREML model with one random-effect component to estimate the proportion of trait variance captured by all DNAm probes. The model is flexible and can be extended to partition the trait variance into components associated with different sets of probes (e.g. a model with 22 components with all the probes on each chromosome as a component). A flexible OREML model can be written as

$\mathbf{y} = \mathbf{C\beta} + \sum_i \mathbf{W}_i \mathbf{u}_i + \mathbf{e}$ with var($\mathbf{y}$) = $\mathbf{V} = \sum_i \mathbf{W}_i \mathbf{W}_i' \sigma_{u_i}^2 + \mathbf{I}\sigma_e^2 = \sum_i \mathbf{A}_i \sigma_{o_i}^2 + \mathbf{I}\sigma_e^2$

where the definitions of all the parameters and variables are similar to those in equations [1] and [2].The variance components can be estimated by REML [33], and the proportion of the trait variance captured by the i-th component can be computed as $\rho_i^2 = \sigma_{o_i}^2/(\sum_i \sigma_{o_i}^2 + \sigma_e^2)$.

The multi-component OREML model can be applied to partition the trait variance into components associated with multiple omic profiles. For example, if SNP genotype, DNAm, and gene expression data are available for all the individuals in a cohort, a multi-component OREML model can be used

to estimate the proportion of trait variance captured by all SNPs (i.e., the SNP-based heritability), the expression levels of all genes, and the DNAm levels at all the CpG sites. The model can be written as $\mathbf{y} = \mathbf{C\beta} + \mathbf{W}_g\mathbf{u}_g + \mathbf{W}_t\mathbf{u}_t + \mathbf{W}_m\mathbf{u}_m + \mathbf{e}$ with $\mathrm{var}(\mathbf{y}) = \mathbf{A}_g\sigma_g^2 + \mathbf{A}_t\sigma_t^2 + \mathbf{A}_m\sigma_m^2 + \mathbf{I}\sigma_e^2$ where $\mathbf{W}_g$, $\mathbf{W}_t$, and $\mathbf{W}_m$ are the matrices of standardised SNP genotypes, gene expression measures, and DNAm levels, respectively, with the corresponding effects $\mathbf{u}_g$, $\mathbf{u}_t$ and $\mathbf{u}_m$; $\mathbf{A}_g = \mathbf{W}_g\mathbf{W}_g'/m_g$ is the genomic relationship matrix (GRM) with m_g being the number of SNPs, $\mathbf{A}_t = \mathbf{W}_t\mathbf{W}_t'/m_t$ is the transcriptomic relationship matrix (TRM) with m_t being the number of transcripts, and $\mathbf{A}_m = \mathbf{W}_m\mathbf{W}_m'/m_m$ is the methylomic relationship matrix (MRM) with m_m being the number of DNAm probes. Note that the model can be reduced by dropping any of the variance components or expanded by including other types of omic profiles (e.g., protein abundance).

Dataset

The LBC cohorts [49, 50] consisted of individuals born in 1921 (LBC1921) and 1936 (LBC1936), mostly living in Edinburgh city and the surrounding Lothian region of Scotland. Blood samples were collected with informed consent. The use of human participants in this study was approved by University of Queensland Human Research Ethics Committee B (approval number: 2011001173). The LBC individuals underwent several waves of SNP genotyping and DNAm measures. DNA methylation levels at 485,512 CpG sites across the genome were measured on 3,191 whole blood samples from 3 waves using the Illumina HumanMethylation450 BeadChip. Duplicates or samples with an excessive proportion of low confidence calls across all probes (>5%) were removed. Probes with an excessive proportion of low confidence calls across all individuals (>5%) or probes located in sex chromosomes were excluded. In addition, probes encompassing SNPs annotated in dbSNP131 using hg19 coordinates or identified as potentially cross-hybridized methylation probes by a previous study [51] were also excluded. After these QC steps, 3,018 samples and 307,360 probes remained (**Supplementary Note 3**). We included in the analysis only the first wave (wave1) of the LBC data consisting of 436 individuals from LBC1921 (average age of 79 years) and 906 individuals from LBC1936 (average age of 70 years) (**Supplementary Table 8**). We finally removed probes with almost invariable beta values across individuals (standard deviation < 0.02) and retained 1,342 individuals and 228,694 probes for further analysis.

There were a number of covariates available in the LBC data including age, sex, batches of the experiment (i.e. plate and position of the sample on a chip), and CTCs. The blood cell counts for different cell types, including basophils, eosinophils, monocytes, lymphocytes, and neutrophils, were

quantified using an LH50 Beckman Coulter instrument on the same day of blood collection. In addition to the covariates, there are a number of traits measured on the LBC individuals including height (measured without shoes), body mass index (BMI), lung function (measured in the highest score of forced expiratory volume in one second), and walking speed (measured in the time taken to walk 6 metres) and smoking status (never smoked, ex-smoker or current smoker) [52, 53]. The numbers of missing measurements are noted in **Supplementary Table 9**. For each trait, we adjusted the phenotype for age in each gender group of each cohort (LBC1921 or LBC1936) and standardised the residuals by rank-based inverse normal transformation. This process removes the age effect and potential difference in mean and variance between two gender groups or cohorts.

The LBC wave1 individuals were also genotyped by Illumina 610-Quadv1 BeadChip. The QC process of the SNP genotype data has been detailed elsewhere [14]. After excluding SNPs from sex chromosomes and SNPs with low allelic frequency (MAF < 0.01), we retained 523,062 genotyped SNPs for analysis.

URLs

OSCA, http://cnsgenomics.com/software/osca;

ReFACTor, https://www.cs.tau.ac.il/~heran/cozygene/software/refactor.html;

EWASher, https://www.microsoft.com/en-us/research/project/fast-lmm-software-papers/;

SVA, https://bioconductor.org/packages/release/bioc/html/sva.html;

LFMM2, https://bcm-uga.github.io/lfmm/;

Data availability

The LBC data are available at https://www.ebi.ac.uk/ega/studies/EGAS00001000910.

References
1. Ritchie MD, Holzinger ER, Li R, Pendergrass SA, Kim D: **Methods of integrating data to uncover genotype-phenotype interactions.** *Nat Rev Genet* 2015, **16:**85-97.
2. Hasin Y, Seldin M, Lusis A: **Multi-omics approaches to disease.** *Genome Biol* 2017, **18:**83.
3. Wu Y, Zeng J, Zhang F, Zhu Z, Qi T, Zheng Z, Lloyd-Jones LR, Marioni RE, Martin NG, Montgomery GW, et al: **Integrative analysis of omics summary data reveals putative mechanisms underlying complex traits.** *Nat Commun* 2018, **9:**918.
4. Consortium GT, Laboratory DA, Coordinating Center -Analysis Working G, Statistical Methods groups-Analysis Working G, Enhancing Gg, Fund NIHC, Nih/Nci, Nih/Nhgri, Nih/Nimh, Nih/Nida, et al: **Genetic effects on gene expression across human tissues.** *Nature* 2017, **550:**204-213.

5. Lloyd-Jones LR, Holloway A, McRae A, Yang J, Small K, Zhao J, Zeng B, Bakshi A, Metspalu A, Dermitzakis M, et al: **The Genetic Architecture of Gene Expression in Peripheral Blood.** *Am J Hum Genet* 2017, **100:**371.

6. Hannon E, Spiers H, Viana J, Pidsley R, Burrage J, Murphy TM, Troakes C, Turecki G, O'Donovan MC, Schalkwyk LC, et al: **Methylation QTLs in the developing brain and their enrichment in schizophrenia risk loci.** *Nat Neurosci* 2016, **19:**48-54.

7. Jaffe AE, Gao Y, Deep-Soboslay A, Tao R, Hyde TM, Weinberger DR, Kleinman JE: **Mapping DNA methylation across development, genotype and schizophrenia in the human frontal cortex.** *Nat Neurosci* 2016, **19:**40-47.

8. Grubert F, Zaugg JB, Kasowski M, Ursu O, Spacek DV, Martin AR, Greenside P, Srivas R, Phanstiel DH, Pekowska A, et al: **Genetic Control of Chromatin States in Humans Involves Local and Distal Chromosomal Interactions.** *Cell* 2015, **162:**1051-1065.

9. Chen L, Ge B, Casale FP, Vasquez L, Kwan T, Garrido-Martin D, Watt S, Yan Y, Kundu K, Ecker S, et al: **Genetic Drivers of Epigenetic and Transcriptional Variation in Human Immune Cells.** *Cell* 2016, **167:**1398-1414 e1324.

10. Battle A, Khan Z, Wang SH, Mitrano A, Ford MJ, Pritchard JK, Gilad Y: **Genomic variation. Impact of regulatory variation from RNA to protein.** *Science* 2015, **347:**664-667.

11. Folkersen L, Fauman E, Sabater-Lleal M, Strawbridge RJ, Franberg M, Sennblad B, Baldassarre D, Veglia F, Humphries SE, Rauramaa R, et al: **Mapping of 79 loci for 83 plasma protein biomarkers in cardiovascular disease.** *PLoS Genet* 2017, **13:**e1006706.

12. Wahl S, Drong A, Lehne B, Loh M, Scott WR, Kunze S, Tsai PC, Ried JS, Zhang W, Yang Y, et al: **Epigenome-wide association study of body mass index, and the adverse outcomes of adiposity.** *Nature* 2017, **541:**81-86.

13. Gusev A, Mancuso N, Won H, Kousi M, Finucane HK, Reshef Y, Song L, Safi A, Schizophrenia Working Group of the Psychiatric Genomics C, McCarroll S, et al: **Transcriptome-wide association study of schizophrenia and chromatin activity yields mechanistic disease insights.** *Nat Genet* 2018, **50:**538-548.

14. Shah S, Bonder MJ, Marioni RE, Zhu Z, McRae AF, Zhernakova A, Harris SE, Liewald D, Henders AK, Mendelson MM, et al: **Improving Phenotypic Prediction by Combining Genetic and Epigenetic Associations.** *Am J Hum Genet* 2015, **97:**75-85.

15. van Kessel KEM, van der Keur KA, Dyrskjot L, Algaba F, Welvaart NYC, Beukers W, Segersten U, Keck B, Maurer T, Simic T, et al: **Molecular Markers Increase Precision of the European Association of Urology Non-Muscle-Invasive Bladder Cancer Progression Risk Groups.** *Clinical Cancer Research* 2018, **24:**1586-1593.

16. Gamazon ER, Wheeler HE, Shah KP, Mozaffari SV, Aquino-Michaels K, Carroll RJ, Eyler AE, Denny JC, Consortium GT, Nicolae DL, et al: **A gene-based association method for mapping traits using reference transcriptome data.** *Nat Genet* 2015, **47:**1091-1098.

17. Gusev A, Ko A, Shi H, Bhatia G, Chung W, Penninx BW, Jansen R, de Geus EJ, Boomsma DI, Wright FA, et al: **Integrative approaches for large-scale transcriptome-wide association studies.** *Nat Genet* 2016, **48:**245-252.

18. Zhu Z, Zhang F, Hu H, Bakshi A, Robinson MR, Powell JE, Montgomery GW, Goddard ME, Wray NR, Visscher PM, Yang J: **Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets.** *Nat Genet* 2016, **48:**481-487.

19.   Liu Y, Aryee MJ, Padyukov L, Fallin MD, Hesselberg E, Runarsson A, Reinius L, Acevedo N, Taub M, Ronninger M, et al: **Epigenome-wide association data implicate DNA methylation as an intermediary of genetic risk in rheumatoid arthritis.** *Nat Biotechnol* 2013, **31:**142-147.

20.   Michels KB, Binder AM, Dedeurwaerder S, Epstein CB, Greally JM, Gut I, Houseman EA, Izzi B, Kelsey KT, Meissner A, et al: **Recommendations for the design and analysis of epigenome-wide association studies.** *Nat Methods* 2013, **10:**949-955.

21.   Jaffe AE, Irizarry RA: **Accounting for cellular heterogeneity is critical in epigenome-wide association studies.** *Genome Biology* 2014, **15.**

22.   Leek JT, Storey JD: **Capturing heterogeneity in gene expression studies by surrogate variable analysis.** *PLoS Genet* 2007, **3:**1724-1735.

23.   Teschendorff AE, Zheng SC: **Cell-type deconvolution in epigenome-wide association studies: a review and recommendations.** *Epigenomics* 2017, **9:**757-768.

24.   Teschendorff AE, Relton CL: **Statistical and integrative system-level analysis of DNA methylation data.** *Nat Rev Genet* 2018, **19:**129-147.

25.   Rakyan VK, Beyan H, Down TA, Hawa MI, Maslau S, Aden D, Daunay A, Busato F, Mein CA, Manfras B, et al: **Identification of type 1 diabetes-associated DNA methylation variable positions that precede disease diagnosis.** *PLoS Genet* 2011, **7:**e1002300.

26.   Teschendorff AE, Menon U, Gentry-Maharaj A, Ramus SJ, Gayther SA, Apostolidou S, Jones A, Lechner M, Beck S, Jacobs IJ, Widschwendter M: **An epigenetic signature in peripheral blood predicts active ovarian cancer.** *PLoS One* 2009, **4:**e8274.

27.   Guintivano J, Aryee MJ, Kaminsky ZA: **A cell epigenotype specific model for the correction of brain cellular heterogeneity bias and its application to age, brain region and major depression.** *Epigenetics* 2013, **8:**290-302.

28.   Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, Wiencke JK, Kelsey KT: **DNA methylation arrays as surrogate measures of cell mixture distribution.** *Bmc Bioinformatics* 2012, **13.**

29.   Gagnon-Bartsch JA, Speed TP: **Using control genes to correct for unwanted variation in microarray data.** *Biostatistics* 2012, **13:**539-552.

30.   Zou J, Lippert C, Heckerman D, Aryee M, Listgarten J: **Epigenome-wide association studies without the need for cell-type composition.** *Nat Methods* 2014, **11:**309-311.

31.   Rahmani E, Zaitlen N, Baran Y, Eng C, Hu D, Galanter J, Oh S, Burchard EG, Eskin E, Zou J, Halperin E: **Sparse PCA corrects for cell type heterogeneity in epigenome-wide association studies.** *Nat Methods* 2016, **13:**443-445.

32.   Caye K, Jumentier B, Francois O: **LFMM 2.0: Latent factor models for confounder adjustment in genome and epigenome-wide association studies.** *bioRxiv* 2018.

33.   Patterson HD, Thompson R: **Recovery of Inter-Block Information When Block Sizes Are Unequal.** *Biometrika* 1971, **58:**545-+.

34.   Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA, Heath AC, Martin NG, Montgomery GW, et al: **Common SNPs explain a large proportion of the heritability for human height.** *Nat Genet* 2010, **42:**565-569.

35.   Yang J, Lee SH, Goddard ME, Visscher PM: **GCTA: a tool for genome-wide complex trait analysis.** *Am J Hum Genet* 2011, **88:**76-82.

36. Kang HM, Sul JH, Service SK, Zaitlen NA, Kong SY, Freimer NB, Sabatti C, Eskin E: **Variance component model to account for sample structure in genome-wide association studies.** *Nat Genet* 2010, **42:**348-354.

37. Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL: **Advantages and pitfalls in the application of mixed-model association methods.** *Nat Genet* 2014, **46:**100-106.

38. Lippert C, Listgarten J, Liu Y, Kadie CM, Davidson RI, Heckerman D: **FaST linear mixed models for genome-wide association studies.** *Nat Methods* 2011, **8:**833-835.

39. Meissner A, Mikkelsen TS, Gu H, Wernig M, Hanna J, Sivachenko A, Zhang X, Bernstein BE, Nusbaum C, Jaffe DB, et al: **Genome-scale DNA methylation maps of pluripotent and differentiated cells.** *Nature* 2008, **454:**766-770.

40. Laurent L, Wong E, Li G, Huynh T, Tsirigos A, Ong CT, Low HM, Kin Sung KW, Rigoutsos I, Loring J, Wei CL: **Dynamic changes in the human methylome during differentiation.** *Genome Res* 2010, **20:**320-331.

41. McGregor K, Bernatsky S, Colmegna I, Hudson M, Pastinen T, Labbe A, Greenwood CM: **An evaluation of methods correcting for cell-type heterogeneity in DNA methylation studies.** *Genome Biol* 2016, **17:**84.

42. Mendelson MM, Marioni RE, Joehanes R, Liu C, Hedman AK, Aslibekyan S, Demerath EW, Guan W, Zhi D, Yao C, et al: **Association of Body Mass Index with DNA Methylation and Gene Expression in Blood Cells and Relations to Cardiometabolic Disease: A Mendelian Randomization Approach.** *PLoS Med* 2017, **14:**e1002215.

43. Gao X, Jia M, Zhang Y, Breitling LP, Brenner H: **DNA methylation changes of whole blood cells in response to active smoking exposure in adults: a systematic review of DNA methylation studies.** *Clin Epigenetics* 2015, **7:**113.

44. Shenker NS, Polidoro S, van Veldhoven K, Sacerdote C, Ricceri F, Birrell MA, Belvisi MG, Brown R, Vineis P, Flanagan JM: **Epigenome-wide association study in the European Prospective Investigation into Cancer and Nutrition (EPIC-Turin) identifies novel genetic loci associated with smoking.** *Hum Mol Genet* 2013, **22:**843-851.

45. Listgarten J, Lippert C, Kadie CM, Davidson RI, Eskin E, Heckerman D: **Improved linear mixed models for genome-wide association studies.** *Nat Methods* 2012, **9:**525-526.

46. Zhou X, Stephens M: **Genome-wide efficient mixed-model analysis for association studies.** *Nat Genet* 2012, **44:**821-824.

47. Peters MJ, Joehanes R, Pilling LC, Schurmann C, Conneely KN, Powell J, Reinmaa E, Sutphin GL, Zhernakova A, Schramm K, et al: **The transcriptional landscape of age in human peripheral blood.** *Nat Commun* 2015, **6:**8570.

48. Zhang Q, Vallerga C, Walker R, Lin T, Henders A, Montgomery G, He J, Fan D, Fowdar J, Kennedy M, et al: **Improved prediction of chronological age from DNA methylation limits it as a biomarker of ageing.** *bioRxiv* 2018.

49. Deary IJ, Gow AJ, Pattie A, Starr JM: **Cohort profile: the Lothian Birth Cohorts of 1921 and 1936.** *Int J Epidemiol* 2012, **41:**1576-1584.

50. Taylor AM, Pattie A, Deary IJ: **Cohort Profile Update: The Lothian Birth Cohorts of 1921 and 1936.** *Int J Epidemiol* 2018, **47:**1042-1042r.

51. Price ME, Cotton AM, Lam LL, Farre P, Emberly E, Brown CJ, Robinson WP, Kobor MS: **Additional annotation enhances potential for biologically-relevant analysis of the**

Illumina Infinium HumanMethylation450 BeadChip array. *Epigenetics Chromatin* 2013, **6:**4.

52. Deary IJ, Whiteman MC, Starr JM, Whalley LJ, Fox HC: **The impact of childhood intelligence on later life: following up the Scottish mental surveys of 1932 and 1947.** *J Pers Soc Psychol* 2004, **86:**130-147.

53. Deary IJ, Gow AJ, Taylor MD, Corley J, Brett C, Wilson V, Campbell H, Whalley LJ, Visscher PM, Porteous DJ, Starr JM: **The Lothian Birth Cohort 1936: a study to examine influences on cognitive ageing from age 11 to age 70 and beyond.** *BMC Geriatr* 2007, **7:**28.

Figure 1 Power and false positive rate for the MWAS methods in simulation scenario 1. The phenotypes were simulated based on the effects from 100 causal probes but no direct effects from the CTCs. (a) Mean genomic inflation factor from a method across 100 simulation replicates with an error bar representing +/- SE of the mean. The dashed line at 1 shows the expected value if there is no inflation. (b) Each dot represents the FPR of a method (with an error bar representing +/- the SE). The dashed line at 0.05 is the expected FPR under the null. (c) Each column represents the mean of the mean chi-squared values of 100 causal probes from a method across 100 simulation replicates with an error bar representing +/- SE of the mean. Note that chi-squared values of the causal probes (on the odd chromosomes) were recalibrated by the genome inflation factor of the null probes on the even chromosomes. (d) Each dot represents the power of a method (with an error bar representing +/- the SE), computed as the proportion of causal probes detected at a genome-wide significance level based on the recalibrated chi-squared values in 100 simulation replicates.

Figure 2 Genomic inflation factor and false positive rate for the MWAS methods in simulation scenario 2 (effects from CTCs but no causal probes). Shown on the horizontal axis are the R^2_{CTCs} values used to simulate the phenotype. (a) Each dot represents the mean λ value from 1000 simulation replicates given a specified R^2_{CTCs} value for a method with an error bar representing +/- the SE of the mean. (b) Each dot represents the false positive rate, calculated as the proportion of simulation replicates with one or more null probes detected at a genome-wide significance level.

Figure 3 QQ plot of p-values from MWAS analysis for 4 quantitative traits in the LBC data. The DNAm measures were adjusted for age, sex, and batches. The phenotypes were stratified into groups by sex and cohort and were adjusted for age and standardised to z-scores by rank-based inverse normal transformation in each group. The phenotypes are (a) BMI, (b) height (c) lung function, and (d) walking speed.

Table 1 Genomic inflation factor and genome-wide significant associations reported by different MWAS methods for the 4 traits in the Lothian Birth Cohorts.

	BMI		Height		Lung function		Walking speed	
	λ	Number of associations	λ	Number of associations	λ	Number of associations	λ	Number of associations
Unadjusted	1.68	14	1.30	0	0.98	13	1.28	1
5PCs	1.11	2	0.96	0	1.06	17	1.04	1
SVA	1.04	1	0.95	0	1.06	6	1.01	0
LFMM2-ridge	1.09	2	1.00	0	1.10	20	1.04	0
LFMM2-lasso	1.08	2	0.99	0	1.09	20	1.03	0
ReFACTor	1.13	2	0.97	0	1.09	14	1.02	0
EWASher	1.11	3	0.96	0	1.09	2	0.96	1
MOA	0.99	2	1.05	0	0.97	14	0.99	1
MOMENT	1.00	1	1.05	0	0.98	0	1.00	1