Hydrogenotrophic methanogens of the mammalian gut: functionally similar, thermodynamically different - A modelling approach

Rafael Muñoz-Tamayo[1*,¶], Milka Popova[2, ¶], Maxence Tillier[2], Diego P. Morgavi [2], Jean-Pierre

Morel [3], Gérard Fonty [3], Nicole Morel-Desrosiers[3]

[1]UMR Modélisation Systémique Appliquée aux Ruminants, INRA, AgroParisTech, Université Paris-

Saclay, 75005, Paris, France

[2]Institute National de la Recherche Agronomique, UMR1213 Herbivores, Clermont Université,

VetAgro Sup, UMR Herbivores, Clermont-Ferrand, France

[3]Université Clermont Auvergne, CNRS, LMGE, F-63000 Clermont-Ferrand, France

Running title: Thermodynamic and kinetic modelling of gut methanogens

* Correspondence: Rafael Muñoz-Tamayo, UMR Modélisation Systémique Appliquée aux

Ruminants, INRA, AgroParisTech, 16 Rue Claude Bernard, Paris, 75005, France. Tel: 00 33

144081759. E-mail : Rafael.Munoz-Tamayo@inra.fr

14
15 ¶ These authors contributed equally to this work

Abstract

Methanogenic archaea occupy a functionally important niche in the gut microbial ecosystem of mammals. Our purpose was to quantitatively characterize the dynamics of methanogenesis by integrating microbiology, thermodynamics and mathematical modelling. For that, *in vitro* growth experiments were performed with pure cultures of key methanogens from the human and ruminant gut, namely *Methanobrevibacter smithii*, *Methanobrevibacter ruminantium* and *Methanobacterium formicium*. Microcalorimetric experiments were performed to quantify the methanogenesis heat flux. We constructed an energetic-based mathematical model of methanogenesis. Our model captured efficiently the dynamics of methanogenesis with concordance correlation coefficients of 0.94 for CO_2, 0.99 for H_2 and 0.97 for CH_4. Together, experimental data and model enabled us to quantify metabolism kinetics and energetic patterns that were specific and distinct for each species despite their use of analogous methane-producing pathways. Then, we tested *in silico* the interactions between these methanogens under an *in vivo* simulation scenario using a theoretical modelling exercise. *In silico* simulations suggest that the classical competitive exclusion principle is inapplicable to gut ecosystems and that kinetic information alone cannot explain gut ecological aspects such as microbial coexistence. We suggest that ecological models of gut ecosystems require the integration of microbial kinetics with nonlinear behaviours related to spatial and temporal variations taking place in mammalian guts. Our work provides novel information on the thermodynamics and dynamics of methanogens. This understanding will be useful to construct new gut models with enhanced prediction capabilities and could have practical applications for promoting gut health in mammals and mitigating ruminant methane emissions.

Introduction

Methanogenic archaea inhabit the gastro-intestinal tract of mammals where they have established syntrophic interactions within the microbial community (1–3) playing a critical role in the energy balance of the host (4,5). In the human gut microbiota, the implication of methanogens in host homeostasis or diseases is poorly studied, but of growing interest (6). *Methanobrevibacter smithii* (accounting for 94% of the methanogen population) and *Methanosphaera stadtmanae* are specifically recognized by the human innate immune system and contribute to the activation of the adaptive immune response (7). Decreased abundance of *M. smithii* was reported in inflammatory bowel disease patients (8), and it has been suggested that methanogens may contribute to obesity (9). In the rumen, the methanogens community is more diverse though still dominated by *Methanobrevibacter* spp., followed by *Methanomicrobium* spp., *Methanobacterium* spp. (10) and *Methanomassillicoccus* spp (11). However, the proportion of these taxa could vary largely, with *Methanomicrobium mobile* and *Methanobacterium formicium* being reported as major methanogens in grazing cattle (12). Though methanogens in the rumen are essential for the optimal functioning of the ecosystem (by providing final electron acceptors), the methane they produce is emitted by the host animal, contributing to global greenhouse gas (GHG) emissions. The livestock sector is responsible for 14.5% of the anthropogenic GHG emissions, of which enteric methane from ruminants contributes to more than 40% of total emissions (13). Methanogens can be separated in two groups based on the presence or absence of cytochromes, which are membrane-associated electron transfer proteins providing energetic advantages for microbial growth (14). However, in the gastrointestinal tract of mammals, the only methanogens with cytochromes are the *Methanosarcinales* which are minor members of the community (4,15). Major rumen methanogens (16) and the dominant human archaeon *M. smithii* (17), are hydrogenotrophic archaea without cytochrome. Cytochrome-lacking methanogens exhibit lower growth yields than archaea with cytochromes (14). However, this apparent energetic disadvantage has been counterbalanced by a greater adaptation to the environmental conditions prevailing in the gastrointestinal tract (15), and by the establishment of syntrophic interactions with feed fermenting

microbes. This syntrophic cooperation centred on hydrogen allows the anaerobic reactions of substrate

conversion to proceed close to the thermodynamic equilibrium (18,19) (that is with Gibbs free energy

change close to zero).

To our knowledge, the impact of thermodynamics on human gut metabolism has been poorly addressed

in existing mathematical models (20–23). For the rumen, thermodynamic principles have been

incorporated already into mathematical research frameworks because of their important role in feed

fermentation, (24–29). Despite these relevant efforts, much work remains to be conducted for attaining

a predictive thermodynamic-based model that allows for quantitative assessment of the impact of the

thermodynamics on fermentation dynamics. Theoretical frameworks have been developed to calculate

stoichiometric and energetic balances of microbial growth from the specification of the anabolic and

catabolic reactions of microbial metabolism (30,31), and advances have been done to link

thermodynamics to kinetics (32–34). These works constitute a solid basis for tackling the

thermodynamic modelling of gut metabolism. In this respect, new knowledge on the extent of

methanogenesis could help to improve existing gut models. Accordingly, our purpose was to

quantitatively characterize the dynamics of hydrogen utilization, methane production, growth and heat

flux of three hydrogenotrophic gut methanogens by integrating microbiology, thermodynamics, and

mathematical modelling. We investigated the rate and extent of methanogenesis by performing *in vitro*

experiments with three methanogenic species representing major human and ruminant genera: *M.*

smithii, M. formicium and *Methanobrevibacter ruminantium*. To interpret and get the most out of the

resulting data, a mathematical model with thermodynamic basis was developed to describe the

dynamics of the methanogenesis. Our findings allowed to quantify metabolism kinetics and energetic

patterns that were specific and distinct for each species despite their use of analogous methane-

producing pathways and their common belonging to the group of cytochrome-lacking methanogenic

archaea.

Material and Methods

In vitro growth experiments

Archaeal strains and growth media

Archaeal strains used in the study were *M. ruminantium* M1 (DSM 1093)*, M. smithii* PS (type strain DSM 861)*,* and *M. formicium* MF (type strain DSM 1535). The growth media was prepared as previously described (35) and composition is summarized in Table S1 of the Supplementary material.

Experimental design and measures

Starter cultures were grown until reaching optical density at 660 nm (OD_{660}) of 0.400 ± 0.030. Optical density was measured on a Jenway spectrophotometer (Bibby Scientific). Then, 0.6 ml were used to inoculate one experimental tube. Commercially prepared high purity H_2/CO_2 (80%/20%) gas mix was added to inoculated tubes by flushing for 1 min at 2.5 Pa. Mean initial OD_{660} and pressure values are summarized in Table S2 of the Supplementary material. Growth kinetics for each strain were followed over 72 h. The experiment was repeated twice. Each kinetics study started with 40 tubes inoculated at the same time. At a given time point (3 h, 5 h, 6.5 h, 24 h, 26 h, 30 h, 47h, 72 h post inoculation), two tubes with similar OD_{660} values were sampled. The tubes were used for measuring gas parameters: pressure was measured using a manometer and composition of the gas phase was analysed by gas chromatography on a Micro GC 3000A (Agilent Technologies, France). One of the tubes was centrifuged 10 min at 13 000 g. The microbial pellet was weighed and stored at -20°C in 2 ml screw-cap tubes containing 0.4 g of sterile zirconia beads (0.3 g of 1 mm and 0.1 g of 0.5 mm).

DNA extraction and qPCR quantification of 16S rRNA genes

One ml of lysis buffer (50mM NaCl, 50 mM TrisHCl pH 7.6, 50 mM EDTA, 5 % SDS) was added

directly to the frozen microbial pellet before homogenizing for 2×30 s at 5100 tours/min in a Precellys

bead-beater (Bertin Instruments). Samples were centrifuged for 3 min at 14 000 g and the liquid phase

transferred to a new tube before adding 600 µl of phenol–chloroform–3-methyl-1-butanol (25:24:1)

solution. After centrifugation at 14 000 g for 3 min, the aqueous phase was transferred to a fresh tube

and 500 µl of chloroform were added. The chloroform-washing step was repeated twice with

centrifugation at 14000 g for 3 min between steps. The final volume of the aqueous phase was measured

and DNA precipitation was initiated by adding 70% of the volume of isopropanol 100% and 10% of

the volume of sodium acetate 3M. Sedimentation at 14 000 g for 30 min was again performed and the

resulting DNA pellet was washed with 500 µl of ethanol 70% and dissolved in 50µl of molecular

biology quality water. The extraction yield was checked on a Nanodrop 1000 Spectrophotometer

(Thermo Fisher Scientific, France) and extracts run on a FlashGel System (Lonza, Rockland, Inc) to

check integrity.

Copies of 16S rRNA genes were quantified using a qPCR approach. Primers used are those of Ohene-

Adjei et al (36) ; reaction assay and temperature cycles were as described previously (37) . Triplicate

qPCR quantification was performed on 20 ng of extracted DNA. Amplifications were carried out using

SYBR Premix Ex Taq (TaKaRa Bio Inc., Otsu, Japan) on a StepOne system (Applied Biosystems,

Courtabeuf, France). Absolute quantification involved the use of standard curves that had been

prepared with gDNA of *Methanobrevibacter ruminantium* DSM 1093. PCR efficiency was of 103%.

Results were expressed as copy numbers per ng of extracted DNA per g of microbial pellet. *M. smithii*

and *M. ruminantium* strains used in this study possess two copies of 16S rRNA genes in their genomes.

The number of cells was computed by dividing 16S copy numbers by 2.

Microcalorimetry

Microcalorimetric experiments were performed to determine the heat flux pattern of each methanogen.

Metabolic activity and microbial growth were monitored by using isothermal calorimeters of the heat-

conduction type (A TAM III, TA Instruments, France) equipped with two multicalorimeters, each

holding six independent minicalorimeters, allowed continuous and simultaneous recording as a

function of time of the heat flux produced by 12 samples. The bath temperature was set at 39°C; its

long-term stability was better than ± 1x10^{-4}°C over 24h. Each minicalorimeter was electrically

calibrated. The specific disposable 4 mL microcalorimetric glass ampoules capped with butyl rubber

stoppers and sealed with aluminium crimps were filled with 1.75 mL of Balch growth media and

overpressed with 2.5 Pa of H_2/CO_2 80%/20% gas mixture for 30 s. There was no significant difference

in pressure at the beginning of the study. They were sterilized by autoclave and stored at 39°C until the

beginning of the microcalorimetric measurements. Actively growing cultures of methanogens (OD_{660}

of 0.280±0.030 for *M. smithii*, 0.271±0.078 for *M. ruminantium* and 0.142±0.042 for *M. formicium)*

were stored at -20°C in order to diminish microbial activity before inoculation. Cultures were thawed

for 30 min at ambient temperature and inoculation was carried out by injecting 0.25 mL of the culture

through the septum of the overpressed microcalorimetric ampoules just before inserting them into the

minicalorimeters. Samples took about two hours to reach the bath temperature and yield a stable zero

baseline. Blank experiments were also carried out by inserting ampoules that were not inoculated and,

as expected, no heat flux was observed confirming the medium sterility. Each experiment was repeated

thrice.

The heat flux $\left(\frac{dQ}{dt}\right)$, also called thermal power output P, was measured for each methanogen and blank

samples with a precision ≥ 0.2 μW. The heat flux data of each sample were collected every 5 minutes

during more than 10 days. The total heat Q was obtained by integrating the overall heat flux time curve

using the TAM Assistant Software and its integrating function (TA Instruments, France).

Classically, the heat flux-time curve for a growing culture starts like the S-shaped biomass curve (a lag

phase followed by an exponential growth phase) but differs beyond the growth phase, the heat flux

being then modulated by transition periods (38). Heat flux data can be used to infer the microbial

growth rate constant provided the existence of a correlation between isothermal microcalorimetry data

and microbiological data (e.g., cell counts) at early growth (39). During the exponential growth phase,

microbial growth follows a first-order kinetics defined by the specific growth rate constant μ_c (h^{-1}).

Analogously, the heat flux follows an exponential behaviour determined by the parameter μ_c as

described by (38,39).

$$\frac{dQ}{dt} = \mu_c \cdot Q \tag{1}$$

167   The growth rate constant μ_c can be determined by fitting the exponential part of the heat flux-time

168   curve using the fitting function of the TAM Assistant Software. In our case study, careful selection of

169   the exponential phase of heat flux dynamics was performed to provide a reliable estimation of the

170   maximum growth rate constant from calorimetric data.

Mathematical model development

Modelling *in vitro* methanogenesis

The process of *in vitro* methanogenesis is depicted in Figure 1. The H_2/CO_2 mixture in the gas phase

diffuses to the liquid phase. The H_2 and CO_2 in the liquid phase are further utilized by the pure culture

to produce CH_4. Methane in the liquid phase diffuses to the gas phase.

Model construction was inspired by our previous dynamic models of human gut (20) and rumen *in*

vitro fermentation (40) followed by certain simplifications. The model considers the liquid-gas transfer

of carbon dioxide. Due to the low solubility of hydrogen and methane (41), the concentration of these

two gases in the liquid phase was not modelled. We assumed that the dynamics of concentrations in

the gas phase are determined by kinetic rate of the methanogenesis. To incorporate thermodynamic

information, instead of using the Monod equation in the original formulation, we used the kinetic rate

function proposed by Desmond-Le Quéméner and Bouchez (33). The resulting model is described by

the following ordinary differential equations

184
$$\frac{dx_{H_2}}{dt} = \mu_{max} \cdot \exp\left(-\frac{K_s \cdot V_g}{n_{g,H_2}}\right) \cdot x_{H_2} - k_d \cdot x_{H_2} \tag{2}$$

185
$$\frac{ds_{CO_2}}{dt} = -\frac{-Y_{CO_2} \cdot \mu_{max}}{Y} \cdot \exp\left(-\frac{K_s \cdot V_g}{n_{g,H_2}}\right) \cdot x_{H_2} - k_L a \cdot \left(s_{CO_2} - K_{H,CO_2} \cdot R \cdot T \cdot n_{g,CO_2}/V_g\right) \tag{3}$$

186
$$\frac{dn_{g,H_2}}{dt} = -\frac{\mu_{max}}{Y} \cdot \exp\left(-\frac{K_s \cdot V_g}{n_{g,H_2}}\right) \cdot V_L \cdot x_{H_2} \tag{4}$$

187
$$\frac{dn_{g,CO_2}}{dt} = V_L \cdot k_L a \cdot \left(s_{CO_2} - K_{H,CO_2} \cdot R \cdot T \cdot n_{g,CO_2}/V_g\right) \tag{5}$$

188
$$\frac{dn_{g,CH_4}}{dt} = \frac{Y_{CH_4} \cdot \mu_{max}}{Y} \cdot \exp\left(-\frac{K_s \cdot V_g}{n_{g,H_2}}\right) \cdot V_L \cdot x_{H_2} \tag{6}$$

where s_{CO_2} is the concentration (mol/L) of carbon dioxide in the liquid phase and x_{H_2} is the biomass

concentration (mol/L) of hydrogenotrophic methanogens. The numbers of moles in the gas phase are

represented by the variables $n_{g,H_2}, n_{g,CO_2}, n_{g,CH_4}$. The gas phase volume $V_g = 20$ mL and the liquid

phase volume $V_L = 6$ mL. Liquid-gas transfer for carbon dioxide is described by a non-equilibria

transfer rate which is driven by the gradient of the concentration of the gases in the liquid and gas

phase. The transfer rate is determined by the mass transfer coefficient $k_L a$ (h^{-1}) and the Henry's law

coefficients K_{H,CO_2} (M/bar). R (bar·(M · K)$^{-1}$) is the ideal gas law constant and T is the temperature (K).

Microbial decay is represented by a first-order kinetic rate with k_d (h^{-1}) the death cell rate constant.

Microbial growth was represented by the rate function proposed by Desmond-Le Quéméner and

Bouchez (33) using hydrogen as the limiting reactant

199
$$\mu = \mu_{\max} \cdot \exp\left(-\frac{K_s \cdot V_g}{n_{g,H_2}}\right) \tag{7}$$

200   where μ is the growth rate (h^{-1}), $\mu_{\max}$ (h^{-1}) is the maximum specific growth rate constant and K_s(mol/L)

201   the affinity constant. Equation (7) is derived from energetic principles following Boltzmann statistics

202   and uses the concept of exergy (maximum work available for a microorganism during a chemical

203   transformation). The affinity constant has an energetic interpretation since it is defined as

204
$$K_s = \frac{E_M + E_{dis}}{v_{harv} \cdot E_{cat}} \tag{8}$$

where E_{dis} (kJ/mol) and E_M (kJ/mol) are the dissipated exergy and stored exergy during growth

respectively. E_{cat} (kJ/mol) is the catabolic exergy of one molecule of energy-limiting substrate, and

v_{harv} is the volume at which the microbe can harvest the chemical energy in the form of substrate

molecules (33). E_{cat} is the absolute value of the Gibbs energy of catabolism ($\Delta G_{r,c}$) when the reaction

is exergonic ($\Delta G_{r,c}<0$) or zero otherwise. The stored exergy E_M is calculated from a reaction (destock)

representing the situation where the microbe gets the energy by consuming its own biomass. E_M is the

absolute value of the Gibbs energy of biomass consuming reaction ($\Delta G_{r,destock}$) when the reaction is

exergonic ($\Delta G_{r,destock}<0$) or zero otherwise. Finally, the dissipated exergy E_{dis} is the opposite of the

Gibbs energy of the overall metabolic reaction, which is a linear combination of the catabolic and

destock reactions. This calculation follows the Gibbs energy dissipation detailed in Kleerebezem and

Van Loosdrecht (31).

In our model, the stoichiometry of methanogenesis is represented macroscopically by one catabolic

reaction (R$_1$) for methane production and one anabolic reaction (R$_2$) for microbial formation. We

assumed that ammonia is the only nitrogen source for microbial formation. The molecular formula of

microbial biomass was assumed to be $C_5H_7O_2N$ (41).

R_1: 4 H_2 + CO_2 → CH_4 + 2 H_2O

R_2: 10 H_2 + 5 CO_2 + NH_3 → $C_5H_7O_2N$ + 8 H_2O

In the model, the stoichiometry of the reactions is taken into account *via* the parameters Y, Y_{CO_2}, Y_{CH_4},

which are the yield factors (mol/mol) of microbial biomass, CO_2 and CH_4. The microbial yield factor

Y was extracted from literature. We assumed that *M. smithii* and *M. ruminantium* have the same yield

(being both Methanobrevibacter). This yield factor was set to 0.006 mol biomass/mol H_2, using the

methane-based molar growth yield of 2.8 g biomass/mol CH_4 estimated for *M. smithii* (42) and the

equations (9) and (11). Similarly, the yield factor for *M. formicium* was set to 0.007 mol biomass/mol

H_2 using the methane-based molar growth yield of 3.5 g biomass/mol CH_4 reported by Schauer and

Ferry (43). The fraction of H_2 utilized for microbial growth (reaction R_2) is defined by the yield factor

Y (mol of microbial biomass/mol of H_2). Now, let f be the fraction of H_2 used for the catabolic reaction

R_1. Reaction R_2 tells us that for every 10 moles of H_2 used in R_2, we get 1 mol of microbial biomass.

Hence, it follows that

$$Y = \frac{1}{10} \cdot (1 - f) \tag{9}$$

If Y is known, the fraction f can be computed from Equation (9).

The yield factors of CO_2 and CH_4 can be expressed as functions of the the fraction f:

$$Y_{CO_2} = \left(\frac{1}{4}\right) \cdot f + \left(\frac{5}{10}\right) \cdot (1 - f) \tag{10}$$

$$Y_{CH_4} = \left(\frac{1}{4}\right) \cdot f \tag{11}$$

The model has two physicochemical parameters $(k_L a, K_{H,CO_2})$ and four biological parameters

(μ_{max}, K_s, Y, k_d). The initial condition for s_{CO_2} is unknown and was also included in the parameter

vector for estimation. The Henry's law coefficients are known values calculated at 39°C using the

equations provided by Batstone et al. (41). If the article is accepted, an implementation of the model in

the Open Source software Scilab (https://www.scilab.org/) will be made available at Zenodo

(https://doi.org/10.5281/zenodo.3271611).

Theoretical model to study interactions among methanogens

The experimental study of microbial interactions requires sophisticated *in vitro* systems under

continuous operation such as the one developed by Haydock *et al.* (44). In our work, we explored by

means of mathematical modelling how the methanogens can interact under *in vivo* conditions. For this

theoretical study, we elaborated a toy model based on the previous model for *in vitro* methanogenesis.

Let us consider the following simple model for representing the consumption of hydrogen by the

methanogenic species i under an *in vivo* scenario of continuous flow

253
$$\frac{dx_{H_2,i}}{dt} = \mu_{max,i} \cdot \exp\left(-\frac{K_{s,i} \cdot V_g}{n_{g,H_2}}\right) \cdot x_{H_2;i} - D_i \cdot x_{H_2,i} \tag{12}$$

254
$$\frac{dn_{g,H_2}}{dt} = q_{H_2} - \frac{\mu_{max,i}}{Y_i} \cdot \exp\left(-\frac{K_{s,i} \cdot V_g}{n_{g,H_2}}\right) \cdot V_L \cdot x_{H_2,i} - b \cdot n_{g,H_2} \tag{13}$$

255   where q_{H_2} (mol/h) is the flux of hydrogen produced from the fermentation of carbohydrates. The kinetic

256   parameters are specific to the species i ($x_{H_2,i}$). The parameter D_i (h^{-1}) is the dilution rate of the

257   methanogens and b (h^{-1}) is an output substrate rate constant. Extending the model to n species with a

258   common yield factor Y, the dynamics of hydrogen is given by

259
$$\frac{dn_{g,H_2}}{dt} = q_{H_2} - \frac{V_L}{Y} \sum_{i=1}^{n} \mu_{max,i} \cdot \exp\left(-\frac{K_{s,i} \cdot V_g}{n_{g,H_2}}\right) \cdot x_{H_2,i} - b \cdot n_{g,H_2} \tag{14}$$

260   where the sub index i indicates the species. In our case study, $n = 3$.

Parameter identification

Before tackling the numerical estimation of the model parameters, we addressed the question of

whether it was theoretically possible to determine uniquely the model parameters given the available

measurements from the experimental setup. This question is referred to as structural identifiability (45).

Structural identifiability analysis is of particular relevance for model whose parameters are biologically

meaningful, since knowing the actual value of the parameter is useful for providing biological insight

on the system under study (46). Moreover, in our case, we are interested in finding accurate estimates

that can be further used as priors in an extended model describing the *in vivo* system.

We used the freely available software DAISY (47) to assess the structural identifiability of our model.

Physical parameters $(k_\text{L}\text{a}, K_{\text{H,CO}_2})$ were set to be known. The model was found to be structurally

globally identifiable. In practice, however, to facilitate the actual identification of parameters and

reduce practical identifiability problems such as high correlation between the parameters (48), we fixed

some model parameters to values reported in the literature. The transport coefficient $k_\text{L}\text{a}$, the Henry's

law coefficient $K_{\text{H,CO}_2}$, and the dead cell rate constant k_d were set to be known and were extracted from

Batstone et al. (41). Therefore, only the parameters $\mu_{\max}, K_\text{s}$ and initial condition of s_{CO_2} were set to

be estimated. To capitalize on the calorimetric data, we further assumed that $\mu_{\max}$ was equal to the

specific rate constant μ_c estimated from the heat flux-time curve. By this, only the affinity constant for

each strain and the initial condition of s_{CO_2} were left to be estimated.

The parameter identification for each methanogen was performed with the IDEAS Matlab® toolbox

(49), which is freely available at http://genome.jouy.inra.fr/logiciels/IDEAS. The measured variables

are the number of moles in the gas phase (H_2, CH_4, CO_2). The Lin's concordance correlation coefficient

(CCC) (50) was computed to quantify the agreement between the observations and model predictions.

Results

Methanogens biomass

285   Archaea-specific primers targeting the 16S rRNA gene were used to enumerate microbial cells in each

286   pure culture. Three hours post inoculation microbial numbers varied from 7.62×10^7 to 2.81×10^8 and

287   reached 10^9 after 72 hours of incubation. Table S3 summarizes microbial numbers at different sampling

288   times.

Calorimetric pattern of methanogens

Figure 2 displays a representative isothermal calorimetric curve for each methanogen. The three

measured heat flux dynamics of each methanogen were found to follow similar energetic patterns. *M.*

smithii and *M. formicium* exhibited a lag phase of a few hours, while *M. ruminantium* was already

metabolically active when introduced into the minicalorimeter though several attempts were made to

obtain a lag phase by changing storage conditions and thawing the culture just before inoculating the

microcalorimetric ampoules. The pattern of heat flux for all tested methanogens is characterized by

one predominant peak which was observed at different times for each methanogen. *M. smithii* exhibited

a second metabolic event occurring at 60 h with an increase of heat flux. The same phenomenon was

observed for *M. formicium* but at a lower intensity that started at 140 h. One possible explanation for

this event is cell lysis (39). The process was considered completed when the heat flux ceased marking

the end of the metabolic activity. It is noted that *M. formicium* produced a small peak at 14 h (Fig. 2).

A similar peak, but of much smaller size, was observed on the other curves obtained with this

methanogen. *M. smithii* also exhibits a small peak (occurrence of 3 out of 3) at 7.4 h shown in the inset

of Figure 2. For *M. ruminantium*, we do not know whether the small peak exists since the initial part

of the curve is missing. This small peak translates in a metabolic activity that remains to be elucidated.

The total heat (Q_m) produced during the methanogenesis process that took place under the present

experimental conditions was, on average, -5.5 ± 0.5 J for the three methanogens (for *M. ruminantium*,

the missing initial part of the heat flux-time curve was approximately estimated by extrapolating the

exponential fit). As we shall see below, this experimental value is consistent with the theoretically

expected value.

Estimation of thermodynamic properties

We defined two macroscopic reactions to represent the catabolism (R1) and anabolism (R2) of the

methanogenesis (see Modelling in vitro methanogenesis section). All thermodynamic properties result

from the contribution of both catabolic and anabolic reactions. In the Supplementary Material, the

calculations of total heat (Q_m), enthalpy (ΔH_m), Gibbs energy (ΔG_m) and entropy (ΔS_m) of the

methanogenesis are detailed. The estimated overall heat produced during the methanogenesis process

under our experimental conditions was in average $Q_m = -5.66$ J. This heat results from the sum of

the heat of the catabolic reaction (Q_c) and the heat of the anabolic reaction (Q_a). From the total heat

of the methanogenesis, the anabolic reaction contributes to 7% of the metabolic heat for *M. smithii* and

M. ruminantium. For *M. formicium*, the contribution of the anabolic reaction to the metabolic heat is

9%. It is also interesting to note that there is a very good agreement between the theoretical value

calculated above and the overall heat experimentally determined by microcalorimetry (-5.5 ± 0.5 J).

Table 1 shows the thermodynamic properties per mole of biomass formed during methanogenesis of

M. ruminantium, M. smithii and *M. formicium* on H_2/CO_2. These properties are compared with values

found in the literature for other methanogens grown on different substrates.

Dynamic description of *in vitro* kinetics

The developed mathematical model was calibrated with the experimental data from *in vitro* growth

experiments in Balch tubes. Table 2 shows the parameters of the dynamic kinetic model described in

Equations 2-6. The reported value of μ_{max} for each methanogen corresponds to the average value

obtained from three heat flux-time curves. From Table 2, it is concluded that *M. smithii* exhibited the

highest growth rate constant, followed by *M. ruminantium* and finally *M. formicium*. In terms of the

affinity constant K_s, while *M. smithii* and *M. ruminantium* are of the same order, the affinity constant

for *M. formicium* is lower in one order of magnitude.

Figure 3 displays the dynamics of the compounds in the methanogenesis for the three methanogens.

Experimental data are compared against the model responses. Table 3 shows standard statistics for

model evaluation. The model captures efficiently the overall dynamics of the methanogenesis.

Hydrogen and methane are very well described by the model with concordance correlation coefficients

(CCC) of 0.99 and 0.97 respectively. For carbon dioxide, CCC = 0.94.

Figure 4 displays the dynamics for the methanogens as measured by the 16S rRNA gene, as well as the

dynamics of biomass as predicted for the model. As observed, the microbes follow a typical Monod-

like trajectory.

Discussion

Our objective in this work was to quantitatively characterize the dynamics of hydrogen utilization,

methane production, growth and heat flux of three hydrogenotrophic methanogens by integrating

microbiology, thermodynamics and mathematical modelling. Our model developments were

instrumental to quantify energetic and kinetic differences between the three methanogens studied,

strengthening the potentiality of microcalorimetry as a tool for characterizing the metabolism of

microorganisms (51). This modelling work provides estimated parameters that can be used as prior

values for other modelling developments of gut microbiota.

Energetic and kinetic differences between methanogens

Methanogenesis appears as a simple reaction with a single limiting substrate (H_2). The

microcalorimetry approach we applied revealed that this simplicity is only apparent and that

hydrogenotrophic methanogens exhibit energetic and kinetic differences. Methanogenesis is indeed a

complex process that can be broken down in several stages. The dominant metabolic phase is

represented by one peak that occurs at different times. The magnitude of the peak differs between the

methanogens and also the slope of the heat flux trajectories. The return time of the heat flux to the zero

baseline was also different. The energetic difference is associated with kinetic differences that translate

into specific kinetic parameters, namely affinity constant (K_s) and maximum growth rate constant

(μ_{max}). Previously, energetic differences between methanogens have been ascribed to the presence or

absence of cytochromes (14). These differences are translated into different yield factors, H_2 thresholds,

and doubling times. The kinetic differences revealed in this study for three cytochrome-lacking

methanogens indicate that factors other than the presence of cytochromes might play a role in the

energetics of methanogenesis. Interestingly, calorimetric experiments showed that *M. ruminantium*

was metabolically active faster than the other methanogens, characteristic that could explain the

predominance of *M. ruminantium* in the rumen (52). Looking at the expression of the affinity constant

(Equation (8)), the differences between the affinity constants among the methanogens can be explained

by the differences between the by the harvest volume v_{harv} and the yield factors. Note that in the kinetic

function developed by Desmond-Le Quéméner and Bouchez (33), the maximum growth rate did not

have any dependency on the energetics of the reaction. Our experimental study revealed that μ_{max} is

species-specific and reflects the dynamics of the heat flux of the reaction at the exponential phase. This

finding suggests that a further extension of the kinetic model developed by Desmond-Le Quéméner

and Bouchez (33) should include the impact of energetics on $\mu_{\max}$. Since our study is limited to three

species, it is important to conduct further research on other methanogens to validate our findings.

Thermodynamic analysis

Regarding the energetic information for different methanogens summarized in Table 1, it is observed

that the thermodynamic behaviour of the three methanogens is analogous to that observed for

Methanobacterium thermoautotrophicum (53). The values reported in Table 1 show indeed that the

methanogenesis on H_2/CO_2 is characterized by large heat production. The growth is highly exothermic,

with a ΔH_m value that largely exceeds the values found when other energy substrates are used. The

enthalpy change ΔH_m, which is more negative than the Gibbs energy change ΔG_m, largely controls the

process. Growth on H_2/CO_2 is also characterized by a negative entropic contribution $T\Delta S_m$ which, at

first sight, may look surprising since entropy increases in most cases of anaerobic growth (54).

However, this can be understood if one remembers that $T\Delta S_m$ corresponds in fact to the balance

between the final state and the initial state of the process, that is

$$T\Delta S_m = \frac{(1-10Y)}{4Y} T\Delta S_c + T\Delta S_a = \frac{(1-10Y)}{4Y} T(S_{final} - S_{initial})_c + T(S_{final} - S_{initial})_a$$

Methanogenesis on H_2/CO_2 is particular because the final state of its catabolic reaction (1 mol CH_4 +

2 mol H_2O) involves a smaller number of moles than the initial state (4 mol H_2 + 1 mol CO_2), which

results in a significant loss of entropy during the process. For spontaneous growth in such a case, the

ΔH_m must not only contribute to the driving force but must also compensate the growth-unfavourable

$T\Delta S_m$, which means that ΔH_m must be much more negative than ΔG_m (55). For this reason,

methanogenesis on H_2/CO_2, which is accompanied by a considerable decrease of entropy and a large

production of heat, has been designed as an entropy-retarded process (53). More generally, von Stockar

and Liu (55) noticed that when the Gibbs energy of the metabolic process is resolved into its enthalpic

and entropic contributions, very different thermodynamic behaviours are observed depending on the

growth type. These thermodynamic behaviours are: aerobic respiration is clearly enthalpy-driven (ΔH_m

$<< 0$ and $T\Delta S_m > 0$), whereas fermentative metabolism is mainly entropy-driven ($\Delta H_m < 0$ and $T\Delta S_m$

$>> 0$). Methanogenesis on H_2/CO_2 is enthalpy-driven but entropy-retarded ($\Delta H_m << 0$ and $T\Delta S_m < 0$) ,

whereas methanogenesis on acetate is entropy-driven but enthalpy-retarded ($\Delta H_m > 0$ and $T\Delta S_m >> 0$).

In the present case, the highly exothermic growth of *M. ruminantium, M. smithii* and *M. formicium* on

H_2/CO_2 is largely due to the considerable decrease of entropy during the process: in fact, 50% of the

heat produced here serves only to compensate the loss of entropy. A proportion of 80% was found for

M. thermoautotrophicum (53)*,* which results from the fact that their $T\Delta S_m$ and ΔH_m values are,

respectively, 2.7 and 1.7 times larger than ours. This difference might be due to the differences in the

temperature of the studies , namely 39°C in our study vs 60°C in the study by Schill et al. (53).

Do our results inform on ecological questions such as species coexistence?

The competitive exclusion principle (56) states that coexistence cannot occur between species that

occupy the same niche (the same function). Only the most competitive species will survive. Recently,

by using thermodynamic principles, Großkopf & Soyer (34) demonstrated theoretically that species

utilizing the same substrate and producing different compounds can coexist by the action of

thermodynamic driving forces. Since in our study the three methanogens perform the same metabolic

reactions, the thermodynamic framework developed Großkopf & Soyer (34) predicts, as the original

exclusion principle (56), the survival of only one species. By incorporating thermodynamic control on

microbial growth kinetics, Lynch et al (57) showed theoretically that differentiation of ATP yields can

explain ecological differentiation of methanogens over a range of liquid turnover rates. This theoretical

work predicts that for a fixed liquid turnover rate, only one species survives. For the continuous culture

of microorganisms, it has been demonstrated that at the equilibrium (growth rate equals the dilution

rate) with constant dilution rates and substrate input rates, the species that has the lowest limiting

substrate concentration wins the competition. From Eq. (12), the number of moles of hydrogen of the

species $n^*_{g,H_2,i}$ at the steady state is

421
$$n^*_{g,H_2,i} = \frac{K_{s,i} \cdot V_g}{\log(\mu_{max,i}/D_i)}$$

Using the model parameters of Table 2, we studied *in silico* three possible competition scenarios,

assuming a constant environment (constant dilution rate D). Two dilution rates were evaluated: $D =$

0.021 h^{-1} (retention time = 48 h) and $D =$ 0.04 h^{-1} (retention time = 25 h). A retention time of 48 h

corresponds to values measured in small ruminants (58) and to humans as we used in our gut model

(20). For higher retention times, the results obtained for 48 h hold. For $D =$ 0.021 h^{-1}, we obtained that

$n^*_{g,H_2,Ms} = 0.32$ mmol, $n^*_{g,H_2,Mr} = 0.68$ mmol, $n^*_{g,H_2,Mf} = 0.28$ mmol, where the subindex Ms, Mr, Mf

stand for *M. smithii*, *M. ruminantium* and *M.formicium*. From these results, it appears that under a

constant environment, *M. formicium* will win the competition. Since $n^*_{g,H_2,Ms} < n^*_{g,H_2,Mr}$, *M.*

ruminantium will be extinguished before *M. smithii*. For $D =$ 0.04 h^{-1}, we obtained that $n^*_{g,H_2,Ms} = 0.49$

mmol, $n^*_{g,H_2,Mr} =$ 1.42 mmol, $n^*_{g,H_2,Mf} =$ 1.57 mmol, and thus *M. smithii* wins the competition. To win

the competition, *M. ruminantium* requires longer retention times than its competitors. Retention times

of digesta longer than 48 h are physiologically uncommon, thus the presence of *M. ruminantium* in the

gut ecosystem can be explained, for example, from known adhesion properties (both *M. ruminantium*

and *M. smithii* genes encode adhesin-like proteins (59,60). To illustrate these aspects, we built a

multiple-species model with the three methanogens using Eq. (12) and Eq. (14). The parameter b was

set to 0.5 h^{-1} and the hydrogen flux production q_{H_2} rate was set to 0.02 mol/min. Figure 5A displays

the dynamics of the three methanogens for the first scenario (D= 0.021 h^{-1}). It is observed that at 50 d

only *M. formicium* survives. This result, however, is not representative of what occurs in the rumen

where the three methanogens coexist (5,61). It is intriguing that in our toy model it is *M. formicium*

that wins the competition, bearing in mind that *M. ruminantium* and *M. smithii* are more abundant

than *M. formicium* (5,52). Figure 5 shows that selective conditions favour the survival of one species.

Similar results can be obtained for the human gut by including the effect of pH on microbial growth

(23) and setting the gut pH to select one of the species. On the basis of the competitive exclusion

principle, it is thus intriguing that having a very specialized function, methanogens are a diverse group

that coexist. Gut ecosystems, therefore, exhibit the paradox of the plankton introduced by Hutchinson

(1961) that presents the coexistence of species all competing for the same substrate in a relatively

isotropic or unstructured environment (62) . In the case of the rumen, our modelling work suggests that

in addition to kinetic and thermodynamic factors, other forces contribute to the ecological shaping of

the methanogens community in the rumen favouring the microbial diversity. Indeed, methanogenic

diversity in the rumen results from multiple factors that include pH sensitivity, the association with

rumen fractions (fluid and particulate material), and the endosymbiosis with rumen protozoa (5,52).

For the human gut, ecological factors enable methanogens to coexist to a competitive environment

where hydrogenotrophic microbes (acetogens, methanogenic archaea and sulfate-reducing bacteria)

utilize H$_2$ *via* different pathways (63–65). Both in the human gut and in the rumen, microbes grow in

association with biofilms that form a polymer-based matrix that provides nutritional and hydraulic

advantages for microbial growth and resistance to shear forces (20,66). Indeed, in our modelling work

of human gut fermentation (20), we suggested that, from the different actions the mucus has on colonic

fermentation, the mechanism of promoting conditions for microbial aggregation appears as the most

relevant factor for attaining the high microbial density and the high level of fibre degradation

characteristic of the human gut. Altogether, these factors result in nonlinear behaviours, spatial and

temporal variations that promote coexistence and diversity, that, as discussed in dedicated literature on

microbial ecology (67–71), render the classical formulation of the competitive exclusion principle

(56,72) inapplicable to gut ecosystems.

Finally, mathematical modelling is expected to enhance our understanding of gut ecosystems (66,73).

It is then key that in addition to metabolic aspects, mathematical models of gut fermentation incorporate

the multiple aspects that shape microbial dynamics to provide accurate predictions and improve insight

on gut metabolism dynamics and its potential modulation. For ruminants, the development of precision

livestock technologies provides promising alternatives for integrating real-time data of key animal

phenotypes such as feeding behaviour with mathematical models for estimating methane emissions

(74) and rumen function indicators at large scale. These tools will be instrumental to support livestock

management decisions and guide timely interventions. Similarly, for humans, mathematical models

coupled with electronic technologies for online monitoring of gut function (75) might facilitate the

diagnosis and the design of personalized therapies for gastrointestinal diseases.

Acknowledgements

477 We are grateful to Dominique Graviou (UMRH, Inra) for her skilled assistance on the *in vitro* growth

478 experiments and qPCR assays. We thank the Inra PHASE department and the Inra MEM

479 metaprogramme for financial support.

Conflict of interest

482 No conflict.

References

References

1.   Miller TL, Wolin MJ, Hongxue Z, Bryant MP. Characteristics of methanogens isolated from

bovine rumen. Appl Environ Microbiol. 1986;51:201–2.

2.   Dridi B, Fardeau ML, Ollivier B, Raoult D, Drancourt M. Methanomassiliicoccus luminyensis

gen. nov., sp. nov., a methanogenic archaeon isolated from human faeces. Int J Syst Evol

Microbiol. 2012;62:1902–7.

3.   Paul K, Nonoh JO, Mikulski L, Brune A. "Methanoplasmatales," thermoplasmatales-related

archaea in termite guts and other environments, are the seventh order of methanogens. Appl

Environ Microbiol. 2012;78:8245–53.

4.   Dridi B, Raoult D, Drancourt M. Archaea as emerging organisms in complex human

microbiomes. Anaerobe. 2011;17:56–63.

5.   Carberry CA, Waters SM, Kenny DA, Creevey CJ. Rumen methanogenic genotypes differ in

abundance according to host residual feed intake phenotype and diet type. Appl Env

Microbiol. 2014;80:586–94.

6.   Borrel G, McCann A, Deane J, Neto MC, Lynch DB, Brugère JF, et al. Genomics and

metagenomics of trimethylamine-utilizing Archaea in the human gut microbiome. ISME J.

2017;11:2059–74.

7.   Bang C, Weidenbach K, Gutsmann T, Heine H, Schmitz RA. The intestinal archaea

Methanosphaera stadtmanae and Methanobrevibacter smithii activate human dendritic cells.

PLoS One. 2014;9:e99411.

8.   Ghavami SB, Rostami E, Sephay AA, Shahrokh S, Balaii H, Aghdaei HA, et al. Alterations of

the human gut Methanobrevibacter smithii as a biomarker for inflammatory bowel diseases.

Microb Pathog. 2018;117:285–9.

9.   Mathur R, Barlow GM. Obesity and the microbiome. Vol. 9, Expert Review of

Gastroenterology and Hepatology. 2015. p. 1087–99.

10.   Hook SE, Wright A-DG, McBride BW. Methanogens: methane producers of the rumen and

mitigation strategies. Archaea. 2010;945785.

11.   Poulsen M, Schwab C, Jensen BB, Engberg RM, Spang A, Canibe N, et al. Methylotrophic

methanogenic Thermoplasmata implicated in reduced methane emissions from bovine rumen.

Nat Commun. 2013;4:1428.

12.   Jarvis GN, Strömpl C, Burgess DM, Skillman LC, Moore ERB, Joblin KN. Isolation and

identification of ruminal methanogens from grazing cattle. Curr Microbiol. 2000;40:327–32.

13.   Gerber P, Steinfeld H, Henderson B, Mottet A, Opio C, Dijkman J, et al. Tackling Climate

Change through Livestock. Food and Agriculture Organization. 2013.

14.   Thauer RK, Kaster AK, Seedorf H, Buckel W, Hedderich R. Methanogenic archaea:

Ecologically relevant differences in energy conservation. Nat Rev Microbiol. 2008;6:579–91.

15.   Morgavi DP, Forano E, Martin C, Newbold CJ. Microbial ecosystem and methanogenesis in

ruminants. Animal. 2010;4:1024–36.

16.   Friedman N, Jami E, Mizrahi I. Compositional and functional dynamics of the bovine rumen

methanogenic community across different developmental stages. Environ Microbiol.

2017;19:3365–73.

17.   Hansen EE, Lozupone CA, Rey FE, Wu M, Guruge JL, Narra A, et al. Pan-genome of the

dominant human gut-associated archaeon, Methanobrevibacter smithii, studied in twins. Proc

Natl Acad Sci. 2011;108(Supplement_1):4599–606.

18.   Jackson BE, McInerney MJ. Anaerobic microbial metabolism can proceed close to

thermodynamic limits. Nature. 2002;415:454–6.

19.   Gonzalez-Cabaleiro R, Lema JM, Rodriguez J, Kleerebezem R. Linking thermodynamics and

kinetics to assess pathway reversibility in anaerobic bioprocesses. Energy Environ Sci.

2013;6:3780–9.

20.   Muñoz-Tamayo R, Laroche B, Walter E, Doré J, Leclerc M. Mathematical modelling of

carbohydrate degradation by human colonic microbiota. J Theor Biol. 2010;266:189–201.

21.   Van Wey AS, Lovatt SJ, Roy NC, Shorten PR. Determination of potential metabolic pathways

of human intestinal bacteria by modeling growth kinetics from cross-feeding dynamics. Food

Res Int. 2016;88:207–16.

22.   Shoaie S, Ghaffari P, Kovatcheva-Datchary P, Mardinoglu A, Sen P, Pujos-Guillot E, et al.

Quantifying Diet-Induced Metabolic Changes of the Human Gut Microbiome. Cell Metab.

2015;22:320–31.

23.   Kettle H, Louis P, Holtrop G, Duncan SH, Flint HJ. Modelling the emergent dynamics and

major metabolites of the human colonic microbiota. Environ Microbiol. 2015;17:1615–30.

24.   Janssen PH. Influence of hydrogen on rumen methane formation and fermentation balances

through microbial growth kinetics and fermentation thermodynamics. Anim Feed Sci Technol.

2010;160:1–22.

25.   Kohn R a, Boston RC. The Role of Thermodynamics in Controlling Rumen Metabolism.

546   Model Nutr Util Farm Anim. 2000;11–24.

547   26.   Offner A, Sauvant D. Thermodynamic modeling of ruminal fermentations. 2006;55:343–65.

548   27.   Ungerfeld EM. A theoretical comparison between two ruminal electron sinks. Front Microbiol.

549   2013;4.

550   28.   Van Lingen HJ, Plugge CM, Fadel JG, Kebreab E, Bannink A, Dijkstra J. Thermodynamic

551   driving force of hydrogen on rumen microbial metabolism: A theoretical investigation. PLoS

552   One. 2016;11:e0161362.

553   29.   Ghimire S, Gregorini P, Hanigan MD. Evaluation of predictions of volatile fatty acid

554   production rates by the Molly cow model. J Dairy Sci. 2014;97:354–62.

555   30.   Heijnen JJ, Dijken JP. In search of a thermodynamic description of biomass yields for the

556   chemotrophic growth of microorgansims. Biotechnol Bioengeneering. 1992;39:833–52.

557   31.   Kleerebezem R, Van Loosdrecht MCM. A Generalized Method for Thermodynamic State

558   Analysis of Environmental Systems. Crit Rev Environ Sci Technol. 2010;40:1–54.

559   32.   Hoh CY, Cord-Ruwisch R. A practical kinetic model that considers endproduct inhibition in

560   anaerobic digestion processes by including the equilibrium constant. Biotechnol Bioeng.

561   1996;51:597–604.

562   33.   Desmond-Le Quemener E, Bouchez T. A thermodynamic theory of microbial growth. Isme J.

563   2014;8:1747–51.

34.   Großkopf T, Soyer OS. Microbial diversity arising from thermodynamic constraints. ISME J.

2016;10:2725–33.

35.   Wolfe RS. Techniques for cultivating methanogens. Methods Enzymol. 2011;494:1–22.

36.   Ohene-Adjei S, Teather RM, Ivan M, Forster RJ. Postinoculation protozoan establishment and

association patterns of methanogenic archaea in the ovine rumen. Appl Environ Microbiol.

2007;73(14):4609–18.

37.   Popova M, Martin C, Eugene M, Mialon MM, Doreau M, Morgavi DP. Effect of fibre- and

starch-rich finishing diets on methanogenic Archaea diversity and activity in the rumen of

feedlot bulls. Anim Feed Sci Technol. 2011;166–67:113–21.

38.   Bricheux G, Bonnet JL, Bohatier J, Morel JP, Morel-Desrosiers N. Microcalorimetry: a

powerful and original tool for tracking the toxicity of a xenobiotic on Tetrahymena pyriformis.

Ecotoxicol Env Saf. 2013;98:88–94.

39.   Braissant O, Bonkat G, Wirz D, Bachmann A. Microbial growth and isothermal

microcalorimetry: Growth models and their application to microcalorimetric data.

Thermochim Acta. 2013;555:64–71.

40.   Muñoz-Tamayo R, Giger-Reverdin S, Sauvant D. Mechanistic modelling of in vitro

fermentation and methane production by rumen microbiota. Anim Feed Sci Technol.

2016;220:1–21.

41.   Batstone DJ, Keller J, Angelidaki I, Kalyuzhnyi S V, Pavlostathis SG, Rozzi A, et al.

Anaerobic Digestion Model No.1 (ADM1). IWA Task Group for Mathematical Modelling of

Anaerobic Digestion Processes. IWA Publishing, London; 2002.

42.   Pavlostathis SG, Miller TL, Wolin MJ. Cellulose Fermentation by Continuous Cultures of

Ruminococcus-Albus and Methanobrevibacter-Smithii. Appl Microbiol Biotechnol.

1990;33(1):109–16.

43.   Schauer NL, Ferry JG. Metabolism of formate in Methanobacterium formicicum. J Bacteriol.

1980;142:800–7.

44.   Haydock AK, Porat I, Whitman WB, Leigh JA. Continuous culture of Methanococcus

maripaludis under defined nutrient conditions. FEMS Microbiol Lett. 2004;

45.   Walter E, Pronzato L. Identification of Parametric Models from Experimental Data. Springer,

London; 1997.

46.   Muñoz-Tamayo R, Puillet L, Daniel JB, Sauvant D, Martin O, Taghipoor M, et al. Review: To

be or not to be an identifiable model. Is this a relevant question in animal science modelling?

Animal. 2018;12:701–12.

47.   Bellu G, Saccomani MP, Audoly S, D'Angio L. DAISY: A new software tool to test global

identifiability of biological and physiological systems. Comput Methods Programs Biomed.

2007;88:52–61.

48.   Vanrolleghem PA, Vandaele M, Dochain D. Practical identifiability of a biokinetic model of

activated-sludge respiration. Water Res. 1995;29:2561–70.

49.   Muñoz-Tamayo R, Laroche B, Leclerc M, Walter E. IDEAS: A parameter identification

toolbox with symbolic analysis of uncertainty and its application to biological modelling. In:

IFAC Proceedings Volumes. 2009. p. 1271–6.

50.   Lin LI. A concordance correlation-coefficient to evaluate reproducibility. Biometrics.

1989;45:255–68.

51.   Ruiz T, Bec A, Danger M, Koussoroplis A, Aguer J, Morel J, et al. A microcalorimetric

608   approach for investigating stoichiometric constraints on the standard metabolic rate of a small

609   invertebrate. Ecol Lett. 2018;21:1714–22.

610   52.   Janssen PH, Kirs M. Structure of the archaeal community of the rumen. Appl Environ

611   Microbiol. 2008;74:3619–25.

612   53.   Schill NA, Liu JS, von Stockar U. Thermodynamic analysis of growth of Methanobacterium

613   thermoautotrophicum. Biotechnol Bioeng. 1999;64:74–81.

614   54.   von Stockar U, Larsson C, Marison IW. Calorimetry and energetic efficiencies in aerobic and

615   anaerobic microbial growth. Pure Appl Chem. 1993;65:1889–92.

616   55.   Von Stockar U, Liu JS. Does microbial life always feed on negative entropy? Thermodynamic

617   analysis of microbial growth. Biochim Biophys Acta - Bioenerg. 1999;1412:191–211.

618   56.   Hardin G. The competitive exclusion principle. Science (80-). 1960;131:1292–7.

619   57.   Lynch TA, Wang Y, van Brunt B, Pacheco D, Janssen PH. Modelling thermodynamic

620   feedback on the metabolism of hydrogenotrophic methanogens. J Theor Biol. 2019;477:14–23.

621   58.   Udén P, Rounsaville TR, Wiggans GR, Van Soest PJ. The measurement of liquid and solid

622   digesta retention in ruminants, equines and rabbits given timothy (Phleum pratense) hay. Br J

623   Nutr. 1982;48:329–339.

624   59.   Ng F, Kittelmann S, Patchett ML, Attwood GT, Janssen PH, Rakonjac J, et al. An adhesin

625   from hydrogen-utilizing rumen methanogen Methanobrevibacter ruminantium M1 binds a

626   broad range of hydrogen-producing microorganisms. Env Microbiol. 2016;18:3010–21.

60.   Samuel BS, Hansen EE, Manchester JK, Coutinho PM, Henrissat B, Fulton R, et al. Genomic

and metabolic adaptations of Methanobrevibacter smithii to the human gut. Proc Natl Acad
Sci. 2007;104:10643–8.

61. Kelly WJ, Leahy SC, Li D, Perry R, Lambie SC, Attwood GT, et al. The complete genome
sequence of the rumen methanogen Methanobacterium formicicum BRM9. Stand Genomic
Sci. 2014;9:15.

62. Hutchinson GE. The paradox of the plankton. Am Nat. 1961;95:137–45.

63. Bernalier A, Lelait M, Rochet V, Grivet JP, Gibson GR, Durand M. Acetogenesis from H2 and
CO2 by methane- and non-methane-producing human colonic bacterial communities. FEMS
Microbiol Ecol. 1996;19:193–202.

64. Nava GM, Carbonero F, Croix JA, Greenberg E, Gaskins HR. Abundance and diversity of
mucosa-associated hydrogenotrophic microbes in the healthy human colon. ISME J.
2012;6:57–70.

65. Flint HJ, Duncan SH, Scott KP. Interactions and competition within the microbial community
of the human colon : links between diet and health. Environ Microbiol. 2007;9:1101–11.

66. Huws SA, Creevey CJ, Oyama LB, Mizrahi I, Denman SE, Popova M, et al. Addressing
global ruminant agricultural challenges through understanding the rumen microbiome: past,
present, and future. Front Microbiol. 2018;9:2161.

67. Pfeiffer T, Schuster S, Bonhoeffer S. Cooperation and competition in the evolution of ATP-
producing Pathways. Science (80-). 2001;292:50–507.

68. Vandermeer J, Evans MA, Foster P, Höök T, Reiskind M, Wund M. Increased competition
may promote species coexistence. Proc Natl Acad Sci. 2002;99:8731–6.

649 69. MacLean RC, Gudelj I. Resource competition and social conflict in experimental populations

650   of yeast. Nature. 2006;441:498–501.

651 70. Rapaport A, Dochain D, Harmand J. Long run coexistence in the chemostat with multiple

652   species. J Theor Biol. 2009;257:252–9.

653 71. Grognard F, Masci P, Benoît E, Bernard O. Competition between phytoplankton and bacteria:

654   exclusion and coexistence. J Math Biol. 2015;70:959–1006.

655 72. Hsu SB, Hubbell S, Waltman P. A mathematical theory for single-nutrient competition in

656   continuous cultures of micro-organisms. SIAM J Appl Math. 1977;32:366–83.

657 73. Widder S, Allen RJ, Pfeiffer T, Curtis TP, Wiuf C, Sloan WT, et al. Challenges in microbial

658   ecology: Building predictive understanding of community function and dynamics. ISME J.

659   2016;10:2557–68.

660 74. Muñoz-Tamayo R, Ramirez Agudelo JF, Dewhurst RJ, Miller G, Vernon T, Kettle H. A

661   parsimonious software sensor for estimating the individual dynamic pattern of methane

662   emissions from cattle. Animal. 2018;

663 75. Ou JZ, Yao CK, Rotbart A, Muir JG, Gibson PR, Kalantar-zadeh K. Human intestinal gas

664   measurement systems: In vitro fermentation and gas capsules. Trends Biotechnol.

665   2015;33:208–13.

669   **Figures**

671   **Figure 1** Schematics of the *in vitro* methanogenesis process by hydrogenotrophic methanogens.

672   Double arrows represent fluxes due to liquid-gas transfer, simple arrows represent metabolic fluxes.

Figure 2 Example of isothermal calorimetric curves for *M. ruminantium* (dashed blue line), *M. smithii* (solid red line) and *M. formicium* (dotted black line). The dominant metabolic phase is represented by one peak. The magnitude of the peak differs between the methanogens and also the slope of the heat flux trajectories. The return of the heat flux to the zero baseline also differs between the three methanogens. The inset zoom displays the peak exhibited by *M. smithii* at 7.4 h.

Figure 3 Top plots: dynamics of methanogenesis by *M. ruminantium* (*), *M. smithii* (o) and *M. formicium* (□). Experimental data (*,o,□) are compared against model predicted responses: dotted blue lines (*M. ruminantium*), solid red lines (*M. smithii*) and dashed black lines (*M. formicium*). Bottom plots: summary observed vs predicted variables. The solid red line is the isocline.

Figure 4 Dynamics of methanogens as measured by 16S rRNA gene copies (circles) and biomass

concentrations (solid line) predicted by the model.

Figure 5 Possible competition scenarios between *M. ruminantium* (blue dashed line)*, M. smithii* (red solid line) and *M. formicium* (black dotted line) in a hypothetical constant environment. A. At constant dilution rate of 0.021 h^{-1}, *M. formicium* displaces the other two methanogens. B. With a constant dilution rate of 0.04 h^{-1}, *M. smithii* wins the competition. At constant environmental conditions, only one species wins and displaces the other methanogens.

Tables

Table 1 Gibbs energies, enthalpies and entropies of metabolic processes involving some methanogens

growing on different energy sources

Microorganism	Energy substrate	Growth conditions	ΔG_m kJ / C-mol	ΔH_m kJ / C-mol	$T\Delta S_m$ kJ / K^{-1} C-mol	Driving force	Reference
M. ruminantium, M. smithii,	H_2/CO_2	anaerobic	-1073	-2132	-1059	Enthalpy-driven but Entropy-retarded	this work
M. formicium	H_2/CO_2	anaerobic	-801	-1605	-804	Enthalpy-driven but Entropy-retarded	this work
M. thermo-autotrophicum	H_2/CO_2	anaerobic	-802	-3730	-2928	Enthalpy-driven but Entropy-retarded	(53)
M. formicium	formate	anaerobic	-880	-613	+267	Enthalpy-driven	(30)
M. barkeri	methanol	anaerobic	-570	-420	+150	Enthalpy-driven	(30)

| M. barkeri | acetate | anaerobic | -366 | +145 | +511 | Entropy-driven but enthalpy-retarded | (76) |

Table 2 Parameters of the model of *in vitro* methanogenesis. The value reported $\mu_{\max}$ for each methanogen is the mean value obtained from three heat flux-time curves

Parameter	Definition	Value		
$k_L a$ (h^{-1})	Liquid–gas transfer constant	8.33		
K_{H,CO_2} (M/bar)	Henry's law coefficient of carbon dioxide	0.0246		
k_d (h^{-1})	Death cell rate constant	8.33x10^{-4}		
		M. smithii	*M. ruminantium*	*M. formicium*
K_s (mol/L)	Affinity constant	0.028	0.042	0.011
$\mu_{\max}$ (h^{-1})	Maximum specific growth rate constant	0.12	0.07	0.046
Y (mol biomass /mol H$_2$)	Microbial biomass yield factor	0.006	0.006	0.007

735   **Table 3** Statistical indicators for model evaluation

Component	CCC*	r^2	CV$_{RMSE}$**
Hydrogen	0.99	0.97	14
Methane	0.97	0.95	18
Carbon dioxide	0.94	0.88	6

736   * CCC: Lin's concordance correlation coefficient.

737   ** CV$_{RMSE}$: coefficient of variation of the root mean squared error.

Supplementary Material

Hydrogenotrophic methanogens of the mammalian gut: functionally similar, thermodynamically different - A modelling approach

Rafael Muñoz-Tamayo[1*,¶], Milka Popova[2, ¶], Maxence Tillier[2], Diego P. Morgavi [2], Jean-Pierre Morel [3], Gérard Fonty [3], Nicole Morel-Desrosiers[3]

[1]UMR Modélisation Systémique Appliquée aux Ruminants, INRA, AgroParisTech, Université Paris-Saclay, 75005, Paris, France

[2]Institute National de la Recherche Agronomique, UMR1213 Herbivores, Clermont Université, VetAgro Sup, UMR Herbivores, Clermont-Ferrand, France

[3]Université Clermont Auvergne, CNRS, LMGE, F-63000 Clermont-Ferrand, France

1. Growth media information and inoculation conditions

Table S1 Methanogens growth media composition (modified from DSMZ medium 119 https://www.dsmz.de/microorganisms/medium/pdf/DSMZ_Medium119.pdf . Growth media was distributed in Balch tubes (6 ml per tube), tubes were sealed and sterilized by autoclaving at 121°C for 20 min. Media preparation and distribution was realized under CO_2 flushing to assure anoxic conditions. Oxygen traces from commercial gases were scrubbed using a heated cylinder containing reduced copper (35).

Composition per 100 ml	Amount
Clarified rumen fluid [1]	30 ml
Dipotassium phosphate (K_2HPO_4) 0.06% (w/v)	5 ml
Balch Mineral solution [3]	5 ml
Tryptone	0.2 g
Yeast extract	0.2 g
Balch oligo-elements solution [4]	1 ml
Balch vitamin solution[5]	1 ml
Resazurin 0.1%[2]	1 ml

Ammonium chloride	0.05 g
Sodium acetate	0.25 g
Sodium formate	0.25 g
Sodium carbonate	0.5g
L-cystein HCl[2]	0.4g
Distilled water	qs 100 ml

---

[1] Rumen fluid, that was the main constituent of the culture medium, was sampled through the rumen cannula from a grazing dairy cow
prior to the beginning of the experiment. Sampled rumen contents were strained through a monofilament cloth, the filtrate was then
centrifuged at 5 000 g for 15 min. The supernatant was autoclaved and centrifuged again at the same conditions as above. The clarified
rumen fluid (decanted supernatant) was stored at -20°C and centrifuged again after thawing prior to media preparation.

[2] Media was boiled to expel dissolved oxygen, a reducing agent (L-cystein) and a redox indicator (resazurin) were added to keep a low
redox potential and indicate the oxidative state of the medium, respectively.

[3] $KH_2PO_4.2H_2O$ (0.6g), $(NH_4)_2SO_4$ (0.6g), NaCl (1.2g), $MgSO_4.7H_2O$ (0.12g), $CaCl_2.2H_2O$ (0.12g), distilled water qs 100 ml

[4] Nitrilotriacetic acid (0.15 g), $MgSO_4.7H_2O$ (0.3g), $MnSO_4.2H_2O$ (0.05g), NaCl (0.1g), $FeSO_4.7H_2O$ (0.01g), $CoSO_4$ (0.01g),
$CaCl_2.2H_2O$ (0.01g), $ZnSO_4.2H_2O$ (0.01g), $CuSO_4.5H_2O$ (0.001g), $AlK(SO_4)_2$ (0.001g), H_3BO_3 (0.001g), $NaMoO_4.2H_2O$ (0.001g),
$NiCl.6H_2O$ (0.01g), Na_2SeO_3 (0.001g), distilled water qs 100 ml

[5] Biotine (0.2 mg), PABA (0.5 mg), Riboflavine (0.5 mg), Pantothenic acid (0.5 mg), Sodium ascorbate (0.5 mg), Folic acid (0.2 mg),
Niacin (0.5 mg), Pyridoxine (0.10 mg), thiamine (0.05 mg), Vitamin B12 0.1mg/ml (0.1 ml), lipoic acid (0.5 mg), Choline chloride (0.5
mg), Inositol (0.5 mg), Nicotinamide (0.5 mg), Pyridoxal (0.5mg), distilled water qs 100 ml

Table S2 Summary of initial OD and pressure measured immediately after primary inoculation. Presented are means (sd) of 31 values per strain.

Strain	*Methanobrevibacter ruminantium*	*Methanobrevibacter smithii*	*Methanobacterium formicium*
Initial OD	0.054 (0.007)	0.099 (0.027)	0.046 (0.01)
Initial pressure (mbar)	2996 (39.956)	2927 (98.985)	2910 (48.883)

2. Quantification of methanogens

Table S3 qPCR quantification of 16S rRNA genes

	Methanobrevibacter smithii						
Time point (hours after inoculation)	3	5	6.5	24	26	30	72
Microbial pellet (g) *	0.204 ($\pm$0.007)	0.16 ($\pm$0.017)	0.147 ($\pm$0.003)	0.194 ($\pm$0.002)	0.182 ($\pm$0)	0.176 ($\pm$0.012)	0.202 ($\pm$0.012)
Extracted DNA ng/g of microbial pellet*	9765.5 ($\pm$3078)	12695.1 ($\pm$838.4)	13752.8 ($\pm$2662.3)	41257.4 ($\pm$25296.9)	31983.2 ($\pm$512.5)	49386.3 ($\pm$3956.8)	28943.7 ($\pm$7300.7)
Total number of 16S rRNA gene copies in the microbial pellet	5.62×10^8	4.34×10^8	2.71×10^8	2.43×10^9	2.68×10^9	2.66×10^9	2.45×10^9
Total number of cells in the microbial pellet **	2.81×10^8	2.17×10^8	1.35×10^8	1.21×10^9	1.34×10^9	1.33×10^9	1.22×10^9
	Methanobrevibacter ruminantium						
Time point (hours after inoculation)	3	5	6.5	24	26	30	72
Microbial pellet (g) *	0.148 ($\pm$0.048)	0.134 ($\pm$0.006)	0.183 ($\pm$0.013)	0.161 ($\pm$0.006)	0.153 ($\pm$0.039)	0.196 ($\pm$0.004)	0.164 ($\pm$0.023)
Extracted DNA ng/g of	8764.7 ($\pm$1345.4)	10864.3 ($\pm$1325.5)	7546.3 ($\pm$1126.5)	22274.2 ($\pm$16)	19412.8 ($\pm$4976.2)	16184.1 ($\pm$578.6)	23463.5 ($\pm$2053.3)

microbial pellet*							
Total number of 16S rRNA gene copies in the microbial pellet	1.52×10^8	1.74×10^8	2.29×10^8	1.82×10^9	9.95×10^8	1.23×10^9	2.06×10^9
Total number of cells in the microbial pellet **	7.62×10^7	8.71×10^7	1.15×10^8	9.10×10^8	4.97×10^8	6.16×10^8	1.03×10^9
Methanobacterium formicium							
Time point (hours after inoculation)	3	5	6.5	24	26	30	72
Microbial pellet (g) *	0.13 (±0.017)	0.167 (±0.017)	0.086 (±0.003)	0.15 (±0.064)	0.167 (±0.044)	0.195 (±0.016)	0.153 (±0.052)
Extracted DNA ng/g of microbial pellet*	19517.7 (±3588.2)	13263.9 (±875)	21530.9 (±7188.2)	63810.6 (±28842.7)	52442.7 (±21309.7)	38602.9 (±2761.3)	60968.8 (±28872.4)
Total number of 16S rRNA gene copies in the microbial pellet	2.05×10^8	2.31×10^8	2.60×10^8	1.81×10^9	2.02×10^9	2.42×10^9	3.19×10^9
Total number of cells in the microbial pellet	2.05×10^8	2.31×10^8	2.60×10^8	1.81×10^9	2.02×10^9	2.42×10^9	3.19×10^9

* Mean and standar deviation of two observations.
** *M. smithii* and *M. ruminantium* possess two copies of 16S rRNA gene in their genomes.

3. Calculation of thermodynamic properties of the methanogenesis

In our model, the methanogenesis is represented macroscopically by one catabolic reaction (R1) for methane production and one anabolic reaction (R2) for microbial formation. We assumed that ammonia is the only nitrogen source for microbial formation. The molecular formula of microbial biomass was assumed to be $C_5H_7O_2N$ (41).

R_1: $4 H_2 + CO_2 \rightarrow CH_4 + 2 H_2O$

R_2: $10 H_2 + 5 CO_2 + NH_3 \rightarrow C_5H_7O_2N + 8 H_2O$

The thermodynamic properties associated to the methanogenesis result from the contribution of both catabolic and anabolic reactions.

3.1 Thermodynamic properties of formation used for the calculations

Table S4 Standard enthalpies ($\Delta H_f°$) and Gibbs energies ($\Delta G_f°$) of formation at 25°C of compounds involved in hydrogenotrophic methanogenesis. Values were extracted from Wagman et al. (77), with the exception of the microbial biomass that was calculated from values for *Methanosarcina barkeri* reported by Liu et al. (76)

Compound (phase)	$\Delta H_f°$ (kJ/mol)	$\Delta G_f°$ (kJ/mol)
H_2O (l)	-285.830	-237.129
H_2 (g)	0	0
CO_2 (g)	-393.509	-394.359
CH_4 (g)	-74.81	-50.72
NH_3 (aq)	-80.29	-26.50
$C_5H_7O_2N$	-511.50[*]	-349.50[*]

[*] These values are five times the values reported by Liu et al. (76) since the biomass formula we used has five carbon molecules, while the biomass formula used by Liu et al. has one carbon.

3.2 Enthalpies

The heat produced during methanogenesis results from the contribution of both catabolic and anabolic reactions. So, first, we calculated the standard enthalpies of the catabolic and anabolic reactions using the standard enthalpies of formation given in Table S4 for the different compounds involved in methanogenesis.

The standard enthalpy of the catabolic reaction $\Delta H_{r,c}°$ was calculated as follows

$$\Delta H_{r,c}° = \Delta H_{f,CH_4}° + 2 \cdot \Delta H_{f,H_2O}° - \left(4 \cdot \Delta H_{f,H_2}° + \Delta H_{f,CO_2}° \right) = -252.96 \frac{kJ}{mol}$$

A similar equation was used for the calculation of the standard enthalpy of the anabolic reaction $\Delta H_{r,a}^{\circ}$

$$\Delta H_{r,a}^{\circ} = \Delta H_{f,C_5H_7O_2N}^{\circ} + 8 \cdot \Delta H_{f,H_2O}^{\circ} - \left(10 \cdot \Delta H_{f,H_2}^{\circ} + 5 \cdot \Delta H_{f,CO_2}^{\circ} + \Delta H_{f,NH_3}^{\circ}\right) = -750.31 \frac{\text{kJ}}{\text{mol}}$$

These results are at 25°C since this is the temperature of the standard enthalpies of formation reported in Table S3. A correction could be made to get results at 39°C but the heat capacities reported by Wagman et al. (77) show that the temperature correction can be neglected. Similarly, in the interest of simplicity, we assumed that the effect of pressure is negligible. Next, we considered the fact that the heat of a given reaction can be calculated at any state along the reaction pathway *via* the determination of the reaction coordinate or degree of advancement ε (78). Under our assumptions, the heat produced or consumed by a particular reaction during a given interval can be calculated as follows

$$Q = \int_{\varepsilon_0}^{\varepsilon_{t_f}} \Delta H_r^{\circ} \, d\varepsilon$$

For our two reactions, at the instant t we have

$$\varepsilon_c(t) = \frac{n_{H_2}(t) - f \cdot n_{H_{2,0}}}{-4}$$

$$\varepsilon_a(t) = \frac{n_{H_2}(t) - (1-f) \cdot n_{H_{2,0}}}{-10}$$

where $n_{H_2}(t)$ is the number of moles of hydrogen at the instant t, $n_{H_{2,0}}$ is the initial number of moles of hydrogen, and f is the fraction of H_2 used for the catabolic reaction. Our calorimetric experiments started with $n_{H_{2,0}} = 8.83 \cdot 10^{-5}$ mol in all cases. At the final time t_f, all the hydrogen was consumed, so that $n_{H_2}(t_f) = 0$. For *M. smithii* and *M. ruminantium*, the microbial yield factor is Y=0.006 (6) which implies that f =0.94. Accordingly, $\varepsilon_c = 2.075 \cdot 10^{-5}$ mol and $\varepsilon_a = 5.30 \cdot 10^{-7}$ mol. It thus follows that the overall heat produced during the methanogenesis process (Q_m) can be calculated using the following equation

$$Q_m = Q_c + Q_a = \varepsilon_c(t_f) \cdot \Delta H_{r,c}^{\circ} + \varepsilon_a(t_f) \cdot \Delta H_{r,a}^{\circ}$$

where Q_c, Q_a are the heat produced during catabolism and anabolism respectively. The previous equation can also be written as

$$Q_m = n_{H_{2,0}} \left[\frac{(1-10Y)}{4} \cdot \Delta H_{r,c}^{\circ} + Y \cdot \Delta H_{r,a}^{\circ}\right]$$

Under the experimental conditions of our study, this yields

$$Q_m = Q_c + Q_a = (-5.25) + (-0.40) = -5.65 \text{ J}$$

This result shows that the anabolic reaction contributes to only 7% of the metabolic heat.

Since the substrate was totally consumed, the enthalpy of the methanogenesis process per mole (or C-mol) of biomass formed, ΔH_m, can be calculated as follows

$$\Delta H_m = \frac{Q_m}{n_{biomass}} = \frac{Q_m}{Y \cdot n_{H_{2,0}}} = \frac{(1 - 10Y)}{4 \cdot Y} \cdot \Delta H_{r,c}^{\circ} + \Delta H_{r,a}^{\circ}$$

which yields

$$\Delta H_m = -10658 \frac{kJ}{mol} = -2132 \frac{kJ}{C - mol}$$

For *M. formicium*, Y=0.007 (7). Applying the same procedure, we obtained $Q_m = -5.66$ J and $\Delta H_m = -1830 \frac{kJ}{C - mol}$

The anabolic reaction contributes to 8% of the metabolic heat.

3.3 Gibbs energies and entropies

Following a procedure analogous to the one used above for the enthalpies, the standard Gibbs energies of the catabolic ($\Delta G_{r,c}^{\circ}$) and anabolic ($\Delta G_{r,a}^{\circ}$) reactions were calculated using the standard Gibbs energies of formation listed in Table S3.

$$\Delta G_{r,c}^{\circ} = \Delta G_{f,CH_4}^{\circ} + 2 \cdot \Delta G_{f,H_2O}^{\circ} - \left(4 \cdot \Delta G_{f,H_2}^{\circ} + \Delta G_{f,CO_2}^{\circ}\right) = -130.62 \frac{kJ}{mol}$$

$$\Delta G_{r,a}^{\circ} = \Delta G_{f,C_5H_2O_7N}^{\circ} + 8 \cdot \Delta G_{f,H_2O}^{\circ} - \left(10 \cdot \Delta G_{f,H_2}^{\circ} + 5 \cdot \Delta G_{f,CO_2}^{\circ} + \Delta G_{f,NH_3}^{\circ}\right) = -248.24 \frac{kJ}{mol}$$

The free energy of the methanogenesis process per mole (or C-mol) of biomass formed, ΔG_m, can then obtained from the following equation

$$\Delta G_m = \frac{(1 - 10Y)}{4 \cdot Y} \cdot \Delta G_{r,c}^{\circ} + \Delta G_{r,a}^{\circ}$$

which yields for *M. smithii* and *M. ruminantium*

$$\Delta G_m = -5364 \frac{kJ}{mol} = -1073 \frac{kJ}{C - mol}$$

Knowing that

$$\Delta G_m = \Delta H_m - T \cdot \Delta S_m$$

it follows that the entropic contribution to the methanogenesis process is equal to

$$T \cdot \Delta S_m = \Delta H_m - \Delta G_m = -5294 \; \frac{kJ}{mol} = -1059 \; \frac{kJ}{C-mol}$$

which gives, at 39°C, the following value for the entropy of the methanogenesis process per mole (or C-mol) of biomass formed

$$\Delta S_m = \frac{(\Delta H_m - \Delta G_m)}{(273.15 + 39)} = -16.96 \; \frac{kJ}{K \, mol} = -3.40 \; \frac{kJ}{K \, C-mol}$$

The same procedure applied to *M. formicium* yields $\Delta G_m = -917 \; \frac{kJ}{C-mol}, \Delta S_m = -2.92 \; \frac{kJ}{C-mol}$

References

1. Wolfe RS. Techniques for cultivating methanogens. Methods Enzymol. 2011;494:1–22.

2. Batstone DJ, Keller J, Angelidaki I, Kalyuzhnyi S V, Pavlostathis SG, Rozzi A, et al. Anaerobic Digestion Model No.1 (ADM1). IWA Task Group for Mathematical Modelling of Anaerobic Digestion Processes. IWA Publishing, London; 2002.

3. Wagman DD, Evans WH, Parker VB, Schumm RH, Halow I, Bailey SM, et al. The Nbs Tables of Chemical Thermodynamic Properties - Selected Values for Inorganic and C-1 and C-2 Organic-Substances in Si Units. J Phys Chem Ref Data. 1982;11:S2.

4. Liu JS, Marison IW, von Stockar U. Microbial growth by a net heat up-take: A calorimetric and thermodynamic study on acetotrophic methanogenesis by Methanosarcina barkeri. Biotechnol Bioeng. 2001;75:170–80.

5. Prigogine I, Defay R. Traité de thermodynamique, conformément aux méthodes de Gibbs et De Donder. Éditions Desoer. Liège; 1950.

6. Pavlostathis SG, Miller TL, Wolin MJ. Cellulose Fermentation by Continuous Cultures of Ruminococcus-Albus and Methanobrevibacter-Smithii. Appl Microbiol Biotechnol. 1990;33(1):109–16.

7. Schauer NL, Ferry JG. Metabolism of formate in Methanobacterium formicicum. J Bacteriol. 1980;142:800–7.