methyl-ATAC-seq measures DNA methylation at accessible chromatin

Spektor R[1], Tippens ND[2], Mimoso CA[3], Soloway PD[4,5]

1 Department of Molecular Biology and Genetics, Field of Genetics, Genomics, and Development, Cornell University, Ithaca, New York 14853, USA.
2 Tri-Institutional Training Program in Computational Biology and Medicine, Cornell University, Ithaca, New York 14853, USA.
3 College of Agricultural and Life Sciences, Cornell University, Ithaca, New York 14853, USA
4 College of Veterinary Medicine, Department of Biomedical Sciences, Cornell University, Ithaca, New York 14853, USA.
5 College of Agriculture and Life Sciences, Division of Nutritional Sciences, Cornell University, Ithaca, New York 14853, USA

ABSTRACT:

Chromatin features are characterized by genome-wide assays for nucleosome location, protein binding sites, 3-dimensional interactions, and modifications to histones and DNA. For example, Assay for Transposase Accessible Chromatin sequencing (ATAC-seq) identifies nucleosome-depleted (open) chromatin, which harbors potentially active gene regulatory sequences; and bisulfite sequencing (BS-seq) quantifies DNA methylation. When two distinct chromatin features like these are assayed separately in populations of cells, it is impossible to determine, with certainty, where the features are coincident in the genome by simply overlaying datasets. Here we describe methyl-ATAC-seq (mATAC-seq), which implements modifications to ATAC-seq, including subjecting the output to BS-seq. Merging these assays into a single protocol identifies the locations of open chromatin, and reveals, unambiguously, the DNA methylation state of the underlying DNA. Such combinatorial methods eliminate the need to perform assays independently and infer where features are coincident.

INTRODUCTION:

Active promoters, enhancers, and other gene regulatory sequences are typically bound by sequence-specific transcription factors (TFs), free of nucleosomes, and these facilitate transcription. Such regulatory sequences can be identified by methods that detect nucleosome-depleted regions (NDRs), including DNase-seq, which identifies NDRs by their hypersensitivity to DNase I (Thurman et al. 2012); FAIRE-seq, which identifies NDRs according to their reduced protein content (Gaulton et al. 2010); and ATAC-seq, which identifies NDRs based on their increased accessibility to Tn5 transposase integration, and accordingly are called Transposase hypersensitive sites (THS) (Buenrostro et al. 2013). There is considerable agreement among the regions identified by each assay. ATAC-seq has received further use recently owing to its simplified workflow, reduced material requirements and lower background signals. Additional advancements such as Omni-ATAC (Corces et al. 2017) and Fast-ATAC (Corces et al. 2016) have further improved the utility of ATAC-seq.

DNA within NDRs may have different modification states, including methylation at the fifth carbon of Cytosine (5mC), and oxidized derivatives. In the mammalian genome, most 5mC is found at CpG dinucleotides, and is generally associated with transcriptionally inactive regions. Bisulfite sequencing (BS-seq) uses selective chemical deamination of unmodified cytosines to uracil, leaving 5mC unchanged. The extent of methylation at a given CpG in a sample is detected after amplification, sequencing, aligning reads to the genome, and then assessing the proportion of aligned reads that retained a C at a CpG, diagnostic of methylation, *vs.* a T, which reports an unmethylated residue.

Two features of BS-seq dramatically increase costs compared to routine sequencing assays. First, bisulfite treatment reduces the yield and complexity of DNA libraries, resulting in fewer reads uniquely aligning to the genome. Second, to reliably quantify the extent of methylation of a given CpG requires high read coverage. For these reasons, Reduced Representation Bisulfite Sequencing (RRBS) (Meissner et al. 2005) and derivatives (Boyle et al. 2012; Chatterjee et al. 2012; Garrett-Bakelman et al. 2015) have been used to focus analysis on CpG dense regions. However, not all gene regulatory sequences are detected by RRBS, and many regions that are detected are not regulatory.

Integrating results from assays for distinct chromatin features have defined novel categories of regulatory elements. These include bivalent promoters (Bernstein et al. 2006), enhancers

70   (Heintzman et al. 2009), and widely observed chromatin states likely to harbor shared regulatory functions (Roadmap Epigenomics et al. 2015). In most of these studies, results from assays for single features are superimposed, and when a given locus has signals for multiple features, the features are inferred to be coincident on the same molecule. Though many inferences might be accurate, there is uncertainty inherent in such approaches, owing to the fact that samples

75   commonly contain multiple sub-populations of cells, each with a characteristic chromatin state. Accordingly, the population-averaged results might report chromatin states found in no individual subpopulation of cells. Methods that combine assays for multiple chromatin features in a single protocol can eliminate this ambiguity for the features assayed. Here, we describe methyl-ATAC-seq (mATAC-Seq), a modification of ATAC-seq that combines ATAC-seq with

80   BS-seq, identifying the locations of open chromatin, and the methylation state of the underlying DNA. In addition to providing more reliable assignments of chromatin states, mATAC-seq can focus DNA methylation analyses to accessible regulatory regions of the genome.

RESULTS:

85 Fig. 1 shows the workflow and sample results for mATAC-seq. It includes two primary modifications during the transposition step of the Omni-ATAC-seq protocol: (1), methylated oligonucleotides are loaded onto Tn5 to generate the transposome (Fig. 1A) which is then used to perform ATAC-seq (Fig. 1B); and (2), 5-methyldeoxycytosine triphosphate (5-mdCTP) is substituted for dCTP during the subsequent end repair step (Fig. 1C). These modifications

90 protect the Nextera adapter sequences during the final step of mATAC-seq library preparation, bisulfite treatment of the tagmented DNA (Fig. 1D). Use of methylated oligonucleotides, and 5-mdCTP during end repair protects cytosines in the adaptors from deamination caused by bisulfite treatment, which is necessary for successful PCR amplification and sequencing of the resulting libraries. Sequenced libraries provide information on both DNA methylation and

95 Transposase hypersensitivity (Fig. 1E).

We applied mATAC-seq to nuclei prepared from HCT116 colorectal carcinoma cells. mATAC-seq reads in peaks were highly reproducible in biological replicates (r^2=0.90, Fig. S1A). To validate that mATAC-seq captured open chromatin domains as well as conventional methods,

100 we compared Transposase Hypersensitive (THS) sites found by mATAC-seq with those we identified using the standard Omni-ATAC-seq protocol (Fig. 2A-D) (Corces et al. 2017). Approximately 92% of called peaks found by Omni-ATAC-seq were found by mATAC-seq (Fig. 2A). There was also strong concordance between mATAC-seq and Omni-ATAC-seq with respect to gene features detected by both assays, with promoter regions being the most

105 commonly identified features (Fig. 2B). In addition, reads in peaks identified by Omni-ATAC-seq and mATAC-seq were well correlated (Fig. S1A, B). Regions of greatest divergence include difficult to map regions such as repetitive elements, low complexity sequences, and simple repeat annotations (Fig. S1C). These analyses demonstrate that mATAC-seq detects open chromatin comparably to traditional Omni-ATAC-seq, and that protocol modifications that enable

110 subsequent bisulfite sequencing do not compromise detection of open chromatin.

To validate that mATAC-seq identified DNA methylation patterns as reliably as conventional methods, we next compared the mATAC-seq methylation data with whole genome bisulfite sequencing (WGBS) data reported for HCT116 cells at THS sites and CpG islands (Blattler et

115 al. 2014). DNA methylation detected by mATAC-seq replicates was highly reproducible at peaks (r^2=0.83) and CpG Islands (r^2=0.95) (Fig. S2A, B); and methylation levels reported by mATAC-seq correlated well with levels reported by WGBS at peaks (r^2=0.68) and CpG Islands (r^2=0.85)

(Fig S2A, B). THS peaks identified by mATAC-Seq in HCT116 were predominantly unmethylated, and this agrees with existing WGBS data (Fig S2C, D). Fig. 2E and Fig. 2F report DNA methylation patterns assayed respectively by mATAC-seq and WGBS across gene bodies spanning from 2kb 5' of transcriptional start sites (TSS) to 2kb 3' of transcriptional end sites (TES). These patterns are consistent with the high correlations described above. We find these high correlations despite the fact that the assays were performed by different labs; also, WGBS and mATAC-seq assays are different in that mATAC-seq queries DNA methylation at open chromatin, whereas WGBS assays the entire genome, regardless of chromatin state. Our mATAC-seq data showed a reciprocal relationship between accessibility and 5mC density. These are in agreement with previous results from NOMe-seq (Kelly et al. 2012), which can also report sites of accessible chromatin and DNA methylation states but requires much greater sequencing depth. Both assays reveal that highly accessible chromatin is depleted of 5mC, and that there is an abundance of methylation in less accessible chromatin over gene bodies (Fig. 2E, F). Having shown that sites of open chromatin and DNA methylation states reported by mATAC-seq, Omni-ATAC-seq, and WGBS are in agreement, we concluded that mATAC-seq can be used to simultaneously identify the locations of the genome with accessible chromatin, and the methylation state of the underlying DNA. Because mATAC-seq measures accessibility and methylation in a single assay, it eliminates the inherent uncertainty about coincidence of chromatin features that can arise when ATAC and bisulfite assays are performed independently, and inferences are made after overlaying the two datasets, and at lower costs.

We extended our analyses of HCT116 cells, performing mATAC-seq on HCT116-derived *DNMT1* and *DNMT3B* double knock-out cells (DKO) (Rhee et al. 2002) to assess the functional significance of these methyltransferases on chromatin accessibility and methylation states in parental HCT116 cells. In DKO cells, there were 23,301 hyper-accessible sites, and 3,166 hypo-accessible sites, compared to parental HCT116 cells (Fig. 3A; | $\log_2$ fold change | > 1, q < 0.01); 16,170 THS sites observed in HCT116 cells were unchanged in DKO cells (| $\log_2$ fold change | < 1, q > 0.8). Compared to unchanged sites, hyper-accessible sites in DKO cells were depleted of DNA methylation (Fig. 3B); these sites were enriched for ATF3, FOSL1, FOSL2, BATF, AP1 and JUNB binding motifs (Fig. 3C). These TFs were previously shown to interact more strongly to their binding motifs when unmethylated (methyl-minus TFs (Yin et al. 2017)). We infer that chromatin hyper-accessibility at these sites in DKO cells was due to enhanced binding of the methyl-minus TFs when methylation was diminished; this had the effect of limiting nucleosome deposition, thus enabling increased chromatin accessibility. Conversely, hypo-accessible sites

in DKO cells were modestly depleted of DNA methylation (Fig. 3B), and enriched for SP1, NFYA, SP5, KLF9, KLF14, and KLF3 binding motifs (Fig. 3D). These TFs were previously shown to exhibit less binding when their sites were unmethylated (methyl-plus TFs (Yin et al.

155   2017)). We infer that chromatin hypo-accessibility at these sites in DKO cells was due to reduced binding of the methyl-plus TFs when methylation was diminished, and that this led to increased nucleosome deposition, and reduced chromatin accessibility. In support of this is the observation that promoters showing the greatest increases in chromatin accessibility in DKO cells were also the promoter that were most extensively hypomethylated (Fig. 3E). These

160   findings and conclusions are consistent with previously described mechanisms whereby TF binding can regulate nucleosome density (Zaret and Carroll 2011). These conclusions may be tempered by the fact that we are assaying methylation at accessible sequences, the same loci, when in an inaccessible state, are underrepresented in our methylation analyses.

165   To assess how promoter accessibility states detected by mATAC-seq relate to gene expression, we queried existing RNA-seq data from HCT116 and DKO cells (Blattler et al. 2014). Promoters that were hypo-accessible in DKO cells exhibited no significant gene expression changes relative to the corresponding promoters in parental HCT116 cells. At promoters that exhibited no differences in accessibility in the two cell types, there were significant, but very modest

170   differences in mean expression levels. At promoters that were hyper-accessible in DKO cells, we observed substantial and significantly higher levels of expression in DKO cells relative to HCT116, with expression differences increasing as accessibility increased (Fig. 3F). These are in accordance with previous findings (Kelly et al. 2012), further validating the utility of mATAC-seq, and demonstrating the concordance between the extent of chromatin accessibility at

175   promoters, and promoter activity.

Our analyses so far have separately examined methylation and chromatin accessibility results from mATAC-seq. We next combined methylation and accessibility data to take advantage of added value of the combined results afforded by mATAC-seq. We first performed *k*-means

180   clustering of DNA methylation levels at THS sites in HCT116 and DKO cells. DNA methylation at mATAC-seq peaks in HCT116 cells formed five distinct clusters (Fig. 4A). In Cluster 1, accessible peaks, and the 1kb intervals flanking the peaks, were hypermethylated in HCT116 relative to DKO cells, with the flanks exhibiting more hypermethylation. Clusters 2 and 3 were hypomethylated at peak centers in both cells; the clusters were respectively hypermethylated in

185   HCT116 cells in one or the other of the two intervals flanking the peaks. Cluster 4 was

hypermethylated over the peaks only in HCT116 cells, and hypomethylated in the peak and flanks in DKO cells. Cluster 5 was hypomethylated in the peaks and flanks of both cell types (Fig. 4A, C, D).

190   When we assessed mRNA expression from promoters within the five clusters, differences between DKO and HCT116 emerged that varied according to cluster. Promoters in DKO cells from Clusters 1 and 4 were significantly more active than the corresponding promoters from the same clusters in HCT116 cells, with respective increases in mRNA of 2.5-, and 3.2- fold (Fig. 4B). Clusters 2 and 3 exhibited a modest change of 1.3-fold between the cell types. Cluster 5,

195   which was both hypomethylated and hyper-accessible in both cell types, showed no difference in expression.

Besides the differences in DNA methylation and expression, the clusters have additional distinguishing features. There are more promoters, CpG islands, and exons in Cluster 1

200   compared to Cluster 4; and more intronic and distal intergenic elements in Cluster 4 compared to Cluster 1 (Fig. S3A, B). One feature is the broad domain of H2A.Z in Cluster 1 that accompanied the loss of DNA methylation in DKO cells (Fig. 4E). This finding is consistent with reports that DNA modification and H2A.Z are mutually antagonistic (Zilberman et al. 2008). In Cluster 4, where hypermethylation in HCT116 cells is largely confined to the mATAC-seq peak,

205   there was also an increase in H2A.Z in DKO cells, with the increase being more modest and confined to a narrower portion of the 2kb window displayed. Additional histone modifications associated with active chromatin (H3K4me1, H3K4me3, H3K27ac) were elevated in DKO cells near Cluster 1 mATAC-seq peaks, but these effects were limited or absent in Cluster 4 (Fig. 4E). Like H2A.Z, H3K27me3 was increased in DKO cells at Cluster 1, with the effects also being

210   more modest at Cluster 4 (Fig. 4F). This is also consistent with antagonism reported between H3K27me3 and DNA methylation (Lindroth et al. 2008). In contrast to these histone modifications and variants, H3K9me3 at mATAC peaks was largely unaffected by DNMT-loss. Cluster 5 shows no DNA methylation changes between the two conditions, and there were little to no changes in deposition of histone modifications and variants.

Motifs for TFs, and CTCF binding also varied by cluster. Cluster 1 is enriched for motifs recognized by DNA methyl-plus TFs such as CTCFL, MYC, and BHLHE40; ZFX and ZNF711 contain similar motifs to ZNF704, a methyl-minus transcription factor (Fig 4H). Of the top five TFs enriched in Cluster 4, three are MEF-family TFs, followed by ARNT, which was previously

220   suggested to be methyl-sensitive (Lay et al. 2015) (Fig. 4I). ARNT motifs share substantial sequence identity with BHLHE40, a methyl-minus TF.

DISCUSSION:

225   ATAC-seq identifies nucleosome-depleted regions of the genome, which are arguably the most relevant for gene regulation within cells. By including bisulfite treatment in the workflow, mATAC-seq targets DNA methylation profiling to open chromatin sites that are enriched for active regulatory regions of the genome. Accordingly, mATAC-seq queries the functional methylome of cells, using relatively few reads compared to WGBS. This is in contrast to other

230   assays for DNA methylation that query the entire genome, or other domains that may not be regulatory.

By applying mATAC-seq to the well-characterized HCT116 cell line, and its DNA methylation-deficient DKO derivative, we demonstrated that mATAC-seq detects DNA methylation patterns

235   that agree with both previously described WGBS results, and with our Omni-ATAC-seq results. These tests validated the fidelity, and compatibility of combining tagmentation and bisulfite treatment steps in the mATAC-seq workflow. DKO cells had many hyper-accessible sites relative to parental HCT116 cells, and these sites exhibited loss of methylation. These same regions were also enriched for methyl-minus TF binding sites, which interact more strongly with

240   DNA when the sites are in an unmethylated context. This highlights the instructive role of TF binding for nucleosome occupancy in the genome. Specifically, our data indicate that when DNA is unmethylated, it facilitates the recruitment of methyl-minus TFs, and that these in turn enable chromatin to assume an open state. Our data also revealed that hyper-accessible and hypomethylated domains in DKO cells were enriched for the histone variant H2A.Z, implicating

245   this factor in limiting DNA methylation, and nucleosome density at sequences where it is recruited. In contrast, regions that displayed no change in methylation showed little change in accessibility. We did not observe a depletion of H3K9me3 at sites with increased accessibility, and decreased DNA methylation, confirming statements in previous studies (Blattler et al. 2014). Such findings were made possible by the combination of DNA methylation, and open

250   chromatin status provided by mATAC-seq.

We envisage that mATAC-seq could be applied to many other systems. For example, HCT116 derivatives carrying single *DNMT* knockouts (Rhee et al. 2000; Rhee et al. 2002) would enable us to identify regulatory elements the different DNMTs individually target for methylation, and

255   their respective influences on nucleosome placement. The various DNMTs have been shown to regulate DNA methylation states by independent, as well as cooperative mechanisms (Liang et al. 2002). Repetitive elements are common targets of the DNMTs, and the resulting DNA

methylation contributes to their silencing. However, silencing can occur when DNMT activities are impaired, indicating that compensating mechanisms can silence transposons, likely
260   involving H3K9 methylation, and possibly other chromatin modifications (Horard et al. 2009; Karimi et al. 2011; Walter et al. 2016; Jorda et al. 2017). Querying the specificities of the DNMTs, and their influences on chromatin accessibility at repetitive elements using mATAC-seq can elaborate mechanisms underlying repeat regulation.

265   Additionally, by using HCT116 DKO cells, we studied the effects of DNA methylation depletion that arose by passive mechanisms due to a lack of DNA methylation maintenance. Active demethylation by TET dioxygenases, and AID/APOBEC deaminases, occurring during differentiation and response to stimuli, is a distinct process. Applying mATAC-seq to stem cell differentiation – including under conditions where these active demethylation mechanisms are
270   altered – can reveal both the combinatorial changes in accessibility and DNA methylation, and the effects active DNA methylation mechanisms have on chromatin state during differentiation.

Pioneer transcription factors have the unique property of binding chromatin that is generally inaccessible to other transcription factors (Zaret and Mango 2016). Application of mATAC-seq
275   to systems where pioneer factor functions are altered can reveal the influences these factors have on both chromatin accessibility, and methylation state of the underlying DNA.

Our protocol for mATAC-seq can potentially be integrated with existing methods for combinatorial detection of other DNA modifications including 5-hydroxymethylcytosine (Yu et al.
280   2012; Booth et al. 2013), 5-formylcytosine (Song et al. 2013), and 5-carboxycytosine (Wu et al. 2016). ChIPmentation uses Tn5 tagmentation in a chromatin immunoprecipitation workflow (Schmidl et al. 2015). This too could be implemented, using steps we developed for mATAC-seq, to identify locations of DNA-bound proteins, and the underlying DNA modification states in a combinatorial detection strategy similar to other methyl-ChIP strategies (Brinkman et al. 2012;
285   Statham et al. 2012). Combinatorial indexing as a low-cost strategy to query single cells can be used to enable the extension of mATAC-seq to a single cell format; specifically, methods such as single-cell combinatorial indexing assay for transposase accessible chromatin using sequencing (sci-ATAC-seq) (Cusanovich et al. 2015), and for methylation analysis (sci-MET) (Mulqueen et al. 2018). Some alterations are necessary to adapt our mATAC-seq protocol for
290   single cell sequencing, including, extending the indexed adapter set to use methylated sci-ATAC-seq adapters during tagmentation, followed by split-pooling, methylated end-repair,

bisulfite conversion in a 96-well format, and PCR. The challenge to this approach is the depletion of reads due to the destructiveness of bisulfite conversion, and the limited sequence complexity in bisulfite converted reads.

MATERIALS AND METHODS:

Cell Culture:

Cultured cells (#28 HCT116 Parental and #343 DKO) were procured from the Genetic
300 Resources Core Facility at Johns Hopkins School of Medicine and cultured in McCoy's Modified
5A Medium containing 10% heat-inactivated FBS and 1× Penn/Strep (Gibco #15140122). Cells
for each experiment were grown apart for at least 2 passages before library preparation.

Genotyping:

305 DNA from each cell line was extracted using EZ-10 Spin Columns (Bio Basic #BS427) following
the manufacturer's protocol. Genotyping PCR was performed on 50ng genomic DNA using
oligos from Table S2 from (Das and Chadwick 2016) for 40 cycles using GoTaq (Promega
#M3001) (94°C 2 minutes, 40 cycles of: [94°C 30 seconds, 60°C 30 seconds, 72°C 30
seconds], 72°C 5 minutes) and run on a 2% agarose gel. Cells were confirmed to be
310 Mycoplasma-free and HeLa-free via PCR (Rahbari et al. 2009; Young et al. 2010) on 50ng
genomic DNA and cell-culture media (Figure S4).

Omni-ATAC-seq:

Cells were trypsinized and subsequently inactivated in cell culture media. Following inactivation,
315 cells were pelleted and resuspended in cold PBS (without Ca++ and Mg++). Cells were stained
with Trypan Blue and counted on a hemocytometer. Lysis and tagmentation were performed
exactly as described (Corces et al. 2017) with modifications to inactivation and size selection.
Briefly, 100,000 HCT116 Parental and DKO cells were lysed on ice for 3 minutes in 50 µL ice-
cold Lysis Buffer (10mM Tris pH 7.4, 10mM NaCl, 3mM $MgCl_2$, 0.1% NP-40, 0.1% TWEEN 20,
320 0.01% Digitonin in DEPC H2O), resuspended in 1mL ice-cold RBS-Wash (10mM Tris pH 7.4,
10mM NaCl, 3mM $MgCl_2$, 0.1% TWEEN 20) and pelleted at 4°C at 500 × g for 10 minutes.
Tagmentation was performed in 1 × Tagmentation Buffer (10mM Tris pH 7.4, 5mM $MgCl_2$, 10%
DMF, 33% PBS, 0.1% TWEEN 20, 0.01% Digitonin) using 100nM Tn5 Transposase for 30
minutes at 37°C. Tagmentation was inactivated with the addition of 5 volumes SDS Lysis Buffer
325 (100mM Tris pH 7.4, 50mM NaCl, 10mM EDTA, 0.5% SDS in H_2O) and 100µg Proteinase K
(Invitrogen #25530049) for 30 minutes at 55°C followed by Isopropanol precipitation using
GlycoBlue (Invitrogen #AM9516) as a carrier. DNA was size selected using Ampure XP beads
(Beckman Coulter # A63880) using a 0.5 × volumes to remove large fragments followed by a
1.8 × final volume according to the manufacturer's instructions. PCR was performed for using
330 Q5 DNA polymerase (NEB #M0491S) with 1 × GC buffer (72°C 5 minutes, 98C 30 seconds, 11
cycles of: [98°C 10 seconds, 65°C 30 seconds, 72°C 30 seconds], 72°C 5 minutes) followed by
a final cleanup using a 1.8 × volumes of Ampure XP beads according to the manufacturer's
instructions.

335 methylATAC-seq:

Cell lysis was performed identically to Omni-ATAC-seq. Tagmentation was performed on
250,000 HCT116 Parental and DKO cells using 700nM Tn5 Transposase assembled using pre-
annealed Tn5ME-A_mC and Tn5ME-B_mC (Table S3) for 30 minutes at 37°C following the
addition of 0.01ng of unmethylated Lambda DNA (Promega #D1521). We recommend

340   performing a titration of Tn5 transposase to nuclei input to assay minimum amounts required as in Figure S5. Inactivation and size-selection were performed identically to our modified Omni-ATAC-seq protocol. Tagmented DNA was End-Repaired for 30 minutes at 37°C (5U Klenow Exo- (NEB #M2012S), 1 × NEB Buffer 2, and 0.5 mM/each dATP, dGTP, dTTP, and 5-mdCTP (NEB #N0365S)) similar to T-WGBS (Lu et al. 2015) and X-WGBS (Suzuki et al. 2018). End

345   repair was cleaned using a 1.8 × volumes of Ampure XP beads according to the manufacturer's instructions. 10% of the product was kept for quality control PCR (Fig. S6A). Bisulfite conversion was performed using EZ DNA Methylation-Lightning (Zymo #D5030T) following the manufacturer's protocol. PCR was immediately performed using PfuTurbo Cx (Agilent #600410) (94°C 2 minutes, 13 cycles of: [98°C 10 seconds, 6°5C 30 seconds, 72°C 30 seconds], 72°C 5

350   minutes) (Fig S6B) followed by a final cleanup using a 1.8 × volumes of Ampure XP beads according to the manufacturer's instructions.

Tn5 Transposase:
Tn5 was produced exactly as described (Picelli et al. 2014) with no modifications. For Omni-

355   ATAC-seq, Tn5 transposase was assembled as described (Adey and Shendure 2012) using pre-Annealed Tn5MEDS-A and Tn5MEDS-B from Table S3. For methylATAC-seq, Tn5 transposase was assembled using pre-Annealed Tn5ME-A_5mC and Tn5MEB_5mC oligonucleotides from Table S3. Oligonucleotides were annealed by combining ME-A or ME-B oligos to Tn5MErev and incubating for 2 minutes at 94°C followed by a 0.1°C/s ramp to 25°C.

360   Enzyme was stored at -80°C.

Data Analysis:
Libraries were quantified using the Qubit dsDNA HS Assay Kit (ThermoFisher #Q32854). High-throughput sequencing was performed by the Cornell University Genomics Facility on the

365   Illumina NextSeq 500 with single-end 75bp reads. Trimming for mATAC-seq and Omni-ATAC-seq was performed using fastp (Chen et al. 2018) -q 20 -l 20 -a CTGTCTCTTATACACATCT. Trimming for ChIP-seq, RNA-seq, and WGBS data was performed using fastp -q 20 -l 20 -a AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC.

370   <u>Alignment to hg19:</u> In this study we used GRCh37 instead of GRCh38 to match previous studies using similar cells and methods. These results would not be affected by this change because we do not study centromeric sequences and predominantly discuss changes at promoters.

    <u>ChIP-seq hg19:</u> Trimmed FASTQ files were aligned using BWA-MEM (Li and Durbin 2010) to

375   hg19. Reads were deduplicated using Picard [http://broadinstitute.github.io/picard/] MarkDuplicates. HCT116 and DKO ChIP-seq data for H2A.Z, H3K4me3, H3K4me1, H3K27Ac, H3K27me3, H3K9me3, and H3K36me3 data (Lay et al. 2015) were downloaded from NCBI GEO database accession GSE58638. HCT116 and DKO ChIP-seq data for CTCF (Maurano et al. 2015) were downloaded from NCBI GEO database accession GSE50610.

    <u>RNA-seq hg19:</u> Pair-end trimmed FASTQ files were aligned using HISAT2 (Kim et al. 2015) to hg19. HCT116 and DKO RNA-seq data (Blattler et al. 2014) were downloaded from NCBI GEO database accessions GSE52429 and GSE60106, respectively.

385   Omni-ATAC and mATAC-Seq: Trimmed FASTQ files were aligned using Bismark (Krueger and
        Andrews 2011) v0.19.0 to hg19 using the following settings: --score_min L,0,-0.6. Bisulfite reads
        to be used for MethylKit were filtered for non-conversion using Bismark's filter_non_conversion
        and deduplicated using deduplicate_bismark. Methylation was extracted using Bismark's
        methylation extractor --gzip --bedgraph --counts --ignore 9 --ignore_3prime 9. Reads used for
390   peak calling and ATAC-seq visualization were deduplicated using deduplicate_bismark without
        filtering for non-conversion. Conversion rate (Table S1) was measured by aligning to the lambda
        genome (GenBank: J02459.1) and filtered as above; percent conversion rate was calculated as
        (1-(Total methylated C's in all contexts)/(Total number of C's analyzed)) × 100.

395   WGBS hg19: Trimmed FASTQ files were aligned using Bismark v0.19.0 to hg19 using the
        following settings: --score_min L,0,-0.6. Bisulfite reads to be used for MethylKit were filtered for
        non-conversion using Bismark's filter_non_conversion and deduplicated using
        deduplicate_bismark. Methylation was extracted using Bismark's methylation extractor --gzip --
        bedgraph --counts. HCT116 and DKO WGBS data (Blattler et al. 2014) were downloaded from
400   NCBI GEO database accession GSE60106.

        Methylation: Differential methylation was quantified using MethylKit (Akalin et al. 2012) at
        merged HCT116 and DKO mATAC-seq peaks extended to 1kb tiles covering at least 3 CpGs.
        Promoters were defined as being within 1kb of a TSS using Genomation (Akalin et al. 2015).
405
        Peak calling: ATAC-seq peaks were called using HOMER (Heinz et al. 2010) findPeaks
        localSize 50000 -size 150 -minDist 50 –fragLength 0 -style dnase. ChIP-seq peaks were called
        using HOMER findPeaks -style histone. CTCF ChIP-seq peaks were called using HOMER
        findPeaks -style factor. Reads were assigned to peaks merged from HCT116 and DKO cells
410   using featurecounts (Liao et al. 2013) on reads filtered for a minimum log_2 CPM of 0.5 in at least
        2 samples. Differential accessibility was called using DESeq2 (Love et al. 2014) lfcShrink.
        Hyper- and hypo-accessible peaks were defined as having a | log_2 FC | > 1 with an adjusted p
        value < 0.01 in DKO compared to HCT116 parental cells. Promoters were defined as being
        within 1kb of a TSS using Genomation. FRiP scores in Table S1 and sample correlation in Fig.
415   S2 were quantified using DiffBind (Stark and Brown 2018) on libraries downsampled to 5M
        reads using Picard DownsampleSam using peaks called by HOMER. Peak overlaps for Fig. 1A
        and Fig. S2E were generated using ChIPpeakAnno (Zhu et al. 2010). Feature overlaps for Fig.
        1B and S3A were generated using ChIPseeker (Yu et al. 2015). Motif enriched in changed
        peaks were called using HOMER findMotifsGenome to the hg19 genome using unchanged
420   peaks as background.

        RNA-seq quantification: Unstranded hg19-aligned reads were assigned to hg19 genes using
        featurecounts inbuilt reference using default settings. Differential expression was quantified
        using DESeq2 lfcShrink on reads filtered for a minimum CPM of 0.5 in at least 2 samples.
425
        Genome browser visualizations: ATAC-seq and mATAC-seq bigWig files were made using
        eepTools (Ramirez et al. 2016) bamCoverage --binSize 1 --normalizeUsing RPKM --

ignoreForNormalization chrM --scaleFactor N and viewed on UCSC genome browser. ChIP-Seq bigWigs were made using deepTools bamCoverage --binSize 10 --normalizeUsing RPKM --

430 ignoreForNormalization chrM --scaleFactor N. Scale factor was determined by coverage of peaks called by HOMER shared between HCT116 and DKO via bedops --intersect where N = (% reads in shared peaks in HCT116)/(% reads in shared peaks in DKO) when N > 1.1. Scaling was applied to the following samples: DKO_mATAC 1.877, DKO_H3K27ac_R2 1.48, H3K4me3_R2 = 1.47.

Gene body heatmaps were produced using deepTools plotheatmap --beforeRegionStartLength 2000 --regionBodyLength 3000 --afterRegionStartLength 2000 to Ensembl hg19 APPRIS PRINCIPAL:1 flagged transcripts (Rodriguez et al. 2013). Heatmaps for differential peaks were centered on peaks called by HOMER. Peaks from HCT116 and DKO were combined using

440 BEDOPS for clustering of DNA methylation at THS sites. Clustering was performed using deepTools plotheatmap --kmeans 5, the output and order of which was used for all subsequent heatmaps.

Data Access:

445 All raw and processed sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; http://www.ncbi.nlm.nih.gov/geo/) under accession number GSE126215.

450 **ACKNOWLEDGEMENTS:**

Jennifer D. Mosher, Ann E. Tate, Jeff C. Mattison, and Peter A. Schweitzer from the Cornell University Biotechnology Resource Center (BRC) for genomic sequencing; Bert Vogelstein for HCT116 cells and derivatives; Önder Kartal, and Erin Chu for proofreading and helpful

455   comments; the Cornell College of Veterinary Medicine and the National Institutes of Health for funding (R01HG006850 and R01GM105243).

REFERENCES:

460   Adey A, Shendure J. 2012. Ultra-low-input, tagmentation-based whole-genome bisulfite sequencing. *Genome Res* **22**: 1139-1143.

Akalin A, Franke V, Vlahovicek K, Mason CE, Schubeler D. 2015. Genomation: a toolkit to summarize, annotate and visualize genomic intervals. *Bioinformatics* **31**: 1127-1129.

Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A, Mason CE. 465   2012. methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biol* **13**: R87.

Bernstein BE, Mikkelsen TS, Xie X, Kamal M, Huebert DJ, Cuff J, Fry B, Meissner A, Wernig M, Plath K et al. 2006. A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* **125**: 315-326.

470   Blattler A, Yao L, Witt H, Guo Y, Nicolet CM, Berman BP, Farnham PJ. 2014. Global loss of DNA methylation uncovers intronic enhancers in genes showing expression changes. *Genome Biol* **15**: 469.

Booth MJ, Ost TWB, Beraldi D, Bell NM, Branco MR, Reik W, Balasubramanian S. 2013. Oxidative bisulfite sequencing of 5-methylcytosine and 5-hydroxymethylcytosine. *Nat* 475   *Protocols* **8**: 1841-1851.

Boyle P, Clement K, Gu H, Smith ZD, Ziller M, Fostel JL, Holmes L, Meldrim J, Kelley F, Gnirke A et al. 2012. Gel-free multiplexed reduced representation bisulfite sequencing for large-scale DNA methylation profiling. *Genome Biol* **13**: R92.

Brinkman AB, Gu H, Bartels SJ, Zhang Y, Matarese F, Simmer F, Marks H, Bock C, Gnirke A, 480   Meissner A et al. 2012. Sequential ChIP-bisulfite sequencing enables direct genome-scale investigation of chromatin and DNA methylation cross-talk. *Genome Res* **22**: 1128-1138.

Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. 2013. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding 485   proteins and nucleosome position. *Nat Methods* **10**: 1213-1218.

Chatterjee A, Rodger EJ, Stockwell PA, Weeks RJ, Morison IM. 2012. Technical considerations for reduced representation bisulfite sequencing with multiplexed libraries. *Journal of biomedicine & biotechnology* **2012**: 741542.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. 490   *Bioinformatics* **34**: i884-i890.

Corces MR, Buenrostro JD, Wu B, Greenside PG, Chan SM, Koenig JL, Snyder MP, Pritchard JK, Kundaje A, Greenleaf WJ et al. 2016. Lineage-specific and single-cell chromatin accessibility charts human hematopoiesis and leukemia evolution. *Nat Genet* **48**: 1193-1203.

495   Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, Satpathy AT, Rubin AJ, Montine KS, Wu B et al. 2017. An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues. *Nat Methods* **14**: 959-962.

Cusanovich DA, Daza R, Adey A, Pliner H, Christiansen L, Gunderson KL, Steemers FJ, 500   Trapnell C, Shendure J. 2015. Multiplex single-cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* **348**: 910-914.

Das S, Chadwick BP. 2016. Influence of Repressive Histone and DNA Methylation upon D4Z4 Transcription in Non-Myogenic Cells. *PLoS ONE* **11**: e0160022.

Garrett-Bakelman FE, Sheridan CK, Kacmarczyk TJ, Ishii J, Betel D, Alonso A, Mason CE, 505   Figueroa ME, Melnick AM. 2015. Enhanced reduced representation bisulfite sequencing for assessment of DNA methylation at base pair resolution. *Journal of visualized experiments : JoVE* doi:10.3791/52246: e52246.

Gaulton KJ, Nammo T, Pasquali L, Simon JM, Giresi PG, Fogarty MP, Panhuis TM, Mieczkowski P, Secchi A, Bosco D et al. 2010. A map of open chromatin in human pancreatic islets. *Nat Genet* **42**: 255-259.

Heintzman ND, Hon GC, Hawkins RD, Kheradpour P, Stark A, Harp LF, Ye Z, Lee LK, Stuart RK, Ching CW et al. 2009. Histone modifications at human enhancers reflect global cell-type-specific gene expression. *Nature* **459**: 108-112.

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass

CK. 2010. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**: 576-589.

Horard B, Eymery A, Fourel G, Vassetzky N, Puechberty J, Roizes G, Lebrigand K, Barbry P, Laugraud A, Gautier C et al. 2009. Global analysis of DNA methylation and transcription of human repetitive sequences. *Epigenetics* **4**: 339-350.

Jorda M, Diez-Villanueva A, Mallona I, Martin B, Lois S, Barrera V, Esteller M, Vavouri T, Peinado MA. 2017. The epigenetic landscape of Alu repeats delineates the structural and functional genomic architecture of colon cancer cells. *Genome Res* **27**: 118-132.

Karimi MM, Goyal P, Maksakova IA, Bilenky M, Leung D, Tang JX, Shinkai Y, Mager DL, Jones S, Hirst M et al. 2011. DNA methylation and SETDB1/H3K9me3 regulate predominantly

distinct sets of genes, retroelements, and chimeric transcripts in mESCs. *Cell Stem Cell* **8**: 676-687.

Kelly TK, Liu Y, Lay FD, Liang G, Berman BP, Jones PA. 2012. Genome-wide mapping of nucleosome positioning and DNA methylation within individual DNA molecules. *Genome Res* **22**: 2497-2506.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory requirements. *Nat Methods* **12**: 357-360.

Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* **27**: 1571-1572.

Lay FD, Liu Y, Kelly TK, Witt H, Farnham PJ, Jones PA, Berman BP. 2015. The role of DNA

methylation in directing the functional organization of the cancer epigenome. *Genome Res* **25**: 467-477.

Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**: 589-595.

Liang G, Chan MF, Tomigahara Y, Tsai YC, Gonzales FA, Li E, Laird PW, Jones PA. 2002.

Cooperativity between DNA methyltransferases in the maintenance methylation of repetitive elements. *Mol Cell Biol* **22**: 480-491.

Liao Y, Smyth GK, Shi W. 2013. The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic Acids Res* **41**: e108.

Lindroth AM, Park YJ, McLean CM, Dokshin GA, Persson JM, Herman H, Pasini D, Miro X,

Donohoe ME, Lee JT et al. 2008. Antagonism between DNA and H3K27 Methylation at the Imprinted Rasgrf1 Locus. *PLoS Genetics* **4**: e1000145.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Lu H, Yuan Z, Tan T, Wang J, Zhang J, Luo HJ, Xia Y, Ji W, Gao F. 2015. Improved

tagmentation-based whole-genome bisulfite sequencing for input DNA from less than 100 mammalian cells. *Epigenomics* **7**: 47-56.

Maurano MT, Wang H, John S, Shafer A, Canfield T, Lee K, Stamatoyannopoulos JA. 2015. Role of DNA Methylation in Modulating Transcription Factor Occupancy. *Cell Rep* **12**: 1184-1195.

Meissner A, Gnirke A, Bell GW, Ramsahoye B, Lander ES, Jaenisch R. 2005. Reduced representation bisulfite sequencing for comparative high-resolution DNA methylation analysis. *Nucleic Acids Res* **33**: 5868-5877.

Mulqueen RM, Pokholok D, Norberg SJ, Torkenczy KA, Fields AJ, Sun D, Sinnamon JR, Shendure J, Trapnell C, O'Roak BJ et al. 2018. Highly scalable generation of DNA methylation profiles in single cells. *Nat Biotechnol* **36**: 428-431.

Picelli S, Björklund ÅK, Reinius B, Sagasser S, Winberg G, Sandberg R. 2014. Tn5 transposase and tagmentation procedures for massively scaled sequencing projects. *Genome Research* **24**: 2033-2040.

Rahbari R, Sheahan T, Modes V, Collier P, Macfarlane C, Badge RM. 2009. A novel L1 retrotransposon marker for HeLa cell line identification. *Biotechniques* **46**: 277-284.

Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dundar F, Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**: W160-165.

Rhee I, Bachman KE, Park BH, Jair KW, Yen RW, Schuebel KE, Cui H, Feinberg AP, Lengauer C, Kinzler KW et al. 2002. DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* **416**: 552-556.

Rhee I, Jair KW, Yen RWC, Lengauer C, Herman JG, Kinzler KW, Vogelstein B, Baylin SB, Schuebel KE. 2000. CpG methylation is maintained in human cancer cells lacking DNMT1. *Nature* **404**: 1003-1007.

Roadmap Epigenomics C, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, Kheradpour P, Zhang Z, Wang J et al. 2015. Integrative analysis of 111 reference human epigenomes. *Nature* **518**: 317-330.

Rodriguez JM, Maietta P, Ezkurdia I, Pietrelli A, Wesselink JJ, Lopez G, Valencia A, Tress ML. 2013. APPRIS: annotation of principal and alternative splice isoforms. *Nucleic Acids Res* **41**: D110-117.

Schmidl C, Rendeiro AF, Sheffield NC, Bock C. 2015. ChIPmentation: fast, robust, low-input ChIP-seq for histones and transcription factors. *Nat Methods* **12**: 963-965.

Song CX, Szulwach KE, Dai Q, Fu Y, Mao SQ, Lin L, Street C, Li Y, Poidevin M, Wu H et al. 2013. Genome-wide profiling of 5-formylcytosine reveals its roles in epigenetic priming. *Cell* **153**: 678-691.

Stark R, Brown G. 2018. DiffBind: Differential binding analysis of ChIP- Seq peak data. In *Bioconductor*.

Statham AL, Robinson MD, Song JZ, Coolen MW, Stirzaker C, Clark SJ. 2012. Bisulfite sequencing of chromatin immunoprecipitated DNA (BisChIP-seq) directly informs methylation status of histone-modified DNA. *Genome Res* **22**: 1120-1127.

Suzuki M, Liao W, Wos F, Johnston AD, DeGrazia J, Ishii J, Bloom T, Zody MC, Germer S, Greally JM. 2018. Whole-genome bisulfite sequencing with improved accuracy and cost. *Genome Res* **28**: 1364-1371.

Thurman RE, Rynes E, Humbert R, Vierstra J, Maurano MT, Haugen E, Sheffield NC, Stergachis AB, Wang H, Vernot B et al. 2012. The accessible chromatin landscape of the human genome. *Nature* **489**: 75-82.

Walter M, Teissandier A, Perez-Palacios R, Bourc'his D. 2016. An epigenetic switch ensures transposon repression upon dynamic loss of DNA methylation in embryonic stem cells. *eLife* **5**.

Wu H, Wu X, Zhang Y. 2016. Base-resolution profiling of active DNA demethylation using MAB-seq and caMAB-seq. *Nature protocols* **11**: 1081-1100.

Yin Y, Morgunova E, Jolma A, Kaasinen E, Sahu B, Khund-Sayeed S, Das PK, Kivioja T, Dave K, Zhong F et al. 2017. Impact of cytosine methylation on DNA binding specificities of human transcription factors. *Science* **356**.

Young L, Sung J, Stacey G, Masters JR. 2010. Detection of Mycoplasma in cell cultures. *Nature protocols* **5**: 929-934.

Yu G, Wang LG, He QY. 2015. ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization. *Bioinformatics* **31**: 2382-2383.

610 Yu M, Hon GC, Szulwach KE, Song CX, Zhang L, Kim A, Li X, Dai Q, Shen Y, Park B et al. 2012. Base-Resolution Analysis of 5-Hydroxymethylcytosine in the Mammalian Genome. *Cell* **149**: 1368–1380.

Zaret KS, Carroll JS. 2011. Pioneer transcription factors: establishing competence for gene expression. *Genes Dev* **25**: 2227-2241.

Zaret KS, Mango SE. 2016. Pioneer transcription factors, chromatin dynamics, and cell fate
615 control. *Curr Opin Genet Dev* **37**: 76-81.

Zhu LJ, Gazin C, Lawson ND, Pages H, Lin SM, Lapointe DS, Green MR. 2010. ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data. *BMC bioinformatics* **11**: 237.

Zilberman D, Coleman-Derr D, Ballinger T, Henikoff S. 2008. Histone H2A.Z and DNA
620 methylation are mutually antagonistic chromatin marks. *Nature* **456**: 125-129.

625 **FIGURES AND FIGURE LEGENDS:**

Figures 1-4

A Isolate nuclei

B Tagmentation

C Methylated end-repair

D Bisulfite conversion and PCR

E Sequencing

Accessibility

Methylation

CpG islands
Genes

GAPDH
IFFO1

CpG

5mCpG

C to T
conversion

nucleosome

methylated
ME-A

methylated
ME-B

Tn5
transposase

Figure 1: Overview of mATAC-seq; (A) Tn5 carrying methylated oligonucleotides (red and blue segments) is used to (B) perform tagmentation on nuclei at THS sites. (C) Tagmented DNA is end-repaired using 5mdCTP + dDTPs, purified, (D) Bisulfite converted, amplified, and (E) sequenced to measure DNA methylation and accessibility simultaneously; sample data are shown for one region in HCT116 cells. Peak height in accessibility track is proportional to read abundance; bar height in methylation track is proportional to extent of methylation at CpGs.

Figure 2: Comparison of methods; (A) Omni-ATAC and mATAC share a majority of peaks (B) Features at peaks are similar for mATAC and Omni-ATAC. (C) Accessibility in mATAC-seq is comparable to (D) Omni-ATAC-seq at gene bodies +/- 2kb, n=21,305. (E) methylation reported by mATAC is comparable to (F) WGBS at gene bodies +/- 2kb, n=21,305, though WGBS includes data absent from mATAC.

Figure. 3: Accessibility and methylation at peaks; (A, B) significantly changed mATAC-seq hyper-accessible (log$_2$ fold change > 1, q < 0.01, n = 23,310 peaks), hypo-accessible (log$_2$ fold change < -1, q < 0.01, n = 3,166 peaks), and unchanged peaks (| log$_2$ fold change | < 1, q > 0.8 n = 16,170). Motifs enriched in (C) hyper- and (D) hypo-accessible sites compared to unchanged sites. (E) DNA methylation changes at promoters binned by accessibility, reported as the change in methylation ratio of DKO cells relative to HCT116 (DKO/HCT116). (F) mRNA expression changes in DKO cells relative to HCT116, reported as log$_2$ CPM, at genes binned by differential accessibility of their promoters as in (E). q values are for Wilcoxon tests with Benjamini-Hochberg correction.

Figure 4: Combined Accessibility and methylation analysis. (A) DNA methylation at mATAC-seq peaks from HCT116 and DKO cells form 5 distinct clusters by DNA methylation. (B) mRNA expression in $\log_2$ CPM at identified clusters. Features in clusters 1, 4 and 5 are depicted according to (C) accessibility, (D) DNA methylation, (E) activating histone modifications, (F)

665 silencing histone modifications, and (G) CTCF. Motifs enriched in cluster 1 (H), and cluster 4 (I), compared to cluster 5. q values are for Wilcoxon tests with Benjamini-Hochberg correction.