Independent mechanisms for acquired salt tolerance versus growth resumption induced

by mild ethanol pretreatment in *Saccharomyces cerevisiae*

Elizabeth A. McDaniel[a,*], Tara N. Stuecker[a], Manasa Veluvolu[a], Audrey P. Gasch[b,c], Jeffrey A.

Lewis[a,#]

[a] Department of Biological Sciences, University of Arkansas–Fayetteville, Fayetteville, AR

72701

[b] Laboratory of Genetics, University of Wisconsin–Madison, Madison, WI 53713

[c] Great Lakes Bioenergy Research Center, University of Wisconsin–Madison, Madison, WI

53713

[*] Present Address: Department of Bacteriology, University of Wisconsin–Madison, Madison, WI

[#]Corresponding author:

Email: lewisja@uark.edu

ABSTRACT

All living organisms must recognize and respond to various environmental stresses throughout their lifetime. In natural environments, cells frequently encounter fluctuating concentrations of different stressors that can occur in combination or sequentially. Thus, the ability to anticipate an impending stress is likely ecologically relevant. One possible mechanism for anticipating future stress is acquired stress resistance, where cells pre-exposed to a mild sub-lethal dose of stress gain the ability to survive an otherwise lethal dose of stress. We have been leveraging wild strains of *Saccharomyces cerevisiae* to investigate natural variation in the yeast ethanol stress response and its role in acquired stress resistance. Here, we report that a wild vineyard isolate possesses ethanol-induced cross-protection against severe concentrations of salt. Because this phenotype correlates with ethanol-dependent induction of the *ENA* genes, which encode sodium efflux pumps already associated with salt resistance, we hypothesized that variation in *ENA* expression was responsible for differences in acquired salt tolerance across strains. Surprisingly, we found that the *ENA* genes were completely dispensable for ethanol-induced survival of high salt concentrations in the wild vineyard strain. Instead, the *ENA* genes were necessary for the ability to resume growth on high concentrations of salt following a mild ethanol pretreatment. Surprisingly, this growth acclimation phenotype was also shared by the lab yeast strain despite lack of *ENA* induction under this condition. This study underscores that cross protection can affect both viability and growth through distinct mechanisms, both of which likely confer fitness effects that are ecologically relevant.

IMPORTANCE

Microbes in nature frequently experience "boom or bust" cycles of environmental stress.

Thus, microbes that can anticipate the onset of stress would have an advantage. One way

microbes anticipate future stress is through acquired stress resistance, where cells exposed to a

mild dose of one stress gain the ability survive an otherwise lethal dose of a subsequent stress.

In the budding yeast *Saccharomyces cerevisiae*, certain stressors can cross protect against

high salt concentrations, though the mechanisms governing this acquisition of higher stress

resistance are not well understood. In this study, we took advantage of wild yeast strains to

understand the mechanism underlying ethanol-induced cross protection against high salt

concentrations. We found that mild ethanol stress allows cells to resume growth on high salt,

which involves a novel role for a well-studied salt transporter. Overall, this discovery highlights

how leveraging natural variation can provide new insights into well-studied stress defense

mechanisms.

INTRODUCTION

All organisms experience stress and must respond to environmental perturbations throughout their lifetime. Unlike animals, unicellular organisms generally lack the ability to escape stressful environments. Thus, microbes have evolved sophisticated stress defense strategies such as genome rearrangements, small molecule synthesis, and dynamic gene expression programs to enable stress acclimation (1-3). The model eukaryote *Saccharomyces cerevisiae* responds to diverse stresses by coordinating the expression of condition-specific genes with a large, common gene expression program called the environmental stress response (ESR) (4, 5). The ESR encompasses ~15% of the yeast genome, including ~600 repressed genes that are enriched for processes related to protein synthesis and growth, and the ~300 induced genes that encode diverse functions related to stress defense.

The discovery of a coordinated common response to stress suggested a possible mechanism for the long-observed phenomenon of acquired stress resistance and cross protection, where cells pretreated with a mild dose of stress are better able to survive an otherwise lethal dose of severe stress (6-9). In yeast, defective ESR expression correlates with diminished acquired stress resistance, suggesting that stress-activated gene expression changes may serve to protect against future challenges (10, 11). Beyond yeast, acquired stress resistance has been observed in diverse organisms including bacteria, plants, and humans, and has major implications for food production, agriculture, and human health. For example, mild stress induces higher resistance of the food-borne pathogens *Listeria monocytogenes* and *Salmonella typhimurium* to food preservatives (12, 13). Acquired stress resistance has also been implicated in host survival in the form of bile acid tolerance and antibiotic resistance (14, 15). Acquired thermotolerance and drought resistance in agriculturally significant plants is increasingly important due to climate change (16, 17). In humans, short-term fasting protects healthy cells, but not cancer cells, from the toxic effects of chemotherapy drugs (18, 19).

Altogether, understanding how cells are able to acquire further resistance has broad

applications ranging from agriculture and biotechnology to human health and disease.

We have been leveraging natural variation in the yeast ethanol response to better

understand the cellular mechanisms governing acquisition to high stress resistance. While

mutagenesis studies in laboratory strains of yeast have identified genes and processes

necessary for acquired stress resistance (20-23), there are inherent limitations to using a single

strain background (24). In the case of yeast, the S288c laboratory strain historically used for

large-scale mutagenesis screening is a genetic and physiological outlier compared to wild yeast

strains (25, 26). We have previously noted that S288c has an aberrant gene expression

response to ethanol (11, 27), which we hypothesized was responsible for the strain's inability to

acquire resistance to any other stresses following pre-treatment with mild ethanol (10, 11). We

subsequently found that defective induction of antioxidant defenses in response to mild ethanol

was responsible for S288c's inability to acquire further hydrogen peroxide (28).

In the present study we found that in contrast to S288c, mild ethanol stress induces

cross protection against severe salt concentrations in the wild vineyard strain M22. This

phenotype correlated with the induction of the Ena sodium efflux pump system by ethanol in a

wild vineyard isolate, which was not induced in the S288c-derived common laboratory strain.

Because the ENA system has been previously implicated in salt tolerance (29-31), we

hypothesized that variation in *ENA* expression was responsible for phenotypic differences

across strains. Surprisingly, we found that the *ENA* genes were completely dispensable for

ethanol-induced survival of high salt concentrations in the wild strain. Instead, the *ENA* genes

were necessary for a novel growth resumption phenotype that we call "ethanol-induced

acclimation," where mild ethanol stress allows cells to eventually resume growth on high

concentrations of salt. More surprisingly, our common laboratory strain also exhibited ethanol-

dependent acclimation to high salt concentrations, even though *ENA* is not induced by ethanol

in this strain background. Overall, this study demonstrates that cross protection can affect both

viability and growth through distinct mechanisms, both of which likely confer fitness effects that

are ecologically relevant.

RESULTS

Ethanol Induces Cross Protection Against Severe Salt in a Wild Vineyard Strain

We previously observed that in our S288c-derived common laboratory strain, mild

ethanol pretreatment could not induce acquired resistance to severe ethanol concentrations or

cross protect against other stresses (10). In contrast, mild ethanol pretreatment did induce

further ethanol resistance in the vast majority of wild yeast strains (11). Additionally, we recently

discovered that mild ethanol stress can cross protect against oxidative stress in a wild oak strain

(28). Because yeast in nature are likely to experience environmental shifts between high

concentrations of sugars and high ethanol concentrations, we hypothesized that the inability of

ethanol to cross protect against osmotic stress in S288c may be another aberrant acquired

stress resistance phenotype in this strain background. We tested this by examining ethanol-

induced cross protection against severe salt in a wild vineyard strain background (M22). Cross

protection assays were performed by exposing cells to a mild dose of ethanol (5% v/v) for 1

hour, and scoring their survival across a panel of 11 increasingly severe doses of NaCl (see

Materials and Methods).

We found that ethanol pretreatment weakly cross protected against severe NaCl in

S288c (Fig. 1), though cross protection was not completely absent as previously reported (10).

In contrast, ethanol strongly improved M22's ability to survive otherwise lethal salt

concentrations. Notably, S288c had intrinsically higher basal resistance to NaCl. However, the

diminished cross protection phenotype of S288c relative to M22 cannot be explained by the

higher baseline resistance, as mild NaCl pretreatment did strongly increase S288c's NaCl

resistance (Fig. 2). Moreover, the levels of acquired NaCl resistance following mild NaCl

pretreatment were similar for both S288c and M22.

Induction of *ENA1* by Ethanol in a Wild Vineyard Isolate

Because acquired stress resistance relies on stress-activated gene expression changes (10, 11), we hypothesized that the phenotypic differences in cross protection between S288c and M22 may be due to differences in ethanol-responsive gene expression. We analyzed our previous ethanol-responsive transcriptome changes (27), specifically looking for candidate salt resistance genes with higher induction by ethanol treatment in M22 compared to S288c. We noticed that *ENA1* encoding a sodium efflux pump known to be involved in salt resistance (29) showed a 4.7-fold induction by ethanol in M22 versus a 1.4-fold decrease in expression in S288c, placing it within the top 25 genes in terms of magnitude of differential ethanol-responsive expression when comparing M22 and S288c.

The Ena P-type ATPase sodium efflux pumps are known to play a critical role in maintaining Na+ ion homeostasis in high salt conditions (30-32). In many yeast strain backgrounds, the *ENA* locus consists of a tandem array of nearly identical genes that can vary in copy number (33). S288c contains three copies (*ENA1, ENA2,* and *ENA5)*, whereas M22 appears to contain a single copy (34). This single copy is somewhat unusual, in that M22 appears to contain a large 3885-bp deletion in *ENA1* relative to that of S288c, which results in a full-length in-frame fusion of *ENA1* and *ENA2* (34).

ENA copy number has been linked to high salt tolerance (25–27), likely explaining why S288c has higher basal NaCl tolerance than M22. In S288c, the *ENA* genes are lowly expressed under standard growth conditions (24), but are highly induced in response to saline or alkaline pH stresses (23). Including our previous studies, there are currently no reports of *ENA* induction by ethanol stress in the S288c, suggesting that this mode of *ENA* regulation may have been lost in the laboratory strain background (see Discussion).

181 **The ENA System is not Required for Ethanol-Induced Survival in Severe Salt, but is**

182 **Required for Growth Resumption During Salt Acclimation**

183 Because variation in *ENA* expression is linked to basal salt resistance, we hypothesized

184 *ENA1* induction by ethanol may protect M22 from otherwise lethal salt stress. To test this, we

185 deleted the entire *ENA* region in M22 and examined its role in ethanol-induced acquired NaCl

186 resistance. Surprisingly, we found no defect in acquired salt resistance in the M22 *ena∆* mutant

187 (Fig. 3A). One possible explanation is that the ENA system is not necessary for short-term

188 exposure to acute NaCl, but is instead required for long-term survival. Thus, we measured

189 acquired NaCl resistance in the M22 *ena∆* strain over 24 hours. Even after 24 hours, the M22

190 *ena∆* strain acquired NaCl resistance equivalently to the wild-type strain, and had at most a mild

191 basal NaCl resistance defect (Fig. 3A). We did notice that for ethanol pre-treated wild-type cells,

192 the plating density significantly increased over time (Fig. 3B). This suggested that the ethanol

193 pretreatment enabled cells to acclimate and resume growth on high concentrations of NaCl, and

194 that this acclimation phenotype may be Ena dependent.

195 To further examine the role of the Ena in acclimation to high salt stress after ethanol

196 pretreatment, we measured growth in liquid media, with or without mild ethanol pretreatment

197 (Fig. 4A). Ethanol pretreatment allowed wild-type M22 cells to acclimate and resume growth in

198 1.25M NaCl, and this growth resumption was completely abolished in the M22 *ena∆* mutant.

199 These data suggest that two cross protection phenotypes—survival vs. growth—have distinct

200 cellular mechanisms.

201 In light of the distinct requirement of *ENA* for growth resumption on salt but not survival

202 in the M22 background, we examined whether S288c was able to acclimate and resume growth

203 on high salt following ethanol pretreatment. Indeed, ethanol pretreatment induced growth

204 resumption in S288c (Fig. 4B), which was abolished in the S288c *ena∆* mutant. This was

205 somewhat surprising, considering that *ENA1* is not induced by ethanol in S288c under these

206 conditions (see Discussion).

DISCUSSION

In this study, we initially sought to understand how ethanol cross protects against severe salt stress. We found that ethanol-induced cross protection against severe salt was weaker in the common lab strain S288c when compared to the wild vineyard strain M22. We examined previous transcriptional profiling of the yeast ethanol response to identify candidate genes induced by ethanol in M22 but not S288c (27). Based on these data, we identified the ENA system as a prime candidate to test. The ENA system uses the hydrolysis of ATP through P-type ATPases to transport sodium out of the cell against the electrochemical gradient (22), and mutants lacking *ENA* function are salt sensitive (29-31). Interestingly, the *ENA* locus of many yeast strains including S288c and M22 appears to be the result of a recent introgression from *S. paradoxus* and shows significant copy number variation across strains (33). Other strains have a single non-S288c-like *ENA6* gene that does not share sequence similarity to the *ENA* genes from *S. paradoxus* (26, 30). Genetic mapping studies have linked both copy number variation and polymorphisms in the *ENA* region to variation in NaCl and LiCl tolerance (26, 35-37).

Thus, we were somewhat surprised to find that the ENA system of M22 was completely dispensable for survival in high salt following ethanol pretreatment. Instead, the ENA system was required for a novel ethanol-induced cross-protection phenotype that allows for acclimation and subsequent growth resumption in the presence of high salt. Notably, the vast majority of studies examining salt sensitivity phenotypes for *ENA* have been performed by growing cells on salt-containing plates, which cannot easily distinguish between viability and growth. Both phenotypes are likely important in natural environments. Wild yeast cells growing on fruit such as the M22 vineyard strain may experience simultaneous or fluctuating hyperosmotic stress and ethanol stress, which could explain the evolution of cross protection.

Because we were able to separately examine survival and growth, we reassessed the role of the ENA system in S288c. Surprisingly, we found that while ethanol only weakly induced higher survival on high salt in the S288c background, ethanol-induced acclimation to high salt

was similar between the two strains. This ethanol-induced acclimation phenotype in S288c was

also *ENA* dependent, despite the lack of induction of *ENA* by ethanol in this strain. Notably,

ENA is known to be induced by NaCl in the S288c background (38), which is likely necessary

for growth resumption on high salt. Additionally, basal *ENA* expression is higher in S288c

compared to M22 (11, 27), likely due to copy number variation (our S288c-derived laboratory

strain contains three *ENA* copies, while M22 contains a single copy (34)). It is likely that other

ethanol-induced genes and processes are necessary for ethanol-induced acclimation to high

salt concentrations.

The striking induction of the ENA system by ethanol in M22 but not S288c implies

regulatory differences between the two strains. Recently, natural variation in the promoter

region of *ENA6* in a sake strain was shown to increase Ena6p expression and thus increase salt

tolerance (37). In this strain, a 33-bp deletion in the promoter eliminates glucose repression by

eliminating repressor binding sites for the Nrg1p and Mig1/2p transcription factors. In contrast,

we hypothesize that the novel regulation of *ENA1* by ethanol in M22 is likely not due to promoter

variation. Comparing the promoters of *ENA1* between the S288c and M22 backgrounds reveals

two SNPs and a 20-bp AT repeat insertion within a 20-bp poly-AT repeat region. However,

these promoter differences do not alter or introduce any predicted transcription factor binding

sites, suggesting promoter variation is unlikely responsible for the observed expression

differences between the two strains. Instead, it is likely that *trans* regulatory variation is

responsible for the novel induction by ethanol in the M22 background. The phenotypic

consequences of this novel induction of *ENA1* by ethanol in the M22 strain remain an

unresolved question, as S288c exhibits a similar growth resumption phenotype. Nonetheless,

these findings expand our knowledge of the ENA system's role in stress defense mechanisms,

and highlight the power of using natural variation to yield new insight into even previously well-

studied aspects of cellular physiology, such as the ENA system.

MATERIALS AND METHODS

Strains and Growth Conditions

Strains and primers used in this study are listed in Tables 1 and 2, respectively. The entire *ENA* region was deleted by homologous recombination and replacement with a KanMX4 drug resistance marker in the haploid MATa strain BY4741. This strain was used as a genomic template for introducing the *ena1-5Δ*::KanMX4 allele into different strain backgrounds. To generate homozygous *enaΔ* diploids in the S288c strain background, the *ena1-5Δ*::KanMX4 region was amplified and transformed into MATa and MATα haploid derivatives of DBY8268, which were then mated together. To generate homozygous *enaΔ* diploids in the M22 background, the *ena1-5Δ*::KanMX4 region was transformed into the diploid M22 strain, resulting in an *enaΔ* heterozygote. M22 is capable of mating-type switching, and thus sporulation and dissection yielded homozygous *enaΔ* diploids. Homozygous deletions were verified by diagnostic PCR.

All strains were grown in YPD (1% yeast extract, 2% peptone, 2% dextrose) at 30°C with orbital shaking (270 rpm). Optical density was recorded using a Unico spectrophotometer. Sporulation was achieved by growing cells to saturation for 2 days in YPD, harvesting by centrifugation, resuspending in 1% potassium acetate, and incubating for up 3-5 days at 25°C with shaking.

Acquired Stress Resistance Assays

The acquired stress resistance assays were performed as described in (28). Briefly, cells were grown to overnight to saturation, sub-cultured into in 30-mL fresh media, and then grown for at least 8 generations into exponential phase (OD_{600} of 0.3-0.6) to reset any cellular memory of starvation stress (39). Each culture was split and pretreated with either a mild "primary" stress or a mock (equivalent concentration of water) control. Primary stresses included either 5% v/v ethanol or 0.4 M NaCl. Cells were incubated with the pretreatment for 1 hour and then collected

by mild centrifugation at 1,500 $x\,g$ for 3 min to remove the primary stress. Cells were

resuspended in fresh media to an OD_{600} of 0.6, and then diluted 3-fold into a microtiter plate

containing a panel of severe NaCl doses ranging from 1.2 M to 3.2 M (0.2 M increments).

Plates were sealed breathable Rayon films (VWR), and incubated with secondary stress at

30ºC with 800 rpm shaking in a VWR Symphony Incubating Microplate Shaker. Secondary

treatments were for 2h unless otherwise noted. Following secondary treatment, 4 µl of a 50-fold

cell dilution was spotted directly onto YPD agar plates and grown for 48 hours at 30°C. Viability

at each dose was scored using a 4-point semi-quantitative scale that compared survival in each

secondary dose against an unstressed (YPD only) control: 100% viability = 3 points, 50-90%

viability = 2 points, 10-50% viability = 1 point, and 0% viability = 0 points. An overall tolerance

score was calculated as the sum of scores across all 11 stress doses. Acquired stress

resistance assays were performed in biological triplicate, and raw phenotypic data can be found

in Table S3. A detailed acquired stress resistance assay protocol can be found on protocols.io

under doi dx.doi.org/10.17504/protocols.io.g7sbzne. Statistical analyses were performed using

Prism 7 (GraphPad Software).

Ethanol-Induced Growth Resumption Analysis

To assess ethanol-induced growth resumption in the presence of salt, cells were given

mild primary ethanol (5% v/v) or mock pretreatments as described for the acquired resistance

assays. Following 1 hour pretreatment, cells were gently centrifuged at room temperature for 3

minutes at 1500 $x\,g$, and then resuspended in YPD containing 1.25 M NaCl at an OD_{600} of 0.1.

Five ml of each sample was transferred to a glass test tube and incubated at 30ºC at 270 rpm.

The OD_{600} of all samples was then manually measured with a Unico spectrophotometer over

approximately 72 hours. Growth assays were performed in biological triplicate.

ACKNOWLEDGEMENTS

This work was supported in part by National Science Foundation Grant No. IOS-1656602, the

Arkansas Biosciences Institute (Arkansas Settlement Proceeds Act of 2000), and startup funds

provided by the University of Arkansas to JAL, University of Arkansas Honors College Grants to

EAM and MV, and the DOE Great Lakes Bioenergy Research Center (DOE Office of Science

BER DE-FC02-07ER64494) The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication.

REFERENCES

1. Chadha S, Sharma M. 2014. Transposable elements as stress adaptive capacitors induce genomic instability in fungal pathogen Magnaporthe oryzae. PLoS One 9:e94415.

2. Boutte CC, Crosson S. 2013. Bacterial lifestyle shapes stringent response activation. Trends Microbiol 21:174-180.

3. Gasch AP, Werner-Washburne M. 2002. The genomics of yeast responses to environmental stress and starvation. Funct Integr Genomics 2:181-92.

4. Gasch AP, Spellman PT, Kao CM, Carmel-Harel O, Eisen MB, Storz G, Botstein D, Brown PO. 2000. Genomic expression programs in the response of yeast cells to environmental changes. Mol Biol Cell 11:4241-57.

5. Causton HC, Ren B, Koh SS, Harbison CT, Kanin E, Jennings EG, Lee TI, True HL, Lander ES, Young RA. 2001. Remodeling of yeast genome expression in response to environmental changes. Mol Biol Cell 12:323-37.

6. Collinson LP, Dawes IW. 1992. Inducibility of the response of yeast cells to peroxide stress. J Gen Microbiol 138:329-335.

7. Lewis JG, Learmonth RP, Watson K. 1995. Induction of heat, freezing and salt tolerance by heat and salt shock in *Saccharomyces cerevisiae*. Microbiology 141 (Pt 3):687-94.

8. Flattery-O'Brien J, Collinson LP, Dawes IW. 1993. *Saccharomyces cerevisiae* has an inducible response to menadione which differs from that to hydrogen peroxide. J Gen Microbiol 139:501-7.

9. Jamieson DJ. 1992. *Saccharomyces cerevisiae* has distinct adaptive responses to both hydrogen peroxide and menadione. J Bacteriol 174:6678-81.

10. Berry DB, Gasch AP. 2008. Stress-activated genomic expression changes serve a preparative role for impending stress in yeast. Mol Biol Cell 19:4580-4587.

11.   Lewis JA, Elkon IM, McGee MA, Higbee AJ, Gasch AP. 2010. Exploiting natural

variation in *Saccharomyces cerevisiae* to identify genes for increased ethanol

resistance. Genetics 186:1197-205.

12.   Lou Y, Yousef AE. 1997. Adaptation to sublethal environmental stresses protects

Listeria monocytogenes against lethal preservation factors. Appl Environ Microbiol

63:1252-5.

13.   Leyer GJ, Johnson EA. 1993. Acid adaptation induces cross-protection against

environmental stresses in *Salmonella typhimurium*. Appl Environ Microbiol 59:1842-7.

14.   Leverrier P, Dimova D, Pichereau V, Auffray Y, Boyaval P, Jan G. 2003. Susceptibility

and adaptive response to bile salts in *Propionibacterium freudenreichii*: physiological

and proteomic analysis. Appl Environ Microbiol 69:3809-18.

15.   Mathieu A, Fleurier S, Frenoy A, Dairou J, Bredeche MF, Sanchez-Vizuete P, Song X,

Matic I. 2016. Discovery and function of a general core hormetic stress response in *E.

coli* induced by sublethal concentrations of antibiotics. Cell Rep 17:46-57.

16.   Baurle I. 2016. Plant heat adaptation: priming in response to heat stress. F1000Res

5:694

17.   Ding Y, Fromm M, Avramova Z. 2012. Multiple exposures to drought 'train'

transcriptional responses in *Arabidopsis*. Nat Commun 3:740.

18.   Raffaghello L, Lee C, Safdie FM, Wei M, Madia F, Bianchi G, Longo VD. 2008.

Starvation-dependent differential stress resistance protects normal but not cancer cells

against high-dose chemotherapy. Proc Natl Acad Sci U S A 105:8215-20.

19.   Raffaghello L, Safdie F, Bianchi G, Dorff T, Fontana L, Longo VD. 2010. Fasting and

differential chemotherapy protection in patients. Cell Cycle 9:4474-6.

20.   Berry DB, Guan Q, Hose J, Haroon S, Gebbia M, Heisler LE, Nislow C, Giaever G,

Gasch AP. 2011. Multiple means to the same end: the genetic basis of acquired stress

resistance in yeast. PLoS Genet 7:e1002353.

21.   Gibney PA, Lu C, Caudy AA, Hess DC, Botstein D. 2013. Yeast metabolic and signaling

388   genes are required for heat-shock survival and have little overlap with the heat-induced

389   genes. Proc Natl Acad Sci U S A 110:E4393-402.

390   22.   Kelley R, Ideker T. 2009. Genome-wide fitness and expression profiling implicate Mga2

391   in adaptation to hydrogen peroxide. PLoS Genet 5:e1000488.

392   23.   Zakrzewska A, van Eikenhorst G, Burggraaff JE, Vis DJ, Hoefsloot H, Delneri D, Oliver

393   SG, Brul S, Smits GJ. 2011. Genome-wide analysis of yeast stress survival and

394   tolerance acquisition to analyze the central trade-off between growth rate and cellular

395   robustness. Mol Biol Cell 22:4435-46.

396   24.   Gasch AP, Payseur BA, Pool JE. 2016. The power of natural cariation for model

397   organism biology. Trends Genet 32:147-154.

398   25.   Kvitek DJ, Will JL, Gasch AP. 2008. Variations in stress sensitivity and genomic

399   expression in diverse *S. cerevisiae* isolates. PLoS Genet 4:e1000223.

400   26.   Warringer J, Zorgo E, Cubillos FA, Zia A, Gjuvsland A, Simpson JT, Forsmark A, Durbin

401   R, Omholt SW, Louis EJ, Liti G, Moses A, Blomberg A. 2011. Trait variation in yeast is

402   defined by population history. PLoS Genet 7:e1002111.

403   27.   Lewis JA, Broman AT, Will J, Gasch AP. 2014. Genetic architecture of ethanol-

404   responsive transcriptome variation in *Saccharomyces cerevisiae* strains. Genetics

405   198:369-382.

406   28.   Stuecker TN, Scholes AN, Lewis JA. 2018. Linkage mapping of yeast cross protection

407   connects gene expression variation to a higher-order organismal trait. PLoS Genet

408   14:e1007335.

409   29.   Haro R, Garciadeblas B, Rodriguez-Navarro A. 1991. A novel P-type ATPase from yeast

410   involved in sodium transport. FEBS Lett 291:189-191.

411   30.   Daran-Lapujade P, Daran JM, Luttik MA, Almering MJ, Pronk JT, Kotter P. 2009. An

412   atypical *PMR2* locus is responsible for hypersensitivity to sodium and lithium cations in

the laboratory strain *Saccharomyces cerevisiae* CEN.PK113-7D. FEMS Yeast Res

9:789-792.

31.   Kinclova-Zimmermannova O, Gaskova D, Sychrova H. 2006. The Na+,K+/H+ -antiporter

Nha1 influences the plasma membrane potential of *Saccharomyces cerevisiae*. FEMS

Yeast Res 6:792-800.

32.   Wieland J, Nitsche AM, Strayle J, Steiner H, Rudolph HK. 1995. The PMR2 gene cluster

encodes functionally distinct isoforms of a putative Na+ pump in the yeast plasma

membrane. EMBO J 14:3870-3882.

33.   Strope PK, Skelly DA, Kozmin SG, Mahadevan G, Stone EA, Magwene PM, Dietrich FS,

McCusker JH. 2015. The 100-genomes strains, an *S. cerevisiae* resource that

illuminates its natural phenotypic and genotypic variation and emergence as an

opportunistic pathogen. Genome Res 25:762-774.

34.   Doniger SW, Kim HS, Swain D, Corcuera D, Williams M, Yang SP, Fay JC. 2008. A

catalog of neutral and deleterious polymorphism in yeast. PLoS Genet 4:e1000183.

35.   Kim HS, Fay JC. 2007. Genetic variation in the cysteine biosynthesis pathway causes

sensitivity to pharmacological compounds. Proc Natl Acad Sci U S A 104:19387-19391.

36.   Treusch S, Albert FW, Bloom JS, Kotenko IE, Kruglyak L. 2015. Genetic mapping of

MAPK-mediated complex traits across *S. cerevisiae*. PLoS Genet 11:e1004913.

37.   Sirr A, Scott AC, Cromie GA, Ludlow CL, Ahyong V, Morgan TS, Gilbert T, Dudley AM.

2018. Natural variation in *SER1* and *ENA6* underlie condition-specific growth defects in

Saccharomyces cerevisiae. G3 (Bethesda) 8:239-251.

38.   Garciadeblas B, Rubio F, Quintero FJ, Banuelos MA, Haro R, Rodriguez-Navarro A.

1993. Differential expression of two genes encoding isoforms of the ATPase involved in

sodium efflux in *Saccharomyces cerevisiae*. Mol Gen Genet 236:363-368.

39.   Guan Q, Haroon S, Bravo DG, Will JL, Gasch AP. 2012. Cellular memory of acquired

stress resistance in *Saccharomyces cerevisiae*. Genetics 192:495-505.

40.   Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, Andre B, Bangham R,

Benito R, Boeke JD, Bussey H, Chu AM, Connelly C, Davis K, Dietrich F, Dow SW, El

Bakkoury M, Foury F, Friend SH, Gentalen E, Giaever G, Hegemann JH, Jones T, Laub

M, Liao H, Liebundguth N, Lockhart DJ, Lucau-Danila A, Lussier M, M'Rabet N, Menard

P, Mittmann M, Pai C, Rebischung C, Revuelta JL, Riles L, Roberts CJ, Ross-

MacDonald P, Scherens B, Snyder M, Sookhai-Mahadeo S, Storms RK, Veronneau S,

Voet M, Volckaert G, Ward TR, Wysocki R, Yen GS, Yu K, Zimmermann K, Philippsen

P, et al. 1999. Functional characterization of the *S. cerevisiae* genome by gene deletion

and parallel analysis. Science 285:901-906.

465 **TABLES**

466 **Table 1. Strains used in this study.**

Strain	Background	Genotype	Source
DBY8268	S288c (lab strain)	*MATa/MATα ura3-52/ura3Δ ho/ho GAL2/GAL2*	David Botstein
JL187	BY4741	*MATa his3Δ leu2Δ met15Δ ura3Δ ena1-5Δ::kanMX*	This study
JL505	DBY8268	*MATa ho GAL2 ura3-52 or ura3Δ*	
JL506	DBY8268	*MATα ho GAL2 ura3-52 or ura3Δ*	
JL1127	DBY8268	*MATa ho GAL2 ura3-52 or ura3Δ ena1-5Δ::KanMX*	This study
JL1128	DBY8268	*MATα ho GAL2 ura3-52 or ura3Δ ena1-5Δ::KanMX*	This study
JL1131	DBY8268	*MATa/MATα ho/ho GAL2/GAL2 ura3-52/ura3Δ ena1-5Δ::KanMX/ena1-5Δ::KanMX*	This study
M22	wild vineyard strain	*MATa/MATα*	Robert Mortimer
JL213	M22	*MATa/MATα ena1-5Δ::KanMX/ena1-5Δ::KanMX*	This study
JL214	M22	*MATa/MATα ena1-5Δ::KanMX/ENA1-5*	This study

468 **Table 2. Primers used in this study.**

Primer Name	Sequence	Notes
ENA1_KanMX_up_484	CATTTATTTCCTACTTCTATGACGTTTGTTAGGGCAGGGATGTAGTACGCTGCAGGTCGAC	Contains 45-bp homology to a region 448-bp upstream of the *ENA1* start codon. Paired with "ENA5_KanMX_down_92" to amplify the KanMX4 for generating *ena1-5ΔKanMX4*.
ENA5_KanMX_down_92	CTCATTACCTAAATTTGTTTATGTTCGGTAGCCCTAAAGGAGCTTCATCGATGAATTCGAGCTCG	Contains 45-bp homology to a region 92-bp downstream of the *ENA5* stop codon. Paired with "ENA5_KanMX_up_448" to amplify the KanMX4 for generating *ena1-5ΔKanMX4*.
kanB	CTGCAGCGAGGAGCCGTAAT	Common MX4 cassette primer from (40). Paired with "ENA 1kb up F" to verify insertion of KanMX4 into the *ENA* locus.
ENA 1kb up F	GTCAATATTTTAGGGTTATCGGT	Anneals 1-kb upstream of the *ENA1* start codon in S288c. Paired with "ENA 1kb down R" to amplify *ena1-5ΔKanMX* (3 kb product).
ENA 1kb down F	GTCAATATTTTAGGGTTATCGGT	Anneals 1-kb downstream of the *ENA5* stop codon in S288c. Paired with "ENA 1kb up F" to amplify ena1-5ΔKanMX (3 kb product).
ENA1_qPCR_up	ACACTGACAGCCCAGTCAAGGAAT	Anneals 2,921-bp downstream of the *ENA1* start codon. Paired with "ENA1_qPCR_down" to amplify 181-bp product. Used to verify loss of *ENA1* during *ena1-5ΔKanMX* mutant construction.
ENA1_qPCR_down	ATTGTGAATGCAATGGCGAGACCC	Anneals 3,198-bp downstream of ENA1 start codon. Paired with "ENA1_qPCR_up" to amplify 181-bp product. Used to verify loss of *ENA1* during *ena1-5ΔKanMX* mutant construction.

470 **FIGURES**

Fig. 1. Ethanol induces strong cross protection against severe salt in a wild vineyard

isolate. A) A representative NaCl cross protection assay. S288c (DBY8268 lab strain) and M22

(wild vineyard strain) were exposed to mild 5% ethanol or a mock (5% water) pretreatment for 1

hour, washed, and then exposed to 11 increasingly severe doses of NaCl for 2 hours before

plating to score viability. **B)** NaCl tolerance scores were calculated from the viability across each

of the 11 doses (see Methods). Error bars denote the standard deviation of four independent

biological replicates. Asterisks indicate significantly higher resistance in ethanol-pretreated

versus mock-pretreated cells (*** *P* < 0.001, ns = not significant (*P* > 0.05), *t*-test).

Fig. 2. NaCl allows acquisition of even higher NaCl resistance in both the lab and

vineyard strains. A) A representative acquired NaCl resistance assay is shown for both S288c

(DBY8268 lab strain) and M22 (wild vineyard strain). Cells were split and exposed to either 0.4

M NaCl or a mock (media only) pretreatment for 1 hour, washed, exposed to 11 doses of severe

NaCl for 2 hours, and then plated to score viability. **B)** Salt tolerance scores across each of the

11 doses are plotted as the mean and standard deviation of four independent biological

replicates. Asterisks indicate significantly higher resistance in NaCl-pretreated versus mock-

pretreated cells (* $P < 0.05$, *** $P < 0.001$, *t*-test).

Fig. 3. The ENA system is not necessary for ethanol-induced survival against severe salt.

A) Representative NaCl cross protection assays are shown with increasing amounts of time in

the severe secondary stress doses for Wild Type M22 and the M22 *ena∆* mutant. **B)** Salt

tolerance scores across each of the 11 doses were calculated for each of the timepoints from

Panel A. Error bars denote the standard deviation of three independent biological replicates.

Asterisks indicate timepoints with significantly higher resistance in M22 compared to the *ena∆*

mutant (*$P < 0.01$, *t*-test).

Fig. 4. The ENA system is required for ethanol-induced growth resumption in severe salt in both S288c and M22. A) M22 and M22 *enaΔ (*JL213) and **B)** S288c (JL505) and S288c *enaΔ* (JL1131) were exposed to mild 5% ethanol or a mock (5% water) pretreatment for 1 hour, washed, and then resuspended in YPD containing 1.2 M NaCl. Growth was then measured over the course of 3 days. Error bars denote the standard deviation of three independent biological replicates.