Glass confers rhabdomeric photoreceptor identity in

Drosophila, but not across all metazoans

- 4 F. Javier Bernardo-Garcia^{1,2}, Maryam Syed¹, Gáspár Jékely^{3,*} and Simon G. Sprecher^{1,*}
- ¹Department of Biology, University of Fribourg, Chemin du Musée 10, CH-1700 Fribourg,
- 7 Switzerland

1

2

3

5

11

- 8 ²Department of Biochemistry and Biophysics, University of California, San Francisco, San
- 9 Francisco, CA 94158, USA
- ³Living Systems Institute, University of Exeter, Stocker Road, Exeter EX4 4QD, United Kingdom
- 12 *corresponding authors: g.jekely@exeter.ac.uk, simon.sprecher@unifr.ch

ABSTRACT

Across metazoans, visual systems employ different types of photoreceptor neurons to detect light. These include rhabdomeric PRs, which exist in distantly related phyla and possess an evolutionarily conserved phototransduction cascade. While the development of rhabdomeric PRs has been thoroughly studied in the fruit fly *Drosophila melanogaster*, we still know very little about how they form in other species. To investigate this question, we tested whether the transcription factor Glass, which is crucial for instructing rhabdomeric PR formation in *Drosophila*, may play a similar role in other metazoans. Glass homologues exist throughout the animal kingdom, indicating that this protein evolved prior to the metazoan radiation. Interestingly, our work indicates that *glass* is not expressed in rhabdomeric photoreceptors in the planarian *Schmidtea mediterranea* nor in the annelid *Platynereis dumerilii*. Combined with a comparative analysis of the Glass DNA-binding domain, our data suggest that the fate of rhabdomeric PRs is controlled by Glass-dependent and Glass-independent mechanisms in different animal clades.

Keywords: Glass, transcription factor, *Drosophila*, *Schmidtea*, *Platynereis*, photoreceptor development, rhabdomeric photoreceptors, eye development, evolutionary conservation

INTRODUCTION

Most animals can detect visual cues, which provide them with detailed information about their environment. This information may include the shape of nearby objects, colours, movements or the day–night cycle, and it is relevant for surviving. As a consequence, animals have evolved various types of photoreceptor neurons (PRs) such as ciliary and rhabdomeric PRs [1, 2], which play different roles in different animal species. For instance, rhabdomeric PRs are critical for image-forming vision (e.g. in *Drosophila* compound eye PRs) and for identifying the direction of a light source (e.g. in the planarian *Schmidtea mediteranea* and in the annelid *Platynereis dumerilii*) [3-5]. Nevertheless, in the case of most metazoan clades, very little is known about how rhabdomeric PRs develop.

Interestingly, all known rhabdomeric PRs appear to use a similar assortment of phototransduction proteins. These PRs possess rhabdomeric-type opsins that can modify their spatial conformation upon light stimulation, which allows them to activate Gaq. Gaq signals through phospholipase C (PLC) causing the opening of cation channels on the cytoplasmic membrane of PRs and thus leading to the formation of action potentials. This light-sensing machinery is present in distantly related animal phyla [1, 6, 7], including vertebrates (due to the 'intrinsically photosensitive retinal ganglion cells', ipRGCs [8]), which poses the question of to what degree the development of rhabdomeric PRs is evolutionarily conserved. Is the acquisition of the rhabdomeric phototransduction cascade regulated by a similar set of transcription factors across metazoans? Using the fruit fly Drosophila melanogaster as a model system we have recently shown that the zinc finger transcription factor Glass acts as critical cell fate selector by directing the maturation of PR precursors into adult, light-sensing PRs. In Drosophila, Glass is required for the expression of virtually all the phototransduction proteins [9], and it regulates the development of all types of rhabdomeric PRs (including those in the Bolwig organ, the ocelli, and the compound eye) [10-12]. Therefore, we investigated whether Glass may also be involved in rhabdomeric PR differentiation in other animal species.

The planarian *Schmidtea* and the annelid *Platynereis* are emerging model organisms whose visual systems have been well characterised [3, 5, 13-19]. Interestingly, by analysing recently published single-cell sequencing data of *Schmidtea* we found that *glass* is not expressed in rhabdomeric PRs in this species. Moreover, using *in situ* hybridisation we could not detect *glass* expression in rhabdomeric PRs in *Platynereis*. Thus, while Glass is critical for the specification of rhabdomeric PR identity in *Drosophila*, the absence of Glass in rhabdomeric PRs in *Schmidtea* and *Platynereis* supports that different genetic programmes are required for controlling rhabdomeric PR cell fate in different animal clades. Therefore, while the initial specification of the eye field appears to be controlled by an evolutionarily conserved group of transcription factors (called the retinal determination network, RDN) [17, 20, 21], the

subsequent steps that diversify distinct cell types, including rhabdomeric PRs, are likely controlled by diverse developmental programmes.

METHODS

Phylogenetic analysis

We searched for protein sequences similar to *Drosophila* Glass [22] and *Platynereis* Glass [23] (see sequences in Additional file 1 and Additional file 2) by using NCBI BLAST [24] and the *Schmidtea mediterranea* Genome Database [25]. Redundant sequences were removed from the collection using CD-HIT with an identity cutoff of 90% [26]. To obtain cluster maps based on all against all pairwise similarity, we used CLANS2 with the BLOSUM62 matrix and a *p*-value cutoff of 1e–60 [27]. For phylogenetic tree construction we selected a non-derived set of sequences from the *glass* cluster and aligned them with MUSCLE [28]. Sequences shorter than 300 amino acids were removed prior to the alignment. We trimmed the alignments with TrimAI in 'Automated 1' mode [29]. We identified the JTT+I+G4 model as the best by IQ-TREE [30]. Maximum likelihood trees and bootstrap analysis were carried out with IQ-TREE. Trees were visualized with FigTree [31] (for the data corresponding to this analysis, see Additional file 3).

Glass-binding site analysis

We examined a subset of Glass-like protein sequences by aligning them with either BLAST [24] or MUSCLE [28], and analysed them with 'DNA-binding site predictor for Cys₂His₂ Zinc Finger Proteins' [32, 33] (for details on the sequences that we used see Fig. 3 and Additional file 4). To investigate the DNA-binding specificity of each of these candidates, we copied its full amino acid sequence as input, and asked the software to search for Cys₂His₂ domains [32]. Then, we predicted the binding sites for those regions that best aligned with the 4th and the 5th zinc fingers of Glass, which are responsible for recognising its targets *in vivo* [34-37]. We used 'expanded linear SVM' as prediction model.

Animal caretaking

Drosophila melanogaster stocks were cultured at 25 °C in a 12:12 hour light–dark cycle, and we fed them with cornmeal medium (which was supplemented with molasses, fructose and yeast). We used Canton-S as a wild-type strain (courtesy of R. Stocker), *glass-Gal4* (courtesy of S. Kim) [38] and *UAS-mCD8::RFP* (Bloomington Stock Center, no. 32219).

Our wild-type *Platynereis dumerilii* were a mixed population of worms, originally captured in the sea in Naples (Italy) and Arcachon (France). We also used *r-opsin1-GFP* worms (courtesy of F. Raible) [14]. These animals were kept in sea water at 22 °C, in a 16:8 hours light–dark cycle. We maintained them synchronised to an artificial moon cycle, induced by slightly increasing the light intensity at night for 1 week per month (using a 10 W light bulb, to simulate the full moon). *Platynereis* had a varied diet that included *Artemia salina*, *Tetraselmis marina*, fish food and spinach leaves. For our experiments (i.e. *in situ* hybridisation and microinjections) we crossed males and females and collected the fertilised eggs, as previously described [39]. The larvae that hatched from these eggs were kept at 18 °C.

Immunohistochemistry and in situ hybridisation

In the case of *Drosophila* antibody stainings, these were performed on cryosections of *glass>mCD8::RFP* flies, as previously described [9, 40]. We dissected heads (removing the proboscis to improve the penetration of our reagents) and fixed them for 20 minutes with 3.7% formaldehyde in 0.01 M phosphate buffer (PB; pH 7.4). Then, we washed our samples with PBT (Triton X-100 0.3% in PB) and incubated them with a cryoprotectant solution (sucrose 25% in PB) overnight at 4 °C. The following day we embedded the fly heads in OCT, froze them with liquid nitrogen, and cut 14 µm cryosections in the transverse plane. Once the samples were dry, we proceeded to immunostain them. For this, we washed the slides with PBT (this buffer was also used in subsequent washing steps) and incubated them in primary antibody (rabbit anti-DsRed, 1:100, Clontech, no. 632496) at 4 °C overnight. Then, we washed the cryosections and incubated them in secondary antibody (goat anti-rabbit conjugated to Alexa Fluor 568, 1:200, Molecular Probes, no. A-11011) at 4 °C overnight, and washed again the next day. We mounted our samples by using Vectashield that contained DAPI (Vector, H-1200) and took images with a Leica SP5 confocal microscope.

To detect the *glass* transcript in *Drosophila*, we used the ViewRNA *in situ* hybridisation kit of Affimetrix (no. QVT0012) – which is a proprietary method – and proceeded according to the instructions of the company. Briefly, we took head cryosections (as described above for antibody stainings) and ordered a mix of labelled RNA probes against *glass* from Affimetrix. Then, we processed the samples by digesting them with protease QF, and washed with PB and with various commercial solutions included in the kit. We incubated our cryosections with the *glass* probes for 2 hours, at 40 °C. After this, we continued with a series of washing and signal amplification steps, followed by a colour reaction (we used Fast Red as a fluorophore). We finished by washing the samples with PB, and used Vectashield containing DAPI (Vector, H-1200) to cover the slides. Imaging was done with a Leica SP5 confocal microscope.

To perform double in situ hybridisation in Platynereis, we followed – with few modifications – a protocol that has been previously used for characterising the expression pattern of r-opsin1 [16, 41]. For the present work, we also produced an RNA probe against the glass transcript (for details on the glass probe, see Additional file 1). We fixed 3-5 day old larvae in 4% formaldehyde, and we subjected them to a mild proteinase K digestion to improve the penetration of our reagents. These larvae were prehybridised at 65 °C by using a hybridisation mix (Hyb-Mix), containing 50% formamide, 5x saline-sodium citrate buffer (SSC), 50 μg/ml heparin, 0.1% Tween 20, and 5 mg/ml torula RNA. Then, we dissolved the probes against ropsin1 and glass (labelled with either fluorescein-UTP or digoxigenin-UTP) in Hyb-Mix, denatured them at 80 °C for 10 minutes, and added this solution to our samples. We hybridised both probes simultaneously by incubating at 65 °C overnight. Then, we washed the samples at 65 °C with a series of solutions that initially contained 50% formamide and 2x SSCT (obtained from a stock solution with Tween 20 0.1% in 4x SSC), and we progressively decreased the concentration of both formamide and SSCT throughout successive washes. After washing, we placed the larvae at room temperature and proceeded to immunostain them. We detected the two probes sequentially, by using peroxidase-conjugated primary antibodies against fluorescein (1:250, Roche) and digoxigenin (1:50, Roche). To detect the first probe, we incubated our samples overnight at 4 °C in one of these antibodies, washed them with Tris NaCl Tween 20 buffer (TNT; 0.1 M Tris-HCl, 0.15 M NaCl, 0.1% Tween 20; pH 7.5), and started the colour reaction by adding a solution that contained fluorescent tyramide (conjugated to either Cy3 or fluorescein). We controlled the development of the signal by using a fluorescence microscope and, when it was ready, we washed in TNT and stopped the peroxidase activity with H₂O₂. To detect the second probe, we repeated these immunostaining steps similarly. We mounted our samples with 90% glycerol, and scanned them in a confocal microscope (example confocal stacks can be found within the Additional file 5).

Microinjection of glass-Tomato

145

146

147

148149

150151

152

153

154

155156

157158

159

160

161

162163

164165

166167

168169

170171

172173

174

175

176

177

178179

180

181

182

We used an unpublished assembly of the *Platynereis* genome (courtesy of D. Arendt, EMBL Heidelberg) for making a *glass-Tomato* reporter (see Additional file 1 and Additional file 2 for details). We PCR-amplified a fragment of the *Platynereis glass* promoter and cloned it into a plasmid in frame with the tandem dimer version of *Tomato* (courtesy of L. A. Bezares-Calderón) by using sticky end ligation with Apal and SgsI [42]. The fragment that we cloned included a 5,789 bp long upstream sequence, and also the beginning of the Glass coding sequence: the first ATG codon was predicted both by aligning the *Platynereis* version of Glass to the Glass homologues of other species and by using the ATGpr software [43, 44]. For details on the plasmid that we injected, see its annotated sequence in Additional file 6.

For microinjections, we collected freshly fertilised *Platynereis* eggs and proceeded as previously described [14]. Briefly, we removed the jelly of the eggs by digesting with proteinase K and washing with abundant sea water, using a sieve. We diluted the *glass-Tomato* plasmid to a final concentration of about 200 ng/µl, and delivered it into 1-cell embryos with a microinjection setup, by using Femtotip II microcapillaries (Eppendorf). Larvae were kept at 18 °C, and we imaged them with a Leica SP8 confocal microscope to study the expression of the reporter (representative confocal stacks are available in the Additional file 5). The expression of this reporter showed some degree of mosaicism, given that it was not integrated into the genome, which allowed us to investigate the morphology of the individual neurons that expressed it. We investigated over 100 surviving, Tomato-positive *Platynereis* larvae.

RESULTS

Glass homologues are present throughout metazoans

Glass plays a fundamental role for the differentiation of rhabdomeric PRs in the fruit fly [9, 11, 45, 46]. To investigate if it provides a similar function across metazoans, we first decided to search for Glass homologues in other species.

To do this, we obtained Glass-like sequences by using NCBI BLAST [24] and the *Schmidtea mediterranea* Genome Database [25]. We analysed these sequences with the CLANS2 software (using the BLOSUM62 matrix and a *p*-value cutoff of 1e–60) to produce a cluster map (Fig. 1A) [27]. In this type of graph, closely related sequences (represented as points) are clustered together and connected by dark lines. Based on their similarities, we were able to identify multiple Glass homologues across distantly related species. Some more derived sequences (e.g. from *Strongylocentrotus* and *Saccoglossus*) were also clearly supported as Glass homologues in our analysis. Using these data, we constructed a maximum likehood phylogenetic tree for Glass, which was visualised with FigTree (Fig. 1B) [31] (for more details on our analysis, see the Methods section and the Additional file 3). Importantly, our data reveal that Glass homologues are widely present throughout the animal kingdom.

Neither vertebrates nor choanoflagellates have clear Glass homologues

Based on the distribution of Glass homologues, it seems this protein was present in the common ancestor of all metazoans, but not in choanoflagellates (the sister group of metazoans). Intriguingly, we could also not find any Glass homologue in vertebrates (Fig. 2). Since we identified Glass across multiple animal clades, we wondered why we could not find its vertebrate homologue. Several species have fully sequenced, well annotated genomes, like

zebrafish, mice, or humans [47-51]. For this reason, we decided to further investigate the evolutionary conservation of Glass by scrutinising its sequence.

221

222

223224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250251

252253

254255

256

257

258

259

Glass homologues share a distinctive cluster of five Cys₂His₂ zinc fingers in most species (one exception is Caenorhabditis, in which it only has four zinc fingers because the first one is missing). Particularly, the 4th and the 5th zinc fingers are especially important because they are responsible for guiding Glass towards its targets, given that they recognise its DNA-binding motif in vivo, GAARCC [34-37]. Therefore, we modified our bait by using the consensus sequence of either the full cluster of five zinc fingers, or only the 4th and 5th zinc fingers, and we repeated our BLAST search against vertebrates and choanoflagellates. By doing this we obtained results like, for example, ZSCAN22, ZNF253, or KOX 26 in humans, which still showed less similarity to Glass than any of those homologues that we identified in other species (Fig. 3, sequences available in Additional file 4). We also considered the human candidates that appeared annotated as putative Glass orthologues in Flybase via DIOPT [22, 52], including ZNF764, ZNF683, or ZNF500, but, likewise, they aligned poorly with the consensus sequence of the Glass zinc fingers (Fig. 3, sequences available in Additional file 4). Next, we analysed if any of these proteins would be able to functionally substitute Glass by recognising its DNAbinding motif, the GAARCC sequence [34, 35, 37]. For this, we used the online tool 'DNAbinding site predictor for Cys₂His₂ Zinc Finger Proteins', which predicts the DNA-binding behaviour of zinc finger proteins [32, 33]. This software indicates that those Glass-like proteins that exist in vertebrates and choanoflagellates cannot recognise the GAARCC motif, in contrast to the clear Glass homologues that we found in other animals (i.e. in Amphimedon, Schmidtea, Platynereis, Aplysia, Caenorhabditis, Drosophila, Strongylocentrotus and Branchiostoma) (Fig. 3). Consequently, it remains unclear what happened to the glass gene during the evolution of vertebrates: it could be that they lost Glass, or that it severely changed its amino acid sequence and its DNA-binding motif. Intriguingly, similar to Drosophila, some cells in the vertebrate retina also use the rhabdomeric phototransduction cascade – the ipRGCs, which detect irradiance [8] - and, based on our data, it seems highly probable that these cells develop through different mechanisms in *Drosophila* and in vertebrates.

glass is not expressed in rhabdomeric PRs in the Schmidtea eye

Given that Glass is an essential transcription factor for activating the expression of phototransduction proteins in all *Drosophila* PRs [9, 10], we investigated whether Glass has a similar function in other organisms. For this, we tested if it is expressed in PRs in the eye of the planarian *Schmidtea mediterranea*. Planarians typically possess one pair of eyes, located in the head, that mediate light avoidance [5, 17, 53]. Importantly, their eyes contain rhabdomeric PRs, which are evolutionarily homologous to *Drosophila* PRs [1, 17].

Recently, a single-cell transcriptome atlas has been published for *Schmidtea*, and it is available online [18, 19, 54]. Using this database, rhabdomeric PRs can be identified because they form a cluster of non-ciliated neurons that express phototransduction proteins, including the *opsin* gene (Fig. 4A) [19]. Surprisingly, these cells do not co-express Glass (Fig. 4B), suggesting that, in contrast to *Drosophila*, Glass is not important for the function of rhabdomeric PRs in the *Schmidtea* eye.

glass is not expressed in rhabdomeric PRs in the Platynereis eye

260

261

262

263

264

265

266267

268269

270

271

272

273

274

275276

277

278

279

280

281

282283

284

285

286287

288

289

290

291

292293

294

295

296

297

298

We next tested whether Glass is expressed in rhabdomeric PRs in the marine ragworm *Platynereis dumerilii*. The visual system of *Platynereis* has been well studied, both from a molecular and a functional point of view. *Platynereis* possesses two types of bilateral eyes containing rhabdomeric PRs, called the dorsal and ventral eyes (also known as adult and larval eyes, respectively). These two eye types are able to detect the direction of light, thus mediating phototaxis [3, 13-16].

In Drosophila, glass is expressed in all rhabdomeric PRs [12, 55]. We could detect glass expression in the compound eye of adult flies both with in situ hybridisation and with a glass-Gal4 line crossed to UAS-mCD8::RFP (Figs. 5A-B'), which confirms previous data [12, 55]. By contrast, in the case of Platynereis, in situ hybridisations performed in 3-5 day old larvae did not show co-expression of the glass transcript with rhabdomeric opsin 1 (r-opsin1), which is a marker for rhabdomeric PRs in both the dorsal and the ventral eyes [14, 16], indicating that glass is not present in these cells (Figs. 5C-C", also see confocal stacks in Additional file 5). In addition, we also generated a Platynereis glass reporter by cloning 5.7 kb of its upstream sequence into a plasmid, where the glass start codon was in frame with Tomato (a red fluorescent protein). We used this plasmid for transient transgenesis by injecting it in 1-cell embryos containing a stable r-opsin1-GFP insertion [14]. r-opsin1-GFP animals consistently showed strong GFP signal in their dorsal eye PRs, and this signal was weaker in the ventral eye PRs. In the case of the dorsal eyes, all PRs project their rhabdomeres into a pigment cup, and their axons form four nerves that innervate the optic neuropil in the brain [3, 14, 16]. After microinjections, we tested 3-8 day old larvae (slightly older than those that we used for in situ, to guarantee that positive cells had enough fluorescence to distinguish them) but we did not observe co-expression of GFP and Tomato. glass-Tomato-expressing neurons were consistently located in the head of *Platynereis*, distant from the ventral eyes. The expression of glass-Tomato showed some degree of mosaicism due to this reporter not being integrated into the genome, which allowed us to observe the morphology of individually labelled cells in detail. Some of these Tomato-positive cells appeared close to the dorsal eyes, but they did not project any rhabdomere-like extension into the pigment cup, and their axons did not innervate the optic neuropil (Figs. 5D-E", confocal stacks are available in Additional file 5), indicating that they were not part of the eye rhabdomeric PRs. We conclude that, while Glass is expressed in all types of rhabdomeric PRs in *Drosophila*, it is not present in known rhabdomeric PRs in *Platynereis*.

Glass is expressed in Platynereis sensory neurons

Since *glass* is predominantly expressed in PRs in *Drosophila*, we wondered what type of cells express *glass* in *Platynereis*. We observed that most of the neurons that were labelled with the *glass-Tomato* reporter innervated the neurosecretory neuropil (which is ventral to the optic neuropil, Figs. 5D–D") [56], and, interestingly, many of them were bipolar neurons (Fig. 6). These two features are relevant because an ongoing electron microscopy (EM) connectome reconstruction shows that, in *Platynereis* larvae, most neurons are either unipolar or pseudounipolar, and there are very few bipolar neurons [3, 56-59]. Based on their position and on their morphology, all bipolar neurons in this EM reconstruction are considered sensory neurons because they possess distinctive membranous specialisations (called sensory dendrites) that project towards the surface [3, 56-59]. Therefore, it is very likely that a subset of *glass*-expressing cells in *Platynereis* are sensory neurons.

The neurosecretory neuropile of *Platynereis* contains multiple sensory neurons, and it has been characterised both from an anatomical and a molecular point of view [56]. However, it is still unknown whether this region is homologous to any structure of the *Drosophila* brain. Given that *glass* is also required for the development of the corpora cardiaca in *Drosophila* [60], it could be possible that Glass has an evolutionarily conserved function in neurosecretoy cells. In addition, it could also be that Glass regulates the formation of other sensory neurons. Notably, the *Caenorhabditis* homologue of Glass (called CHE-1) is expressed in ASE chemosensory neurons, and it regulates their development [34, 61].

CONCLUSIONS

Remarkably, the earliest steps of eye development are controlled by a group of transcription factors, called the 'retinal determination network' (RDN), which is both required and sufficient for eye formation in distantly related species [20, 62-68]. RDN members, such as Eyeless, Sine oculis, or Eyes absent are important for inducing eye field specification. To achieve this, they establish complex epistatic interactions with each other. These interactions occur similarly across model organisms, suggesting that this is an evolutionarily conserved process [20, 69]. In contrast to the early steps of eye field specification, subsequent mechanisms that specify the cell fate of PRs are not well understood. Here we provide evidence that, during the late stages of eye development, rhabdomeric PRs mature through different mechanisms in different species.

In *Drosophila*, we have recently shown that Sine oculis (a core component of the RDN) directly activates the expression of the transcription factor *glass*, which is crucial for activating the expression of virtually all the phototransduction proteins in all types of *Drosophila* PRs [9, 10, 70]. Based on similarities in their light-sensing machinery, *Drosophila* PRs are considered homologous to the ipRGCs of vertebrates, and also to the rhabdomeric PRs that exist in *Schmidtea* and *Platynereis* [1, 6, 7, 15, 17, 19]. Intriguingly, while we did identify Glass homologues in most metazoans, we could not find a clear Glass homologue in vertebrates. Moreover, our data indicate that *glass* is not expressed in the rhabdomeric PRs of *Schmidtea* or *Platynereis*. This suggests that metazoans have evolved alternative transcriptional pathways to direct the formation of rhabdomeric PRs. One of these pathways requires Glass (e.g. in *Drosophila*), while others do not, (e.g. in vertebrates, *Schmidtea*, or *Platynereis* larvae).

It could be possible that Glass started being expressed in rhabdomeric PRs at some point during the evolution of ecdysozoans and that it became specialised in regulating the differentiation of these cells. Therefore, comparing the differentiation of Glass-expressing and non-Glass-expressing PRs provides a valuable entry point to dissect shared and dissimilar aspects of the developmental programme. Additionally, it would also be interesting to know the identity of Glass-expressing cells for understanding the ancestral function of Glass. The glass transcript is rare and lowly expressed in the Schmidtea single cell transcriptome data that we have currently available [18, 19], and it was also lowly expressed in the single cell transcriptome datasets of Platynereis, to the point of being removed from the analyses of the two papers in which the sequencing was published [71, 72], which makes it impossible to compare the function of glass-expressing cells between different species at this moment. It could be possible that this is because only a few cells in the brain express Glass, and these may not have been included in the samples that were sequenced. Therefore we expect that, in the near future, increasing both the number and the quality such single cell transcriptomes for these and other species will be useful to address several questions about the evolution of specific cell fates. For example, some opsins may have other functions apart from light-sensing [73], and it would be relevant to know if glass regulates the expression of any such opsin outside the *Platynereis* eye (for example), at any stage.

The absence of Glass in rhabdomeric PRs in the eye of some species argues for other transcription factors being capable of activating the expression of phototransduction proteins, however the underlying mechanism remains unknown. Our data support a rather complex scenario for the evolution of rhabdomeric PRs, but future works on the targets of the RDN may help to better understand how rhabdomeric PR identity is regulated.

ABBREVIATIONS ac-Tub: acetylated Tubulin, EM: electron microscopy, PB: phosphate buffer, PBT: phosphate buffer with Triton X-100, PR: photoreceptor neuron, RDN: retinal determination network, ropsin1: rhabdomeric opsin 1, SSC: saline-sodium citrate buffer, SSCT: saline-sodium citrate buffer with Tween 20. **DECLARATIONS** Ethics approval and consent to participate Not applicable. Consent for publication Not applicable. Availability of data and material All data presented in this article are accessible online, as indicated in either the main text or in the Additional file section. **Competing interests** The authors do not declare competing or financial interests. **Funding** This work was funded by the Swiss National Science Foundation (31003A_149499 to S.G.S.) and the European Research Council (ERC-2012-StG 309832-PhotoNaviNet to S.G.S.). Authors' contributions F.J.B.-G., G.J., and S.G.S. conceived the study. F.J.B.-G., G.J. and M.S. performed the experiments. F.J.B.-G., G.J., and S.G.S. wrote the manuscript. **Acknowledgements**

We thank the Bloomington Stock Center, R. Stocker, and S. Kim for fly stocks; F. Raible for the *r-opsin1-GFP Platynereis* strain; D. Arendt for providing access to the draft *Platynereis* genome; and L. A. Bezares-Calderón for plasmids. We are also grateful to C. Adler, J. Cleland, and to our colleagues of the Sprecher and Jékely labs for valuable discussions.

FIGURES

- **Fig. 1: Glass phylogeny**. To identify Glass homologues we searched for Glass-like sequences with BLAST and obtained a cluster map by using all against all pairwise similarity. In this graph, those sequences that are most similar appear clustered together, and connected by a darker line (A). Based on these data, we built a maximum likehood tree for Glass (B) (for further details see the Methods section, the tree file and the sequences that we used for it are included in the Additional file 3).
- **Fig. 2: Glass homologues exist in most animal groups**. Based on sequence comparison (Additional file 4, also see Fig. 3), we infer that *glass* appeared in the common ancestor of all metazoans, and that it has been transmitted to most present-day animals (shown in green on the phylogenetic tree [74]). However, we were not able to identify *glass* in vertebrates.
- Fig. 3: Analysis of the Glass zinc fingers. Generally, Glass homologues possess a cluster of five Cys₂His₂ zinc fingers, each of them containing the following motif: Cys-X_{2.4}-Cys-X₁₂-His- $X_{3.4.5}$ -His. Of these, we compared the sequences of the 4th and the 5th zinc fingers, which are responsible for recognising the DNA Glass-binding motif in PRs in vivo [34-37], from the following species: Amphimedon (Porifera), Schmidtea (Platyhelminthes), Platynereis (Annelida), Aplysia (Mollusca), Caenorhabditis (Nematoda), Drosophila (Arthropoda), Strongylocentrotus (Echinodermata) and Branchiostoma (Cephalochordata). In the table, those amino acids that match the Glass consensus sequence (deduced by aligning the homologues of different species, in the first column) appear on black background. The 3D structure of the DNA-bound Cys₂His₂ domain has been resolved [75], and it is expected that four amino acids per zinc finger directly recognise three base pairs. These amino acids are well evolutionarily conserved across different Glass homologues and, in the sequences that we show, they are no. 10 (D), 12 (S), 13 (T), and 16 (K) in the 4th zinc finger, and no. 38 (Q), 40 (G), 41 (N), and 44 (R) in the 5th zinc finger. Other residues and neighbouring zinc fingers are also expected to contribute to the DNAbinding specificity of Glass [76]. Similarly, we aligned Glass-like proteins from vertebrates (e.g. human) and choanoflagellates (e.g. Salpingoeca) with BLAST [24] and MUSCLE [28], but they showed little similarity to the Glass consensus sequence (shown on the second column). Furthermore, a 'DNA-binding site predictor for Cys₂His₂ Zinc Finger Proteins' has been developed and is available online [32, 33]. This software predicts that, based on their amino acid sequence, all Glass homologues (in the first column) can bind to the same DNA motif:

GAAGCC, which was expected from experimental works in *Drosophila* and *Caenorhabditis* [34, 35]. By contrast, it appears that the Glass-like proteins of vertebrates and choanoflagellates (on the second column) would not be able to recognise this motif. All sequences are available in the Additional file 4.

Fig. 4: *glass* is not expressed in rhabdomeric PRs in *Schmidtea*. These graphs were obtained from the Planarian Digiworm atlas, a single-cell transcriptome database for *Schmidtea mediterranea* [19, 25]. Each point corresponds to one single cell, and they are clustered according to the similarity of their transcriptome. One of the clusters shown, corresponding to non-ciliated neurons, is formed by 14 rhabdomeric PRs which can be identified because of the expression of the *opsin* gene (dd_Smed_v4_15036_0_1, A). However, these PRs do not appear to express the *Schmidtea glass* homologue (annotated as dd_Smed_v4_75162_0_1 in this website [19, 54], B).

Fig. 5: glass is not expressed in rhabdomeric PRs in Platynereis. (A, B) glass is present in all Drosophila rhabdomeric PRs, including those in the compound eye [12, 55]. This can be observed in head cryosections, either by using in situ hybridisation (magenta in A, grevscale in A') or with glass>mCD8::RFP flies (magenta in B, greyscale in B'). In both cases samples were counterstained with DAPI (green). (C-E) In contrast to Drosophila, double in situ hybridisation against the glass (red) and r-opsin1 (green) transcripts shows that glass is not present in Platynereis rhabdomeric PRs. Samples were counterstained with antibodies against acetylated Tubulin (ac-Tub, blue), which is a neuropil marker (C, transversal view of a whole-mounted, 5 day old larva). To the right, close-ups of the dorsal (arrow in C; C',C") and ventral eyes (arrowhead in C; C"", C"") show that glass (in magenta/greyscale) is not expressed in either of these visual organs. Similarly, we found that a microinjected glass-Tomato reporter (magenta/greyscale) was not co-expressed with a stable r-opsin1-GFP insertion (green). Brightfield (BF, greyscale) was imaged as a reference (D-D", dorsal view of a whole mounted, 8 day old larva). The positions of the dorsal and ventral eyes are shown with an arrow and an arrowhead, respectively. Close-ups to the right show how the axons of Tomato and GFPpositive neurons project into two different areas in the brain (D', D"; orthogonal views taken along the Z segment are shown below). As a control, we also imaged an 8 day old, wild-type, uninjected larva to test its autofluorescence (using two excitation laser wavelengths: 552 nm, same as for Tomato; and 488 nm, same as for GFP). Scale bars: 10 µm in C', C'"; 20 µm in D-E; and 50 µm in A, B. Axes: D, dorsal; M, medial; P, posterior; V, ventral.

Fig. 6: Glass-expressing cells in *Platynereis* **include sensory neurons**. When we injected our *glass-Tomato* reporter, we observed that many of the neurons that appeared labelled in the *Platynereis* head were bipolar, located close to the surface, and they often possessed membranous specialisations resembling sensory dendrites (arrows) (A–D). Scale bars: 5 μm.

ADDITIONAL FILES

- Additional file 1: supplementary methods.
- 497 Additional file 2: Platynereis glass supplementary nucleotide sequences (both genomic and
- 498 transcriptomic).

492493

494 495

496

499

501

504

506

509

510511

- Additional file 3: Glass phylogenetic tree and sequences data.
- Additional file 4: subset of glass-like sequences for which the DNA-binding affinity was
- 503 investigated.
- 505 Additional file 5: example confocal stacks.
- 507 Additional file 6: annotated sequence of the glass-tomato plasmid that was used for
- 508 Platynereis microinjections.

REFERENCES

- 512 1. Fain GL, Hardie R, Laughlin SB: Phototransduction and the evolution of photoreceptors.
- 513 Current Biology 2010;20:R114–124.
- 514 2. Nilsson D-E: Photoreceptor evolution: Ancient siblings serve different tasks. Current
- 515 Biology 2005;15:R94-96.
- 516 3. Randel N, Asadulina A, Bezares-Calderon LA, Veraszto C, Williams EA, Conzelmann
- 517 M, Shahidi R, Jekely G: Neuronal connectome of a sensory-motor circuit for visual navigation.
- 518 Elife 2014;3.
- 519 4. Borst A: Drosophila's view on insect vision. Current Biology 2009;19:R36–47.
- 520 5. Paskin TR, Jellies J, Bacher J, Beane WS: Planarian Phototactic Assay Reveals
- 521 Differential Behavioral Responses Based on Wavelength. PloS one 2014;9:e114708.
- 522 6. Provencio I, Warthen DM: Melanopsin, the photopigment of intrinsically photosensitive
- retinal ganglion cells. WIREs: Membrane Transport and Signaling 2012;1:228–237.
- 524 7. Montell C: *Drosophila* visual transduction. Trends in neurosciences 2012;35:356–363.

- 525 8. Hankins MW, Hughes S: Vision: melanopsin as a novel irradiance detector at the heart
- 526 of vision. Current Biology 2014;24:R1055–1057.
- 9. Bernardo-Garcia FJ, Fritsch C, Sprecher SG: The transcription factor Glass links eye
- 528 field specification with photoreceptor differentiation in *Drosophila*. Development 2016;143:1413–
- 529 1423.
- 530 10. Bernardo-Garcia FJ, Humberg T-H, Fritsch C, Sprecher SG: Successive requirement of
- 531 Glass and Hazy for photoreceptor specification and maintenance in Drosophila. Fly
- 532 2017;11:112-120.
- 533 11. Moses K, Ellis MC, Rubin GM: The glass gene encodes a zinc-finger protein required by
- 534 Drosophila photoreceptor cells. Nature 1989;340:531–536.
- 535 12. Ellis MC, O'Neill EM, Rubin GM: Expression of Drosophila glass protein and evidence
- for negative regulation of its activity in non-neuronal cells by another DNA-binding protein.
- 537 Development 1993;119:855-865.
- 538 13. Jékely G, Colombelli J, Hausen H, Guy K, Stelzer E, Nédélec F, Arendt D: Mechanism
- of phototaxis in marine zooplankton. Nature 2008;456:395–399.
- 540 14. Backfisch B, Veedin Rajan VB, Fischer RM, Lohs C, Arboleda E, Tessmar-Raible K,
- 541 Raible F: Stable transgenesis in the marine annelid *Platynereis dumerilii* sheds new light on
- 542 photoreceptor evolution. Proceedings of the National Academy of Sciences of the United States
- 543 of America 2013;110:193–198.
- 544 15. Arendt D, Tessmar-Raible K, Snyman H, Dorresteijn AW, Wittbrodt J: Ciliary
- 545 photoreceptors with a vertebrate-type opsin in an invertebrate brain. Science 2004;306:869-
- 546 871.
- 547 16. Randel N, Bezares-Calderón LA, Gühmann M, Shahidi R, Jékely G: Expression
- 548 dynamics and protein localization of rhabdomeric opsins in *Platynereis* larvae. Integrative and
- 549 comparative biology 2013;53:7–16.
- 550 17. Lapan SW, Reddien PW: Transcriptome analysis of the planarian eye identifies ovo as
- a specific regulator of eye regeneration. Cell reports 2012;2:294-307.
- 552 18. Plass M, Solana J, Wolf FA, Ayoub S, Misios A, Glazar P, Obermayer B, Theis FJ,
- Kocks C, Rajewsky N: Cell type atlas and lineage tree of a whole complex animal by single-cell
- transcriptomics. Science 2018;360.
- 555 19. Fincher CT, Wurtzel O, de Hoog T, Kravarik KM, Reddien PW: Cell type transcriptome
- atlas for the planarian Schmidtea mediterranea. Science 2018;360.

- 557 20. Silver SJ, Rebay I: Signaling circuitries in development: insights from the retinal
- determination gene network. Development 2005;132:3–13.
- 559 21. Arendt D, Tessmar K, Medeiros de Campos-Baptista M-I, Dorresteijn A, Wittbrodt J:
- 560 Development of pigment-cup eyes in the polychaete Platynereis dumerilii and evolutionary
- conservation of larval eyes in Bilateria. Development 2002;129:1143–1154.
- 562 22. Flybase. http://flybase.org/.
- 563 23. Max Planck Institute for Developmental Biology: Platynereis dumerilii transcriptome.
- http://jekely-lab.tuebingen.mpg.de/blast/.
- 565 24. NCBI BLAST service. http://blast.ncbi.nlm.nih.gov/.
- 566 25. Schmidtea mediterranea Genome Database. http://smedgdkc03.stowers.org.
- 567 26. Li W, Godzik A: Cd-hit: a fast program for clustering and comparing large sets of protein
- or nucleotide sequences. Bioinformatics 2006;22:1658-1659.
- 569 27. Frickey T, Lupas A: CLANS: a Java application for visualizing protein families based on
- pairwise similarity. Bioinformatics 2004;20:3702-3704.
- 571 28. Edgar RC: MUSCLE: Multiple sequence alignment with high accuracy and high
- throughput. Nucleic acids research 2004;32:1792–1797.
- 573 29. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T: trimAl: a tool for automated
- alignment trimming in large-scale phylogenetic analyses. Bioinformatics 2009;25:1972-1973.
- 575 30. IQ-TREE software for phylogenomic inference. http://www.igtree.org/.
- 576 31. FigTree viewer of phylogenetic trees. http://tree.bio.ed.ac.uk/software/figtree/.
- 577 32. DNA-binding site predictor for Cys₂His₂ zinc finger proteins. http://zf.princeton.edu/.
- 578 33. Persikov AV, Singh M: De novo prediction of DNA-binding specificities for Cys₂His₂ zinc
- finger proteins. Nucleic acids research 2014;42:97–108.
- 580 34. Etchberger JF, Lorch A, Sleumer MC, Zapf R, Jones SJ, Marra MA, Holt RA, Moerman
- 581 DG, Hobert O: The molecular signature and cis-regulatory architecture of a C. elegans
- 582 gustatory neuron. Genes & development 2007;21:1653–1674.
- 583 35. Naval-Sánchez M, Potier D, Haagen L, Sánchez M, Munck S, Van de Sande B,
- 584 Casares F, Christiaens V, Aerts S: Comparative motif discovery combined with comparative

- 585 transcriptomics yields accurate targetome and enhancer predictions. Genome research
- 586 2013;23:74-88.
- 587 36. O'Neill EM, Ellis MC, Rubin GM, Tjian R: Functional domain analysis of glass, a zinc-
- 588 finger-containing transcription factor in Drosophila. Proceedings of the National Academy of
- Sciences of the United States of America 1995;92:6557–6561.
- 590 37. Enuameh MS, Asriyan Y, Richards A, Christensen RG, Hall VL, Kazemian M, Zhu C,
- 591 Pham H, Cheng Q, Blatti C et al: Global analysis of Drosophila Cys₂-His₂ zinc finger proteins
- 592 reveals a multitude of novel recognition motifs and binding determinants. Genome research
- 593 2013;23:928-940.
- 594 38. Park S, Bustamante EL, Antonova J, McLean GW, Kim SK: Specification of Drosophila
- 595 corpora cardiaca neuroendocrine cells from mesoderm is regulated by Notch signaling. PLoS
- 596 genetics 2011;7.
- 597 39. Hauenschild C, Fischer A: Platynereis dumerilii: Mikroskopische Anatomie,
- 598 Fortpflanzung, Entwicklung. Stuttgart, Germany: Gustav Fischer Verlag; 1969.
- 599 40. Wolff T: Histological techniques for the *Drosophila* eye. Part II: Adult. In: Sullivan W,
- 600 Ashburner M, Hawley RS, editors. Drosophila protocols. Cold Spring Harbor, New York: Cold
- Spring Harbor Laboratory Press; 2000: p. 229–243.
- 602 41. Tessmar-Raible K, Steinmetz PR, Snyman H, Hassel M, Arendt D: Fluorescent two-
- 603 color whole mount in situ hybridization in Platynereis dumerilii (Polychaeta, Annelida), an
- emerging marine molecular model for evolution and development. BioTechniques 2005;39:460-
- 605 462.
- 606 42. Day RN, Davidson MW: The fluorescent protein palette: tools for cellular imaging.
- 607 Chemical Society reviews 2009;38:2887–2921.
- 608 43. ATGpr service. http://atgpr.dbcls.jp/.
- 609 44. Nishikawa T, Ota T, Isogai T: Prediction whether a human cDNA sequence contains
- 610 initiation codon by combining statistical information and similarity with protein sequences.
- 611 Bioinformatics 2000;16:960-967.
- 612 45. Liang X, Mahato S, Hemmerich C, Zelhof AC: Two temporal functions of Glass:
- ommatidium patterning and photoreceptor differentiation. Developmental biology 2016;414:4-
- 614 20.


- 615 46. Morrison CA, Chen H, Cook T, Brown S, Treisman JE: Glass promotes the
- differentiation of neuronal and non-neuronal cell types in the *Drosophila* eye. PLoS genetics
- 617 2018;14.
- 618 47. Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M,
- 619 Evans CA, Holt RA et al: The sequence of the human genome. Science 2001;291:1304–1351.
- 48. Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K,
- 621 Doyle M, FitzHugh W et al: Initial sequencing and analysis of the human genome. Nature
- 622 2001;409:860-921.
- 623 49. Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, Agarwala R,
- 624 Ainscough R, Alexandersson M, An P et al: Initial sequencing and comparative analysis of the
- 625 mouse genome. Nature 2002;420:520–562.
- 626 50. Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE,
- 627 Humphray S, McLaren K, Matthews L et al: The zebrafish reference genome sequence and its
- relationship to the human genome. Nature 2013;496:498–503.
- 629 51. Ensembl genome browser. http://www.ensembl.org/.
- 630 52. Hu Y, Flockhart I, Vinayagam A, Bergwitz C, Berger B, Perrimon N, Mohr SE: An
- 631 integrative approach to ortholog prediction for disease-focused and other functional studies.
- 632 BMC bioinformatics 2011;12:357.
- 633 53. Arees EA: Absence of light response in eyeless planaria. Physiology & behavior
- 634 1986:36:445-449.
- 635 54. Planarian Digiworm atlas. https://digiworm.wi.mit.edu.
- 636 55. Moses K, Rubin GM: Glass encodes a site-specific DNA-binding protein that is
- 637 regulated in response to positional signals in the developing Drosophila eye. Genes &
- 638 development 1991;5:583-593.
- 639 56. Williams EA, Veraszto C, Jasek S, Conzelmann M, Shahidi R, Bauknecht P, Mirabeau
- 640 O, Jekely G: Synaptic and peptidergic connectome of a neurosecretory center in the annelid
- 641 brain. Elife 2017;6.
- 642 57. Bezares-Calderon LA, Berger J, Jasek S, Veraszto C, Mendes S, Guhmann M, Almeda
- R, Shahidi R, Jekely G: Neural circuitry of a polycystin-mediated hydrodynamic startle response
- 644 for predator avoidance. Elife 2018;7.

- 645 58. Randel N, Shahidi R, Veraszto C, Bezares-Calderon LA, Schmidt S, Jekely G: Inter-
- 646 individual stereotypy of the Platynereis larval visual connectome. Elife 2015;4:e08069.
- 647 59. Veraszto C, Ueda N, Bezares-Calderon LA, Panzera A, Williams EA, Shahidi R, Jekely
- 648 G: Ciliomotor circuitry underlying whole-body coordination of ciliary activity in the Platynereis
- 649 larva. Elife 2017;6.
- 650 60. De Velasco B, Shen J, Go S, Hartenstein V: Embryonic development of the Drosophila
- 651 corpus cardiacum, a neuroendocrine gland with similarity to the vertebrate pituitary, is controlled
- by sine oculis and glass. Dev Biol 2004;274:280-294.
- 653 61. Uchida O, Nakano H, Koga M, Ohshima Y: The C. elegans che-1 gene encodes a zinc
- 654 finger transcription factor required for specification of the ASE chemosensory neurons.
- 655 Development 2003;130:1215-1224.
- 656 62. Hoge MA: Another gene in the fourth chromosome of Drosophila. The American
- 657 Naturalist 1915;49:47-49.
- 658 63. Quiring R, Walldorf U, Kloter U, Gehring WJ: Homology of the eyeless gene of
- 659 Drosophila to the Small eye gene in mice and Aniridia in humans. Science 1994;265:785–789.
- 660 64. Halder G, Callaerts P, Flister S, Walldorf U, Kloter U, Gehring WJ: Eyeless initiates the
- expression of both sine oculis and eyes absent during Drosophila compound eye development.
- 662 Development 1998;125:2181-2191.
- 663 65. Halder G, Callaerts P, Gehring WJ: Induction of ectopic eyes by targeted expression of
- the eyeless gene in *Drosophila*. Science 1995;267:1788–1792.
- 665 66. Pignoni F, Hu B, Zavitz KH, Xiao J, Garrity PA, Zipursky SL: The eye-specification
- 666 proteins So and Eya form a complex and regulate multiple steps in Drosophila eye
- development. Cell 1997;91:881–891.
- 668 67. Chow RL, Altmann CR, Lang RA, Hemmati-Brivanlou A: Pax6 induces ectopic eyes in a
- 669 vertebrate. Development 1999;126:4213–4222.
- 670 68. Loosli F, Winkler S, Wittbrodt J: Six3 overexpression initiates the formation of ectopic
- 671 retina. Genes & development 1999;13:649–654.
- 672 69. Donner AL, Maas RL: Conservation and non-conservation of genetic pathways in eye
- specification. The International journal of developmental biology 2004;48:743–753.
- 674 70. Jusiak B, Karandikar UC, Kwak S-J, Wang F, Wang H, Chen R, Mardon G: Regulation
- of Drosophila eye development by the transcription factor Sine oculis. PloS one 2014;9.

- 676 71. Achim K, Eling N, Vergara HM, Bertucci PY, Musser J, Vopalensky P, Brunet T, Collier
- 677 P, Benes V, Marioni JC et al: Whole-Body Single-Cell Sequencing Reveals Transcriptional
- Domains in the Annelid Larval Body. Mol Biol Evol 2018;35:1047-1062.
- 679 72. Achim K, Pettit JB, Saraiva LR, Gavriouchkina D, Larsson T, Arendt D, Marioni JC:
- High-throughput spatial mapping of single-cell RNA-seq data to tissue of origin. Nat Biotechnol
- 681 2015;33:503-509.
- 682 73. Leung NY, Montell C: Unconventional Roles of Opsins. Annu Rev Cell Dev Biol
- 683 2017;33:241-264.

691

- 684 74. Dunn CW, Giribet G, Edgecombe GD, Hejnol A: Animal Phylogeny and Its Evolutionary
- 685 Implications. Annu Rev Ecol Evol S 2014;45:371-+.
- 686 75. Pavletich NP, Pabo CO: Zinc finger-DNA recognition: Crystal structure of a Zif268-DNA
- 687 complex at 2.1 A. Science 1991;252:809–817.
- 688 76. Garton M, Najafabadi HS, Schmitges FW, Radovani E, Hughes TR, Kim PM: A
- 689 structural approach reveals how neighbouring C2H2 zinc fingers influence DNA binding
- 690 specificity. Nucleic acids research 2015;43:9147–9157.



XP_014779223.1_Octopus_bimaculoides

AHA51257.1_Beroe_forskalii

AHA51336.1_Euplokamis_dunlapae

comp416373_c0_seq8_Platynereis_dumerilii glass

64

100

0.2

Figure 2




Figure 3

stid >Amphimedon, https://blast.ncbi.nhm.nih.gov/ XP 019864430.1 CDVGCKCEAETSTITKERETESCERENEGETONLCESCENERENKTE > Schmidtea, http://smedgdkc03.spwers.org/ EGI57499.1 CHESKAESESSTITKETENESCERENEGETOLEESCENERENERENER > Platynereis, http://seety-lab.tuebingen.mpg.de/blast. COMP416373.c0.seq8 CRNGKKAESESSTITKETENESCERENCOKICLLESSCENENERENKTE > Aplysia, https://blast.ncbi.nlm.nih.gov/ XP 005091155.1 CRNGKKAESESSTITKETENESCERENCOKICLLESSCENENERENKTE > Caenorhabditis, https://blast.ncbi.nlm.nih.gov/ NP 001076597.1, CHE-1 CACENORHEESESSTITKETENESTESCERENCOKICLLESSCENENERENKTE > Drosophila, http://flybase.org/ Glass CANGRKAESESESSTITKETENESTESCERENCOKICLLESSCENENERENKTE > Strongylocentrotus, https://blast.ncbi.nlm.nih.gov/ XP 790671.3 CANGRKAERESSESSTITKETENESTESCERENCOKICLLESSCENERENKENEKE oioRxit preprint doi: Preprint

Glass

stid

stid

stid

stid

oflagellates) choan and (vertebrates proteins ass-like

https://blast.ncbi.nlm.nih.gov/

>Human,

CDACGRAFSDCSALIRHIRTHSGEKPYQCKVCPKAFAQSSSTIEHQ 011525220.1, ZSCAN22

https://blast.ncbi.nlm.nih.gov/ >Human,

EAW84863.1, ZNF253

CEECGKAFNRSTDITTHKIVHTGEKPYKCEECGKAFNQPSTISKFE

https://blast.ncbi.nlm.nih.gov/

CKECGEAF SOSSTITKHIKVYTGEKPYTCKDCRKAFSOSSITQHQ CAA36583.1, KOX 26 >Human,

stid

http://www.uniprot.org/ >Human, **ZNF764**

ROSSEMAAHRRTHSGEKPYPCPOGGRRFGOKSAVAKHO CPDCGRCE

atid

http://www.uniprot.org/ >Human, ZNF683

http://www.uniprot.org/ >Human,

SSSSNIKTHLRIHSGARPFQCSVCRSRFTQHIHFKLHH

REVCHKRE

stid 0.0-1

CTOCGKRENNSSHESAHRRTHTGEKPYTCPACGRGERRGTDIHKHO

ZNF500

RT

> Salpingoeca, https://blast.ncbi.nlm.nih.gov/


XP 004993762.1


CSICGKSFITSSHVRMHERTHDDARHFSCPLCSKPFKTKPG KQHI


> Salpingoeca, https://blast.ncbi.nlm.nih.gov/

XP 004989424.1

CDQCSFKTAYKHILTTHKRIHTGEKPYKCDQCEYKTAYKRNMPRHKRIH










Figure 4




Figure 5




Figure 6

