Molecularly distinct models of zebrafish *Myc*-induced B cell leukemia

Chiara Borga[1*], Clay A. Foster[1*], Sowmya Iyer[2,3*], Sara P. Garcia, David M. Langenau[2,3]†, J. Kimble Frazer[1]†

[1]Section of Pediatric Hematology-Oncology, Department of Pediatrics, University of Oklahoma Health Sciences Center, Oklahoma City, OK 73104; [2]Department of Pathology and Center for Cancer Research, Massachusetts General Hospital Research Institute, Boston, MA 02114; [3]Harvard Stem Cell Institute, Boston, MA 02114

[*]Co-first authorship contribution

†Equal senior authorship contribution

Corresponding authors:

J. Kimble Frazer, University of Oklahoma Health Sciences Center, Basic Sciences Education Bldg., 941 Stanton L Young, Rm. 229 Oklahoma City, OK 73104, USA. Phone: 405-271-5311 ext. 42375. Fax: 405-271-1025. Email: Kimble-Frazer@ouhsc.edu

David M. Langenau, Harvard Medical School and Massachusetts General Hospital, 149 13th Street, office #6012, Charlestown, Massachusetts 02129, USA. Phone: 617-643-6508. Email: dlangenau@mgh.harvard.edu

Conflict of interest disclosure statement: Authors declare no conflict of interests.

Abstract:

Zebrafish models of T cell acute lymphoblastic leukemia (T-ALL) have been studied for over a decade, but curiously, robust zebrafish B cell ALL (B-ALL) models had not been described. Recently, our laboratories reported two seemingly closely-related models of zebrafish B-ALL. In these genetic lines, the primary difference is expression of either murine or human transgenic c-MYC, each controlled by the zebrafish *rag2* promoter. Here, we compare ALL gene expression in both models. Surprisingly, we find that B-ALL arise in different B cell lineages, with $ighm^+$ vs. $ighz^+$ B-ALL driven by murine *Myc* vs. human *MYC*, respectively. Moreover, these B-ALL types exhibit signatures of distinct molecular pathways, further unexpected dissimilarity. Thus, despite sharing analogous genetic makeup, the ALL types in each model are markedly different, proving subtle genetic changes can profoundly impact model organism phenotypes. Investigating the mechanistic differences between mouse and human c-MYC in these contexts may reveal key functional aspects governing MYC-driven oncogenesis in human malignancies.

Zebrafish are a valuable leukemia model due to highly conserved hematopoietic and oncogenic pathways, facile genetics, and ease of use in chemical genetic screens. However, until recently, robust zebrafish B-cell leukemia models had not been described[1, 2]. The first transgenic zebrafish leukemia model was created 15 years ago and targeted murine c-*Myc* (*mMyc*) to thymocytes of AB strain zebrafish, leading to the rapid development of T-cell acute lymphoblastic leukemia (T-ALL)[3]. Additional genetic models were subsequently developed that result in induction of T-ALL[4], but B-cell leukemia models lagged behind[5, 6].

In two recent reports published in *Leukemia*, our groups independently demonstrated the development of zebrafish B-ALL using transgenic expression of *mMyc* or human c-*MYC* (*hMYC*) controlled by the zebrafish recombination activating gene 2 (*rag2*) promoter[1, 2]. Both models shared genetic and phenotypic features, but there were also key differences including strain background and species differences in the MYC transgene that was used to generate each model. Here, we compare and contrast these models, making the important finding that zebrafish develop at least four molecularly distinct ALL types, including cortical thymocyte-arrested $cd4^+/cd8^+$ T-ALL, $ighm^+$ B-ALL, $ighz^+$ B-ALL, and biphenotypic T/B-ALL.

Thirteen ALLs were purified from *rag2:hMYC;lck:eGFP* double-transgenic fish[2]. As previously reported, these leukemias had heterogeneous GFP expression, with T-ALL being exclusively GFP^{hi}, B-ALL exclusively GFP^{lo}, and other fish harboring mixed-ALL with both GFP^{hi} and GFP^{lo} cells, representing simultaneous T- and B-ALL, respectively[2]. Leukemias were subjected to RNA-seq transcriptomic profiling and compared with transplanted leukemias generated from single ALL clones described by Garcia et al., 2018[1]. Principal Component Analysis clearly distinguished *mMyc*-induced T-ALL from B-ALL, with the single *mMyc*-induced biphenotypic B/T-ALL clustering between these samples (Fig. 1A). The eight *hMYC-*

induced ALLs that were largely GFP^{hi} clustered with known T-ALLs (2, 6, 8-11, 13, 14), while

three primarily GFP^{lo} ALL clustered near the *mMyc* B-ALLs (3-5). Two *hMYC*-induced ALLs

with substantial populations of both GFP^{lo} and GFP^{hi} cells grouped near the *mMyc* biphenotypic

B/T leukemia. Hierarchical clustering using the top 100 positively- and negatively-correlated

genes from PC2 confirmed that these genes defined B and T lymphocytes, respectively (Fig. 1B),

with B-ALLs expressing *cd79b*, *syk*, *pax5*, *blnk* and *efb1*, while T-ALLs expressed *cd8b*, *lck*,

runx3 and *gata3*. As previously reported, the biphenotypic B/T-ALL and mixed *hMYC*-induced

ALLs expressed both T- and B-cell lineage genes[1, 2]. PC2 up-regulated genes were enriched for

B cell signaling pathways when independently assessed by GSEAsig (Supplemental Tables 1 and

2).

To examine clonality and maturation in *mMyc*- and *hMYC*-induced ALL, we next

analyzed the expression of constant and variable regions of the T cell receptor β (*tcrβ*) and

immunoglobulins μ and ζ (*ighm*, *ighz*; Fig. 2A). Every *mMyc* and *hMYC* T-ALL exhibited *tcrβ*

expression, with V(D)J recombination occurring in most samples as determined by expression of

specific variable regions[1]. Conversely, *mMyc* and *hMYC* B-ALL did not recombine or express

tcrβ, but expressed constant regions of *ighm* or *ighz*. Ig variable regions were not detected,

indicating V(D)J rearrangement likely had not occurred in these leukemias and suggesting B-

ALLs arrest at the early pro-B cell stage. As expected, *hMYC* mixed-ALL contained distinct T-

and B-ALL clones expressing both *tcr* and *ig* mRNAs, with their relative expression correlating

well with the percentage of GFP^{hi}/T-ALL vs. GFP^{lo}/B-ALL cells found in each sample (Fig. 2A).

Intriguingly, *mMyc* B-ALL expressed exclusively *ighm* while *hMYC* B-ALL favored *ighz*

expression, indicating that *mMyc* and *hMYC* might be oncogenic in distinct B cell lineages.

To further explore differences between these models, we next identified genes uniquely-

expressed by T-ALL, *mMyc/ighm*⁺ B-ALL, or *hMYC/ighz*⁺ B-ALL (Fig. 2B). As expected T-

ALLs expressed known T cell lineage markers, yet *mMyc/ighm*⁺ and *hMYC/ighz*⁺ B-ALLs were

transcriptionally distinct. *mMyc/ighm*⁺ B-ALL expressed *gfi1ab, zfhx3, notch1a, nf1b,* and

gtf3aa. By contrast, *hMYC/ighz*⁺ B-ALL expressed higher *cd79a*, *cd83, mef2cb*, and *jak2a* levels.

To further test for differences in these two molecular subtypes of B-ALL, we next performed

GSEAsig using these same differentially-regulated genes. From this analysis, we uncovered that

mMyc/ighm⁺ B-ALLs exhibited significant enrichment for pathways regulating ribosome

biogenesis and RNA binding (Fig. 2C and Supplemental Tables 3 and 4). By contrast,

hMYC/ighz⁺ B-ALLs were enriched for intracellular signaling, protein binding, and germinal

center B cell maturation pathways. In support of our findings, Liu et al. recently reported the

identification of molecularly and biological distinct *ighz*⁺ and *ighm*⁺ B cell lineages using

rag2:mCherry; cd79b:GFP transgenic zebrafish[7]. In the context of normal B cell development,

ighz⁺ B cells were mCherry^hi/GFP^lo while *ighm*⁺ B cells were mCherry^hi/GFP^hi. Overall, these

results demonstrate that *mMyc/ighm*⁺ and *hMYC/ighz*⁺ B-ALLs are not subtle B cell leukemia

variants, but rather distinct malignancies that arise in different B cell types with vastly different

molecular pathway signatures.

In summary, although zebrafish B cell leukemia models were lacking for many years, our

analyses reveal two highly-divergent types of B-ALL. This is surprising, as both models utilize

the same promoter (*rag2*) to regulate a near-identical oncoprotein, c-Myc/MYC, with the only

differences being the MYC transgene species of origin and the genetic backgrounds upon which

the models were developed. Yet, despite the high molecular similarity of both models, these B-

ALL subtypes also show unique gene expression signatures when compared to one another,

which likely reflects differences in both their lineage (*ighm* vs. *ighz*) and potential differences in

MYC transcriptional targets expressed by the early developmental stages of these distinct pro-B

cell populations. Our new analysis of these models reconciles the perceived differences in the

manuscripts published by our groups, identifying four molecularly distinct ALL subtypes in

zebrafish: cortical $cd4^+/cd8^+$ T-ALL, biphenotypic B/T ALL, $ighm^+$ B-ALL, and $ighz^+$ B-ALL.

Developing a wider array of leukemia models and refining mechanisms that drive their growth,

aggression, and stem cell frequency will surely lead to new insights into human disease.

Figure Legends:

Figure 1. Gene expression differences define T- vs. B-lineages in ALL of both the *rag2:mMyc* and *rag2:hMYC;lck:eGFP* models. (**A**) Principal component analysis of RNAseq expression profiles of previously-classified *mMyc*-induced T- (blue, n=8), B- (red, n=2), or biphenotypic ALL (pink, n=1) and compared with 13 unknown *hMYC*-induced ALL (black). (**B**) Heat map and hierarchical clustering using the top 100 positively- and negatively-correlated genes from PC2. B cell-specific genes are denoted in blue and T cell-specific genes in red at the right.

Figure 2. Identification of two molecularly-distinct B-ALL types arising independently in either the *ighm*$^+$ or *ighz*$^+$ B cell lineages. (**A**) T cell receptor beta and Ig heavy chain expression in individual ALLs. *tcrβ* recombination is denoted by grey-shaded boxes (left column), with percentage of GFP^{hi} cells in each *rag2:hMYC;lck:eGFP* ALL noted in right column. Histograms depict expression of *tcrβ* and *igh* constant regions by each ALL. Not available (NA). (**B**) Heatmap showing expression of genes differentially expressed in T-ALL, *mMyc*/*ighm*$^+$ B-ALL, and *hMYC*/*ighz*$^+$ B-ALL. (**C**) Gene set enrichment analysis using genes positively correlated with each ALL molecular subtype. T-ALL (blue), *mMyc*/*ighm*$^+$ B-ALL (red), and *hMYC*/*ighz*$^+$ B-ALL (orange). Complete geneset and GSEAsig results are provided in Supp. Tables 3 and 4.

130 **References:**

131   1.   Garcia EG, Iyer S, Garcia SP, Loontiens S, Sadreyev RI, Speleman F, *et al.* Cell of origin dictates
132   aggression and stem cell number in acute lymphoblastic leukemia. *Leukemia* 2018 2018/04/18.

133
134   2.   Borga C, Park G, Foster C, Burroughs-Garcia J, Marchesin M, Shah R, *et al.* Simultaneous B and T
135   cell acute lymphoblastic leukemias in zebrafish driven by transgenic MYC: implications for
136   oncogenesis and lymphopoiesis. *Leukemia* 2018 2018/08/15.

137
138   3.   Langenau DM, Traver D, Ferrando AA, Kutok JL, Aster JC, Kanki JP, *et al.* Myc-induced T cell
139   leukemia in transgenic zebrafish. *Science* 2003 Feb 7; **299**(5608): 887-890.

140
141   4.   Frazer JK, Meeker ND, Rudner L, Bradley DF, Smith AC, Demarest B, *et al.* Heritable T-cell
142   malignancy models established in a zebrafish phenotypic screen. *Leukemia* 2009 Oct; **23**(10):
143   1825-1835.

144
145   5.   He S, Jing CB, Look AT. Zebrafish models of leukemia. *Methods Cell Biol* 2017; **138**: 563-592.

146
147   6.   Squiban B, Frazer JK. Danio rerio: Small Fish Making a Big Splash in Leukemia. *Curr Pathobiol Rep*
148   2014 Jun; **2**(2): 61-73.

149
150   7.   Liu X, Li YS, Shinton SA, Rhodes J, Tang L, Feng H, *et al.* Zebrafish B Cell Development without a
151   Pre-B Cell Stage, Revealed by CD79 Fluorescence Reporter Transgenes. *J Immunol* 2017 Sep 1;
152   **199**(5): 1706-1715.

A

	tcrβ recomb.	lck:GFP	

rag2:mouse cMyc CG1-strain (transplant)

9.2 — ighm+ B-ALL
10.2
12.2 — T/B bi-phenotypic
15.2
14.1.1.1
13.1.1
8.4.1.1
8.3
7.2.1
4.3.1
2.1 — T-ALL

rag2:human cMYC-ER WIK/AB-strain (primary)

hMYC-10 98
hMYC-2 81
hMYC-6 96
hMYC-8 92
hMYC-9 88
hMYC-11 97
hMYC-14 NA
hMYC-12 50 — Mixed
hMYC-13 77
hMYC-4 24
hMYC-3 6
hMYC-5 1 — ighz+ B-ALL
hMYC-1 NA

Legend:
- tcrβ-C1
- tcrβ-C2
- ighm
- ighd
- ighz

Expression (read counts/million): 0 100 200 300 400 500

B

gfi1ab
zfhx3
notch1a
tle3b

nf1b
gtf3aa

hopx
mef2cb
cd83
jak2a

lyn
cd79a

bcl11b
cd8a
cd226
cd4-1
lck
tcf7

9.2A 9.2B 10.2A 10.2B 12.2A 12.2B hMYC-5 hMYC-3 hMYC-4 hMYC-1 hMYC-12 hMYC-13 hMYC-14 hMYC-11 hMYC-8 hMYC-2 hMYC-9 hMYC-6 hMYC-10 8.3A 8.3B 7.2.1A 7.2.1B 4.3.1A 4.3.1B 8.4.1.1 2.1A 2.1B 14.1.1.1 15.2A 15.2B 13.1.1A 13.1.1B

row z-score: -2 0 2

C

- GO T CELL DIFFERENTIATION
- GSE10325 CD4 T CELL VS MYELOID UP
- LEE DIFFERENTIATING T LYMPHOCYTE
- GO PRERIBOSOME
- GO POLY-A RNA BINDING
- GO RNA BINDING
- PASQUALUCCI LYMPHOMA BY GC STAGE DN
- GO PROTEIN DOMAIN SPECIFIC BINDING
- GO INTRACELLULAR SIGNAL TRANSDUCTION

Legend:
- T-ALL
- Igm+ B-ALL
- Igz+ B-ALL

FDR q-value: 1×10^{-1} 1×10^{-3} 1×10^{-5} 1×10^{-7} 1×10^{-9} 1×10^{-11} 1×10^{-13} 1×10^{-15} 1×10^{-17}