The winner takes it all: how semelparous insects can become periodical*

Odo Diekmann[1] and Robert Planqué[2]

Abstract The aim of this short note is to give a simple explanation for the remarkable periodicity of *Magicicada* species, which appear as adults only every 13 or 17 years, depending on the region. We show that a combination of two types of density dependence may drive, for large classes of initial conditions, all but one year class to extinction. Competition for food leads to negative density dependence in the form of a uniform (i.e., affecting all age classes in the same way) reduction of the survival probability. Satiation of predators leads to positive density dependence within the reproducing age class. The analysis focuses on the full life cycle map derived by iteration of a semelparous Leslie matrix.

Keywords Age-structured population models · Nonlinear dynamics · Nonlinear Leslie matrix models · Single year class dynamics · Semelparous insects.

Mathematics Subject Classification (2010) 37N25 · 92B25

Dedicated to Hans Othmer, a winner who shares rather than deprives.

1 Periodical insects: a conundrum

According to M.G. Bulmer (Bulmer 1977), "An insect population is said to be periodical if the life cycle has a fixed length of k years ($k > 1$) and if the adults do not appear every year but only every kth year." He then provides several examples, one by quoting Lloyd and Dybas (1966)

> "Periodical cicadas, *Magicicada* spp., have the longest life cycles known for insects. In any one population, all but a tiny fraction are the same age. The nymphs suck juices from the roots of deciduous forest trees in eastern United States. Mature nymphs finally emerge from the ground, become adults, mate, lay their eggs, and die within the same few weeks of every seventeenth (or, in the South, every thirteenth) year. Not one species does this, but three. There are three separate and distinct species that occur together over most of the range of periodical cicadas and, wherever the species coexist, they are invariably synchronized. In different regions, different "broods" of periodical cicadas may be out of synchrony by several years, but the species (in a given region) never are. Finally, the same three species—the same as nearly as anyone can tell by looking at them or listening to the songs of their males—exist both as 17 and as 13-year periodical cicadas! [Pp. 133 - 134]"

Odo Diekmann
Department of Mathematics, Utrecht University, Budapestlaan 6, 3584 Utrecht, The Netherlands. email: o.diekmann@uu.nl. Tel: +31 30 253 1487.
Robert Planqué
Department of Mathematics, Vrije Universiteit Amsterdam, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands. email: r.planque@vu.nl. ORCID: 0000-0002-0489-5425.

In the wake of Bulmer's pioneering paper a rich literature on discrete time models for semelparous organisms developed, see (Behncke 2000, Cushing 2009; 2015, Cushing and Henson 2012, Diekmann et al. 2005, Kon 2012; 2017, Webb 2001, Wikan 2017) and the references given there. The population splits into year classes according to the year of birth (equivalently: the year of reproduction) counted modulo k. As a year class is reproductively isolated from the other year classes, it forms a population by itself. Accordingly, competition may lead to exclusion.

Despite this rich literature, the spatio-temporal pattern displayed by periodical cicada population dynamics is, we think, still an enigma. The interest of one of the authors in the subject was rekindled while refereeing an early version of (Machta et al. 2018). The model introduced and studied in that paper is characterized by

– uniform (i.e., age-class unspecific) negative density dependence during development;
– positive density dependence for reproduction.

The first of these captures competition for food, the second satiation of predators to eat adults in a given year (Williams et al. 1993).

The paper (Machta et al. 2018) considers the limit $k \to \infty$ and employs a continuous time description of the competition for resources (whence the word "hybrid" in the title). The main result establishes the instability of any steady state with more than one year class present.

Here we focus on the "full life cycle" map, i.e., the k-times iterated Leslie matrix, cf. (Davydova et al. 2003; 2005). The full life cycle map is represented by a diagonal matrix. If competition is uniform, in the sense that in any year any of the survival probabilities is the product of a fixed age-class specific factor and a year-specific factor that is the same for all age-classes, the diagonal elements all have the same overall survival factor. In other words, negative density dependence results over a full life cycle for all year classes in exactly the same multiplication factor. Each diagonal element has an additional reproduction factor. Positive density dependence causes this factor to be highest for the year class that had, in the year that it reproduced, the highest density. In this way, a numerical advantage is reinforced and asymptotically there remains, "generically", only one year class: all the others go extinct. See Figure 1 for an example.

Note that this holds independently of the resulting single year class (SYC) dynamics in the sense that the winners may exhibit ultimately steady state, periodic or even chaotic dynamics. Also note that it depends on the initial condition which one of the k year classes will win the competition. Domains of attraction (or, dominance) are separated by sets of special initial conditions for which several year classes coexist forever (e.g., steady coexistence states and their stable manifolds). In principle these separatrices may have an intricate structure (e.g., see (Hofbauer et al. 2004)).

The upshot is that the combination of uniform negative density dependence and concentrated (in one point of the life cycle) positive density dependence, as assumed in (Machta et al. 2018), leads, as a rule, to exclusion of all but at most one year class. This does not prove, of course, that this is the mechanism underlying the observed phenomenon of SYC dynamics in *Magicicada*. But in the spirit of Occam's Razor, this probably currently offers the simplest, and hence most plausible, potential explanation.

Anyhow, as far as we are aware, we provide below the first analytical demonstration of one year class driving all other year classes to extinction without severe restrictions on the initial conditions and without any condition on the resulting SYC dynamics, except for boundedness.

2 Model formulation

Let k be an integer bigger than one. Let $\boldsymbol{N}(t)$ be the k-vector such that $N_j(t)$ is the sub-population density of individuals that at the census moment in year t have age j. In view of linear algebra we take $j = 1, 2, \ldots, k$, even though biologically $j = 0, 1, \ldots, k-1$ would perhaps make more sense if, as we assume, the census moment is in the early autumn, so after reproduction took place. We assume that

$$\boldsymbol{N}(t+1) = L(\boldsymbol{h}(\boldsymbol{N}(t))\boldsymbol{N}(t). \tag{1}$$

where

$$L(\boldsymbol{h}) = \begin{pmatrix} 0 & 0 & \cdots & 0 & h_k \\ h_1 & 0 & & 0 & 0 \\ 0 & h_2 & & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & h_{k-1} & 0 \end{pmatrix} \tag{2}$$

is a "semelparous" Leslie matrix (here the adjective "semelparous" indicates that in the first row only the last element is non-zero, reflecting that individuals reproduce at age k and then die). The dependence of the matrix elements h_i on $\boldsymbol{N}$ captures density dependence. We assume that

$$h_i(\boldsymbol{N}) = \sigma_i \pi(\boldsymbol{N}) \quad \text{for } i = 1, \ldots, k-1, \tag{3}$$
$$h_k(\boldsymbol{N}) = \sigma_k \pi(\boldsymbol{N}) \beta(\sigma_k \pi(\boldsymbol{N}) N_k). \tag{4}$$

where

$$\sigma_i \in (0, 1], \quad i = 1, \ldots, k, \tag{5}$$
$$\pi : \mathbb{R}_+^k \to [0, 1], \tag{6}$$
$$\beta : [0, \infty) \to [0, \infty) \text{ is continuous, strictly increasing and bounded.} \tag{7}$$

Here the σ_i are survival probabilities under "standard" conditions and the (uniform, i.e., i-independent) factor $\pi(\boldsymbol{N})$ reflects the reduction of the survival probability as a result of crowding and competition for food. The last age group too needs to survive the winter before they can emerge as adults, so the population density of emerging adults equals $\sigma_k \pi(\boldsymbol{N}) N_k$. The function β is a composite model ingredient. It describes the number of offspring at the census moment per emerging adult. The monotonicity of β reflects that the per adult capita risk of falling victim to predation decreases with adult population density. This is the effect of predators becoming satiated (or even over-satiated) when prey density is very high, and that the time interval between emergence and death of the adults is too short for a numerical increase in the number of predators (mainly birds).

3 The main results

So far we did not specify any property of π that justifies the description "negative" density dependence. But now we require that population densities remain bounded and implicitly this is an assumption concerning the function π.

Hypothesis 1 *(dissipativity (Hale 1988)) There exists $R > 0$ such that*

$$B_R^+ := \{\boldsymbol{N} \in \mathbb{R}_+^k : \sum_{i=1}^{k} N_i \le R\} \tag{8}$$

is invariant and attracts all orbits.

By "attracts all orbits" we mean that for every $\boldsymbol{N}_0 \in \mathbb{R}_+^k$ there exists $m = m(\boldsymbol{N}_0)$ such that the solution of (1) with initial condition $\boldsymbol{N}(0) = \boldsymbol{N}_0$ satisfies $\boldsymbol{N}(t) \in B_R^+$ for $t \ge m(\boldsymbol{N}_0)$. We adopt Hypothesis 1 for the rest of the paper. In Section 4 below we provide easily verifiable conditions on π and β that guarantee that Hypotheses 1 is satisfied.

At the end of Section 1 we already observed that (due to symmetry, see (Davydova et al. 2005, Diekmann and van Gils 2003) for a bit more detail) any year class is a candidate for becoming the sole inhabiter of the world. It turns out to be relatively easy to describe a set of initial conditions such that the year class that has age k at the initial time will outcompete all the other year classes.

Theorem 1 *The set*
$$\Omega_k := \{\boldsymbol{N} \in \mathbb{R}_+^k : N_k > N_i \text{ for } i = 1, \ldots, k-1\} \tag{9}$$

is invariant under the full life cycle map, and if $\boldsymbol{N}(0) \in \Omega_k$ and for $i \in \{1, \ldots, k-1\}$, $N_i(0) > 0$, then

$$\frac{N_i(mk)}{N_k(mk)}$$

is a strictly decreasing function of m. This function has limit zero for $m \to \infty$ in case $\limsup_{m \to \infty} N_k(mk) > 0$.

The set Ω_3 is depicted in Figure 1. Note that *all* year classes go extinct if $N_k(mk) \to 0$ for $m \to \infty$. Hypothesis 1 rules out the possibility that both $N_i(mk)$ and $N_k(mk)$ grow beyond any bound for $m \to \infty$ while their ratio goes to zero. So we conclude that for initial conditions belonging to Ω_k only the k-th year class can possibly persist.

Proof In order to avoid overburdening the reader with notational detail, we focus on the proof of the case $k = 3$. The proof for general k does not require new arguments.

From (1) it follows that

$$\boldsymbol{N}(3) = L(\boldsymbol{h}(\boldsymbol{N}(2)))L(\boldsymbol{h}(\boldsymbol{N}(1)))L(\boldsymbol{h}(\boldsymbol{N}(0)))\boldsymbol{N}(0). \tag{10}$$

The product of the three Leslie matrices is a diagonal matrix. We claim that the three elements of the diagonal of this matrix have a factor

$$\theta := \sigma_1 \sigma_2 \sigma_3 \pi(\boldsymbol{N}(0))\pi(\boldsymbol{N}(1))\pi(\boldsymbol{N}(2)) \tag{11}$$

in common and are in fact of the form $\beta(\cdot)\theta$ with the argument of β being θN_1 for the first element, $\sigma_2 \sigma_3 \pi(\boldsymbol{N}(0))\pi(\boldsymbol{N}(1))N_2(0)$ for the second element and $\sigma_3 \pi(\boldsymbol{N}(0))N_3(0)$ for the third element.

To verify the claim, we elaborate (10) from right to left, using π_1 to denote $\pi(\boldsymbol{N}(0))$, π_2 to denote $\pi(\boldsymbol{N}(1))$ and π_3 to denote $\pi(\boldsymbol{N}(2))$. All factors in the products in the vectors below follow the order of events over time.

$$\begin{pmatrix} N_1 \\ N_2 \\ N_3 \end{pmatrix} \mapsto \begin{pmatrix} \beta(\sigma_3 \pi_1 N_3)\sigma_3 \pi_1 N_3 \\ \sigma_1 \pi_1 N_1 \\ \sigma_2 \pi_1 N_2 \end{pmatrix} \mapsto \begin{pmatrix} \beta(\sigma_3 \pi_2 \sigma_2 \pi_1 N_2)\sigma_3 \pi_2 \sigma_2 \pi_1 N_2 \\ \sigma_1 \pi_2 \beta(\sigma_3 \pi_1 N_3)\sigma_3 \pi_1 N_3 \\ \sigma_2 \pi_2 \sigma_1 \pi_1 N_1 \end{pmatrix} \tag{12}$$

$$\mapsto \begin{pmatrix} \beta(\sigma_3 \pi_3 \sigma_2 \pi_2 \sigma_1 \pi_1 N_1)\sigma_3 \pi_3 \sigma_2 \pi_2 \sigma_1 \pi_1 N_1 \\ \sigma_1 \pi_3 \beta(\sigma_3 \pi_2 \sigma_2 \pi_1 N_2)\sigma_3 \pi_2 \sigma_2 \pi_1 N_2 \\ \sigma_2 \pi_3 \sigma_1 \pi_2 \beta(\sigma_3 \pi_1 N_3)\sigma_3 \pi_1 N_3 \end{pmatrix}.$$

Noting that $\theta = \sigma_1 \sigma_2 \sigma_3 \pi_1 \pi_2 \pi_3$, inspection reveals that the claim is correct. It follows that

$$\frac{N_1(3)}{N_3(3)} = \frac{\beta(\theta N_1(0))}{\beta(\sigma_3 \pi(\boldsymbol{N}(0))N_3(0))} \frac{N_1(0)}{N_3(0)}. \tag{13}$$

Now for $\boldsymbol{N}(0) \in \Omega_3$, $N_1(0) < N_3(0)$, so that

$$\begin{aligned} \theta N_1(0) &= \sigma_3 \pi(\boldsymbol{N}(0))\Big(\sigma_2 \sigma_1 \pi(\boldsymbol{N}(1))\pi(\boldsymbol{N}(2))\Big)N_1(0) \\ &\leq \sigma_3 \pi(\boldsymbol{N}(0))N_1(0) \\ &< \sigma_3 \pi(\boldsymbol{N}(0))N_3(0), \end{aligned}$$

and we conclude that, since β is strictly increasing,

$$\beta(\theta N_1(0)) < \beta(\sigma_3 \pi(\boldsymbol{N}(0))N_3(0)).$$

Hence,

$$\begin{aligned} (i) \quad & N_1(3) < N_3(3), \\ (ii) \quad & \frac{N_1(3)}{N_3(3)} < \frac{N_1(0)}{N_3(0)} \quad \text{when } N_1(0) > 0. \end{aligned}$$

Essentially the same argumentation yields that, when $\boldsymbol{N}(0) \in \Omega_3$

$$\begin{aligned} (iii) \quad & N_2(3) < N_3(3), \\ (iv) \quad & \frac{N_2(3)}{N_3(3)} < \frac{N_2(0)}{N_3(0)} \quad \text{when } N_2(0) > 0. \end{aligned}$$

It follows from (i) and (iii) that Ω_3 is invariant under the full life cycle map and from (ii) and (iv) that for $i = 1, 2$ the sequences

$$\frac{N_i(3m)}{N_3(3m)}$$

are decreasing in m, strictly when positive. It remains to be shown that either their limit is zero or all year classes go extinct.

Assume that $N_1(3m)/N_3(3m) \to \alpha$ for $m \to \infty$. By iteration (13) leads to

$$\frac{N_1(3(m+1))}{N_3(3(m+1))} = \frac{\beta(\sigma_1\sigma_2\sigma_3\pi(\boldsymbol{N}(3m))\pi(\boldsymbol{N}(3m+1))\pi(\boldsymbol{N}(3m+2))N_1(3m))}{\beta(\sigma_3\pi(\boldsymbol{N}(3m))N_3(3m))} \frac{N_1(3m)}{N_3(3m)}. \tag{14}$$

Passing to the limit $m \to \infty$ in (14), it follows that if $\alpha > 0$ then necessarily, since β is continuous,

$$\frac{\beta(\sigma_1\sigma_2\sigma_3\pi(\boldsymbol{N}(3m))\pi(\boldsymbol{N}(3m+1))\pi(\boldsymbol{N}(3m+2))\alpha N_3(3m)}{\beta(\sigma_3\pi(\boldsymbol{N}(3m))N_3(3m))} \to 1$$

for $m \to \infty$. This is very well possible if $N_3(3m) \to 0$ for $m \to \infty$ (and in that case, since Ω_3 is invariant, also $N_1(3m)$ and $N_2(3m)$ converge to zero for $m \to \infty$, so all year classes go extinct). But if $\limsup_{m\to\infty} N_3(3m) > 0$, this is *not* possible, since $\alpha < 1$, $\sigma_1\sigma_2 \le 1$, $\pi(\boldsymbol{N}(3m+1))\pi(\boldsymbol{N}(3m+2)) \le 1$ and β is strictly increasing.

Theorem 1 does neither restrict nor predict the resulting SYC dynamics and allows for a large class of initial conditions. We now present a kind of local version in which we incorporate more restrictions, but also describe more precisely the ultimate dynamics.

The full life cycle map acts on $\mathbb{R}^k$. Any subspace characterised by $k - 1$ components being zero is invariant. The restriction to such an invariant subspace corresponds to a map from $\mathbb{R}$ to $\mathbb{R}$ that we shall call a SYC full life cycle map. The k maps are different but equivalent, see [8, chapter V, 9, Section 7]. To stay in line with Theorem 1 we work primarily with the map corresponding to $N_1 = N_2 = \ldots = N_{k-1} = 0$.

Theorem 2 *If a single year class fixed point or periodic orbit of the SYC full life cycle map is linearly stable with respect to that map, so with respect to within year class perturbations, it is automatically also linearly stable with respect to perturbations that involve small introductions of other year classes.*

Proof We again give a proof for the case $k = 3$. The proof for general k uses no new arguments.

The full life cycle map given by (12) is represented by a diagonal 3×3 matrix M acting on a 3 vector,

$$M = \begin{pmatrix} \theta\beta(\theta N_1) & 0 & 0 \\ 0 & \theta\beta(\sigma_2\sigma_3\pi_1\pi_2 N_2) & 0 \\ 0 & 0 & \theta\beta(\sigma_3\pi_1 N_3) \end{pmatrix}, \quad \text{where } \theta = \sigma_1\sigma_2\sigma_3\pi_1\pi_2\pi_3.$$

(Of course, all the π_i are functions of N_1, N_2, and N_3.) We first consider a SYC fixed point and choose the phase such that only the third component of the fixed point vector is positive. Then necessarily $M_{3,3} = \theta\beta(\sigma_3\pi_1 N_3) = 1$ when evaluated in the fixed point. N_3 is itself a fixed point of the SYC map

$$N_3 \mapsto \sigma_1\sigma_2\sigma_3\pi_1\pi_2\pi_3\beta(\sigma_3\pi_1 N_3)N_3,$$

and is, by assumption, stable as such.

The linearisation of the full life cycle map in the fixed point is represented by a matrix as well, of course, L, say. Since the first two components of the fixed point are zero, $L_{1,2} = L_{1,3} = L_{2,3} = L_{2,1} = 0$, making L a lower-diagonal matrix, with eigenvalues equal to the diagonal elements. Direct computation shows that L must be of the form

$$L = \begin{pmatrix} \theta\beta(0) & 0 & 0 \\ 0 & \theta\beta(0) & 0 \\ L_{3,1} & L_{3,2} & L_{3,3} \end{pmatrix}.$$

Since β is increasing, $0 \le \theta\beta(0) < \theta\beta(\sigma_3\pi_1 N_3) = M_{3,3} = 1$.

The diagonal element in the third and last row is complicated, but is equal to the linearisation of the SYC full life cycle map in the fixed point. Hence, since we have assumed that fixed point to be linearly stable, $|L_{3,3}| < 1$. So all eigenvalues of L are less than one in absolute value and the fixed point of the three dimensional full life cycle map is linearly stable.

Fig. 1 Example dynamics for the map (1)–(2), with parameters $\sigma_1 = \sigma_2 = \sigma_3 = 1$, $\pi(\boldsymbol{N}) = e^{-0.4(N_1+N_2+N_3)}$ and $\beta(N_3) = 17.5N_3/(1+N_3)$. Left: full dynamics of all cohorts; middle: the dynamics of the full life cycle map, for all cohorts, shown from the year at which the orbit enters Ω_3; right: dynamics generated by the full life cycle map, with Ω_3 in black, with red points of the orbit outside Ω_3 and blue those inside. We observe convergence to a single year class fixed point inside Ω_3 that lies on the N_3-axis.

For an orbit of period p, we consider the p times iterated full life cycle map. The structure is exactly the same as before, the only difference is that diagonal elements have p factors $\theta\beta$. The Jacobi matrix is again lower diagonal, and of the form

$$K = \begin{pmatrix} \theta(p-1)\cdots\theta(0)\beta^p(0) & 0 & 0 \\ 0 & \theta(p-1)\cdots\theta(0)\beta^p(0) & 0 \\ K_{3,1} & K_{3,2} & K_{3,3} \end{pmatrix}.$$

By the same arguments as before, $\theta(p-1)\cdots\theta(0)\beta^p(0) < 1$, and $|K_{3,3}| < 1$ by the assumption that the p-periodic orbit is linearly stable. So exactly as before we reach the conclusion that all eigenvalues are less than one in absolute value.

4 Model ingredients

To obtain more insight which choices of model ingredients π and β ensure persistence of populations and induce dissipative dynamics as defined in Hypothesis 1, we scale the variables in the following way. Let

$$N_2^{\text{old}} = \sigma_1 N_2^{\text{new}}, \quad N_3^{\text{old}} = \sigma_1\sigma_2 N_3^{\text{new}},$$

and define functions β^{new} and π^{new} according to

$$\beta^{\text{new}}(x) = \sigma_1\sigma_2\sigma_3\beta^{\text{old}}(\sigma_1\sigma_2\sigma_3 x),$$
$$\pi^{\text{new}}(\boldsymbol{N}^{\text{new}}) = \pi^{\text{old}}(\boldsymbol{N}^{\text{old}}).$$

In terms of these new variables, and dropping the superscripts for convenience, the map is seen to be simplified to

$$\boldsymbol{N}(t+1) = \pi(\boldsymbol{N}(t)) \begin{pmatrix} 0 & 0 & \beta(\pi(\boldsymbol{N}(t))N_3(t)) \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \boldsymbol{N}(t). \tag{15}$$

Let us now choose the following form for π:

$$\pi(\boldsymbol{N}) = p(E), \quad \text{with } E := c_1N_1 + c_2N_2 + c_3N_3. \tag{16}$$

Fig. 2 Example graphs of the SYC full life cycle map (17) for different choices of model ingredients. The blue line shows the diagonal; for both full life cycle map graphs, $\pi(E) = e^{-E}$, and β is defined as shown. Further parameters chosen in (17) are $c_1 = c_2 = 0.1$, $c_3 = 1$. Note that both maps show an Allee effect.

So $\pi(\boldsymbol{N})$ is determined by first computing a weighted population size E and next applying a scalar map p that assigns to E a component of the survival probability, i.e., a number between zero and one. Assume $c_3 > 0$, and define

$$E^{\text{new}} = \frac{1}{c_3} E^{\text{old}} = c_1^{\text{new}} N_1 + c_2^{\text{new}} N_2 + N_3,$$

and

$$p^{\text{new}}(E^{\text{new}}) = p^{\text{old}}(E^{\text{old}}).$$

Let us now consider the SYC dynamics. To facilitate the description of that case, we introduce

$$f(x) = \beta(p(x)x)p(x)x.$$

Dropping again the "new" superscripts, the SYC dynamics are given by

$$
\begin{aligned}
N_3 &\mapsto f(N_3) \\
&\mapsto p(c_1 f(N_3)) f(N_3) \\
&\mapsto p(c_2 p(c_1 f(N_3)) f(N_3)) p(c_1 f(N_3)) f(N_3).
\end{aligned}
$$

To ensure boundedness of the population consisting of just one year class, the graph of the SYC full life cycle map,

$$N_3 \mapsto p(c_2 p(c_1 f(N_3)) f(N_3)) p(c_1 f(N_3)) f(N_3) \tag{17}$$

should be below the 45° line for large values of N_3. Our model ingredients β and p must be chosen accordingly. Figure 2 gives an impression of the form of the graph of the SYC full life cycle map, for different choices of the model ingredients.

The derivative at the trivial fixed point $N_3 = 0$ corresponds to multiplication by

$$\beta(0)(p(0))^3.$$

If this quantity exceeds 1, the population cannot go extinct, whereas there is an Allee effect when this quantity is less than one and yet part of the graph lies above the 45° line (see the green and orange curves in Figure 2 for examples).

Some possible choices for p and β include

$$p(E) = \frac{1}{1+E}, \quad p(E) = e^{-E},$$

$$\beta(x) = \beta_0 \left(1 - \frac{\zeta}{x}\right)_+$$

This choice for β is strictly increasing only where it takes strictly positive values and equals zero in a neighbourhood of $x = 0$, so there is certainly an Allee effect. It corresponds to the total population size

of adults being reduced at a constant rate during a fixed time window. Alternatively, we can solve, with P, the predator density, as a parameter,

$$\frac{dx}{dt} = -\frac{x}{1 + \zeta x}P,$$

for a fixed period of time. This leads to an implicitly defined function β. For bifurcation diagrams of Single Year Class Maps, see Chapter V of (Davydova 2004. https://dspace.library.uu.nl/handle/1874/891).

Concerning dissipativity (Hypothesis 1), let

$$|\boldsymbol{N}| := N_1 + N_2 + N_3.$$

Then (15) implies that

$$|\boldsymbol{N}(t+1)| \leq \pi(\boldsymbol{N}(t))\beta(\infty)|\boldsymbol{N}(t)|. \tag{18}$$

If for some choice of $R > 0$ and $\epsilon > 0$ the inequality

$$\pi(\boldsymbol{N})\beta(\infty) \leq 1 - \epsilon$$

holds for all $|\boldsymbol{N}| \geq R$, then the ball

$$\{\boldsymbol{N} \; : \; |\boldsymbol{N}| \leq \max\{\beta(\infty), 1\}R\}$$

is invariant and attracts all orbits. Indeed, any orbit starting outside the ball $\{\boldsymbol{N} \; : \; |\boldsymbol{N}| \leq R\}$ has to enter this ball and once inside this ball we can use (18) and the fact that $\pi(\boldsymbol{N}) \leq 1$ to conclude that, if $\beta(\infty) > 1$, it may again leave the ball with radius R but not the ball with radius $\beta(\infty)R$.

5 Discussion

The discrete time population dynamics of semelparous species with one reproduction opportunity per year, is described by a special kind of Leslie matrices, viz. those for which in the first (= reproduction) row only the last element is non-zero. This reflects that an individual born in a certain year reproduces, if at all, exactly k years later, where k is the length of the life cycle. So the population splits into k reproductively isolated year classes. Mathematically this shows up in the fact that the k times iterated matrix is diagonal, so fully reducible.

Although they are reproductively isolated, the year classes do interact by competition for resources. When resource availability is constant in time, each year class does as well or bad as any other year class. But when resource availability fluctuates in time, the phase of the life cycle relative to resource peaks and troughs can be decisive for success or failure. Thus one year class can drive other year classes to extinction by inducing, for instance, periodic environmental conditions. The work of Davydova and co-workers (Davydova 2004. https://dspace.library.uu.nl/handle/1874/891, Davydova et al. 2003; 2005, Diekmann et al. 2005) focused on this phenomenon.

The present paper is inspired by (Machta et al. 2018) and analyses a model such that, over a full life cycle, competition for food is neutral with respect to the year class distinction. But the model includes a second form of density dependence: it assumes that, due to predator satiation effects, per capita reproduction success of adults is larger when the cohort of adults is larger. So individuals belonging to a large cohort get much offspring while the negative impact of cohort size on survival probabilities is affecting individuals of all other year classes equally strongly, when measured over a full life cycle. Thus the 'strongest' year class becomes even stronger and eventually drives all other year classes to extinction and SYC (single year class) dynamics (Davydova et al. 2003, Mjølhus et al. 2005) gets established. The idea that a combination of predator satiation and intraspecific competition among larvae gives rise to single year class dynamics goes at least back to Hoppensteadt and Keller (1976) and Bulmer (1977), and has been demonstrated in many of the models in the literature by way of numerical experiments. The assumption of uniform competition introduced in (Machta et al. 2018) allows, as we have shown above, to actually prove that SYC dynamics results for large classes of initial conditions. In our opinion, the precise nature of the relationship between mechanisms and phenomena is better revealed by a proof than by simulations.

To explain, in the context of the model, that three species coexist in synchrony would probably require the assumption that the functions π and the functions β for the three species are proportional. So this is still rather puzzling.

We now briefly consider some of the ecological evidence supporting the two chief modelling ingredients.

The long larval stages associated with periodical insects likely result from development constrained by certain abiotic factors such as low temperature, poor food availability, and large adult body size (Danks 1992, Hellövaara et al. 1994). *Magicicada* larvae feed underground on xylem found in plant roots. Xylem is a transport liquid in plants and is poor in nutrients. The competition for this shared food resource thus likely affects all cohorts feeding on them. Cicada nymphs have been shown to be uniformly spatially distributed, suggesting that cohorts do indeed compete for xylem (Williams and Simon 1995).

Predator satiation is well-documented for periodical cicadas (Williams and Simon 1995, and many references therein). The first cicadas to emerge still face a high predation pressure, with as much as 40% eaten within the first days. As numbers increase dramatically in the days following the start of the event, with up to 3,5 million adult cicadas per hectare, per capita predation pressure drops practically to zero (Williams et al. 1993). The predators, mainly birds such as cuckoos, woodpeckers, jays and other larger birds, do not increase much in number during the year of the outbreak, but several show increased population sizes in the one to three years to follow (Koenig and Liebhold 2005). It has also been shown that for several of these species, predator population sizes are in fact lower right before an outbreak event, thus decreasing predation pressure and allowing the insects to benefit even more from their numerical dominance (Koenig and Liebhold 2013).

In this paper we have focussed purely on the problem of elimination of year classes and how a single cohort arises from a starting situation with multiple cohorts. In our modelling framework we have not allowed for so-called stragglers, insects that have a longer developmental time and thus emerge in the year after an outbreak. This has been investigated recently using predominantly numerical simulations by Blackwood et al. (2018).

The enigma of periodical insect species is not confined to *Magicicada*, but is found also in several other insect orders, notably among *Xestia* moths, and in some well-known beetle species such as *Melonontha* cockchafer beetles (Hellövaara et al. 1994). Life spans may vary between species, and even between populations of the same species. *Magicicada* are exceptional, however, in their life spans, which are either 13 or 17 years, and are among the longest of all insect life spans.

There are several intriguing open theoretical questions regarding periodical insects. The adult insects, having developed under ground over a period of 13 or 17 years, all emerge from their burrows within an extremely short time span, usually between 7 and 10 days (Williams and Simon 1995, and references therein). Just how they synchronise so precisely remains unclear (Williams and Simon 1995), although temperature cues have been suggested.

In some outbreak years, as much as half of all individuals fail to develop into adults, and eclose the next year (White and Lloyd 1979). This has been associated with extremely poor nutrition conditions. Differences in the developmental rate of individual larvae are apparently common, with late-developing nymphs 'catching up' before emerging as adults with the rest. To incorporate such phenomena would require developing a physiologically structured population model, with developmental rate directly influenced by food availability and competition (de Roos and Persson 2013).

References

H. Behncke. Periodical cicadas. *J. Math. Biology*, 40:413–431, 2000.

J. C. Blackwood, J. Machta, A. D. Meyer, A. E. Noble, A. Hastings, and A. M. Liebhold. Competition and stragglers as mediators of developmental synchrony in periodical cicadas. *Am. Naturalist*, 192(4): 479–489, 2018.

M. G. Bulmer. Periodical insects. *Am. Naturalist*, 111:1099–1117, 1977.

J. M. Cushing. Three stage semelparous Leslie models. *J. Math. Biology*, 59:75–104, 2009.

J. M. Cushing. On the fundamental bifurcation theorem for semelparous Leslie models. In J. P. Bourguignon, R. Jeltsch, A. Pintoa, and M. Viana, editors, *Mathematics of Planet Earth: Dynamics, Games and Science*, CIM Mathematical Sciences Series, chapter 11. Springer Verlag, 2015.

J. M. Cushing and S. M. Henson. Stable bifurcations in nonlinear semelparous Leslie models. *J. Biol. Dynamics*, 6:80–102, 2012.

H. V. Danks. Long life cycles in insects. *Can. Entomologist*, 124(1):167–187, 1992.

N. V. Davydova. *Old and young. Can they coexist?* PhD thesis, Utrecht University, 2004. https://dspace.library.uu.nl/handle/1874/891.

N. V. Davydova, O. Diekmann, and S. A. van Gils. Year class coexistence or competitive exclusion for strict biennials? *J. Math. Biology*, 46:95–131, 2003.

N. V. Davydova, O. Diekmann, and S. A. van Gils. On circulant populations. i. the algebra of semelparity. *Lin. Algebra and its Appl.*, 398:185–243, 2005.

A. M. de Roos and L. Persson. *Population and Community Ecology of Ontogenetic Development.* Monographs in Population Biology 51. Princeton University Press, Princeton, NJ, 2013.

O. Diekmann and S. A. van Gils. Invariance and symmetry in a year-class model. In J. Buescu, S. Castro, A.P. Dias, and I. Labouriau, editors, *Bifurcation, Symmetry and Patterns*, Birkhäuser Trends in Mathematics. Birkhäuser, 2003.

O. Diekmann, N. V. Davydova, and S. A. van Gils. On a boom and bust year class cycle. *J. Difference Eq. and Appl.*, 11(4):327–335, 2005.

J. K. Hale. *Asymptotic Behavior of Dissipative Systems.* American Mathematical Society, Providence, RI, 1988.

K. Hellövaara, R. Väisänen, and C. Simon. Evolutionary ecology of periodical insects. *Trends Ecol. Evol.*, 9(12):475–480, 1994.

F. Hofbauer, J. Hofbauer, P. Raith, and T. Steingberger. Intermingled basins in a two species system. *J. Math. Biology*, 49(3):293–309, 2004.

F. C. Hoppensteadt and J. B. Keller. Synchronization of periodical cicada emergences. *Science*, 194 (4262):335–337, 1976.

W. D. Koenig and A. M. Liebhold. Effects of periodical cicada emergences on abundance and synchrony of avian populations. *Ecology*, 86(7):1873–1882, 2005.

W. D. Koenig and A. M. Liebhold. Avian predation pressure as a potential driver of periodical cicada cycle length. *Am. Naturalist*, 181(1):145–149, 2013.

R. Kon. Permanence induced by life-cycle resonances: the periodical cicada problem. *J. Biol. Dynamics*, 6(2):855–890, 2012.

R. Kon. Non-synchronous oscillations in four-dimensional nonlinear semelparous Leslie matrix models. *J. Difference Eq. and Appl.*, 23(10):1747–1759, 2017.

M. Lloyd and H. S. Dybas. The periodical cicada problem. *Evolution*, 20:133–149, 1966.

Jonathan Machta, Julie C Blackwood, Andrew Noble, Andrew M Liebhold, and Alan Hastings. A hybrid model for the population dynamics of periodical cicadas. *Bull. Math. Biol.*, to appear, 2018.

E. Mjølhus, A. Wikan, and T. Solberg. On synchronization in semelparous populations. *J. Math. Biology*, 50:1–21, 2005.

G. F. Webb. The prime number periodical cicada problem. *Disc. Cont. Dyn. Sys. B*, 1:387–399, 2001.

J. White and M. Lloyd. Seventeen year cicadas emerging after 18 years: a new brood? *Evolution*, 33 (1193-1199), 1979.

A. Wikan. An analysis of a semelparous population model with density-dependent fecundity and density-dependent survival probabilities. *J Appl. Math.*, 2017:Article ID 8934295, 14 pages, 2017.

K. S. Williams and C. Simon. The ecology, behavior, and evolution of periodical cicadas. *Ann. Rev. Entomol.*, 40:269–295, 1995.

K. S. Williams, K. G. Smith, and F. M. Stephen. Emergence of 13-yr periodical cicadas (Cicadidae: Magicicada): Phenology, mortality, and predators satiation. *Ecology*, 74(4):1143–1152, 1993.