6 May 2019

Nitrogen addition pulse has minimal effect in big sagebrush *(Artemisia tridentata)*

communities on the Pinedale Anticline, Wyoming (USA)

Running head: sagebrush nitrogen fertilization

CHRISTOPHER W. BELTZ[1,2]**,** *Yale University School of Forestry & Environmental Studies,*

195 Prospect St., New Haven, CT 06511, USA

MEGAN L. MOBLEY, *Department of Crop and Soil Science, Oregon State University, Oregon*

State University, Corvallis, OR 97331, USA (meg.mobley@alumni.duke.edu)

INGRID C. BURKE, *Yale University School of Forestry & Environmental Studies, 195*

Prospect St., New Haven, CT 06511, USA (indy.burke@yale.edu)

[1]*E-mail: christopher.beltz@yale.edu*

[2]*Present address: 370 Prospect St., Yale University School of Forestry & Environmental Studies,*

Greeley Memorial Laboratory, New Haven, CT 06511, USA

KEYWORDS *Artemisia tridentata*, *Odocoileus hemionus*, mule deer, dryland, energy

mitigation, sagebrush, nitrogen, fertilization

ABSTRACT

Nitrogen additions are known to elicit variable responses in semi-arid ecosystems, with

responses increasing with precipitation. The response of semi-arid ecosystems to nitrogen are

important to understand due to their large spatial extent worldwide and the global trend of

increasingly available nitrogen. In this study, we evaluated the impact of a single nitrogen

addition pulse on a semi-arid big sagebrush (*Artemisia tridentata*) ecosystem in western

Wyoming. This is important given that sagebrush ecosystems are poorly understood, despite

their prevalence in the western US. In addition, large-scale nitrogen additions have begun on

sagebrush landscapes in Wyoming in order to mitigate population declines in mule deer

(*Odocoileus hemionus*). The study objectives were (1) to evaluate the effectiveness of a nitrogen

fertilization pulse in increasing sagebrush biomass and forage quality, and (2) to assess effects of

nitrogen addition on soil biogeochemistry and vegetation community structure. We fertilized 15

plots across 5 locations in western Wyoming using a single pulse of urea ($5.5g$ N m^{-2}). In

addition, we immobilized available nitrogen through surface hay treatments ($250g$ hay/m^2).

Nitrogen additions failed to increase growth of sagebrush, alter nitrogen content of sagebrush

leaders, or alter greenhouse gas efflux from soils. The plant community also remained

unchanged; total cover, species richness, and community composition were all unaffected by our

treatment application. Over the two years of this study, we did not find indications of nitrogen

limitation of ecosystem processes, despite a wet growing season in 2014. Thus, we have found a

general lack of response to nitrogen in sagebrush ecosystems and no treatment effect of a single

pulse of N to sagebrush biomass or forage quality.

INTRODUCTION

Nitrogen (N) additions are known to elicit varying responses in semi-arid ecosystems (1).

Ecosystems respond to N in multiple ways, including increased net primary production and plant

growth (2,3), plant community composition changes (4), and increased greenhouse gas (GHG)

emissions from soils (5). Most semi-arid and arid systems (henceforth 'drylands') are thought to

be primarily limited by water, though this limitation can be affected by both intra- and inter-

annual variation in precipitation (6). If drylands were only restricted by water-limitation, N

additions to the system should have no effect without a concurrent increase in precipitation.

However, most dryland systems show a response to N additions alone, indicating at least a

partial limitation by N (1,7).

In a meta-analysis of research studies on the effects of N fertilization studies in water-

limited drylands worldwide, Yahdjian et al. (8) found that in almost all cases the addition of N

increased net primary productivity (NPP), showing that nitrogen limitation – even if minor – is a

global phenomenon. The ability to predict the effects of increased N on ecosystem process and

structure is important because of the global trend of increasing N deposition (9). Given that

drylands cover 40% of the Earth's terrestrial surface (10,11) and 20% of the US (12), it is critical

to investigate the effects of increasing N availability in dryland systems, due to their co-

limitation by both water and N. In addition, dryland systems are home to 44% of all cultivated

land (13,14), making understanding the effects of increasing N particularly important due to the

high use of N-based fertilizers.

In the western US, big sagebrush (*Artemisia tridentata* spp.) ecosystems are the dominant

dryland system (15,16), covering 48 million hectares (ha) and 13 states (17). Despite its current

extent in the US, up to 60% of the historic, pre-settlement big sagebrush range has been partially

or completely converted to exotic annual grasslands because of increased disturbance to these

systems and conversion to grazing and agricultural lands (18). In addition, significant portions of

the big sagebrush range continue to be affected by energy development, agriculture, and

expansion of urban areas (19–22). These drivers of the loss of big sagebrush — unsustainable

use of natural resources, population expansion, globalization, as well as climate change — are

the same as those exacerbating desertification in drylands globally (13). Big sagebrush

ecosystems are expected to continue to decline in coverage over the next decade with 2.3-5.5

million hectares projected to be impacted due to oil and gas development (23). While agricultural

additions of N account for a majority of the increase in N availability globally, energy

development also affects N deposition locally, through the production and release of NO_x

compounds (9,24,25).

The effects of nitrogen additions on big sagebrush ecosystems are poorly understood due

to both the small number of studies and the complexities associated with potential co-limitation

of ecosystem processes by water and N in dryland systems (26). Although current knowledge

gaps exist regarding N limitation in dryland ecosystems, increased N availability is widely

known to have environmental consequences in other ecosystems (5). These include: (1)

increasing the amount of carbon stored in terrestrial biomass in both soil and aboveground plant

biomass (27,28), (2) increasing soil efflux of GHGs (29), and (3) changing the species

composition of plant communities (30). Over the past decade, it has become increasingly

important to understand the role of nitrogen in sagebrush ecosystems. Our research will help to

close the gap in knowledge on the effects of N fertilization in sagebrush.

Mule Deer: A Case Study in the Nitrogen Limitation of Drylands

The Pinedale Anticline of western Wyoming is one of the most disturbed regions of big

sagebrush. Land managers are particularly concerned with large population declines in mule deer

(*Odocoileus hemionus*) over the 15 years since the onset of natural gas development (31). The

direct loss of 615 ha of big sagebrush habitat, and the associated winter forage, appear to be

associated with the increased mortality in the Sublette Mule Deer Herd (32). By 2010, this

population decline had been of sufficient magnitude that the Pinedale Anticline Project Office

(PAPO) was required to initiate on-site mitigation.(26). Land managers hypothesized that

fertilizing the area with N would increase habitat quality (33) through increases in big sagebrush

biomass (i.e., forage production) and N content within the forage (i.e., forage quality).

The Pinedale Anticline is over 12,000 ha, with 320 ha in the pilot fertilization area

(26,32). A single pulse of N fertilizer was applied by helicopter in 2010 and 2011. In 2010, two

100-acre plots were fertilized at 45 and 90 kg/ha. In 2011, a separate 400-acre plot was fertilized

at the lower of the two rates (45 kg/ha). Each fertilization application was projected to cost

US$135/ha – totaling greater than US$1.6m per year. Due to the high cost, potential

environmental effects, and the expected dominance of water limitation, some ecologists argued

that fertilization of big sagebrush was unlikely to be effective in offsetting forage loss for mule

deer (26). The eventual monitoring report of the fertilization project reached a similar conclusion

(34), however, as the study was conducted on a previously applied and un-replicated treatment,

no generalizable conclusions could be reached.

N availability is likely to increase through the end of the century, largely due to the

chronic addition of N to environments through N deposition (9). The chronic application (i.e.,

"press") of N is likely to yield different effects than a single pulse, which we applied in this

study to mimic the PAPO protocol (35,36). However, the development of a N fertilization

project on the Pinedale Anticline provides a rare opportunity to ask basic research questions

about N limitation in drylands, while simultaneously helping to inform decisions about a regional

environmental issue.

Project Goals & Hypotheses

Inspired by the actions on the Pinedale Anticline, the goals of this study were (1) to

evaluate the efficacy of the addition of a N pulse for increasing big sagebrush biomass and

forage quality, and (2) to assess effects of N addition on soil biogeochemistry, plant biomass, and

plant community composition. We hypothesized that the N fertilization pulse would increase N

availability in the soil, however it would not increase growth, biomass, or forage quality of big

sagebrush due to primary limitation by water. In addition, we expected changes in plant

community composition and a high potential for increased GHG emission.

MATERIALS AND METHODS

Study area and site selection

We conducted this study between 2012 and 2015 in semi-arid big sagebrush plant

communities in western Wyoming, near the town of Pinedale (42°51′58″ N 109°51′53″ W). The

area surrounding Pinedale is consistent with sagebrush steppe in other parts of Wyoming; plant

communities are co-dominated by Wyoming big sagebrush (*Artemisia tridentata wyomingensis*)

and mountain big sagebrush (*Artemisia tridentata vaseyana*). Other common plant species

present include Sandberg bluegrass (*Poa secunda*), western wheatgrass (*Pascopyrum smithii*),

chaffweed (*Antennaria microphylla*), squirreltail (*Elymus elymoides*), and longleaf phlox (*Phlox*

longifolia).

The soils in this area are categorized as Typic Haplocryalfs, Typic Dystrocryepts, and

Typic Haplocryolls with aridic moisture regimes (37). Soil textures at our sites were categorized

as loam, sandy loam, and sandy clay loam (Table 1). Mean annual precipitation across the

region averages 290 mm and mean annual temperature is 2.6 ℃ (Table 2) (38). The precipitation

in this region falls predominantly in the spring and summer months.

Table 1. Soil texture classes and sand/silt/clay content for each exclosure.

Ranches are the first letter in the ID (i.e., B, P, or S), site is the number in the second position,

and the third letter corresponds to an individual plot (i.e., A, B, or C).

Plot ID	Sand (%)	Clay (%)	Silt (%)	Soil Class
1-1A	26	24	49	loam
1-1B	43	24	32	loam
1-1C	43	16	40	loam
1-2A	41	16	42	loam
1-2B	47	14	38	loam
1-2C	35	20	44	loam
2-1A	54	14	30	sandy loam
2-1B	66	14	18	sandy loam
2-1C	54	16	28	sandy loam
2-2A	50	16	32	loam
2-2B	58	20	21	sandy clay loam
2-2C	63	10	26	sandy loam
3-1A	67	8.5	24	sandy loam
3-1B	67	10	22	sandy loam
3-1C	65	12	22	sandy loam

138 **Table 2. Annual precipitation (mm) for the three Pinedale, WY area ranches over the**

139 **course of the experiment, 2012-2015.**

140 Precipitation values were derived using interpolated values from the PRISM Climate Group,

141 Oregon State University, http://prism.oregonstate.edu, created 13 Feb 2019.

	precipitation (mm)			
Ranch	**2012**	**2013**	**2014**	**2015**
1	174	257	381	286
2	209	295	423	354
3	185	253	378	294

We used NRCS Soil Survey data to identify five site locations across three privately

owned cattle ranches, all within 25 km of the Pinedale Anticline and 10-20 km of each other.

Sites were distributed to represent the spatial extent of the proposed mitigation and fertilization

treatments (geographic coordinates for specific sites are not provided due to privacy concerns).

We obtained permission from three local land-owners to use their privately-owned lands because

of constraints associated with imposing manipulations on public areas adjacent to the Pinedale

Anticline. We identified site locations with similar soil types, textures, and landscape positions,

with all sites positioned on a slight slope. Each site had previously been moderately grazed for

over a century; this grazing history is typical of the region, big sagebrush plant communities

throughout the western U.S., and - most importantly – consistent with the history of the Pinedale

Anticline, for which we are trying to characterize potential effects of N fertilization.

Experimental design and nitrogen fertilization

156   To address our hypotheses, we applied three treatments (nitrogen fertilization, nitrogen

removal through immobilization, and a no-addition control) in a generalized randomized

complete block design with split-plots. Treatments were randomly applied to subplots and

replicated in three main plots; this split-plot setup was replicated at five sites (blocks). The

hierarchy of locations is site > main-plot > subplot. The five replicate sites were located on three

privately-owned ranches. Within each site, we created three plots (10m x 10m). Each main-plot

was enclosed by cattle-proof fencing and contained three subplots (2.5m x 8.5m) each randomly

assigned one of the above treatments. Subplots were fertilized with N in the form of urea (4.5g N

m^{-2}) and surface hay (well-aged for at least one year) was applied (referred as 'nitrogen removal'

here; 250g hay m^{-2}) for the above respective treatments in September 2012. The addition of hay

to the surface immobilizes soil inorganic N, through incorporation of N into microbial biomass

during decomposition of the hay (39–41). All subplots had a 0.56 - 0.625m buffer between

subplots and between subplot edge and fence.

Mineralized Soil Nitrogen

171   Inorganic nitrogen availability, in the form of nitrate (NO_3^-), and ammonium (NH_4^+),

were estimated *in situ* using Plant Root Simulator (PRS®) probes. The PRS® probes are ion

exchange resin membranes held within a rigid plastic frame (Western Ag Innovations, Inc.,

Saskatoon, Canada) and provide an index of nitrogen availability to plants. PRS probes provide

an index of soil N availability that integrates across the time of their deployment (i.e. soil N

availability from Oct-May for this study). PRS probes are placed in pairs, one anion probe

(collecting NH_4^+) and one cation probe (collecting NO_3^-). Multiple pairs were placed within a

single treatment. All probes were washed with DI water following the manufacturer's protocol

and shipped to Western Ag Innovations, Inc., (Saskatoon, Canada) for analysis.

In order to estimate available soil nitrogen (N) during the first year of the experiment

(Sept 2012 – May 2013), we placed PRS® probes within each treatment block at all fifteen plots.

We installed probes over winter to catch the spring snowmelt, the largest pulse of nitrogen for

the year (42), as plots are inaccessible from winter until soils dry out in the spring.

During the first year they were placed on the edge (i.e., cusp) of the canopy of a big

sagebrush individual. During the third year of the study (Oct 2014 – May 2015), probes were

placed at two locations to measure soil N available under both the big sagebrush canopy and

within the interspace between individual shrubs. For example, if an individual sagebrush had a

canopy that extended 20cm from its main stem (i.e., trunk-like base) then the Year 1 probes

would be placed at 20cm (i.e., "cusp"), while the Year 3 probes would be placed at 10cm (i.e.,

"under") and 30cm (i.e., "interspace" or 10cm outside the canopy) from the stem. Probe

placement was stratified by shrub location in an effort to represent a key component of soil N

variability in drylands, as N availability is known to be elevated under shrub canopies (43). This

was of particular concern as time progressed and the effects of our pulse of N fertilizer on soil N

were likely to decrease.

The sets of probes from each sampling campaign (i.e., Year 1 and Year 3) were retrieved

from the field and analyzed after their 7-8 month deployment. A limitation of PRS probes is that

temperature and soil moisture affect the ability of the anion and cation resin to adsorb NH_4^+ and

NO_3^-. Thus, PRS probes provide an index of N availability that can be compared among

treatments within a sampling campaign, but cannot be compared between sampling campaigns.

As such, our analyses are limited to comparisons within a campaign (i.e., within Year 1 or within

Year 3).

Sagebrush Growth & Forage Quality

We estimated big sagebrush annual growth by measuring leader growth (i.e., big

sagebrush stems) and the ephemeral branch growth (i.e., inflorescences) in 2014. Leaders and

ephemeral branches were collected at the end of the growing season in late October 2014 and

dried in the laboratory for five days at 55°C and then at 35°C until mass remained constant over

a 5-day period. We then weighed and measured the length of each individual leader and branch.

The dried leaders were ground in a ball mill and submitted to the Stable Isotope Facility at the

University of Wyoming for total carbon and nitrogen analysis (44), as a proxy for forage quality.

Plant Community

We sampled the plant communities at each site over two days in August 2014. We

surveyed the plant community using 20 randomly placed Daubenmire quadrats (0.1 m^2) within

each subplot. The canopy cover of each plant species was estimated using a modified

Daubenmire (45) method with the following cover categories: <1%, 1-5%, 6-15%, 26-40%, 41-

60%, and >60%. Unknown species were marked as unknown in the field and identified using the

Rocky Mountain Herbarium (Univ. of Wyoming). If the unknown species were unable to be

identified they were not included in the analyses of the plant community.

Soil Trace Gas Emission

We sampled soil trace gas emissions twice in July 2013; each was at least 48 hours after

any precipitation because high amounts of water vapor can be can confound measurement of

CO_2. All sites were sampled within a single day for each sampling period. We deployed 90 soil

gas flux chambers, 2 per subplot across all 15 sites, and arranged them on the cusp of a random

interspace and adjacent big sagebrush canopy, as it is well-recognized that a shrub's location can

create important environmental microsites in mountain big sagebrush ecosystems (43).

Sections of polyvinyl chloride (PVC) pipe (10 cm high x 20 cm diameter) were

permanently buried in the ground to a depth of 6 ± 2cm, which then acted as bases for the

chamber lid. Chamber lids were round, 20cm inside diameter and 8cm deep, made of opaque

PVC and vented following Hutchinson & Mosier (46). The chamber base and lid fit snugly

together and were sealed using a rubber gasket for the 30-minute sampling period. Once the

chamber was sealed, 30 ml samples of air were drawn from the chamber headspace every 15

minutes (t = 0, 15, and 30 mins). Each sample was collected by drawing 60 ml of air from the

headspace into a 60 ml syringe (Allison Medical, Inc.), expelling 30ml into the air outside the

chamber, and then injecting as much as possible of the remaining 30ml sample into a 12ml flat

bottomed, pre-evacuated glass vial (Exetainer, Labco Ltd.) sealed with a rubber septum.

Ambient air samples were collected at each plot following the same procedure above and stored

with the sample vials to account for any systemic change during the sample period. Standards of

known chemical composition were also carried along on field sampling campaigns and analyzed

along with the field samples to account for any systemic change or contamination. Gas samples

were analyzed in the laboratory using gas chromatography on a Shimadzu GC-2014 to measure

concentrations of CO_2, CH_4, and N_2O.

Data Analysis

We conducted all data analysis and figure creation using R version 3.5.3 (R core Team,

2018), using the following libraries: knitr, tidyverse, lubridate, plotrix, emmeans, ecodist, labdsv,

nlme, Rmisc, vegan, soiltexture, and here.

We used a linear mixed effects model (*lme* function, 'nlme' package v 3.1-139 (47)) to

examine the fixed effects of treatment with the nested, random effects of site and plot (i.e., y ~

treatment + 1 | site / plot). Treatment effects were first evaluated using a one-way analysis of

variance (ANOVA), followed by an examination of contrasts to determine the effectiveness of

individual N and hay treatments. Finally, we conducted a post-hoc evaluation of the 95%

confidence intervals for the response to each treatment using estimated marginal means

(*emmeans* function, 'emmeans' package v1.3.4 (48)).

The mixed effects model was used to evaluate the effects of our N and hay treatments on

inorganic soil N availability (total inorganic N, NO_3^-, NH_4^+), sagebrush growth (leader length,

inflorescence length, leader dry mass, and leader dry density), and forage quality (percent C,

percent N, C:N ratio, leader N mass, leader C mass).

Soil trace gas fluxes (CO_2, CH_4, N_2O) were calculated for each treatment and plot using

the rate of change for each of the gases analyzed over the 30-minute period, calculated as the

slope values of a linear regression. These fluxes were then similarly analyzed using the mixed

effects model previously described.

We examined the effects of N and hay treatments on the plant community by evaluating

differences in total plant cover and species richness using the same mixed effects model and

associated processes. To explore the effects of treatment applications on plant community

composition, we used a permutational multivariate analysis of variance (PERMANOVA). Total

cover and abundance by species was assessed by summing the cover class midpoint for each

species by site to obtain the total percent cover for each species and then standardized these

values to relative abundance (*decostand* function, 'Vegan' package v2.5-4 for R (49)). We then

used a PERMANOVA to evaluate differences in community composition between treatments

using a Bray-Curtis dissimilarity index and 9999 permutations (*adonis* function, 'Vegan'

package v2.5-4 for R).

We also explored the effects of total soil N on leader length through calculating a one-

way analysis of covariance (ANCOVA) with site and plot as nested, random factors. This is

similar to the original model described above except treatment is removed, as it is confounded

with soil N availability. We then obtained fitted values at the population and hierarchical level

(i.e., individual sites).

RESULTS

Total Inorganic N in Soil

The addition of N significantly increased the availability of total inorganic N in soils ($p<0.001$;

Fig 1) during the first year of the experiment, while hay addition significantly reduced the

availability of N (p= 0.035; Fig 1). No differences were found for availability of NH_4^+ between

treatments, however NO_3^- was significantly more available in the N fertilized treatments

(p<0.001) than either the control or N removal (p=0.035). By the end of second year of the

experiment, no significant differences remained between treatments for total inorganic N, NO_3^-,

or NH_4^+. During year one, NO_3^- represented 99.0% ± 0.4 of the total available inorganic N across

all treatments/locations, while in year two it was 92.0% ± 1.7.

Fig 1. Mean total soil inorganic N availability within a big sagebrush plant community in

Pinedale, WY.

N fertilization occurred in 2012 with soil available inorganic N measured via PRS probes in

fall/winter/spring of 2012-2013 and 2014-2015 (n=5 sites). The 2013 data were collected from

soil on the *cusp* of the canopy and interspace, while 2015 data were collected separately from

within the *interspace* and *under the canopy*. Significant differences from the control, within each

year and location, are indicated by asterisk based on $\alpha = 0.05$ (* indicates p<0.05, *** indicates

p<0.001). Whiskers indicate the standard error based on among-site variation.

Leader Growth & Nitrogen Content

The addition of N and the application of hay had no treatment effect on the growth of big

sagebrush leaders; they were unchanged in length, mass, and density. While the treatment term

was significant in predicting leader length (p=0.006), neither the nitrogen additions (p=0.103)

nor hay treatment (p=0.079) were significantly different than the control when examining

contrasts. Ephemeral branches (i.e., inflorescences) were unchanged by the addition of N

fertilizer, though the addition of hay reduced their mean length by approximately 8% or 1.1cm

(p=0.029; Fig 2). Leader C:N ratio and N content (i.e., forage quality) were also unchanged by

treatment application.

Fig 2. Growth of sagebrush leaders was unaffected by treatment application, while

inflorescences decreased in length with hay addition.

Treatment is indicated by color (gray=control, khaki=hay addition, blue=nitrogen addition).

a) Leader length, b) leader dry mass, c) inflorescence length, d) leader C:N ratio, e) leader N

content in percent, and f) leader N content in grams.

Exploring the relationship between soil total N availability (Fig 3), we found that the

coefficient for soil N was significantly different from zero (slope=0.00122 ± 0.0003259 SE,

t=3.74365, p=0.0008).

Fig 3. Relationship between sagebrush leader growth and total inorganic N for each of five

sites in Pinedale, WY

Ranch location is indicated by color, while sites within a ranch differ in shape (note: Ranch 3 has

only a single site). Solid lines correspond to the site with the solid color shapes and dashed lines

correspond to the hollow/un-filled shapes. Data presented is across all treatments.

Plant Community

The three ranches each had similar big sagebrush density, typically ranging from 20-30%

of cover. The most common understory species, by cover, across the controls plots within the

three ranches included *Chrysothamnus viscidiflorus*, *Pascopyrum smithii*, *Poa secunda*,

Achnatherum pinetorum, *Hesperostipa comata*, and *Phlox hoodii*. All common understory

species were found at each site, though their prevalence did vary slightly between ranches.

We found no treatment effect on total plant cover or species richness (Fig 4). The plant

community also remained unchanged, regardless of the treatment application.

Fig 4. Total plant cover and species richness were unaffected by treatment application.

Treatment is indicated by color (gray=control, khaki=hay addition, blue=nitrogen addition).

a) species richness, and b) plant abundance

Trace Gas Emission from Soil

Soil trace gas emission displayed no effect from treatments for any gas species (Fig 5).

Neither the addition of hay or N altered the efflux of CO_2, CH_4, or N_2O from the soil surface.

Fig 5. Trace gas emissions (CO_2, CH_4, & N_2O) were unaffected by treatment application.

Treatment is indicated by color (gray=control, khaki=hay addition, blue=nitrogen addition).

DISCUSSION

N fertilization significantly increased the availability of N in soil for the first-year post-treatment,

however there was no significant difference in N availability by the second year. Differences in

control values between year one and two are not surprising given the different locations; length

of deployment, soil temperature, and soil moisture can all affect the adsorption of anion and

cations to their counter-ion resins, making comparisons between years (and conditions)

inappropriate to a certain extent. Precipitation between the two years, 2012 and 2014, varied by

approximately 200mm depending on the ranch examined and could account for a portion of the

variability seen in the control plots between years. Due to differing conditions between years, we

have focused on comparing PRS probe measurements of total inorganic N within single years.

The lack of significant increase in soil N availability in the second year in fertilized plots

is likely due to 'normal' N losses through N mineralization, gaseous loses (volatilization and

microbially-mediated), and leaching. Additions of N were conducted at 4.5 g N m^{-2} (or 45 kg N

ha^{-1}), which is 150-900% of annual N mineralization (43). Trace gas emissions of N-containing

gases (N_2O and NO_x) accounted for an additional 10-20 kg N ha^{-1} yr^{-1} loss from the soil, which

could contribute to the lack of increase observed in inorganic soil N among treatments in year

two (5,26). While we found lower fluxes of N_2O compared to these studies one year after

fertilization, it may be that fluxes were greater immediately after fertilization or during the spring

when water was more available, therefore not captured in our sampling period.

Increased N availability had minimal effect on the length of big sagebrush leaders over

the two years of the study. Additional N did not result in an increase in forage quality; C:N ratio

and mass of N were the same between the control and N fertilized plots. While the addition of

hay minimally decreased the length of ephemeral branches, the addition of N had no effect. It is

possible that the decrease in soil N caused by the hay treatments decreased growth by increasing

N limitations. These limitations on the growth of inflorescences by N may have been more

apparent during 2014, when water was more available.

Our prediction that there would be no differences in plant growth between treatments was

generally supported. Conversely, we also found that leader length was weakly positively

correlated with inorganic soil N availability at each of our sites. On the face of it, these two

statements are contradictory. While we have shown that our treatments significantly affected the

availability of total N in soil across all sites, there was no associated treatment effect on plant

growth. Leader growth correlated with N availability at the site level. It is possible that other

factors may drive or mediate plant growth at these sites. Site-specific factors such as soil texture

(i.e., clay), precipitation, and leaching can affect the conversion of urea fertilizer to N, the

retention of that in soil, and/or to the uptake and allocation of that N by plants to increased leader

growth.

379   Interestingly, differences in leader length between ranches was strong. While this could

380   be due to many factors, the ranch with highest precipitation in each year from 2012-2015 also

381   had the longest sagebrush leaders. However, we did not measure precipitation at any of our sites

382   or ranches, making it impossible to assert this with any confidence.

383   Contrary to our predictions, the addition of a single pulse of N did not alter plant

community composition. *P. smithii* (western wheatgrass) is known to be nitrophilic and respond

to additions of N (50,51), however no significant change was observed. In addition, treatments

did not affect either total plant cover or species richness. This is not surprising as soil N

availability was only increased for the first year after treatment. In addition, drylands – and big

sagebrush ecosystems – are primarily water limited, making short-lived increases of N unlikely

to affect cover or richness.

390   There was no effect of N additions on GHG efflux (i.e., CO_2, CH_4, and N_2O) from soil.

Fluxes of CO_2 and N_2O fell into similar ranges for dry, unwetted sagebrush reported by Norton

et al. (52). The lack of significant treatment effect could be due to a number of reasons: (1) trace

gas measurements are highly variable and sensitive to a number of different environmental

factors; (2) increased GHG fluxes occurred during a time that was not measured, potentially

closer to fertilization or during the spring when water is more available; or (3) the ability of the

microbial community to produce CO_2, CH_4, and N_2O may be limited primarily by water

availability, not N availability.

Our findings are not consistent with other studies, which have generally found a positive

response of plant productivity to N fertilization (1,8) indicating some N limitation. However

given that the mean annual precipitation of the Pinedale, WY area is 290mm, it is not surprising

that we found no response of leader growth to N additions; Hooper and Johnson (1) found that

the response to N was greater for areas with 450-900 mm of precipitation than for 300 mm. It is

important to note that while most studies find a primary water limitation in drylands (53), the

response of dryland ecosystems to N can be expected to increase with increased precipitation

(54).

Our results indicate that a one-time application or single pulse of N does not affect soil N

availability in the long term within the sagebrush communities of the Pinedale Anticline. We

suggest that annual fertilization would be necessary to maintain high levels of soil N availability

within big sagebrush systems, as the levels of soil N were the same between the fertilized and

control plots by the second year. Yet increases in N availability over the long term may have

negative consequences, including higher volatile N losses, and increases in invasive plants (26).

It should be noted that the chronic additions of N fertilizer would likely have different effects

than that of the single pulse of N applied during this study. This study has addressed the effects

of a single pulse of N, such as the PAPO pilot study on the Pinedale Anticline. The increased

levels of N found in the first year after fertilization only minimally increased the length of big

sagebrush leaders, and continued application at that level would be unlikely to increase forage

quality. Ecosystem processes in drylands are primarily water limited and do not change with

short-term N additions, at least over the two-year duration of this study.

CONCLUSIONS

This study was conducted to investigate the role of N limitation in dryland sagebrush ecosystems

and to consider the effect of a pulse of N addition to increase forage and forage quality for mule

deer. We evaluated responses of soil N availability, annual growth rate of big sagebrush, and

forage N concentration in response to N additions. In addition, we investigated the response of

trace gas production and plant community structure to increased N. This study demonstrates that

a single addition of N, in the form of urea fertilizer, does increase N availability in soil for one

year after fertilization. However, this increased soil N availability is only loosely associated with

increased growth of big sagebrush and not likely to impact forage availability and quality for

mule deer. The results of this study suggest that the addition of N in a single pulse is likely a

poor choice for mitigation of mule deer population declines on the Pinedale Anticline due to the

minimal gains in forage, if any, and associated high cost.

ACKNOWLEDGEMENTS

Thank you to B. Amgalan, O. Avirmed, D. Bell, M. Cleary, T. Martyn, K. Palmquist, V.

Pennington, B. Peterson, and C. Rottler for assistance with field work, data collection, lab work,

and manuscript review. Additional thanks to D. Schlaepfer and T. Gregoire for help with data

analysis. Thank you to our three private landowners for making this research possible, including

Freddie Botur and the Cottonwood Ranch in Big Piney, as well as the Sommers Ranch in

Pinedale. An earlier poster version of this study was presented at the Ecological Society of

America conference in 2015. Many thanks to those attendees that provided feedback and

encouragement. Financial support for this research came from the University of Wyoming and

the Wyoming Excellence Fund.

REFERENCES

1. Hooper DU, Johnson L. Nitrogen limitation in dryland ecosystems: Responses to geographical and temporal variation in precipitation. Biogeochemistry. 1999 Jul 1;46(1–3):247–93.

2. Lauenroth WK, Dodd JL, Sims PL. The effects of water- and nitrogen-induced stresses on plant community structure in a semiarid grassland. Oecologia. 1978 Jan 1;36(2):211–22.

3. LeBauer DS, Treseder KK. Nitrogen Limitation of Net Primary Productivity in Terrestrial Ecosystems Is Globally Distributed. Ecology. 2008 Feb 1;89(2):371–9.

4. Reich PB. Elevated CO_2 Reduces Losses of Plant Diversity Caused by Nitrogen Deposition. Science. 2009 Dec 3;326(5958):1399.

5. Vitousek PM, Aber JD, Howarth RW, Likens GE, Matson PA, Schindler DW, et al. Human Alteration of the Global Nitrogen Cycle: Sources and Consequence. Ecol Appl. 1997;7(3):737–50.

6. Sala OE, Gherardi LA, Reichmann L, Jobbágy E, Peters D. Legacies of precipitation fluctuations on primary production: theory and data synthesis. Philos Trans R Soc B Biol Sci. 2012 Nov 19;367(1606):3135–44.

7. Mobley ML, Cleary MJ, Burke IC. Inorganic Nitrogen Supply and Dissolved Organic Nitrogen Abundance across the US Great Plains. PLoS ONE. 2014 Sep;9(9):1–5.

8. Yahdjian L, Gherardi L, Sala OE. Nitrogen limitation in arid-subhumid ecosystems: A meta-analysis of fertilization studies. J Arid Environ. 2011 Aug 1;75(8):675–80.

9. Galloway JN, Townsend AR, Erisman JW, Bekunda M, Cai Z, Freney JR, et al. Transformation of the Nitrogen Cycle: Recent Trends, Questions, and Potential Solutions. Science. 2008 May 16;320(5878):889–92.

10. Wang L, D'Odorico P, Evans JP, Eldridge DJ, McCabe MF, Caylor KK, et al. Dryland ecohydrology and climate change: critical issues and technical advances. Hydrol Earth Syst Sci. 2012 Aug 9;16(8):2585–603.

11. S. Feng, Q. Fu. Expansion of global drylands under a warming climate. Atmospheric Chem Phys. 2013 Oct;13(19):10081–94.

12. Sobecki TM, Moffitt DL, Stone J, Franks CD, Mendenhall AG. Chapter 2: A Broad-Scale Perspective on the Extent, Distribution, and Characteristics of U.S. Grazing Lands. In: Follet RF, Kimble JM, Lal R, editors. The Potential of US Grazing Lands to Sequester Carbon and Mitigate the Greenhouse Effect. Boca Raton, FL: CRC Press; 2001.

13. Millenium Ecosystem Assessment. Ecosystems and Human Well-being: Desertification Synthesis. Washington, DC: World Resources Institute; 2005.

14. Reynolds JF, Smith DMS, Lambin EF, Turner BL, Mortimore M, Batterbury SPJ, et al.
Global Desertification: Building a Science for Dryland Development. Science. 2007 May
11;316(5826):847–51.

15. West NE. Intermountain deserts, shrub steppes, and woodlands. North Am Terr Veg.
1989;209–30.

16. Palmquist KA, Schlaepfer DR, Bradford JB, Lauenroth WK. Mid-latitude shrub steppe
plant communities: climate change consequences for soil water resources. Ecology. 2016
Sep 1;97(9):2342–54.

17. Connelly JW, Knick ST, Schroeder MA, Stiver SJ. Conservation Assessment of Greater
Sage-grouse and Sagebrush Habitats. Cheyenne, Wyoming: Western Association of Fish
and Wildlife Agencies; 2004.

18. Knick ST, Dobkin DS, Rotenberry JT, Schroeder MA, Haegen WMV, van Riper C.
Teetering on the Edge or Too Late? Conservation and Research Issues for Avifauna of
Sagebrush Habitats. The Condor. 2003;105(4):611–34.

19. Walston LJ, Cantwell BL, Krummel JR. Quantifying spatiotemporal changes in a sagebrush
ecosystem in relation to energy development. Ecography. 2009 Dec 1;32(6):943–52.

20. Davies KW, Bates JD, Nafus AM. Vegetation Response to Mowing Dense Mountain Big
Sagebrush Stands. Rangel Ecol Manag. 2012 May 1;65(3):268–76.

21. Manier DJ, Wood DJA, Bowen ZH, Donovan RM, Holloran MJ, Juliusson LM, et al.
Summary of science, activities, programs, and policies that influence the rangewide
conservation of Greater Sage-Grouse (Centrocercus urophasianus): U.S. Geological Survey
Open-File Report 2013–1098 [Internet]. 2013 p. 170. Available from:
http://pubs.usgs.gov/of/2013/1098/.

22. Schlaepfer DR, Lauenroth WK, Bradford JB. Natural Regeneration Processes in Big
Sagebrush (Artemisia tridentata). Rangel Ecol Manag. 2014 Jul 1;67(4):344–57.

23. Copeland HE, Doherty KE, Naugle DE, Pocewicz A, Kiesecker JM. Mapping Oil and Gas
Development Potential in the US Intermountain West and Estimating Impacts to Species.
PLoS ONE. 2009 Oct 14;4(10):e7400.

24. Naftz DL, Schuster PF, Johnson CA. A 50-year record of NOx and SO2 sources in
precipitation in the Northern Rocky Mountains, USA. Geochem Transit. 2011;12(4):1–10.

25. McMurray JA, Roberts DW, Fenn ME, Geiser LH, Jovan S. Using Epiphytic Lichens to
Monitor Nitrogen Deposition Near Natural Gas Drilling Operations in the Wind River
Range, WY, USA. Water Air Soil Pollut. 2013 Mar 1;224(3):1487.

26. Korfanta NM, Mobley ML, Burke IC. Fertilizing western rangelands for ungulate
conservation: An assessment of benefits and risks. Wildl Soc Bull. 2015 Mar 1;39(1):1–8.

27. Frey SD, Ollinger S, Nadelhoffer K, Bowden R, Brzostek E, Burton A, et al. Chronic
nitrogen additions suppress decomposition and sequester soil carbon in temperate forests.
Biogeochemistry. 2014 Nov 1;121(2):305–16.

28. Coskun D, Britto DT, Kronzucker HJ. Nutrient constraints on terrestrial carbon fixation:
The role of nitrogen. J Plant Physiol. 2016 Sep 20;203:95–109.

29. Gomez-Casanovas N, Hudiburg TW, Bernacchi CJ, Parton WJ, DeLucia EH. Nitrogen
deposition and greenhouse gas emissions from grasslands: uncertainties and future
directions. Glob Change Biol. 2016 Apr 1;22(4):1348–60.

30. Milchunas DG, Lauenroth WK. Inertia in Plant Community Structure: State Changes After
Cessation of Nutrient-Enrichment Stress. Ecol Appl. 1995 May 1;5(2):452–8.

31. Sawyer H, Korfanta NM, Nielson RM, Monteith KL, Strickland D. Mule deer and energy
development—Long-term trends of habituation and abundance. Glob Change Biol
[Internet]. 2017; Available from: http://dx.doi.org/10.1111/gcb.13711

32. U.S. Department of the Interior Bureau of Land Management. Environmental Assessment
for Mesa Mule Deer Winter Habitat Improvement Project. Cheyenne, Wyoming (USA):
Bureau of Land Management, Wyoming State Office; 2011a. Report No.: WY-100-EA10-
303.

33. U.S. Department of the Interior Bureau of Land Management. Finding of no significant
impact and decision record for mesa sagebrush fertilization project. Cheyenne, Wyoming
(USA): Bureau of Land Management, Wyoming State Office; 2011b.

34. Dahlke J, Holloran M. Mesa Fertilization Data Collection. Pinedale, WY (USA): Wyoming
Wildlife Consultants; 2015. Report No.: WGFD Service Contract No. 07SC0403657.

35. Smith MD, Knapp AK, Collins SL. A framework for assessing ecosystem dynamics in
response to chronic resource alterations induced by global change. Ecology.
2009;90(12):3279–89.

36. Collins SL, Ladwig LM, Petrie MD, Jones SK, Mulhouse JM, Thibault JR, et al. Press–
pulse interactions: effects of warming, N deposition, altered winter precipitation, and fire
on desert grassland community structure and dynamics. Glob Change Biol.
2017;23(3):1095–108.

37. Munn LC, Arneson CS. Soils of Wyoming. University of Wyoming, Agricultural
Experiment Station; 1998.

38. NOAA. 1981-2010 Climate Normal [Internet]. National Oceanic and Atmospheric
Administration National Centers for Environmental Information; n.d. [cited 2017 Jul 17].
Available from: https://www.ncdc.noaa.gov/cdo-web/datatools/normals

39. Seastedt TR, Parton WJ, Ojima DS. Mass loss and nitrogen dynamics of decaying litter of
grasslands: the apparent low nitrogen immobilization potential of root detritus. Can J Bot.
1992 Feb 1;70(2):384–91.

40. Frey SD, Elliott ET, Paustian K, Peterson GA. Fungal translocation as a mechanism for soil
nitrogen inputs to surface residue decomposition in a no-tillage agroecosystem. Soil Biol
Biochem. 2000 May 1;32(5):689–98.

41. Hart SC, Firestone MK, Paul EA, Smith JL. Flow and fate of soil nitrogen in an annual
grassland and a young mixed-conifer forest. Soil Biol Biochem. 1993 Apr 1;25(4):431–42.

42. Brooks PD, Williams MW, Schmidt SK. Inorganic nitrogen and microbial biomass
dynamics before and during spring snowmelt. Biogeochemistry. 1998 Oct 1;43(1):1–15.

43. Burke IC. Control of Nitrogen Mineralization a Sagebrush Steppe Landscape. Ecology.
1989;70(4):1115–26.

44. Chen J, Carrillo Y, Pendall E, Dijkstra FA, Evans RD, Morgan JA, et al. Soil Microbes
Compete Strongly with Plants for Soil Inorganic and Amino Acid Nitrogen in a Semiarid
Grassland Exposed to Elevated CO2 and Warming. Ecosystems. 2015 Aug 1;18(5):867–80.

45. Daubenmire RF. Plants and environment: a textbook of plant autecology. New York;
London: John Wiley and Sons, Inc..; Chapman and Hall, Ltd.; 1959. 422 pp.

46. Hutchinson GL, Mosier AR. Improved Soil Cover Method for Field Measurement of
Nitrous Oxide Fluxes1. Soil Sci Soc Am J. 1981;45(2):311–6.

47. Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. _nlme: Linear and Nonlinear
Mixed Effects Models_ [Internet]. 2018. Available from: https://CRAN.R-
project.org/package=nlme

48. Length R. emmeans: Estimated Marginal Means, aka Least-Squares Means. 2019.

49. Oksanen JF, Blanchet G, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package. [Internet]. 2019. Available from: https://CRAN.R-
project.org/package=vegan

50. Lowe PN, Lauenroth WK, Burke IC. Effects of Nitrogen Availability on the Growth of
Native Grasses Exotic Weeds. J Range Manag. 2002 Jan 1;55(1):94–8.

51. Samuel MJ, Hart RH. Nitrogen Fertilization, Botanical Composition and Biomass
Production on Mixed-Grass Rangeland. J Range Manag. 1998;51(4):408–16.

52. Norton U, Mosier AR, Morgan JA, Derner JD, Ingram LJ, Stahl PD. Moisture pulses, trace
gas emissions and soil C and N in cheatgrass and native grass-dominated sagebrush-steppe
in Wyoming, USA. Soil Biol Biochem. 2008 Jun 1;40(6):1421–31.

53.   Seagle SW, McNaughton SJ. Simulated Effects of Precipitation and Nitrogen on Serengeti
578   Grassland Productivity. Biogeochemistry. 1993;22(3):157–78.

579   54.   Burke IC, Lauenroth WK, Parton WJ. Regional and temporal variation in net primary
580   production and nitrogen mineralization in grasslands. Ecology. 1997;78(5):1330–40.