Title

Assembly and seasonality of core phyllosphere microbiota on perennial biofuel crops

Authors

Keara L Grady[1,2,a], Jackson W. Sorensen[1,2,a], Nejc Stopnisek[2,3,a], John Guittar[1,4], and Ashley

Shade[1,2,3,5,6]*

1. Department of Microbiology and Molecular Genetics, Michigan State University, East

Lansing MI 48840 USA

2. The DOE Great Lakes Bioenergy Research Center, Michigan State University, East

Lansing, MI 48840

3. Program in Ecology, Evolutionary Biology and Behavior, Michigan State University, East

Lansing MI 48840

4. Kellogg Biological Station, Michigan State University, Hickory Corners, MI 43060

5. The Plant Resilience Institute, Michigan State University, East Lansing MI 48840

6. Department of Plant, Soil and Microbial Sciences, Michigan State University, East

Lansing MI 48840

[a]Contributed equally

*correspondence, shadeash@msu.edu

Keywords

23 Miscanthus, switchgrass, temporal dynamics, microbiome, agroecosystems, bioenergy,

24 sustainability, plant-microbe interactions, leaf, phytobiome

Abstract

Perennial grasses are promising feedstocks for biofuel production, and there is potential to

leverage their native microbiomes to increase their productivity and resilience to

environmental stress. Here, we characterize the 16S rRNA gene diversity and seasonal

assembly of bacterial and archaeal microbiomes of two perennial cellulosic feedstocks,

switchgrass (Panicum virgatum L.) and miscanthus (Miscanthus x giganteus). We sampled

leaves and soil every three weeks from pre-emergence through senescence for two

consecutive switchgrass growing seasons and one miscanthus season, and identified core leaf

taxa based on abundance and occupancy. Virtually all leaf taxa are also detected in soil;

source-sink modeling shows non-random, ecological filtering by the leaf, suggesting that soil

is important reservoir of phyllosphere diversity. Core leaf taxa include early, mid, and late

season groups that were consistent across years and crops. This consistency in leaf

microbiome dynamics and core members is promising for microbiome manipulation or

management to support biofuel crop production.

The phyllosphere (aerial parts of plants) represents the largest environmental surface

area of microbial habitation on the planet [1–3], and much of that surface area is cultivated

agriculture, including an estimated $1.5 \times 10^7 \, km^2$ of cropland [4]. Phyllosphere microorganisms

may provide numerous benefits to plants, including increased stress tolerance [5–7], promotion of

growth and reproduction [8–10], protection from foliar pathogens [11], and, with soil microbes,

control of flowering phenology [12]. Phyllosphere microorganisms are also thought to play

important roles in Earth's biogeochemical cycles by moderating methanol emissions from

plants [13,14] and contributing to global nitrogen fixation [15]. Despite this importance, knowledge

of phyllosphere microbiomes remains relatively modest, especially for agricultural crops [3,16–18].

To leverage plant microbiomes to support productivity and resilience both above and below

ground [19–21], there is a need to advance foundational knowledge of phyllosphere microbiome

diversity and dynamics.

Biofuel crops like miscanthus and switchgrass are selected to have extended growing

seasons, to produce ample phyllosphere biomass, and to maintain high productivity when

grown on marginal lands that are not optimal for food agriculture [22–25]. In the field, these

grasses provide extensive leaf habitat, with a seasonal maximum leaf area index (LAI) of 6.2 for

switchgrass, and 10 m^2 leaf surface per m^2 land for miscanthus [22], as compared to a maximum

LAI of 3.2 for corn [26]. Upon senescence, the aboveground biomass is harvested for conversion

to biofuels and related bioproducts. Improved understanding of the phyllosphere microbiome is

expected to advance goals to predict or manage changes in biomass quality in response to

abiotic stress like drought [27–31] or biotic stress like foliar pathogens [32–34].

Leveraging the Great Lakes Bioenergy Research Center's Biofuel Cropping System

Experiment (BCSE; a randomized block design established at Michigan State's Kellogg Biological

Station in 2008), we asked two questions of the bacterial and archaeal communities

(henceforth: "microbiomes") inhabiting the leaf surfaces and the associated soils of switchgrass

and miscanthus: 1) Are there seasonal patterns of phyllosphere microbiome assembly? If so,

are these patterns consistent across fields of the same crop, different crops, and years? 2) To

what extent might soil serve as a reservoir of phyllosphere diversity?

Results and Discussion

Sequencing summary and alpha diversity

In total, we sequenced 373 phyllosphere epiphyte (leaf surface) and soil samples across

the two growing seasons in 2016 and 2017. The number of sequences per sample after our 97%

OTU (operational taxonomic unit) clustering pipeline ranged from 20,647 to 359,553. The

percentage of sequences belonging to chloroplasts and mitochondria per sample range

between 0.2-99.8%, but 235 of the samples (63%) had fewer than 10% chloroplasts and

mitochondria reads. After removing sequences that were attributed to chloroplasts and

mitochondria or that had unassigned taxonomic classification, we filtered samples that

contained fewer than 1000 reads and rarefied the remaining samples to 1000 reads for

comparative analyses. While this number of reads is not sufficient to fully capture soil diversity,

it does capture phyllosphere diversity (**Figure 1A**). The majority of the switchgrass and

miscanthus phyllosphere communities were exhaustively sequenced, and approached richness

asymptotes with their associated soils.

As reported for other plants [3,35], switchgrass and miscanthus phyllosphere communities

had relatively low richness, with 1480 total taxa observed across both crops and consistently

fewer than 150 taxa per time point, though there was modest seasonal variability in richness

(**Figure S1**). Cumulative richness increased most between the two earliest time points, and then

tapered gradually upward until senescence (**Figure 1B**), showing that the contributions of new

taxa to community richness were low but consistent over time.

Seasonal microbiome dynamics

To perform the most complete temporal analyses of phyllosphere microbiome

seasonality, we also subsampled the amplicon sequencing dataset to include the maximum

number of time points, resulting in inclusion of 51 discrete leaf and soil samples collected over

18 total time points. The overarching patterns in beta diversity were consistent and statistically

indistinguishable from those derived from the same dataset to include more reads per sample

but fewer time points (Mantel tests all p < 0.001; **Table S1, Table S2**, **Figure S2**). For

consistency, we report the patterns from 1000 reads per sample in the main text, but for

transparency and comparison, we report results from the minimum reads per sample, inclusive

of the complete time series, in supporting materials.

There were directional seasonal changes in the structures of switchgrass and

miscanthus phyllosphere bacterial and archaeal communities (**Figure 2A**, **Table 1**), and these

could be attributed to changes in both soil and leaf properties, as well as to weather (**Table S3**).

Over the 2016 season, miscanthus and switchgrass phyllosphere communities were

synchronous (changed at the same pace and to the same extent, Procrustes m12= 0.349, R =

0.807, p = 0.021), and community structure became less variable as the growing season

progressed (**Figure S3**). Switchgrass 2016 and 2017 leaf communities were highly synchronous,

suggesting a predictable, interannual assembly (Procrustes m12= 0.011, R= 0.994, p = 0.008).

The switchgrass community structures were overall equivalent between 2016 and 2017, with

the exception of the final time points that were collected post-senescence. Together with the

species accumulation analysis (**Figure 1B**), these data suggest that these phyllosphere

communities are not stochastically assembled, nor are they a linear accumulation over seasonal

leaf exposure to whatever taxa are dispersed. The communities follow a directional assembly

over the growing season, and the assembly was highly consistent over two years in the

switchgrass.

Contribution of soil microorganisms to phyllosphere assembly

The major sources of microorganisms to the phyllosphere are soils [2], the vascular tissue

of the plant or its seed [36], and the atmosphere or arthropod vectors [3]. As several studies have

shown that soil microbes contribute to the phyllosphere microbiome [35,37], we wanted to

understand the potential for soil as a reservoir of microorganisms inhabiting switchgrass and

miscanthus phyllospheres. We hypothesized that the intersect of shared soil and phyllosphere

taxa would be highest early in the season, after the young grasses emerged through the soil.

Our deep sequencing effort also provided the opportunity to investigate differences in taxon

relative abundances between soil and leaf communities, and to understand what contributions,

if any, the soil rare biosphere has for leaf assembly.

First, we interrogated the 2016 time series to determine the influence of soil-detected

taxa on leaf microbial communities for both crops. As expected, the structures of leaf

communities were highly distinct from soils (**Figure 2B**, **Figure S2B**, **Table 1**). Though soil

communities also changed seasonally, they experienced less overall change than the

phyllosphere (**Table 1**, **Figure 2B**, **Figure S2C**). While fertilization had no impact on phyllosphere

communities, it did have small but significant influence on soil communities (**Table 1**). These

seasonal and fertilization treatment patterns were reproduced in 2017 for switchgrass (**Table

1**).

To better understand the relationship between soil and phyllosphere communities,

including the influence of rare members of the soil, we searched for phyllosphere taxa within

the full soil dataset (not subsampled). Approximately 90% of phyllosphere OTUs were present

in soil samples, with negligible differences between the two crops and modest variability over

time (**Figure 3A**). When considering the relative abundances of taxa, the mean fraction of

phyllosphere communities found in soil samples was even higher at 98% and exhibited no clear

trend over time (**Figure 3B**). Our results show that the majority of abundant, commonly

detected taxa in the phyllosphere are also present in the soil – albeit often at very low

abundances (see below) – highlighting a potentially important role of the soil in harboring

phyllosphere taxa between plant colonization events.

Given the large proportion of phyllosphere taxa present in the soil, we explored the

potential role that immigration from the soil may have in shaping phyllosphere community

composition and seasonal patterns. On balance, many abundant and persistent phyllosphere

taxa were in low abundances in the soil, though there was a positive association between soil

and leaf abundances for soil taxa > 1 e04 relative abundance (**Figure 3C**). This result indirectly

supports the presence of an ecological filter operating on the phyllosphere that favors some

taxa while disfavoring others. To further investigate how ecological filtering may vary over the

growing season, we compared observed trends in OTU richness with those predicted by a

source-sink null model which simulated demographic stochasticity and random immigration

from the soil between subsequent sampling points (see Methods, **Figure 3D-F**). Observed

phyllosphere communities were dramatically less rich than null model predictions, again

supporting the presence of a strong ecological filter. Such a filter could be due to host plant

selection, environmental filtering, competition exclusion among microbial taxa, or a

combination of all three. According to this model, the strength of filtering did not trend

consistently over the growing season.

Finally, other studies have found that soil microbes contribute more to early-season

phyllosphere communities [38], and we observed similar patterns: the most abundant soil taxa

that were also detected on leaves were more prominent in the early season and then became

rare and transient on leaves in the late season (**Figure S4**).

We conclude from these results that soil is a major reservoir of leaf microorganisms for

these perennial crops and note that deep sequencing was required of the soils to observe many

of the prominent leaf taxa. This is in contrast to the studies of other plants that have suggested

that the phyllosphere is comprised largely of passively dispersed and stochastically assembled

microbes from the atmosphere [39–41]. Notably, our analysis cannot inform directionality or

mechanism of dispersal, which could have occurred between soil and leaf via wind, insects, or

through grass emergence, etc. While 133 leaf OTUs (9%) observed in the phyllosphere could

not be detected in the soil (using the unrarified soil dataset) and may be attributable to non-soil

reservoirs, the vast majority of leaf microbes were detectable in local soils and non-neutral

assembly patterns suggest both determinism and habitat filtering.

Core members of the switchgrass and miscanthus phyllosphere

There was high overlap between switchgrass and miscanthus phyllosphere communities

and a trend towards increased intra-crop similarity during senescence (**Figure S5**). There was

also a modest influence of host crop in 2016 (**Table 1**). Therefore, we defined a core

microbiome for each crop and season (**Dataset 1**). We applied an established macroecological

approach [42] to consider both the occupancy and abundance patterns of these taxa (**Figure 4A-**

C); abundance-occupancy relationships have been previously explored for microbial

communities [43,44] and we utilize it here for ecologically informing a core microbiome.

Occupancy is an ecological term that considers how consistently a taxon is detected across

samples in the dataset, expressed as a proportion of occurrences given the total samples

collected (e.g., 1.0 or 100%). Occupancy provides a dataset-aggregated term describing taxon

persistence, which can be informative for defining core taxa when datasets include a time

series [45].

In contrast to the taxa unique to crops and years, which were rare and not persistent,

most of the highly abundant and prevalent taxa were shared (**Figure 4A-C**). We first quantified

the abundance and occupancy distributions of OTUs, and then identified OTUs that were

consistently detected across replicate plots at one sampling time (occupancy of 1) to include in

the core. We found that these 44, 51 and 42 core taxa (as highlighted in **Figure 4A-C**) contribute

84.4%, 79.5% and 79.4% to the total beta diversity in miscanthus 2016, switchgrass 2016 and

2017, respectively (**Figure 4D-F**). While these core taxa were highly abundant and persistent on

these crops' leaves, their functions are yet unknown and additional members could also

transiently contribute. However, we suggest that the core taxa identified here should be

prioritized for follow-up study of functionality and potential plant benefits.

Notably, if we had defined the core as those taxa uniquely detected on each crop (as in

a Venn diagram analysis, **Figure S6**), we would have instead identified rare and transient taxa

(**Figure 4A-C**). Thus, a core analysis based on presence and absence, instead of on abundance

and occupancy, would have provided a different, and arguably less ecologically relevant, core

for these perennial crops. The approach used here provides a reproducible and conservative

option for longitudinal series, and allows for systematic discovery [46,47] of a replicated core over

time.

The core taxa included several Proteobacteria (*Methylobacterium*, *Sphingomonas*, and

Pseudomonas spp.) and Bacteroidetes (*Hymenobacter* spp.). The taxonomic affiliations of these

core taxa are consistent with the literature for other phyllosphere communities [2,3,18,35,48,49],

providing new support for their seasonal importance in the phyllosphere.

We then performed a hierarchical clustering analysis of standardized (e.g., z-score)

dynamics to explore seasonal trends of core taxa (**Figure 4 G-I, Figure S7**)[50]. This analysis

identified several discrete, seasonally-defined groups of core taxa in switchgrass and

miscanthus, respectively. Seasonal groups were taxonomically consistent across crops and

years (**Figure 4 J-L**). This finding suggests potential for functional redundancy because closely

related taxa are hypothesized to have substantial overlap in their functional repertoire [51]. The

early-season groups (**Figure S7, Figure 4G-I** red traces**)** included several *Gammaproteobacteria*

(**Figure 4J-L**). The late-season groups (**Figure S7, Figure 4G-I** blue traces) was comprised of

Alphaproteobacteria to *Cytophagia* and *Actinobacteria* and was pronounced in switchgrass

(**Figure 4J-L**). The third groups (**Figure S7, Figure 4G-I** gray traces**)** included taxa that peaked in

relative abundance mid-season, including *Alphaproteobacteria* and few taxa belonging to *Beta-*

and *Gamma-proteobacteria*, *Cytophagia, Sphingobacteria* and *Actinobacteria* (**Figure 4J-L**).

Our data suggest a compensatory relationship between members within the

Proteobacteria, where members of *Gammaproteobacteria* and *Alphaproteobacteria* replace

one another over time (**Figure 5A**). Such community transitions have been observed on the

phyllosphere of crops such as sugarcane [37], common beans, soybeans, and canola [38]. A study of

endophytic bacteria of prairie grasses, including switchgrass, showed the same trend in

abundance of *Gamma-* and *Alphaproteobacteria* [52] suggesting that these phyllosphere taxa are

facultative endophytes or are similarly affected by the plant development. The benefits plants

may gain from these taxa are well characterized (see review from [53], however it remains

unknown what drives the exclusion of *Pseudomonas* and gives rise to *Alphaproteobacteria*

(predominantly *Methylobacteria*) in the phyllopshere and endopshere. One possible

explanation would be nutrient availability regulated by the plant development which would

selectively influence the abundances of these taxa. Delmotte and colleagues [54] hypothesized

that *Pseudomonas* specialize on monosaccharides, disaccharides and amino acids, whereas

Sphingomonas and *Methylobacteria* are generalist scavengers that can subsist on a variety of

substrates present at low amounts.

Despite similarity in the membership and dynamics of the core microbiota on both crop

plants, there were differences in the relative abundances of the same taxa across crops,

suggesting some microbiome adaptation to or selectivity by the host plant (**Figure 5B**). There

were other core OTUs that had consistent dynamics across both crops, but these examples

demonstrate, first, that dynamics can be crop specific, and second, that abiotic filtering of an

OTU to a particular crop could be manifested as differences in dynamics in addition to the more

extreme scenarios of taxon exclusion or crop specificity that are the hypotheses posed given

unique detection on a particular crop. Indeed, because the crop-unique taxa were generally

rare and transient (**Figure 4A-C**), the dynamics of core taxa with crop-distinct dynamics may

harbor clues as to the competitive landscape and microbially important changes in the host leaf

environment across crops.

The contributions of abiotic variables, space, time, and crop on phyllosphere assembly

We summarize our analyses of the contributions of crop (host plant), space, time, and

abiotic variables to the assembly of the core phyllosphere community in order of least to most

important. Spatial distance between the plots had no explanatory value (assessed by distance-

decay of beta diversity using a Mantel test with a spatial distance, r: 0.013, p = 0.256). This

finding is different from a recent study of annual crops (common beans, canola and soybean)

that showed an influenced of sampling location on leaf microbiome structure [38]. Here, among

those that variables were significant, crop (switchgrass or miscanthus) had the lowest

explanatory value (**Table 1**). However, our work agrees with previous research that has shown a

relationship between plant species/genotype and the leaf microbiota of perennial plants such

as wild mustard [55], sugar cane [37], and tree species like birch, maple, and pine [56]. Time and

measured abiotic factors had highest explanatory value (**Table 1**, **Table S3**). Relatedly, Copeland

et al. 2015 showed that stage in plant development can influence leaf microbiome structure in

annual crops.

To conclude, we investigated the assembly and seasonal dynamics of the phyllosphere

and soil microbes of two perennial grasses, switchgrass and miscanthus, and found consistent

community trajectories and memberships across growing seasons, suggesting that their key

players are predictable and that most of them can be detected in associated soils.

Understanding the seasonal patterns of these key taxa could be used to improve biomass

production, plant health, or facilitate conversion. As seen in [57], the introduction or control of a

few key microbial species can have significant impact on the host plant phenotype. Next steps

should be to interrogate core members for functionality and direct interactions with the plant,

including investigations of the interactions among core members and with the host crop. This

exploration lays the foundation for an approach to biofuel grass production that incorporates

an understanding of host-microbe and microbe-microbe interactions.

Methods

Site description & sampling scheme

Our study system is located within the Great Lakes Bioenergy Research Center (GLBRC)

Biofuel Cropping System Experiment (BCSE) in Hickory Corners, Michigan (42^{o}23'41.6" N,

85^{o}22'23.1" W). We collected samples from two biofuel crops within the BCSE, switchgrass

(*Panicum virgatum* L. cultivar "Cave-in-rock") and miscanthus (*Miscanthus x giganteus)*. Both

crops had been continuously grown since 2008, in replicate 30 x 40 m plots arrayed in a

randomized complete block design. Within each plot, nitrogen-free (no fertilizer) subplots were

maintained in the western-most 3 m of each plot. We sampled replicate plots 1-4 in both the

main and the nitrogen free subplots. We collected leaf and bulk soil samples every three weeks

across the 2016 growing season, including bare soil in April through senescence in October and

November. In total, we collected 152 soil samples (72 switchgrass and 80 miscanthus) and 136

leaf samples (64 switchgrass and 72 miscanthus). At each sampling time, leaves were collected

and pooled at three flags along a standardized path within each plot. Leaves were removed

from the plant stem using ethanol sterilized gloves, then stored in sterile whirl-pak bags until

processing. Bulk soil cores (2 x 10 cm) were collected at the same three locations within a plot,

sieved through 4 mm mesh, then pooled and stored in whirl-pak bags. All samples were kept

on wet ice for transport, then stored at -80 °C.

Soil physico-chemical characteristics (pH, lime, P, K, Ca, Mg, organic matter, NO_3-N, NH_4-

N, and percent moisture) were measured by the Michigan State University (MSU) Soil and Plant

Nutrient Lab (East Lansing, MI, USA, http://www.spnl.msu.edu/) according to their standard

protocols. From each plot, 10 switchgrass leaves or 5 miscanthus leaves were processed for

leaf dry matter content according to [58]. Dried leaves were ground to a fine powder using a

Sampletek 200 vial rotator and iron roll bars (Mavco Industries, Lincoln, NE, USA), then carbon

and nitrogen were measured on an elemental analyzer (Costech ECS 4010; Costech Analytical

Technologies Inc, Valencia, CA, USA). Weather data was collected from the MSU Weather

Station Network, for the Kellogg Biological Station location (https://mawn.geo.msu.edu) for

each sampling day, and plant height and soil temperature were measured on a per-plot basis.

Nucleic acid extraction & sequencing

Throughout, we use "microbiome" to refer to the bacterial and archaeal members as

able to be assessed with 16S rRNA gene sequence analysis. Soil microbial DNA was extracted

using a Powersoil microbial DNA kit (MOBio Inc. Carlsbad, California, USA) according to

manufacturer's instructions. Phyllosphere epiphytic DNA was extracted from intact leaves using

a benzyl chloride liquid:liquid extraction, followed by an isopropanol precipitation [59], using

approximately 5 g of leaves (5-10 switchgrass leaves, or a minimum of 2 miscanthus leaves).

Metagenomic DNA from both soil and phyllosphere was quantified using a qubit 2.0

fluorometer (Invitrogen, Carlsbad, CA, USA), and DNA concentrations were normalized between

all samples prior to sequencing. Paired-end amplicon sequencing was completed by the

Department of Energy's Joint Genome Institute (JGI) using an Illumina MiSeq sequencer, and

using the 16S-V4 (515F-804R) primer set [60], according to the JGI's standard operating protocols,

and incorporating chloroplast- and mitochondria-blocking peptide nucleic acids to prevent co-

amplification of plastid 16S rRNA gene as described in [61].

316   .

318   *Sequence quality control and defining operational taxonomic units*

319   BBDuk (v 37.96) was used to remove contaminants and trim adaptor sequences from

reads. Reads containing 1 or more 'N' bases, having an average quality score of less than ten or

less than 51 bases were removed. Common contaminants were removed with BBMap (v 37.96).

Primers were trimmed using cutadapt (v1.17). Reads were merged, dereplicated, clustered into

97% identity with usearch (v10.0.240), and classified against version 123 of the Silva Database

[62] using sintax [63]. All reads classified as mitochondria, chloroplast or unclassified were removed

before the analysis. Additionally, reads from 4371 OTUs assigned only to the domain level were

extracted and reclassified using SINA online aligner (https://www.arb-silva.de/aligner/) [62]. 696

unclassified reads subsequently were confirmed to be Bacteria could then be classified to more

resolved taxonomic levels. Remaining reads were BLASTed against the entire NCBI nucleotide

database and specifically against the switchgrass genome to check for non-specific binding, but

no hits could be found.

Alpha and beta diversity

For alpha and beta diversity analyses, we performed analyses to datasets subsampled to

the minimum observed quality-filtered reads per sample (141), as well as to 1000 reads per

sample. We did this to enable comparison of the most complete time series to the most

complete comparative view of diversity. We report richness as total number of OTUs clustered

at 97% sequence identity. We used the protest function in the vegan package in R [64] to test for

synchrony in patterns across crops and years. To calculate beta dispersion, we used the

betadisper function in the vegan package in R [64], which is a multivariate analogue of Levene's

test for homogeneity of variances. PERMANOVA was used to test hypothesis of beta diversity

using adonis function in the vegan package in R [64].

Source-sink models and contributions of soil taxa to leaf communities

Given that virtually all phyllosphere taxa were present in the soil, we evaluated the degree to

which observed seasonal patterns in phyllosphere community composition could be explained

with a null model. The null model assumed functional equivalence among taxa and random

source-sink dynamics from the soil. For each crop, we simulated community dynamics between

each pair of sequential samples. This involved: (1) calculating the total number of 0.1 %

incremental increases/decreases in OTU relative abundances observed between the two

sampling points; (2) randomly and iteratively selecting OTUs, weighting their probability of

selection by their relative abundances, to increase or decrease by 0.1 % increments of relative

abundance until reaching the total number of observed increases/decreases for the sample

pair; (3) counting the number of immigrant OTUs which appeared only in the second sample of

the pair; (4) randomly selecting OTUs in the simulated community to decrease by 0.1 %

increments until the total decrease equaled the observed number of arriving OTUs multiplied

by their median initial abundance (0.1 %); and then (5) randomly selecting, again weighting

their probability of selection by their relative abundance, the predicted number of immigrant

OTUs from the soil community, such that the final simulated community abundance was equal

to 1000 sequences. The source soil community was generated by pooling all soil samples from

both years. We used this process to simulate demographic stochasticity. In the model, it was

assumed that 0.1 % incremental changes in relative abundance realistically reflects

phyllosphere population dynamics, and that an immigration event results in the initial relative

abundance of an OTU at 0.1%. Importantly, even if these assumptions are imprecise, the

simulation provides a consistent baseline of community composition against which to compare

observations over the growing season.

Core taxa selection

To infer the core phyllosphere taxa and prioritize them for further inquiry, we calculated the

abundance-occupancy distributions of taxa, as established in macroecology (e.g. Shade et al.

2018). For each OTU, we calculated occupancy and mean relative abundance at each time point

by crop and year. Only OTUs with occupancy of 100% (found in all samples at a particular time

point) were prioritized as core members. Using this conservative threshold for occupancy, we

included all OTUs that had strong temporal signatures; these taxa also were in high abundance

and were persistent as indicated by their abundance-occupancy distributions. These core taxa

also represent potentially important players in plant development, as they were detected at

least at one time point in all sampled fields.

We quantified the explanatory value of the core members to community temporal

dynamics using a previously published method of partitioning community dissimilarity [65]:

379
$$C = \frac{BC_{core}}{BC_{all}}$$

Where *C* is the relative contribution of community Bray Curtis (*BC*) dissimilarity

attributed to the core OTUs.

Hierarchical clustering

To understand the seasonal abundance patterns of the core taxa we performed

hierarchical clustering. We used a z-scored relative abundance matrix subset to contain only

core taxa to generate a w complete linkage distance matrix using the R function hclust() [50].

Groups of core taxa with similar dynamics were defined from the dendrogram using the

function cutree() in R with number of desired groups (k=) to be close to the number of sampling

time points; 8 for miscanthus 2016, 5 for switchgrass 2016 and 7 for switchgrass 2017.

Availability of data, workflows, and material

The datasets generated and/or analyzed during the current study are available in the

Joint Genomes Institute, Integrated Microbial Genomes repository with JGI Projects designated

by year and sample type (Project ID 1139694, 1139696 for 2016 season phyllosphere and soil,

and 1191516 and 1191517 for 2017 season phyllosphere and soil sequences, respectively). Our

sequence analyses and statistical workflows are available at

https://github.com/ShadeLab/PAPER_GradySorensenStopnisek_InPrep

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AS designed the study. AS, KLG, JS, and NS conducted field work. KLG executed lab work. JS,

NS, JG, and AS analyzed the data. All authors discussed and revised the manuscript.

Acknowledgements

We thank SH Lee, M Sleda, S Wu and M Nunez for technical assistance in the field and

laboratory. This material is based upon work supported by the Great Lakes Bioenergy Research

Center, U.S. Department of Energy, Office of Science, Office of Biological and Environmental

Research under Award Numbers DE-SC0018409 and DE-FC02-07ER64494. The work conducted

by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility,

is supported under Contract No. DE-AC02-05CH11231. This work was supported in part by

Michigan State University through computational resources provided by the Institute for Cyber-

Enabled Research. NS acknowledges support from the Michigan State Plant Resilience Institute.

Figure 1. Sequencing effort and alpha diversity for switchgrass and miscanthus phyllosphere

and soils. Operational taxonomic units (OTUs) were defined at 97% sequence identity of 16S

rRNA gene amplicons. (A) Rarefaction curves of quality-controlled reads. The vertical line is the

maximum number of sequences observed in a sample, and the horizontal line is the richness of

that sample. (B) Phyllosphere richness accumulation over time, using a dataset subsampled to

1000 sequences per sample.

Figure 2. Seasonal patterns in the structures of bacterial and archaeal communities

inhabiting the phyllosphere and associated soils of the biofuel feedstocks switchgrass and

miscanthus. (A) Principal coordinates analysis (PCoA) of switchgrass and miscanthus

phyllosphere communities (Bray-Curtis dissimilarity), error bars show 1 deviation around the

centroid (n = 1 to 8 replicate plots/time point). (B) PCoA of the phyllosphere communities

relative to the soil. For both A and B, subsampling depth was 1000 reads per sample and

environmental vectors were fitted when $r^2 > 0.4$ and $p < 0.05$.

Figure 3. The majority of phyllosphere taxa were also present in the soil. (**A**) Circles represent

the mean number of OTUs found in up to eight replicate phyllosphere samples, subsampled to

1000 sequences, for each crop at each time point. An OTU was considered present in the soil if

it occurred at any abundance in any of the 202 unrarefied soil samples over two years. **(B)** The

fractions of the phyllosphere communities present in the soil were even greater when

considering the relative abundances of taxa; each circle represents the mean total relative

abundance of leaf taxa present in the soil in up to four replicate phyllosphere samples. **(C)** The

relative abundances of taxa in pooled phyllosphere samples and pooled soil samples were

positively correlated among taxa that were present at greater than 0.01% total abundance in

the soil. Each black circle represents an OTU present in both phyllosphere and soil

communities; a LOESS smoothing function is shown as a red line. Core members are shown as

large green circles. **(D-F)** Source-sink models of phyllosphere community assembly from soils.

Violin plots show the numbers of observed taxa in the phyllosphere were consistently lower

than the richness values predicted by model simulations. The model assumed random increases

and decreases in taxon abundances between time points and random immigration from the soil

community (see Methods). Each circle represents a single phyllosphere sample.

Figure 4. **Selection and dynamics of core phyllosphere members.** Abundance-occupancy of

leaf taxa for **(A)** miscanthus 2016, **(B)** switchgrass 2016, and **(C)** switchgrass 2017, and their

inclusion in their respective cores. Each point is an OTU. Abundance-occupancy distributions

were calculated at each time point, and taxa that had 100% occupancy at any time point (e.g.,

were detected in all replicate plots at one sampling date) were included in the core (green).

Non-core taxa that were detected in both crops (white/open circles), and crop-specific taxa

(grey) are also indicated. **(D-F)** Contributions of the core taxa to changes in beta diversity over

time. **(G-I)** Patterns of core taxa that share similar temporal changes, as determined by

hierarchical clustering of standardized dynamics. Colors correspond to the dendrograms in

Figure S7. **(J-L)** Patterns of core taxa summed by relative abundances within bacterial class.

"c:" is class and "p:" is phylum.

Figure 5. Compensatory patterns of Protobacteria classes over crops and season in the

phyllosphere of switchgrass and miscanthus. Proteobacteria OTUs contributed 35.2% of the

total taxa detected in the phyllosphere and contributed 116,760 total reads (34.1% of the leaf

reads). **(A)** Changes in the relative contributions of all 521 leaf-detected Proteobacteria OTUs by

class, over time. **(B-D)** Vignettes showing different dynamics of Proteobacteria OTUs that were

detected within the phyllosphere core microbiome, over time and across crops. Class is "c:"

and genus is "g:".

472 Table

473 **Table 1**. Permuted multivariate analysis of variance (PERMANOVA) tables for all hypothesis

474 tests for differences in community structure (beta diversity).

Supporting Figures

Figure S1. Seasonal patterns in the number of observed phyllosphere taxa (richness).

Operational taxonomic units (OTUs) were defined at 97% amplicon sequence identity. Richness

is provided at subsampling depths of 1000 reads (top) and 141 reads (bottom).

Figure S2. Seasonal patterns in the structures of bacterial and archaeal communities

inhabiting the phyllosphere and associated soils of the biofuel feedstocks switchgrass and

miscanthus. (A) Principal coordinates analysis (PCoA) of switchgrass and miscanthus

phyllosphere communities (Bray-Curtis dissimilarity), error bars show 1 deviation around the

centroid (n = 3 to 8 replicate plots/time point). Subsampling depth was 141 reads per sample

and environmental vectors are fitted when $r^2 > 0.4$ and $p < 0.05$. **(B)** PCoA of the phyllosphere

communities relative to the soil, subsampled to 141 sequences per sample. **(C)** PCoA of the soil

communities associated with miscanthus and switchgrass, subsampled to 19,967 sequences per

sample.

Figure S3. Phyllosphere communities become less variable over time. Distance to median was

calculated by analysis of beta-dispersion. Variability in phyllosphere microbiome structure over

time miscanthus 2016, switchgrass 2016, and switchgrass 2017 field seasons. Betadispersion

was calculated from data series subsampled to 1000 reads (top) and 141 reads (bottom).

Figure S4. **Decreases in the contributions of soil-dominating taxa to the phyllosphere**

microbiome over time. Heatmaps represent the 50 top-ranked OTUs from the soil that were

also detected in the phyllosphere. The cell colors are the z-scored relative abundances of the

OTUs in the phyllosphere. The ranking is from top to bottom (e.g., the most abundant soil-

dominant taxon that was also detected in the phyllosphere is represented by the top row in the

heatmap). The left bar shows the classification the taxon as either a core member (green) or

not (pink). **(A)** Miscanthus 2016; **(B)** switchgrass 2016; **(C)** switchgrass 2017.

Figure S5. Intra-crop similarity in leaf communities over time in 2016. We show changes in

Bray-Curtis dissimilarities between switchgrass and miscanthus phyllosphere microbiomes per

time point, inclusive of the maximum number of replicated blocks (up to 8) per time point. **(A)**

"Full" time series, subsampled to 141 reads per sample. **(B)** Time series subsampled to 1000

reads per sample.

Figure S6. Venn diagram of taxa shared across the switchgrass and miscanthus phyllosphere,

in 2016 and 2017. Data were rarefied to 1000 reads per sample.

Figure S7. Hierarchical clustering of standardized (z-scored) dynamics of core phyllosphere

taxa on the phyllosphere of miscanthus 2016 (A), switchgrass 2016 (B) and switchgrass 2017

(C). Seasonally discrete clusters coincide with plant phenology, including groups that achieved

highest relative abundance during early (red), mid (gray) and late (blue) plant growth. Clusters

are labeled in order of temporal occurrence in **Figure 4G-I**. Circles on the dendrogram tips are

color coded by OTU taxonomic classification and labeled with the OTU ID.

Supporting Tables

Table S1. **Sequencing summary of phyllosphere microbial communities characterized in this**

study, categorized by crop and year.

Table S2. **Comparison of overarching patterns of beta diversity across the same dataset**

rarefied to different sequencing depths. We compared all pairs of 141, 500, 1000, 5000, or

10000 reads per sample. All Mantel tests were significant at p < 0.001 on 1000 permutations.

Table S3. Fitted environmental variables that explain changes in microbiome community

structure. Values in which EnvFit R^2 > 0.40 were plotted as vectors in **Figure 2**.

Datasets

Dataset 1. Operational taxonomic unit identifiers and representative 16S rRNA gene amplicon

sequences of the core switchgrass and miscanthus phyllosphere microbiota.

References

1.   Peñuelas, J. & Terradas, J. The foliar microbiome. *Trends Plant Sci.* **19**, 278–280 (2014).

2.   Lindow, S. E. & Brandl, M. T. Microbiology of the phyllosphere. *Appl. Environ. Microbiol.*

542   **69**, 1875–1883 (2003).

543   3.   Vorholt, J. A. Microbial life in the phyllosphere. *Nat. Rev. Microbiol.* **10**, 828–840 (2012).

544   4.   Foley, J. A. *et al.* Solutions for a cultivated planet. *Nature* **478**, 337–342 (2011).

545   5.   Hamilton, C. E., Gundel, P. E., Helander, M. & Saikkonen, K. Endophytic mediation of

546   reactive oxygen species and antioxidant activity in plants: a review. *Fungal Divers.* **54**, 1–

547   10 (2012).

548   6.   Lindow, S. E. & Leveau, J. H. J. Phyllosphere microbiology. *Curr. Opin. Biotechnol.* **13**,

549   238–243 (2002).

550   7.   Redman, R. S., Sheehan, K. B., Stout, R. G., Rodriguez, R. J. & Henson, J. M.

551   Thermotolerance generated by plant/fungal symbiosis. *Science* **298**, 1581 (2002).

552   8.   Canto, A. & Herrera, C. M. Micro-organisms behind the pollination scenes: Microbial

553   imprint on floral nectar sugar variation in a tropical plant community. *Ann. Bot.* **110**,

554   1173–1183 (2012).

555   9.   Doty, S. L. *et al.* Diazotrophic endophytes of native black cottonwood and willow.

556   *Symbiosis* **47**, 23–33 (2009).

557   10.   Taghavi, S. *et al.* Genome survey and characterization of endophytic bacteria exhibiting a

558   beneficial effect on growth and development of poplar trees. *Appl. Environ. Microbiol.*

559   **75**, 748–757 (2009).

11.   Lee, D. W., Hong, J. S., Kim, S. H., Kim, J. W. & Kim, B. S. First Report of Pseudomonas

lurida Causing Bacterial Leaf Spot on Miscanthus sinensis. *J. Phytopathol.* **162**, 195–200

(2014).

12. Wagner, M. R. *et al.* Natural soil microbes alter flowerig phenology and the intensity of

selection on flowering time in a wild *Arabidopsis* relative`. *Ecol. Lett.* **17**, 717–726 (2014).

13. Iguchi, M., Yamanaka, S. & Budhiono, A. Bacterial cellulose - a masterpiece of nature's

arts. *J. Mater. Sci.* **35**, 261–270 (2000).

14. Galbally, I. E. & Kirstine, W. The production of methanol by flowering plants and the

global cycle of methanol. *J. Atmos. Chem.* **43**, 195–229 (2002).

15. Fürnkranz, M. *et al.* Nitrogen fixation by phyllosphere bacteria associated with higher

plants and their colonizing epiphytes of a tropical lowland rainforest of Costa Rica. *ISME*

J. **2**, 561–570 (2008).

16. Weyens, N., van der Lelie, D., Taghavi, S., Newman, L. & Vangronsveld, J. Exploiting plant-

microbe partnerships to improve biomass production and remediation. *Trends in*

Biotechnology **27**, 591–598 (2009).

17. Hacquard, S. & Schadt, C. W. Towards a holistic understanding of the beneficial

interactions across the *Populus* microbiome. *New Phytologist* **205**, 1424–1430 (2015).

18. Kinkel, L. L. Microbial Population Dynamics on Leaves. *Annu. Rev. Phytopathol.* **35**, 327–

347 (1997).

19. Lebeis, S. L. The potential for give and take in plant-microbiome relationships. *Front.*

Plant Sci. **5**, 1–6 (2014).

20. Vandenkoornhuyse, P., Quaiser, A., Duhamel, M., Le Van, A. & Dufresne, A. The

importance of the microbiome of the plant holobiont. *New Phytol.* **206**, 1196–1206

(2015).

21. Turner, T. R., James, E. K. & Poole, P. The plant microbiome. *Genome Biol.* **14**, (2013).

22. Heaton, E. A., Dohleman, F. G. & Long, S. P. Meeting US biofuel goals with less land: The

potential of Miscanthus. *Glob. Chang. Biol.* **14**, 2000–2014 (2008).

23. Tornqvist, C. E. *et al.* Transcriptional Analysis of Flowering Time in Switchgrass. *Bioenergy*

Res. **10**, 700–713 (2017).

24. Robertson, G. P. *et al.* Cellulosic biofuel contributions to a sustainable energy future:

Choices and outcomes. *Science.* **356**, 1-9 (2017).

25. Stoof, C. R. *et al.* Untapped Potential: Opportunities and Challenges for Sustainable

Bioenergy Production from Marginal Lands in the Northeast USA. *BioEnergy Res.* **8**, 482–

501 (2015).

26. Johnson, D. R. & Tanner, J. W. Comparisons of Corn (*Zea mays* L .) Inbreds and Hybrids

Grown at Equal Leaf Area Index , Light Penetration , and Population is generally accepted

that corn hybrids (*Zea raays* I T L .) are superior to inbred lines in terms of overall size

and yield . At a. 482–485 (1972).

27. Wang, B., Seiler, J. R. & Mei, C. A microbial endophyte enhanced growth of switchgrass

under two drought cycles improving leaf level physiology and leaf development. *Environ.*

Exp. Bot. **122**, 100–108 (2016).

28. Ong, R. G. *et al.* Inhibition of microbial biofuel production in drought-stressed

switchgrass hydrolysate. *Biotechnol. Biofuels* **9**, 1–14 (2016).

29. Emerson, R. *et al.* Drought effects on composition and yield for corn stover, mixed

grasses, and Miscanthus as bioenergy feedstocks. *Biofuels* **5**, 275–291 (2014).

30.   Fitzpatrick, C. R. *et al.* Assembly and ecological function of the root microbiome across

606   angiosperm plant species. *Proc. Natl. Acad. Sci.* 201717617 (2018).

607   31.   Santos-Medellín, C., Edwards, J., Liechty, Z., Nguyen, B. & Sundaresan, V. Drought Stress

608   Results in a Compartment-Specific Restructuring of. *MBio* **8**, 1–15 (2017).

609   32.   Cox, C. M., Bockus, W. W., Holt, R. D., Fang, L. & Garrett, K. A. Spatial connectedness of

610   plant species: Potential links for apparent competition via plant diseases. *Plant Pathol.*

611   **62**, 1195–1204 (2013).

612   33.   Alexander, H. M., Bruns, E., Schebor, H. & Malmstrom, C. M. Crop-associated virus

613   infection in a native perennial grass: reduction in plant fitness and dynamic patterns of

614   virus detection. *J. Ecol.* **105**, 1021–1031 (2017).

615   34.   Sattler, S. E. & Funnell-Harris, D. L. Modifying lignin to improve bioenergy feedstocks:

616   strengthening the barrier against pathogens? *Front. Plant Sci.* **4**, 1–8 (2013).

617   35.   Bodenhausen, N., Horton, M. W. & Bergelson, J. Bacterial Communities Associated with

618   the Leaves and the Roots of *Arabidopsis thaliana*. *PLoS One* **8**, (2013).

619   36.   Barret, M. *et al.* Emergence shapes the structure of the seed microbiota. *Appl. Environ.*

620   *Microbiol.* **81**, 1257–1266 (2015).

621   37.   Hamonts, K. *et al.* Field study reveals core plant microbiota and relative importance of

622   their drivers. *Environ. Microbiol.* **20**, 124–140 (2018).

623   38.   Copeland, J. K., Yuan, L., Layeghifard, M., Wang, P. W. & Guttman, D. S. Seasonal

624   Community Succession of the Phyllosphere Microbiome. *Mol. Plant-Microbe Interact.* **28**,

625   274–285 (2015).

39.   Ottesen, A. R. *et al.* Using a Control to Better Understand Phyllosphere Microbiota. *PLoS*

One **11**, 1-16 (2016).

40.   Maignien, L., DeForce, E. A., Chafee, M. E., Murat Eren, A. & Simmons, S. L. Ecological

succession and stochastic variation in the assembly of *Arabidopsis thaliana* phyllosphere

communities. *MBio* **5**, (2014).

41.   Vokou, D. *et al.* Exploring biodiversity in the bacterial community of the mediterranean

phyllosphere and its relationship with airborne bacteria. *Microb. Ecol.* **64**, 714–724

(2012).

42.   Gaston, K. J. *et al.* Abundance - occupancy relationships. *J. Appl. Ecol.* **37**, 39–59 (2000).

43.   Shade, A. *et al.* Macroecology to unite all life, large and small. *Trends Ecol. Evol.* (2018).

44.   Burns, A. R. *et al.* Contribution of neutral processes to the assembly of gut microbial

communities in the zebrafish over host development. *ISME J.* **10**, 655–664 (2016).

45.   Shade, A. & Handelsman, J. Beyond the Venn diagram: The hunt for a core microbiome.

Environ. Microbiol. **14**, 4–12 (2012).

46.   Astudillo-García, C. *et al.* Evaluating the core microbiota in complex communities: A

systematic investigation. *Environ. Microbiol.* **19**, 1450–1462 (2017).

47.   Shade, A. *et al.* Culturing captures members of the soil rare biosphere. *Environmental

Microbiology* **14**, 2247–2252 (2012).

48.   Knief, C., Ramette, A., Frances, L., Alonso-Blanco, C. & Vorholt, J. A. Site and plant species

are important determinants of the *Methylobacterium* community composition in the

plant phyllosphere. *ISME J.* **4**, 719–728 (2010).

49.   Rastogi, G., Coaker, G. L. & Leveau, J. H. J. J. New insights into the structure and function

of phyllosphere microbiota through high-throughput molecular approaches. *FEMS*

Microbiol. Lett. **348**, 1–10 (2013).

50.   R Core Team. R: A language and environment for statistical computing. R Foundation for

651   Statistical Computing, Vienna, Austria. https://www.R-project.org/.(2018).

652   51.   Martiny, J. B. H., Jones, S. E., Lennon, J. T. & Martiny, A. C. Microbiomes in light of traits:

653   A phylogenetic perspective. *Science (80).* **350**, (2015).

654   52.   Ding, T. & Melcher, U. Influences of plant species, season and location on leaf endophytic

655   bacterial communities of non-cultivated plants. *PLoS One* **11**, 1–13 (2016).

656   53.   Bringel, F. & Couée, I. Pivotal roles of phyllosphere microorganisms at the interface

657   between plant functioning and atmospheric trace gas dynamics. *Front. Microbiol.* **6**,

658   (2015).

659   54.   Delmotte, N. *et al.* Community proteogenomics reveals insights into the physiology of

660   phyllosphere bacteria. *Proc. Natl. Acad. Sci.* **106**, 16428–16433 (2009).

661   55.   Wagner, M. R. *et al.* Host genotype and age shape the leaf and root microbiomes of a

662   wild perennial plant. *Nat. Commun.* **7**, 12151 (2016).

663   56.   Laforest-Lapointe, I., Messier, C. & Kembel, S. W. Tree phyllosphere bacterial

664   communities: exploring the magnitude of intra- and inter-individual variation among host

665   species. *PeerJ* **4**, e2367 (2016).

666   57.   Agler, M. T. *et al.* Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome

667   Variation. *PLoS Biol.* **14**, 1–31 (2016).

668   58.   Cornelissen, J. H. C. *et al.* A handbook of protocols for standardised and easy

669   measurement of plant functional traits worldwide. *Aust. J. Bot.* **51**, 335–380 (2003).

59.   Suda, W., Oto, M., Amachi, S., Shinoyama, H. & Shishido, M. A Direct Method to Isolate

DNA from Phyllosphere Microbial Communities without Disrupting Leaf Tissues.

Microbes Environ. **23**, 248–252 (2008).

60.   Caporaso, J. G. *et al.* Moving pictures of the human microbiome. *Genome Biol.* **12**,

(2011).

61.   Lundberg, D. S., Yourstone, S., Mieczkowski, P., Jones, C. D. & Dangl, J. L. Practical

innovations for high-throughput amplicon sequencing. *Nat. Methods* **10**, 999–1002

(2013).

62.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: Improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, (2013).

63.   Edgar, R. C. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS

sequences. *Preprint at* doi:10.1101/074161 (2016).

64.   Oksanen, J. *et al.* vegan: Community Ecology Package. R Package version 2.5-2. (2018).

65.   Shade, A. *et al.* Conditionally rare taxa disproportionately contribute to temporal

changes in microbial diversity. *MBio* **5**, 1–9 (2014).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Table 1. Permuted multivariate analysis of variance (PERMANOVA) tables for all hypothesis

tests for differences in community structure (beta diversity).

Dataset	Variable Tested	Degrees of Freedom	PseudoF	R squared	p-value
2016 All	Microbiome habitat (soil v. leaf)	1	166.09	0.415	0.001
2017 All	Microbiome habitat (soil v. leaf)	1	74.66	0.418	0.001
2016 Phyllosphere	Time	1	21.02	0.183	0.001
2016 Soil	Time	1	7.31	0.050	0.001
2017 Phyllosphere	Time	1	29.09	0.409	0.001
2017 Soil	Time	1	3.05	0.048	0.001
2016 Phyllosphere	Crop	1	14.50	0.134	0.001
2016 Soil	Crop	1	6.76	0.047	0.001
2016 Phyllosphere	Fertilization status	1	0.40	0.004	0.95
2016 Soil	Fertilization status	1	4.77	0.033	0.001
2017 Phyllosphere	Fertilization status	1	0.87	0.020	0.438
2017 Soil	Fertilization status	1	3.41	0.054	0.001

A

B

C

Table S1. **Sequencing summary of phyllosphere microbial communities characterized in this study**, categorized by crop and year. QC is quality controlled.

	Miscanthus 2016	Switchgrass 2016	Switchgrass 2017
Raw Read Pairs	7336923	7120682	9271169
QC Reads	7142900	6883631	7123783
% Chloroplast/ Mitochondria of QC Reads	77.27%	67.16%	12.32%
Samples >= 141 Reads	65	62	44
Samples >= 1,000 Reads	53	43	44
Samples >= 10,000 Reads	36	27	44
OTUs (rarefied to 141 Reads)	540	685	232
OTUs (rarefied to 1,000 Reads)	1010	859	769
OTUs (rarefied to 10,000 Reads)	1121	896	2320

Table S2. **Comparison of overarching patterns of beta diversity across the same dataset rarefied to different sequencing depths**. We compared all pairs of 141, 500, 1000, 5000, or 10000 reads per sample. All tests were significant at p < 0.001 on 1000 permutations.

	141 Reads	500 Reads	1000 Reads	5000 Reads
500 Reads	0.932			
1000 Reads	0.872	0.982		
5000 Reads	0.752	0.931	0.976	
10000 Reads	0.725	0.916	0.967	0.999

Table S3. Fitted environmental variables that explain changes in microbiome community structure. All are p < 0.05 unless designated as not significant (NS). Values in bold (EnvFit R^2 > 0.40) were plotted as vectors in **Figure 2**.

Variable Tested	Axis 1	Axis 2	R squared	P -value
precipitation	-0.127	0.146	0.038	0.703
Air_temp_mean	-0.298	-0.078	0.095	0.384
air_temp_max	-0.212	-0.056	0.048	0.633
Air_Temp_Min	-0.242	0.139	0.078	0.439
Air_Pressure	-0.022	0.217	0.048	0.653
RH	0.474	0.444	0.422	0.008
AH	-0.081	0.102	0.017	0.834
Wind_Speed_Mean	-0.703	-0.213	0.539	0.001
Solar_Radiation	-0.258	-0.368	0.202	0.118
PAR	-0.300	-0.288	0.173	0.162
soil_temp_5_cm_bare_avg	-0.408	-0.061	0.170	0.163
Week	0.730	0.356	0.659	0.001
LDMC_mg_per_g	0.418	0.177	0.206	0.001
nitrogen_percent	-0.392	-0.416	0.327	0.001
carbon_percent	0.345	0.038	0.121	0.001
carbon_per_nitrogen	0.274	0.134	0.093	0.003
height_mean_cm	0.730	-0.230	0.586	0.001
pH	-0.027	0.219	0.049	0.047
P_ppm	-0.120	-0.097	0.024	0.187
K_ppm	-0.445	0.187	0.233	0.001
Ca_ppm	0.047	0.059	0.006	0.659
Mg_ppm	-0.366	-0.125	0.149	0.001
organic_matter	-0.348	-0.036	0.123	0.001
NO3N_ppm	0.003	-0.080	0.006	0.646
NH4_ppm	-0.461	0.090	0.220	0.001
soil_moisture_percent	0.609	0.015	0.371	0.001
soil_temp_10cm	-0.364	-0.028	0.133	0.001