Interaction of modified oligonucleotides with nuclear proteins, formation of novel

nuclear structures and sequence-independent effects on RNA processing

Loren L Flynn #, Ruohan Li # May T Aung-Htut, Ianthe L Pitout, Jack A L Cooper, Alysia

Hubbard, Lisa Griffiths, Charlie Bond, Steve D Wilton, Archa H Fox* and Sue Fletcher*

 # authors contributed equally

corresponding authors

Abstract (max 150 words)

Oligonucleotides and nucleic acid analogues that alter gene expression are showing

therapeutic promise for selected human diseases. The modification of synthetic nucleic acids

to protect against nuclease degradation and to influence drug function is common practice,

however, such modifications may also confer unexpected physicochemical and biological

properties. Here we report backbone-specific effects of modified oligonucleotides on

subnuclear organelles, altered distribution of nuclear proteins, the appearance of novel

structured nuclear inclusions, and modification of RNA processing in cultured cells

transfected with antisense oligonucleotides on a phosphorothioate backbone. Phosphodiester

and phosphorodiamidate morpholino oligomers elicited no such consequences. Disruption

of subnuclear structures and proteins elicit severe phenotypic disturbances, revealed by

transcriptomic analysis of fibroblasts exhibiting such disruption. These data suggest that the

toxic effects and adverse events reported after clinical evaluation of phosphorothioate

nucleic acid drugs may be mediated, at least in part, by non-specific interaction of nuclear

components with the phosphorothioate backbone.

24
25
Key Words

Antisense oligonucleotides, RNA analogue, RNA processing, paraspeckle proteins, nuclear

organelles

Abbreviations

antisense oligonucleotide (AO), spinal muscular atrophy (SMA), Duchenne muscular

dystrophy (DMD) phosphorodiamidate morpholino oligomer (PMO) amyotrophic lateral

sclerosis (ALS)

Introduction

Antisense oligonucleotide drugs are a class of therapeutics designed to alter gene expression and function, and are reported to have delivered therapeutic benefit to spinal muscular atrophy (SMA) type 1 patients (Paton 2017) and a subset of Duchenne muscular dystrophy (DMD) cases (Mendell *et al.* 2013, Mendell *et al.* 2016). *Nusinersen* (*Spinraza*) is a 2′O-methoxyethyl antisense oligonucleotide (AO) on a phosphorothioate backbone, targeting a splice silencer (*ISS-N1*) in *SMN2* intron 7 that promotes exon 7 selection during pre-mRNA splicing (Singh *et al.* 2006) and received Food and Drug Administration (USA) (FDA) approval in December 2016 for the treatment of SMA. *Exondys51* is a phosphorodiamidate morpholino oligomer (PMO) that targets the *DMD* pre-mRNA to exclude exon 51, in order to re-frame the dystrophin transcript around frame shifting deletions flanking exon 51, as a treatment for DMD (Mendell *et al.* 2013). *Exondys51* received accelerated approval from the FDA on September 19[th] 2016. Other drugs exploiting the antisense concept include Kynamro[®] (mipomersen sodium) (Wong and Goldberg 2014), an adjunctive treatment to reduce LDL-C levels in familial hypercholesteremia, approved in January 2013, followed by an FDA hepatotoxity black box warning; and a number of compounds at various stages of clinical development, predominantly siRNA and oligodeoxyribonucleotide analogues designed to induce RNAse H degradation of target transcripts (for review see (Bennett *et al*. 2017, Stein and Castanotto 2017)). Recently, Alnylam announced the first FDA approval of an RNAi Therapeutic, ONPATTRO™ (patisiran) for the treatment of adult patients with polyneuropathy of hereditary transthyretin-mediated amyloidosis. (https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm616518.htm)

Strategies to improve biological stability and confer pharmaceutical properties to nucleic acid drugs include chemical modifications of the bases and nucleic acid backbone to increase resistance to endogenous nucleases; modifications that influence specific oligomer activity and conjugates that enhance cellular uptake and improve tissue distribution. While the phosphorothioate backbone is the most widely applied chemical modification, changes to the ribose moiety (eg, 2′ O-methylation, 2′ O-methoxyethyl, 2′ O-fluoro), nucleobases and other backbone modifications, including peptide nucleic acids and phosphorodiamidate morpholino oligomers can confer specific characteristics and mechanisms of action (for review see (Veedu 2015, Wilton *et al.* 2015, Tri Le *et al.* 2016)).

Many laboratories, including our own, routinely use 2′ O-methyl phosphorothioate oligonucleotides to evaluate antisense sequences for targeted splicing modulation, and different modified bases on a phosphorothioate backbone for various other molecular interventions (Lipi *et al.* 2016, Chakravarthy *et al.* 2017). These compounds can be economically prepared in-house and efficiently transfected into a range of cells as cationic liposomes or dendrimer complexes, using commercially available reagents. However, for sustained splice modification, *in vivo* use and clinical application, PMOs have proved safe and effective (Gebski *et al.* 2003, Fletcher *et al.* 2006, Kinali *et al.* 2009, Mendell *et al.* 2016). We previously reported that antisense exon skipping sequences prepared and optimized *in vitro* as 2′ O-methyl phosphorothioate oligonucleotides generally perform comparably when synthesized as PMO (Adams *et al.* 2007). An extended antisense sequence targeting the *ISS-N1* region of *SMN2* was validated *in vitro* and *in vivo* using 2′ O-methyl phosphorothioate and PMO chemistries (Mitrpant *et al.* 2013).

Phosphorothioate backbone modification of nucleic acid drugs is common practice for protection from rapid degradation by circulating and intracellular nucleases, but these modified AOs can interact undesirably with endogenous cellular components. Early work indicated that negatively charged phosphorothioate AOs bind non-specifically to heparin-like proteins, laminin and collagen (Dias and Stein 2002) and more recently, phosphorothioate AOs were reported to interact with nuclear paraspeckle-associated proteins, sequestering them away from their endogenous target, the long non-coding RNA *NEAT1,* and also seeding paraspeckle like-structures in the absence of *NEAT1* (Shen *et al.* 2014, Shen *et al.* 2015). Paraspeckles and paraspeckle proteins are involved in transcriptional regulation, transport and splicing pathways (Bond and Fox 2009, Naganuma *et al.* 2012). Beyond the paraspeckle-associated subset (Shen *et al.* 2014), phosphorothioate AOs were also reported to bind an additional approximately 50 intracellular proteins (Liang *et al.* 2015) however, these interactions were studied in the context of evaluating the modest impact these interactions had on AO function (Liang *et al.* 2015). A broader question relates to the global consequences on cellular processes arising as a result of the interaction of AOs with nuclear proteins.

Results of *in vivo* studies investigating the consequences of non-specific phosphorothioate AO binding are particularly relevant when considering therapeutic applications. The

sequestration of paraspeckle proteins, in particular proteins of the drosophila

behaviour/human splicing (DBHS) family, was associated with acute hepatotoxicity,

inflammation and apoptosis after application of 2′ fluoro-modified phosphorothioate

antisense oligonucleotides with a 5-10-5 gapmer configuration in mice (Shen *et al.* 2018)

and while this study focused on the severity of 2′ fluoro-modified phosphorothioate AOs, all

2′ modifications evaluated have been shown to sequester paraspeckle proteins, albeit to

varying degrees (Shen *et al.* 2014, Liang *et al.* 2015, Shen *et al.* 2018). The backbone-

dependent binding of phosphorothioate-modified olionucleotides to platelets *in vitro* and *in*

vivo, mediated by the platelet-specific receptor glycoprotein VI (GPVI) is also of note (Flierl

et al. 2015), considering the broad range of nucleic acid therapeutics currently under

investigation (for review see (Bennett *et al.* 2017). Oligonucleotides on a phosphorothioate-

modified backbone elicited strong platelet activation, signaling, reactive oxygen species

generation, adhesion, spreading, aggregation and thrombus formation (Flierl *et al.* 2015). Of

particular relevance to current clinical usage, 2′ O-methyl phosphorothioate AOs were

reported to activate innate immunity when administered directly to the central nervous

system (Toonen *et al.* 2018).

Here we report that transfection of cultured cells with 2′ O-methyl phosphorothioate

antisense sequences resulted in numerous novel, large nuclear inclusions in the form of

highly structured fibril-like aggregates that co-stained for the paraspeckle proteins, NONO,

SFPQ, PSPC1 and FUS. Other nuclear proteins showed altered distribution in 2′ O-methyl

phosphorothioate transfected cells. Intranuclear inclusions begin to form within four hours of

transfection, and become dominant structures throughout the nucleus within 24 hours. The

inclusions appear stable once formed and may remain evident on the culture substrate, even

after death and disintegration of the cell. Transmission electron microscopy on transfected

cells revealed numerous large, regular structures reminiscent of amyloid deposits, with

electron dense regions. Furthermore, gene ontology analyses following RNA sequencing

demonstrated significant disruptions to chromatin silencing; regulation of autophagy;

nucleotide excision repair; membrane and organelle organization; apoptosis; signalling and

protein transmembrane transport, following 2′ O-methyl phosphorothioate transfection.

Materials and Methods

Antisense oligonucleotides

Phosphorodiamidate morpholino oligomers (PMOs) were purchased from Gene-Tools LLC

(Philomath, OR, USA), and 2′ O-methyl phosphorothioate AOs were synthesised by TriLink

BioTechnologies (San Diego CA, USA). The following sequences were evaluated; an AO

encompassing the *SMN* intron 7 *ISS-N1* target, but with a longer sequence *SMN7D(-10-29)*

(5′AUUCACUUUCAUAAUGCUGG 3′); the Gene-Tools standard control oligomer, only

known to be biologically active in reticulocytes carrying a splice mutation in the human beta-

globin pre-mRNA, as an unrelated sham control AO sequence

(5′CCUCUUACCUCAGUUACAAUUUAUA 3′), and identified as 'control AO' throughout

this study; *Smn* (5′CAC CUU CCU UCU UUU UGA UU 3′) designed to induce exon mouse

Smn exon skipping and a *SMN* sense AO (5′ CCAGCAUUAUGAAAGUGAAU 3′)

complementary to *SMN7D(-10-29)*.

Transfection

The 2′O-methyl phosphorothioate AOs were transfected into dermal fibroblasts as lipoplexes

using 3 µl of Lipofectamine 3000 (Life Technologies, Melbourne, Australia) per 1 ml of

OptiMEM, according to the manufacturer's protocol. The transfection mix was applied to

cells seeded at 10,000 per cover slip for immunofluorescence and 15,000 cells per well in 24

well plates for RNA extraction and incubated (37°C) for 24 hours prior to RNA and protein

analysis. SMA patient fibroblasts were used for the experiments shown in Figures 1f and 1g

(Coriell Cell Repositories, GM03813) and fibroblasts from our in-house biobank, obtained

from a healthy volunteer, with informed consent (Murdoch University Human Research

Ethics Committee Approval #2013/156) were used for all other experiments, except live-cell

imaging (see below).

PMOs were delivered to cells either uncomplexed (10 µM), annealed to a complementary

sense DNA oligonucleotide (phosphodiester leash) and transfected as a lipoplex (200 nM), or

by nucleofection using a Nucleofector™ X Unit (Lonza, Melbourne, Australia) according to

the manufacturer's instructions. PMOs were delivered using the P2 nucleofection kit and CA-

137 program at 1 µM, as determined by the final transfection volume. All PMO transfections

were incubated for 72 hours prior to RNA and protein analysis.

RT-PCR on AO transfected cells

RNA was extracted using the MagMAX-96 Total RNA Isolation Kit, including DNase

treatment (Life Technologies), according to the manufacturer's instructions. RT-PCRs were

performed using the One-step Superscript III RT-PCR kit with Platinum Taq polymerase (Life

Technologies) according to the manufacturer's instructions. All primer sequences and PCR

conditions used in this study are detailed in Supplementary File 1**, Table 1**.

cDNA synthesis and quantitative PCR

cDNA was synthesised from (~500 ng) RNA extracted from treated and untreated cell cultures

using the Superscript IV first-strand synthesis system (Life Technologies) as per the

manufacturer's instructions. Prior to qPCR amplification, cDNA was diluted in RNase/DNase

free water (1:5).

The qPCR reactions were performed using Fast SYBR™ Green Master Mix (ThermoFisher

Scientific, Melbourne, Australia) and primers for ribosomal RNA subunits *5S* (100% primer

efficiency), *18S* (95% primer efficiency), *45S* (100% primer efficiency) and housekeeping Tata

box protein (*TBP;*100% primer efficiency) and beta Tubulin (*TUBB;*94% primer efficiency)

transcripts. All primer sequences and cycling conditions used in this study are detailed in

Supplementary File 1, **Table 1.** Transcript abundance was measured using the CFX384 Touch™

Real Time PCR detection system (Bio-Rad Laboratories Pty., Ltd., Gladesville, Australia). The

relative expression of each ribosomal RNA subunit to *TBP* and *TUBB* mRNA was calculated

using the $2^{-DD CT}$ method and presented as a fold change compared to untreated cells.

Immunofluorescence
192
193 Approximately 10,000 fibroblasts were seeded onto 22 mm x 22 mm coverslips in

194 6- well plates and incubated for 24 hours, prior to transfection. Following transfection,

195 the cells were fixed using acetone: methanol (1:1) on ice for 4 minutes and air- dried.

196
197
Fixed cells were washed in PBS containing 1% Triton X-100 to permeabilise the

nuclear membrane, and then in PBS to remove excess Triton X-100. Primary antibodies

were diluted in PBS containing 0.05% Tween20 and applied to cells for 1 hour at room

temperature. All antibody details and staining conditions are listed in Supplementary File

1 **Table 2**. Primary antibodies were detected using AlexaFluor488 anti- mouse (cat no.

A11001), anti-rabbit (cat. No A11008) or AlexaFluor568 anti- mouse (cat. No A11004) or

anti-rabbit (cat. No A11011) (1:400) after incubation for 1 hour at room temperature, and

counterstained with Hoechst 33342 (Sigma-Aldrich) for nuclei detection (1 mg/ml, diluted

1:125).

Western blotting

Cell lysates were prepared with 125 mM Tris/HCl pH 6.8, 15% SDS, 10% Glycerol, 1.25 µM

PMSF (Sigma-Aldrich, NSW, Australia and 1x protease inhibitor cocktail (Sigma-Aldrich)

and sonicated 6 times (1 second pulses) before adding bromophenol blue (0.004%) and

dithiothreitol (2.5 mM). Samples were heated at 94°C for 5 minutes, cooled on ice and

centrifuged at 14,000 x g for 2 min before loading onto the gel.

Total protein (10 µg), measured by BCA, was loaded per sample on a NuPAGE Novex 4-12%

BIS/Tris gel (Life Technologies) and separated at 200 V for 55 minutes. Proteins were

transferred onto a Pall Fluorotrans polyvinylidene fluoride (PVDF) membrane at 350 mA for

2 hours. Following blocking for 1 hour, the membrane was incubated in 5% skim milk

powder in 1x TBST containing the primary antibody diluted as shown in Supplementary File

1, **Table 2.** Immunodetection was performed using the Western Breeze Chemiluminescent

Immunodetection System (Life Technologies) according to the manufacturer's instructions.

Western blot images were captured on a Vilber Lourmat Fusion FX system using Fusion

software and Bio- 1D software was used for image analysis.

Live cell imaging

U2OS cells, previously modified to tag the endogenous SFPQ gene with GFP were used for

live-cell imaging (Li *et al.* 2017). Cells were seeded at 1.5 x 10^5 cells per well in a 12-well

plate, 24 hours before transfection, with high glucose DMEM (Life Technologies, Cat

No.11995065) supplemented with 10% fetal calf serum and 1% penicillin-streptomycin. On

the following day, cells in each well were transfected with 100 nM 2′ O-methyl

phosphorothioate AOs complexed with 2.2 µl Lipofectamine 3000 following the

manufacturer's instructions. Immediately after transfection, the plate was transferred to an

Incucyte S3 (Essen Bioscience) for live-cell imaging at 1 hour intervals under 10 x optical

zoom. SFPQ-GFP and activated caspase-3/7 were visualised under standard green and red

channels. Incucyte Caspase-3/7 Red Apoptosis Assay Reagent (Essen Bioscience, Cat No.

4704) was used to reveal cell apoptosis events.

High resolution microscopy

SIM imaging was performed using an N-SIM microscope (Nikon Corporation, Tokyo, Japan),

with SR Apochromat TIRF 100 x 1.49 NA oil immersion objective. Spherical aberration was

reduced using the Ti2 automated correction collar at the beginning of the imaging session.

Images were acquired using 405 nm, 488 nm and 561 nm lasers, with stacks of step size 0.12

μm (with top and bottom of samples determined visually), using 3D-SIM mode. Images were

reconstructed with NIS Elements software (Nikon Corporation, Tokyo, Japan).

Transmission electron microscopy

Following transfection, cells were washed in PBS and fixed in cold 2.5% phosphate buffered

glutaraldehyde overnight. The fixed cells were scraped and centrifuged, then embedded in 4%

agarose. The pellets were processed using a Leica tissue processor, moving through 1%

aqueous osmium tetroxide, increasing graded alcohols, propylene oxide, propylene

oxide/araldite mix, and finally pure araldite resin. The osmicated cells, surrounded by resin in

a beem capsule were polymerised overnight at 80ºC to form hard blocks.

Semi-thin (0.5 micron) sections were taken from the blocks using glass knives, on an RMC

ultratome, and stained with methylene blue (0.1% aqueous Methylene Blue with 0.1%

Borax) to visualise available cells. Ultrathin sections were then cut from selected blocks,

using a Diatome diamond knife, at approximately 95 nm thickness and mounted on copper

mesh grids and stained with uranyl acetate (5% uranyl acetate solution in 5% aqueous acetic

acid) and Reynold's lead citrate. The grids were viewed using a JEOL 1400 TEM, at 80 kV

and images captured by an 11-megapixel GATAN digital camera at varying magnifications.

RNA sequencing and analysis

RNA quality was confirmed using a Bioanalyser (Perkin Elmar, MA, USA) prior to

RNAseq. Samples were sent to the Australian Genome Research Facility (AGRF, Perth and

Melbourne, Australia) for whole transcriptome library preparation using the TruSeq Stranded

Total RNA Library Prep Kit (Illumina, CA, USA) and ribosomal RNA depletion with the

Ribo-Zero-Gold kit (Illumina, CA, USA). Sequencing was performed using an Illumina

HiSeq 2500 (Illumina, CA, USA) to generate 100 base pair single end reads, resulting in an

average 25 million reads per sample. Raw sequencing files were quality checked using

FastQC (0.11.7), with all files passing. No adapter contamination was identified, and reads

were not trimmed. Transcript quantification was performed with salmon (0.8.2), using an

index constructed from the Ensembl GRCh38.93 annotation. Transcript abundance was

summarised to gene-level counts and imported into R using tximport (1.4.0). Differential

expression analysis was performed using DESeq2 (1.16.1) and the default parameters (alpha

= 0.1). The heatmap of differentially expressed genes ($p_{adj} < 0.1$) was constructed using the

heatmap.2 function from R package gplots (3.0.1). Gene ontology analysis was performed

using GSEA (3.0) with 1000 gene set permutations. The gene ontology network was

constructed using the Enrichment Map plugin for Cytoscape (3.5.1), with cut-offs: $p < 0.05$,

FDR < 0.1 and edge > 0.3.

Results

Formation of nuclear inclusions after 2′ O-methyl phosphorothioate AO transfection

The phosphorothioate backbone can interact non-specifically with both intra- and extra-

cellular proteins, and recruits paraspeckle proteins and forms paraspeckle-like structures

when transfected into Hela cells (Shen *et al.* 2014). However, the dynamics of the formation

of these structures, as well as the long-term consequences of these structures on cell

physiology remain unknown.

In this study, we largely focused on two 2′ O-methyl phosphorothioate AOs, one that

promotes *SMN2* exon 7 retention during splicing by targeting the *ISS-N1* intronic silencer

motif (Singh *et al.* 2006) and the other a control oligomer sequence reported by Genetools

LLC and widely used as a transfection control, although additional AO sequences were also

tested in some experiments, as indicated below. In order to follow nuclear inclusion

formation, we first transfected 2′ O-methyl phosphorothioate AO as lipoplexes (100 nM)

into U2OS cells expressing genome engineered GFP-labelled endogenous SFPQ (Li *et al.*

2017) and performed live cell imaging over 48 hours (**Figure 1a**). The AO transfection

induced formation of nuclear inclusions within 5 hours, and over a period of 12 hours, all of

the SFPQ within the nucleus is observed sequestered within the inclusions, as revealed by

decreased SFPQ staining of the nucleoplasm. Ultimately, only the inclusions are evident, at

which time, the inclusions then coalesce into discrete filaments.

Immunofluorescent staining of the paraspeckle protein, NONO, showed colocalization in

nuclear inclusions in fibroblasts transfected with 2′ O-methyl phosphorothioate *SMN7D(-10-*

29) sequence at 100 nM concentration, but were also evident at AO concentrations as low as

12.5 nM within 24 hours of transfection (**Figure 1b**). Of note, the untreated cells show

diffuse nucleoplasmic NONO signal, as well as naturally occurring punctate NONO inside

endogenous paraspeckle nuclear bodies (Fox *et al.* 2018) (eg. **Figure 1b** (ii)). Nuclear

inclusions were also observed following 2′O-methyl phosphorothioate AO transfection into

normal human myoblasts, transformed mouse *H2K* myoblasts, and the neuroblastoma SH-

SY5Y cell line (Supplementary File 1, **Figure 1).** Once formed, the inclusions appear to be

highly stable, and remain evident on the culture substrate following cell death and nuclear

fragmentation and blebbing **(Figure 1c (i-ii)).** The nuclear inclusions appear as either long

'filaments', punctate foci, or both (**Figure 1c (iii)**). In the past we have routinely observed

marked cell death associated with transfection of 2′ O-methyl phosphorothioate AOs into

primary cells and cell lines, and hence explored any correlation between cell death and the

formation of AO-induced nuclear inclusions. Indeed, we observed here that transfection with

2′ O-methyl phosphorothioate AO sequences induced nuclear inclusions that correlated with

reduced cell survival (**Figure 1d-e**).

While transfection of AOs with phosphorothioate and phosphodiester backbones have been

demonstrated to result in nuclear inclusions (Shen *et al.* 2014), PMOs are yet to be evaluated

in this context. We therefore transfected fibroblasts with the AO sequences described above,

synthesized as PMOs. Transfection of the PMO sequences into fibroblasts did not alter the

apparent distribution of NONO, nor of any other nuclear proteins studied (**Figure 1f).** In

order to demonstrate that poor cellular uptake of these compounds was not a factor, PMOs

were transfected using three established delivery techniques known to lead to altered target

gene expression; uncomplexed PMO (10 µM), nucleofection (1 µM), and annealed to a

complementary DNA "leash" and transfected as a lipoplex (200 nM), all incubated for 72

hours after transfection (**Figure 1f (i-iii),** respectively). Immunofluorescent detection of

NONO showed no PMO-induced intranuclear inclusion formation or altered distribution of

NONO following transfection, under any of the conditions used, and was reproducible,

irrespective of the PMO sequence. Transfection efficiency and altered exon selection in the

target *SMN2* transcript was assessed by RT-PCR across *SMN* exons 4-8, confirming that the

absence of intranuclear inclusions in PMO transfected cells was not a consequence of poor

uptake of PMO (**Figure 1g**).

Figure 1 – **Formation of nuclear inclusions**. Showing **(a)** live cell imaging time-course of U2OS cells with endogenous SFPQ-GFP showing the formation of nuclear inclusions over 24 hours following 2′ O-methyl phosphorothioate control AO transfection (100 nM), inclusions are indicated by the white arrow; **(b)** fibroblasts transfected with 12.5 nM of 2′ O-methyl phosphorothioate *SMN7D(-10-29)* stained for NONO (i), compared to untreated fibroblasts (ii); **(c)** fibroblasts stained for NONO show, (i) nuclear inclusions without evidence of nuclear staining after 2′ O-methyl phosphorothioate *SMN7D(-10-29)* transfection, (ii) nuclear blebbing after 2′ O-methyl phosphorothioate control AO transfection and (iii) nuclear inclusions in the form of filaments or foci induced by transfection with a 2′ O-methyl phosphorothioate *SMN7D(-10-29)* sense sequence AO; **(d)** graph showing the number of viable fibroblasts following 2′ O-methyl phosphorothioate AO transfection (n=4), error bars represent standard error of the mean and P-values were calculated using an unpaired T-test; **(e)** graph showing the percentage of fibroblasts containing nuclear inclusions following 2′ O-methyl phosphorothioate transfection with a minimum of 200 cells counted per sample; **(f)** fibroblasts stained for NONO after transfection with PMO *SMN7D(-*

10-29) at (i) high dose un-complexed (10 µM), (ii) using a complementary leash and transfection reagent (200
nM) and (iii) using electroporation (1 µM), all incubated for 72 hours; and **(g)** RT-PCR of full length (FL) and exon
7 skipped (Δ7) *SMN2* transcripts of RNA collected following transfections as in **(f)**. All scale bars = 10 µm.

To further explore whether AO-induced nuclear inclusions are sequence independent, ninety

2′ O-methyl phosphorothioate AOs targeting structural gene transcripts, transcription factors,

splicing factors and enzymes, were transfected into fibroblasts and the cells were

immunostained for SFPQ (Supplementary File 1, **Table 3)**. All AOs tested disrupted SFPQ

distribution and all but one AO formed nuclear inclusions by 24 hours following AO

transfection at 100 nM, irrespective of length (18-30 bases) and nucleotide composition

(Supplementary File 1, **Table 3**), indicating that with optimal transfection efficiency, any 2′

O-methyl phosphorothioate sequence is likely to induce nuclear inclusions. One transfected

AO induced NONO localization to the nuclear envelope (Supplementary File 1, **Figure 2a**),

while a number of AO sequences induced cytoplasmic SFPQ aggregates in addition to

nuclear inclusions (Supplementary File 1, **Figure 2b**). The AO-mediated cytoplasmic

aggregate formation was also reported by Liang *et al.,* 2014, who observed cytoplasmic

structures as a consequence of TCP1-beta subunit interaction with phosphorothioate-AOs in

the cytoplasm, in addition to formation of nuclear 'phosphorothioate bodies', and concluded

that upon transfection, the TCP1 proteins interact with phosphorothioate AOs and enhance

antisense activity (in this case, activation of the RNase H-mediated target degradation)

(Liang *et al.* 2015).

Composition of nuclear inclusions induced by 2′ O-methyl phosphorothioate AO transfection

The structure and composition of nuclear inclusions was further investigated by

immunostaining of additional selected nuclear proteins found in paraspeckles, the nuclear

envelope, nucleoli, nuclear speckles, Cajal bodies and nuclear stress bodies. The paraspeckle

proteins PSPC1 and FUS were evident in nuclear inclusions (Supplementary File 1, **Figure**

3a and 3b) while the proteins TDP-43 and hnRNP-A1 were not observed to be components

of nuclear inclusions (Supplementary File 1, **Figure 3c and 3d).**

Coilin, a component of nuclear Cajal bodies, shows altered distribution in 2′ O-methyl

phosphorothioate control AO transfected cells and is distributed evenly through the

nucleoplasm and cytoplasm of transfected cells showing NONO-positive nuclear inclusions,

 Figure 2 – Staining of nuclear bodies following 2´ O-methyl phosphorothioate transfection (100 nM, 24
hours), with AOs as indicated in fibroblasts. **(a)** transfected with the control AO showing coilin (red) and
NONO (green) overlayed with hoechst (blue), and graph showing the percentage of abnormal and normally
stained cells in cultures transfected with *SMN7D(-10-29)* and control AOs; **(b)** transfected with *SMN7D(-10-29)*
showing nucleolin (green) and SFPQ (red) overlayed with hoechst (blue), and graph showing the percentage of
abnormal and normally stained cells in cultures transfected with *SMN7D(-10-29)* and control AOs*;* **(c)** transfected
with the control AO showing fibrillarin (red) and NONO (green) overlayed with hoechst (blue), and graph showing
the percentage of abnormal and normally stained cells in cultures transfected with with *SMN7D(-10-29)* and
control AOs and **(d)** transfected with *SMN7D(-10-29)* showing SC35 (green) and SFPQ (red) overlayed with
hoechst (blue), and graph showing the percentage of abnormal and normally stained cell in cultures transfected
with *SMN7D(-10-29)* and control AOs. A minimum of 100 cells were counted for each sample. All scale bars = 10
µm.

whereas it is present almost exclusively in the nucleus of untreated cells (**Figure 2a).**
Nucleolin and fibrillarin, both markers for the nucleolus, show altered distribution in nuclear
inclusion-containing cells, changing from nucleolar to diffuse nucleoplasmic, as shown in
Figures 2b and **2c**, respectively. The splicing factor and component of nuclear speckles,
SC35, showed enhanced cytoplasmic localisation in cells that also contained nuclear
inclusions, revealed here by SFPQ immunofluorescence (**Figure 2d**).

The percentage of AO transfected cells showing disrupted staining or abnormal localization
of coilin, nucleolin, fibrillarin and SC35 staining correlated with cells containing
intranuclear inclusions, as indicated in the graphs (**Figure 2a-d**). Line intensity profiling
shows altered distribution of these nuclear proteins in individual cells that have nuclear
inclusions, and is presented in Supplementary File 1, **Figure 4.** Since components of
subnuclear organelles were disorganised following transfection with phosphorothioate AOs,
the levels of these proteins in the transfected cells were investigated. Western blot analysis
showed that the overall abundance of each protein studied to be essentially unchanged
following AO transfection. Representative western blots probed for SFPQ, NONO, TDP43,
hnRNP-A1, NCL, HSF1 and beta actin, are shown in Supplementary File 1, **Figure 5**.

Morphology of 2´ O-methyl phosphorothioate AO-induced nuclear inclusions
Detailed analysis of the 2′ O-methyl phosphorothioate *SMN7D(-10-29)*AO-induced nuclear
inclusions was performed using transmission electron microscopy (TEM) on ultra-thin,
osmium stained sections (**Figure 3a(i-vi))**. The nuclear inclusions appeared to be
microfibrillar or amyloid-like, approximately 200-250 nm in diameter, occurring mostly in
groups or bundles. For reference, the width of a DNA double helix is ~2 nm. If captured in a

suitable plane of section, the structures appear to have very electron dense termini. Such

nuclear inclusions have never before been revealed by transmission electron microscopy in

the investigating laboratory. The electron dense regions are reminiscent of perichromatin in

size and electron density, the inclusions revealed by TEM extend to ~ 2000 nm in length.

Numerous small, partially formed inclusions are also evident in nuclei of transfected cells

when viewed at higher magnification (**Figure 3a (iii)**).

Since transmission electron microscopy can only capture segments of the presumed large

fibril-like nuclear inclusions, super resolution microscopy was used to examine 2′ O-methyl

phosphorothioate *SMN7D(-10-29)* AO transfected and untreated fibroblasts (**Figure 3b-c**).

AO transfected fibroblasts co-stained for NONO and SFPQ show a network of many nuclear

inclusions characterised by co-localisation of the two proteins, whereas the untreated cells

show more even, punctate distribution within paraspeckles (**Figure 3b (i, ii),** respectively).

Transfected fibroblasts co-stained for SFPQ and FUS reveal large interconnected fibril-like

structures decorated with FUS at mid-points and at the termini (**Figure 3c (i)**). SFPQ, NONO

and FUS are all sequestered to the nuclear inclusions, unlike the even nuclear distribution of

these proteins in untreated cells (**Figure 3c (ii)**).

The nuclear envelope protein, lamin B1 showed altered distribution in a proportion of 2′ O-

methyl phosphorothioate *SMN7D(-10-29)* transfected cells that exhibited nuclear inclusions

(**Figure 3d (i-ii)**). In these cells, the nuclear envelope labelled by immunostaining of lamin

B1 appears multi lobular and distorted, reminiscent of those found in the premature ageing

disease progeria and in E145K cells (Taimen *et al.* 2009). A single z-frame image is shown

in **Figure 3d (i)** that clearly illustrates lamin B1 localized in 4 distinct nuclear lobules.

Figure 3d (ii and iii) shows the composite image and an image of an immunostained,

untreated fibroblast. In all images (**Figure 3b-d**), the nuclear inclusions appear to reside

between areas that may reflect chromosomal territories.

 Figure 3 – Structural analysis of nuclear inclusions in 2´ O-methyl phosphorothioate transfected

fibroblasts. Fibroblasts were transfected with the 2´ O-methyl phosphorothioate AO, *SMN7D(-10-29)* at 100 nM,

for 24 hours). Transfected and untreated control cells were processed for transmission electron microscopy (**a**)

and super resolution fluorescence microscopy (**b-d**). Showing (**a**) transmission electron microscopy of (i) nucleus

with filament-like nuclear inclusions (scale = 5 µm), (ii) higher magnification of (i) (scale = 500 nm), (iii) nucleus

with foci-like nuclear inclusions (scale = 2 µm), (iv) higher magnification of (iv) (scale = 200 nm), (v) partially

formed nuclear inclusions (scale = 200 nm), and (vi) untreated nucleus (scale = 2 µm); (**b**) super resolution

fluorescence microscopy of SFPQ (green) and NONO (red) co-staining following 2´ O-methyl phosphorothioate

(*SMN7D(-10-29)*) transfection (i) and an untreated cell (ii); (**c**) super resolution fluorescence microscopy of SFPQ

(red) and FUS (green) co-staining following transfection (i) and an untreated cell (ii); (**d**) super resolution

fluorescence microscopy of NONO (red) and LAMIN-B1 (green) co-staining following transfection (i-ii) and an

untreated cell (iii). A single z-frame is shown in (i) and a maximum intensity z-stack in (ii). For all SIM images

scale bar = 5 µm.

2´ O-methyl phosphorothioate AOs affect ribosomal RNA processing and maturation

Since Cajal bodies and fibrillarin are important in ribosomal RNA processing and assembly,

we assessed rRNA levels in 2´ O-methyl phosphorothioate AO transfected cells and

untreated cells. qPCR and bioanalyser data show unprocessed rRNA was markedly increased

by 2´ O-methyl phosphorothioate AO transfection in fibroblasts (**Figure 4**). A Bioanalyser

RNA trace from 2´ O-methyl phosphorothioate control AO transfected fibroblasts (**Figure**

4a) showed low levels of 18S and 28S ribosomal subunits, as well as a number of

intermediate peaks compared to that on RNA from untreated fibroblasts (**Figure 4b**),

consistent with incomplete rRNA processing. Poor RNA quality was ruled out, given a

similar pattern was observed in all experimental repeats (n=6). This observation was

supported by qPCR analysis of rRNA levels showing a 2 to 3.5-fold increase in the level of

the unprocessed 45S rRNA in 2´ O-methyl phosphorothioate transfected compared to

untreated fibroblasts (**Figure 4c**). Unprocessed rRNA accumulated over time, following the

formation of nuclear inclusions (**Figure 4d**).

Figure 4 – Analysis of ribosomal RNA processing. Showing bioanalyser trace of rRNA from **(a)** 2´ O-methyl phosphorothioate control AO transfected cells, and **(b)** untreated cells; **(c)** qPCR analysis of 5S, 18S and 45S rRNA levels following 2´ O-methyl phosphorothioate transfection (24 hours). rRNA levels were normalised against *TBP* and compared to those in untreated cells where untreated = 1 (n=3). Error bars represent the standard error of the mean, and P-values were calculated comparing each AO treatment group to the untreated group using an unpaired T-test; and **(d)** qPCR analysis of rRNA levels over 8, 16 and 24 hours; analysis as in **(c)** (n=1)**.**

2´ O-methyl phosphorothioate AO transfection promotes apoptosis

Following phosphorothioate AO transfection of U2OS cells at 100 nM, live cell imaging was undertaken to evaluate caspase activation (**Figure 5a**). Nuclear inclusions were observed

(a)

0h   4h   5h   19h

21h   22h   23h   25h

31h   32h   33h   34h

(b)

2´O-methyl-PS   PMO

SMN Ctrl UT   SMN Ctrl UT

p53 — 53 kDa

Beta-Actin — 42 kDa

(d)

2´O-methyl-PS

Ctrl   SMN   UT

BCL-xL
BCL-xS

(c)

P=0.001

P=0.009

Relative p53 expression

1.20
1.00
0.80
0.60
0.40
0.20
0.00

SMN7D (-10-29) | Control AO | Untreated | SMN7D (-10-29) | Control AO | Untreated

2'O-methyl-PS   PMO

Figure 5 – Analysis of cellular toxicity following 2´ O-methyl phosphorothioate AO transfection. (a) live
cell imaging over time of U2OS cells with endogenous SFPQ-GFP (green) transfected with the 2´ O-methyl
phosphorothioate control AO and stained for caspase 3/7 activation (red). Two sister cells are annotated with
arrows, and scale bar = 10 μm; **(b)** western blot of P53 and β-actin levels in fibroblasts following 2´ O-methyl
phosphorothioate AO (n=3) and PMO transfection (n=1); **(c)** densitometry analysis of **(b)** where P53 levels are
normalized to β-actin and compared to those in untreated cells where untreated = 1. Error bars represent the
standard error of the mean and P-values were calculated using an unpaired T-test; and **(d)** RT-PCR analysis of

BCL2 transcripts in fibroblasts following 2′O-methyl phosphorothioate AO transfection, as indicated at 100 and 50
nM (24 hours).

after 5 hours in two sister cells. Nuclear inclusions became larger and more compact

overtime, followed by caspase activation after 22 hours. No caspase activation was observed

in the absence of intranuclear inclusions, identifying nuclear inclusion as complicit in

caspase activated apoptosis, leading to cell death. The tumour suppressor protein p53 was

decreased, following transfection with 2′ O-methyl phosphorothioate AOs, relative to

untreated control cells (**Figure 5b and c**), suggesting that the cell death observed was p53

independent, contrary to the findings by Shen *et al.,* 2018 (Shen *et al.* 2018) when using a 2′

fluoro-modified AO on a phosphorothioate backbone in the mouse. In comparison to

phosphorothioate AO transfection, p53 levels remained unchanged following PMO

transfection (**Figure 5b and c**).

The mitochondrial membrane and apoptosis regulating, *BCL2*-like gene encodes two

alternatively spliced, functional protein isoforms, a longer anti-apoptotic isoform BCL-xL,

and a shorter pro-apoptotic isoform BCL-xS. Increased abundance of the BCL-xS isoform

induces cytochrome C release from the mitochondria, initiating caspase activated apoptosis.

RT-PCR across the *BCL* transcripts (**Figure 5d**) shows 2′ O-methyl phosphorothioate AO

transfection induces non-specific alternative splicing, increasing the levels of the pro-

apoptotic *BCL-xS* transcript, potentially initiating the caspase activation shown in **Figure 5a**.

Transcriptome analysis of 2′ O-methyl phosphorothioate transfected cells reveals global

cellular disruptions

We next evaluated the effect of 2′ O-methyl phosphorothioate transfection and nuclear

inclusion formation on the transcriptome by RNA-seq. We prepared RNA from fibroblasts

transfected with the two 2′ O-methyl phosphorothioate AOs used throughout this study as

well as two control samples (untreated cells and cells transfected with a DNA phosphodiester

AO (DNA control) version of the same sequence as the control AO, 100 nM for 24 hours) for

RNAseq analysis (n=3). The RNAseq data is available

at https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE121713 (accession number:

GSE121713). An RNAseq heatmap comparing the expression of transcripts in 2′ O-methyl

phosphorothioate AO transfected fibroblasts to that in cells transfected with the DNA

phosphodiester oligonucleotide and untreated cells showed stark differences in expression

Figure 6 – RNA sequencing analysis of 2´ O-methyl phosphorothioate AO transfected fibroblasts. (a) heatmap of all differentially expressed transcripts in control (1, 4, 5-untreated and 2, 3, 6, DNA AO) and 2´ O-methyl phosphorothioate AO *SMN7D(-10-29)* and control AO) transfected (100 nM, 24 hours) cells, whereby red

541 represents overexpressed transcripts, and blue represents under-expressed transcripts; and **(b)** gene ontology
542 (GO) network of significantly affected cellular pathways in samples from transfected, compared to untreated and
543 DNA AO transfected cells. GO terms are represented as dots being either overexpressed (red) or under-
544 expressed (blue), and groups with greater than 50% of genes in common are linked.
545

between the groups (**Figure 6a**). Transcripts that are overexpressed in 2´ O-methyl

phosphorothioate AO transfected cells compared to controls are indicated in red, while those

that are underexpressed are indicated in blue.

Gene ontology (GO) analyses showed significant disruptions in transcripts involved in a

number of major cellular pathways. The GO network (**Figure 6b**) revealed cellular processes

and pathways that are significantly over-expressed include signaling pathways, chromatin

silencing, and metabolic pathways, while those that are significantly under-expressed

include negative regulation of apoptosis and autophagy, membrane and organelle

organisation, protein transmembrane transport and nucleotide excision repair. Gene ontology

pathways included in each cellular process are listed in **Supplementary File 2.** Thus, overall

these data show dramatic and widespread changes to gene expression in human fibroblasts

as a result of the uptake of the 2´ O-methyl phosphorothioate AOs.

Discussion

AOs have therapeutic potential as modulators of gene expression in many different diseases,

and can do so through several different mechanisms. Antisense drugs that are in current

clinical use alter exon selection during pre-mRNA processing, or induce RNaseH degradation

of the target mRNA, the mechanism of action determined by the nature of the AO chemistry

(for review see (Tri Le *et al.* 2016)). However, emerging reports of off-target effects

conferred by synthetic oligonucleotides, identified *in vitro* (Shen *et al.* 2014, Shen *et al.* 2015)

and *in vivo* (Shen *et al.* 2014, Shen *et al.* 2018)*,* demand careful examination of the molecular

effects of these compounds. Here we report that transfection of cultured human dermal

fibroblasts, U2OS osteosarcoma cells, SH-SY5Y neuroblastoma cells and primary myogenic

cells with 2´ O-methyl phosphorothioate oligonucleotides causes the formation of large

structured nuclear inclusions, decorated with proteins that are normally components of

nuclear organelles, in particular, paraspeckles. We show evidence that 2´ O-methyl

phosphorothioate AO transfection alters the distribution of proteins normally associated with

subnuclear bodies, impacts on global transcript expression and ribosomal RNA processing,

among other critical cellular processes, and induces cell stress, followed by apoptosis.

Interaction of phosphorothioate oligonucleotides with nuclear proteins, including certain

paraspeckle proteins, and the formation of paraspeckle-like structures, independent of

NEAT1, has been reported (Shen *et al.* 2014, Shen *et al.* 2015, Shen *et al.* 2018). While Shen

et al. (Shen *et al.* 2014) focused on the effect of phosphorothioate AOs on NONO and

reported altered paraspeckle structure, this was in the context of AO-mediated modulation of

gene expression, not the broader cell biology. Other off-target effects and non-specific

binding of proteins, initiated by the negatively charged phosphorothioate backbone have

been reported elsewhere (Dias and Stein 2002, Winkler *et al.* 2010, Liang *et al.* 2015).

Hence, our experience and extensive research usage of AOs prompted further investigation of

the global consequences of 2′ O-methyl phosphorothioate AO transfection on cell biology,

showing many significant changes, as described above. In contrast, the transfection of

(charge-neutral) PMOs in the current study did not disrupt subnuclear structures, or result in

any alteration in the distribution or staining intensity of the nuclear proteins studied. This

result was reproducible using multiple delivery techniques, including high dose, uncomplexed

PMO transfection, transfection of PMO annealed to a DNA leash and complexed with

liposome reagent, as well as nucleofection. In comparison, all 2′ O-methyl phosphorothioate

AO sequences tested induced novel, structured intranuclear inclusions, revealed by

transmission electron microscopy to have amyloid aggregate-like appearance.

The nuclear inclusions observed in phosphorothioate transfected cells are associated with

altered nuclear architecture, disturbed gene expression and ultimately, apoptosis. While

formation of such large, and apparently irreversible structures would be expected to impact on

nuclear biology, the likely consequences of aberrant sequestration of paraspeckle and other

nuclear proteins on RNA processing and post transcriptional regulation are of prime concern.

Paraspeckles are dynamic, RNA-protein nuclear organelles that occur in the interchromatin

space in mammalian cells and are now known to contain over 40 different proteins associated

with one architectural long non-coding RNA, *NEAT1_2* (for review see (Fox *et al.* 2018)).

Observed in most cultured cells-other than various stem cells, and cells experiencing stress,

paraspeckles regulate gene expression through sequestration of proteins and RNAs (Fox *et al.*

2018), but are also implicated in microRNA biogenesis (Jiang *et al.* 2017). Paraspeckle

proteins can undergo rapid, but normally reversible aggregation from monomers to form

amyloid-like structures, seeded by *NEAT1_2*, and likely mediated through prion-like domains

that are enriched in uncharged polar amino acids (Lancaster *et al.* 2014). The mechanism by

which phosphorothioate oligonucleotides induce some paraspeckle proteins to form aberrant

aggregates is unknown, however, we do know that sub cellular localization and

phosphorothioate AO behaviour in transfected cells is influenced by protein interactions and

the 2′ sugar AO modifications (Bailey *et al.* 2017, Crooke *et al.* 2017), and that synthetic

double stranded RNA is also able to seed *de novo* paraspeckle formation (Shelkovnikova *et*

al. 2018). How other nuclear proteins become incorporated into the phosphorothioate AO

induced nuclear inclusions, and why TDP43, a major component of paraspeckles does not

appear in these inclusions in our study, remains elusive.

Immunofluorescent staining of 2′ O-methyl phosphorothioate transfected fibroblasts revealed

the paraspeckle components NONO, PSPC1, SFPQ and FUS co-localised with large nuclear

structures in excess of 2000 nm in length, with FUS decorating the structures in an ordered

manner. All four of these proteins include prion-like domains (for review see (Fox *et al.*

2018)) and we speculate that the known propensity of phosphorothioate backbone compounds

to interact with and bind proteins then alters the liquid-liquid phase properties of the

paraspeckle proteins, shifting them towards insoluble, amyloid-like aggregations. However,

not all paraspeckle components with prion-like domains investigated were found to colocalise

with the inclusions, and in addition, the localisation and distribution of some proteins

associated with other subnuclear bodies was altered in cells with nuclear inclusions.

Coilin, an integral component of Cajal bodies, nucleolin, one of the most abundant proteins in

the nucleolus but also found in the cytoplasm and on the cell membrane, fibrillarin, located in

the dense fibrillar component of the nucleolus and SC35 (SRSF2), an essential splicing factor

found in nuclear speckles all showed altered distribution in cells that have phosphorothioate

AO -induced nuclear inclusions, but did not co-localize with these inclusions. Cajal bodies

assemble spliceosomal and nucleolar ribonucleoproteins required for pre-mRNA and pre-

rRNA processing, and are recently proposed to contribute to genome organization, with global

effects on gene expression and RNA splicing (Wang *et al.* 2016). The nucleolus is the most

prominent nuclear structure, and is where synthesis and processing of ribosomal transcripts to

yield the mature rRNAs *5.8S, 18S* and *28S* from the *45S* pre- rRNA takes place. We show that

ribosomal RNA processing is greatly impaired by 2′ O-methyl phosphorothioate AO

transfection, and speculate that this is a likely manifestation of improper localization of major

protein components of the nucleolus and Cajal bodies and consequent disruption to their

functions. In addition, the aberrant inclusions that we speculate occupy interchromosomal

spaces may well impose physical constraints upon nuclear organisation, and prevent proper

localisation of Cajal bodies and the nucleoli close to their normal chromosomal sites. Tissues

with high demand for transcript splicing and ribosome biogenesis, and neurons in particular,

have prominent Cajal bodies, juxtaposed to nucleoli (for review see (Lafarga *et al.* 2017)),

and disruption or loss of Cajal bodies is associated with severe neuronal dysfunction (Tapia *et*

al. 2017). Indeed, disruption or depletion of Cajal bodies was seen as the earliest nuclear sign

of motor neuron degeneration in a spinal muscular atrophy mouse model and induced a

progressive nucleolar dysfunction in ribosome biogenesis (Tapia *et al.* 2017).

Paraspeckle biology has gained increasing interest due to the association of paraspeckle

proteins with neurodegenerative disease, in particular amyotrophic lateral sclerosis (ALS),

and the observation that *NEAT1_2* is up-regulated in early-stage motor neurons from the

spinal cords of ALS patients (Shelkovnikova *et al.* 2014, Yamazaki and Hirose 2015).

Paraspeckle formation, not observed in healthy spinal motor neurons, is enhanced in spinal

cords of patients with early stage sporadic and familial ALS, and mutations in many

paraspeckle proteins (eg. TDP-43, FUS, NONO, SFPQ) are associated with ALS (Nishimoto

et al. 2013, Shelkovnikova *et al.* 2018). Whether paraspeckles are protective or causative in

ALS molecular pathology is not known at this time (Yamazaki and Hirose 2015), nor is the

role of paraspeckles in disease well understood. Nevertheless, phosphorothioate AO-

mediated dysregulation of paraspeckle formation and altered nucleoli and Cajal body biology

shown in our study, together with the building body of evidence that nuclear organelle

dysfunction (Lafarga *et al.* 2017, Tapia *et al.* 2017) has implications for central nervous

system, and probably all, clinical applications of these compounds.

Not surprisingly, considering the altered nuclear architecture and distribution of proteins

implicated in RNA biology, transcriptome sequencing showed significant global effects on the

expression of transcripts and revealed disturbances to many critical cellular processes in

phosphorothioate AO transfected cells. Of particular concern, pathways involved in apoptosis,

chromatin silencing, cellular metabolism and a number of signalling pathways, autophagy and

nucleotide excision repair were disturbed. While we acknowledge that these *in vitro* studies in

replicating cells may not fully reflect potential off-target treatment effects in tissues *in vivo,*

Toonan et al., 2018 (Toonen *et al.* 2018) report significant upregulation of immune system-

associated genes in brains of mice treated by intracerebroventricular injection of 2′ O-methyl

phosphorothioate AO. The upregulation of immune system associated genes was detectable for

at least 2 months after the last AO administration.   Here, the exaggerated sequestration of

nuclear proteins as a result of 2′ O-methyl phosphorothioate AO transfection dramatically

disturbs RNA processing, disrupts nuclear architecture, and induces apoptosis, and it seems

reasonable to consider that injection site reactions and adverse effects reported after clinical

evaluation of the 2′ O-methyl phosphorothioate *Drisapersen* for the treatment of Duchenne

muscular dystrophy (Mendell *et al.* 2017) and *Kynamro*® (*Mipomersen*) for the treatment of

familial hypercholesterolemia (Wong and Goldberg 2014) may be mediated at least in part, by

non-specific interactions of nuclear components with the AO backbone. It might also be

prudent to deliberate on the reported effects of antisense drugs, attributed to the AO action on

the target transcript, and consider whether some level of the apparent antisense effect on

splicing, in particular, could perhaps be a consequence of disturbance of RNA processing

pathways more broadly.

We speculate that, unlike endogenous paraspeckles and other dynamic nuclear bodies, the

formation of the aberrant nuclear inclusions seen *in vitro* here does not appear to be

reversible. While 2′ O-methyl phosphorothioate transfected cultures showed reduced cell

numbers, whether this was due to cell death or impaired replication, or both, is uncertain, and

whether the nuclear inclusions *per se* or reduced availability of RNA processing are primarily

responsible requires further investigation. However, any effects resulting in perturbation of

nuclear proteins may be compounded by the formation of the nuclear amyloid-like

aggregates and likely disturbance of protein homeostasis, termed 'proteostasis' (Yerbury *et*

al. 2016). Although other descriptions of exogenously induced nuclear amyloid-like

aggregates are limited, Arnhold *et al.,* 2015 (Arnhold *et al.* 2015) identified large amyloid-

like aggregates in SH-SY5Y cells, treated with mercury, a notorious neurotoxicant, in a study

that explains the mechanism of heavy metal neurotoxicity and identified amyloid protein

aggregation in the cell nucleus as causative. Mass spectrometry of the purified protein

aggregates identified a subset of spliceosomal components and the nuclear envelope protein

lamin B1 (Arnhold *et al.* 2015). In our study, we also detected changes in lamin B1, although

here we did not detect lamin B1 in the nuclear inclusions, we nevertheless observed that

nuclear membrane and lamin B1 organization was distorted in fibroblasts containing nuclear

inclusions. Interestingly, the multi-lobulated nuclei and lamin B1 staining are reminiscent of

cells from progeria patients carrying the lamin A 433G>A mutation (E145K) (Taimen *et al.*

2009).

In summary, we report phosphorothioate backbone-specific effects of modified

oligonucleotides on the distribution and localization of nuclear proteins, appearance of novel

nuclear structures composed in part of a subset of paraspeckle protein components, and

sequence-independent effects on nascent RNA processing. While the *in vivo*, longer term

repercussions of exogenous oligonucleotide-induced nuclear protein aggregates that include

many paraspeckle components and cause sub-nuclear disorganisation are yet to be

determined, our observations suggest that phosphorothioate backbone antisense compounds

destined for clinical application would benefit from further scrutiny.

Acknowledgments

The authors would like to acknowledge technical advice from Russell Johnsen. This work was

performed in part during LLF's and ILP's PhD candidatures. Laboratory studies and ILP were

supported by funding from the Parry Foundation through the Spinal Muscular Atrophy

Association of Australia and LLF received a Team Spencer and Muscular Dystrophy WA PhD

stipend. The authors received research support from the NHMRC Project Grants

APP1147496, APP1086311, APP1144791 and the MNDi foundation, Western Australia.

Author Contributions

Conceived and designed experiments: LLF, RL, SF, MTA, ILP and AF. Performed

experiments: LLF, RL, MTA, ILP, AH, and LG. Analysis of RNA-seq data: JC. Prepared and

edited manuscript: LLF, SF, RL, MTA, ILP, JC, AH, LG, SDW and AF.

Declaration of interests: SF and SDW act as consultants to Sarepta Therapeutics and are

named inventors of patents licensed through the University of Western Australia to Sarepta

Therapeutics. As such, they are entitled to milestone and royalty payments that may be

generated from licensing agreements and are involved in ongoing collaborative research

projects with Sarepta Therapeutics.

738
739
740

References:

AM Adams, PL Harding, PL Iversen, C Coleman, S Fletcher and SD Wilton (2007).
Antisense oligonucleotide induced exon skipping and the dystrophin gene
transcript: Cocktails and chemistries. *BMC Mol Biol* **8**: 57

F Arnhold, KH Guhrs and A von Mikecz (2015). **Amyloid domains in the cell nucleus**
controlled by nucleoskeletal protein lamin b1 reveal a new pathway of mercury
neurotoxicity. *PeerJ* **3**: e754.10.7717/peerj.754

JK Bailey, W Shen, XH Liang and ST Crooke (2017). **Nucleic acid binding proteins affect the subcellular distribution of phosphorothioate antisense oligonucleotides**. *Nucleic Acids Res* **45**(18): 10649-10671.10.1093/nar/gkx709

CF Bennett, BF Baker, N Pham, E Swayze and RS Geary (2017). **Pharmacology of antisense drugs**. *Annu Rev Pharmacol Toxicol* **57**: 81-105.10.1146/annurev-pharmtox-010716-104846

CS Bond and AH Fox (2009). **Paraspeckles: Nuclear bodies built on long noncoding rna**. *J Cell Biol* **186**(5): 637-644.10.1083/jcb.200906113

M Chakravarthy, S Chen, PR Dodd and RN Veedu (2017). **Nucleic acid-based theranostics for tackling alzheimer's disease**. *Theranostics* **7**(16): 3933-3947.10.7150/thno.21529

ST Crooke, S Wang, TA Vickers, W Shen and XH Liang (2017). **Cellular uptake and trafficking of antisense oligonucleotides**. *Nat Biotechnol* **35**(3): 230-237.10.1038/nbt.3779

N Dias and CA Stein (2002). **Potential roles of antisense oligonucleotides in cancer therapy. The example of bcl-2 antisense oligonucleotides**. *Eur J Pharm Biopharm* **54**(3): 263-269

S Fletcher, K Honeyman, AM Fall, PL Harding, RD Johnsen and SD Wilton (2006). **Dystrophin expression in the mdx mouse after localised and systemic administration of a morpholino antisense oligonucleotide**. *J Gene Med* **8**(2): 207-216

U Flierl, TL Nero, B Lim, JF Arthur, Y Yao, SM Jung, E Gitz, AY Pollitt, MT Zaldivia, M Jandrot-Perrus, A Schafer, B Nieswandt, RK Andrews, MW Parker, EE Gardiner and K Peter (2015). **Phosphorothioate backbone modifications of nucleotide-based drugs are potent platelet activators**. *J Exp Med* **212**(2): 129-137.10.1084/jem.20140391

AH Fox, S Nakagawa, T Hirose and CS Bond (2018). **Paraspeckles: Where long noncoding rna meets phase separation**. *Trends Biochem Sci* **43**(2): 124-135.10.1016/j.tibs.2017.12.001

BL Gebski, CJ Mann, S Fletcher and SD Wilton (2003). **Morpholino antisense oligonucleotide induced dystrophin exon 23 skipping in mdx mouse muscle**. *Hum Mol Genet* **12**(15): 1801-1811

L Jiang, C Shao, QJ Wu, G Chen, J Zhou, B Yang, H Li, LT Gou, Y Zhang, Y Wang, GW Yeo, Y Zhou and XD Fu (2017). **Neat1 scaffolds rna-binding proteins and the microprocessor to globally enhance pri-mirna processing**. *Nat Struct Mol Biol* **24**(10): 816-824.10.1038/nsmb.3455

M Kinali, V Arechavala-Gomeza, L Feng, S Cirak, D Hunt, C Adkin, M Guglieri, E Ashton, S Abbs, P Nihoyannopoulos, ME Garralda, M Rutherford, C McCulley, L Popplewell, IR Graham, G Dickson, MJ Wood, DJ Wells, SD Wilton, R Kole, V Straub, K Bushby, C Sewry, JE Morgan and F Muntoni (2009). **Local restoration of dystrophin expression with the morpholino oligomer avi-4658 in duchenne muscular dystrophy: A single-**

799   **blind, placebo-controlled, dose-escalation, proof-of-concept study**. *Lancet Neurol*
800   **8**(10): 918-928.S1474-4422(09)70211-X [pii] 10.1016/S1474-4422(09)70211-X
801

802 M Lafarga, O Tapia, AM Romero and MT Berciano (2017). **Cajal bodies in neurons**. *RNA*
803   *Biol* **14**(6): 712-725.10.1080/15476286.2016.1231360
804

805 AK Lancaster, A Nutter-Upham, S Lindquist and OD King (2014). **Plaac: A web and**
806   **command-line application to identify proteins with prion-like amino acid**
807   **composition**. *Bioinformatics* **30**(17): 2501-2502.10.1093/bioinformatics/btu310
808

809 R Li, AR Harvey, SI Hodgetts and AH Fox (2017). **Functional dissection of neat1 using**
810   **genome editing reveals substantial localization of the neat1_1 isoform outside**
811   **paraspeckles**. *RNA* **23**(6): 872-881.10.1261/rna.059477.116
812

813 S Li, FJ Shu, Z Li, L Jaafar, S Zhao and WS Dynan (2017). **Cell-type specific role of the**
814   **rna-binding protein, nono, in the DNA double-strand break response in the mouse**
815   **testes**. *DNA Repair (Amst)* **51**: 70-78.10.1016/j.dnarep.2017.02.002
816

817 XH Liang, H Sun, W Shen and ST Crooke (2015). **Identification and characterization of**
818   **intracellular proteins that bind oligonucleotides with phosphorothioate linkages**.
819   *Nucleic Acids Res* **43**(5): 2927-2945.10.1093/nar/gkv143
820

821 F Lipi, S Chen, M Chakravarthy, S Rakesh and RN Veedu (2016). **In vitro evolution of**
822   **chemically-modified nucleic acid aptamers: Pros and cons, and comprehensive**
823   **selection strategies**. *RNA Biol* **13**(12): 1232-1245.10.1080/15476286.2016.1236173
824

825 JR Mendell, N Goemans, LP Lowes, LN Alfano, K Berry, J Shao, EM Kaye, E Mercuri, G
826   Eteplirsen Study and DMDIN Telethon Foundation (2016). **Longitudinal effect of**
827   **eteplirsen versus historical control on ambulation in duchenne muscular dystrophy**.
828   *Ann Neurol* **79**(2): 257-271.10.1002/ana.24555
829

830 JR Mendell, LR Rodino-Klapac, Z Sahenk, K Roush, L Bird, LP Lowes, L Alfano, AM
831   Gomez, S Lewis, J Kota, V Malik, K Shontz, CM Walker, KM Flanigan, M Corridore, JR
832   Kean, HD Allen, C Shilling, KR Melia, P Sazani, JB Saoud, EM Kaye and G Eteplirsen
833   Study (2013). **Eteplirsen for the treatment of duchenne muscular dystrophy**. *Ann*
834   *Neurol* **74**(5): 637-647.10.1002/ana.23982
835

836 JR Mendell, Z Sahenk and L Rodino-Klapac (2017). **Clinical trials of exon skipping in**
837   **duchenne muscular dystrophy**. *Expert Opinion on Orphan Drugs* **5**(9): 683-690.DOI:
838   10.1080/21678707.2017.1366310
839

840 C Mitrpant, P Porensky, H Zhou, L Price, F Muntoni, S Fletcher, SD Wilton and AH Burghes
841   (2013). **Improved antisense oligonucleotide design to suppress aberrant smn2 gene**
842   **transcript processing: Towards a treatment for spinal muscular atrophy**. *PLoS One*
843   **8**(4): e62114.10.1371/journal.pone.0062114
844

845 T Naganuma, S Nakagawa, A Tanigawa, YF Sasaki, N Goshima and T Hirose (2012).
846   **Alternative 3'-end processing of long noncoding rna initiates construction of nuclear**
847   **paraspeckles**. *EMBO J* **31**(20): 4020-4034.10.1038/emboj.2012.251
848

Y Nishimoto, S Nakagawa, T Hirose, HJ Okano, M Takao, S Shibata, S Suyama, K Kuwako, T Imai, S Murayama, N Suzuki and H Okano (2013). **The long non-coding rna nuclear-enriched abundant transcript 1_2 induces paraspeckle formation in the motor neuron during the early phase of amyotrophic lateral sclerosis**. *Mol Brain* **6**: 31.10.1186/1756-6606-6-31

DM Paton (2017). **Nusinersen: Antisense oligonucleotide to increase smn protein production in spinal muscular atrophy**. *Drugs Today (Barc)* **53**(6): 327-337.10.1358/dot.2017.53.6.2652413

TA Shelkovnikova, MS Kukharsky, H An, P Dimasi, S Alexeeva, O Shabir, PR Heath and VL Buchman (2018). **Protective paraspeckle hyper-assembly downstream of tdp-43 loss of function in amyotrophic lateral sclerosis**. *Mol Neurodegener* **13**(1): 30.10.1186/s13024-018-0263-7

TA Shelkovnikova, HK Robinson, C Troakes, N Ninkina and VL Buchman (2014). **Compromised paraspeckle formation as a pathogenic factor in fusopathies**. *Human Molecular Genetics* **23**(9): 2298-2312.10.1093/hmg/ddt622

L Shen, A Frazer-Abel, PR Reynolds, PC Giclas, A Chappell, MK Pangburn, H Younis and SP Henry (2014). **Mechanistic understanding for the greater sensitivity of monkeys to antisense oligonucleotide-mediated complement activation compared with humans**. *J Pharmacol Exp Ther* **351**(3): 709-717.10.1124/jpet.114.219378

W Shen, CL De Hoyos, H Sun, TA Vickers, XH Liang and ST Crooke (2018). **Acute hepatotoxicity of 2' fluoro-modified 5-10-5 gapmer phosphorothioate oligonucleotides in mice correlates with intracellular protein binding and the loss of dbhs proteins**. *Nucleic Acids Res* **46**(5): 2204-2217.10.1093/nar/gky060

W Shen, XH Liang and ST Crooke (2014). **Phosphorothioate oligonucleotides can displace neat1 rna and form nuclear paraspeckle-like structures**. *Nucleic Acids Res* **42**(13): 8648-8662.10.1093/nar/gku579

W Shen, XH Liang, H Sun and ST Crooke (2015). **2'-fluoro-modified phosphorothioate oligonucleotide can cause rapid degradation of p54nrb and psf**. *Nucleic Acids Res* **43**(9): 4569-4578.10.1093/nar/gkv298

NK Singh, NN Singh, EJ Androphy and RN Singh (2006). **Splicing of a critical exon of human survival motor neuron is regulated by a unique silencer element located in the last intron**. *Mol Cell Biol* **26**(4): 1333-1346.10.1128/MCB.26.4.1333-1346.2006

CA Stein and D Castanotto (2017). **Fda-approved oligonucleotide therapies in 2017**. *Mol Ther* **25**(5): 1069-1075.10.1016/j.ymthe.2017.03.023

P Taimen, K Pfleghaar, T Shimi, D Moller, K Ben-Harush, MR Erdos, SA Adam, H Herrmann, O Medalia, FS Collins, AE Goldman and RD Goldman (2009). **A progeria mutation reveals functions for lamin a in nuclear assembly, architecture, and chromosome organization**. *Proc Natl Acad Sci U S A* **106**(49): 20788-20793.10.1073/pnas.0911895106

O Tapia, JO Narcis, J Riancho, O Tarabal, L Piedrafita, J Caldero, MT Berciano and M Lafarga (2017). **Cellular bases of the rna metabolism dysfunction in motor neurons of a murine model of spinal muscular atrophy: Role of cajal bodies and the nucleolus**. *Neurobiol Dis* **108**: 83-99.10.1016/j.nbd.2017.08.004

LJA Toonen, J Casaca-Carreira, M Pellise-Tintore, H Mei, Y Temel, A Jahanshahi and WMC van Roon-Mom (2018). **Intracerebroventricular administration of a 2'-o-methyl phosphorothioate antisense oligonucleotide results in activation of the innate immune system in mouse brain**. *Nucleic Acid Ther* **28**(2): 63-73.10.1089/nat.2017.0705

B Tri Le, RN Veedu, S Fletcher and SD Wilton (2016). **Antisense oligonucleotide development for the treatment of muscular dystrophies**. *Expert Opinion on Orphan Drugs* **4**(2): 139-152.10.1517/21678707.2016.1122517

RN Veedu (2015). **Medicinal chemistry of aptamers**. *Curr Top Med Chem* **15**(12): 1065

Q Wang, IA Sawyer, MH Sung, D Sturgill, SP Shevtsov, G Pegoraro, O Hakim, S Baek, GL Hager and M Dundr (2016). **Cajal bodies are linked to genome conformation**. *Nat Commun* **7**: 10966.10.1038/ncomms10966

SD Wilton, RN Veedu and S Fletcher (2015). **The emperor's new dystrophin: Finding sense in the noise**. *Trends Mol Med* **21**(7): 417-426.10.1016/j.molmed.2015.04.006

J Winkler, M Stessl, J Amartey and CR Noe (2010). **Off-target effects related to the phosphorothioate modification of nucleic acids**. *ChemMedChem* **5**(8): 1344-1352.10.1002/cmdc.201000156

E Wong and T Goldberg (2014). **Mipomersen (kynamro): A novel antisense oligonucleotide inhibitor for the management of homozygous familial hypercholesterolemia**. *P T* **39**(2): 119-122

T Yamazaki and T Hirose (2015). **The building process of the functional paraspeckle with long non-coding rnas**. *Front Biosci (Elite Ed)* **7**: 1-41

JJ Yerbury, L Ooi, A Dillin, DN Saunders, DM Hatters, PM Beart, NR Cashman, MR Wilson and H Ecroyd (2016). **Walking the tightrope: Proteostasis and neurodegenerative disease**. *Journal of Neurochemistry* **137**(4): 489-505.10.1111/jnc.13575