Intracellular *Burkholderia* Symbionts Induce Extracellular Secondary Infections;

Driving Diverse Host Outcomes that Vary by Genotype and Environment

Niloufar Khojandi [1,2], Tamara S. Haselkorn [3], Madison N. Eschbach [1], Rana A.

Naser [1], Susanne DiSalvo [1]

[1] Department of Biological Sciences, Southern Illinois University Edwardsville,

Edwardsville, IL 62026

[2] Department of Molecular Microbiology and Immunology, St. Louis University, St. Louis,

MO 63104 (current address)

[3] Department of Biology, University of Central Arkansas, 201 Donaghey Avenue,

Conway, AR 72035

* Correspondence:

Susanne DiSalvo

sdisalv@siue.edu

Keywords: Burkholderia, Amoebae, Dictyostelium, Symbiosis, Infection, Host,

Symbiont

Abstract

Symbiotic associations impact and are impacted by their surrounding ecosystem. The association between *Burkholderia* bacteria and the soil amoeba *Dictyostelium discoideum* is a tractable model to unravel the biology underlying symbiont-endowed phenotypes and their impacts. Several *Burkholderia* species stably associate with *D. discoideum* and typically reduce host fitness in food-rich environments while increasing fitness in food-scarce environments. *Burkholderia* symbionts are themselves inedible to their hosts but induce co-infections with secondary bacteria that can serve as a food source. Thus, *Burkholderia* hosts are "farmers" that carry food bacteria to new environments, providing a benefit when food is scarce. We examined the ability of specific *Burkholderia* genotypes to induce secondary co-infections and assessed host fitness under a range of co-infection conditions and environmental contexts. Although all *Burkholderia* symbionts intracellularly infected *Dictyostelium*, we found that co-infections are predominantly extracellular, suggesting that farming benefits are derived from extracellular infection of host structures. Furthermore, levels of secondary infection are linked to conditional host fitness; *B. agricolaris* infected hosts have the highest level of co-infection and have the highest fitness in food scarce environments. This study illuminates the phenomenon of co-infection induction across *Dictyostelium* associated *Burkholderia* species and exemplifies the contextual complexity of these associations.

Introduction

Symbiotic interactions can alter the fitness and evolutionary trajectory of both partners

(1–4). Clearly detrimental or mutualistic associations have been investigated for obvious

reasons: to eliminate infectious disease, boost health, and restore ecosystems.

However, many symbiotic associations evade simple characterization and related

mechanisms can underlie opposing outcomes (5,6). Invasion and replication strategies

employed by mutualists and pathogens often resemble each other, while genotypes and

external factors modify subsequent outcomes (7). Genotype pairing determines the

outcome of plant-mycorrhizae interactions (8) and amplification of a genomic region in a

normally beneficial *Wolbachia* symbiont leads to over-replication at the hosts expense

(9). Light mediates pathogenicity of a fungal plant endosymbiont (10), temperature

affects reproductive fitness of aphids hosting *Buchnera* (11), and parasitoid pressure

determines whether *Hamiltonella defensa* is beneficial to host aphids (12). These

examples demonstrate that even canonically beneficial or detrimental associations may

produce alternative effects in alternative contexts (4,13–17).

Eukaryotic microbes, such as amoebae, are attractive models for exploring eukaryote-

prokaryote interactions. Amoebae are ubiquitous and efficient phagocytic predators of

bacterial prey, making them important shapers of the microbial community (18). This

pressures prey microbes to evolve virulence strategies that enable evasion of

phagocytosis or subsequent digestion (19). Amoebae are thereby potential training

grounds and environmental reservoirs for bacterial pathogens. Amoebae phagocytosis

also enables bacteria to gain easy access to an attractive intracellular niche, bypassing

the requirement for evolving specialized cell-entry mechanisms. After invasion, bacteria

can be retained in an environmentally resistant cyst or spore (20). A number of bacterial

pathogens, such as *Francisella tularenis* and *Legionellae pneumophila* (21), are

harbored in different species of amoebae and there is a growing list of recently identified

amoebae symbionts (22,23).

The social amoebae *Dictyostelium discoideum* has been appreciated as a model host

for studying bacterial pathogens for some time (24,25). Recently, work with wild isolates

has emphasized its power for exploring naturally occurring microbial symbioses (26,27).

As a social amoeba *Dictyostelium* exhibits a unique life cycle, transitioning between

single- and multi-cellular forms. Under favorable conditions, it lives as a unicellular

amoeba, consuming bacteria and dividing by binary fission. When bacterial food is

depleted, amoebae secrete cAMP, which triggers the transition to multi-cellularity.

During this phase, amoebae aggregate to form a multicellular slug that seeks out a

location for fruiting body formation (such as the soil surface). Fruiting bodies are

comprised of a spherical sorus containing hardy spore cells resting atop a long stalk of

dead cells. This positioning of spore cells likely aids in their dispersal (28). Once

dispersed, spores germinate, and the cycle continues.

D. discoideum grown with a variety of bacterial food traditionally form germ-free sori,

clearing residual bacteria from the multicellular state during development. Microbial

clearance is aided by immune-like sentinel cells which engulf debris and slough off the

migrating slug (29,30). However, approximately one third of wild *D. discoideum* isolates

are naturally and stably colonized by *Burkholderia* bacteria (31,32). *Burkholderia* can be

easily eliminated from host populations with antibiotic treatment and new associations

can be readily initiated through co-culture. These *Burkholderia* symbionts establish

intracellular infections which persist through host development, resulting in sori

containing both extracellular and intracellular bacteria (32). *Burkholderia* symbionts

thereby remain associated with host populations during spore dispersal and can be

acquired through vertical and horizontal transmission routes. This mixed mode of

transmission has interesting implications for the fitness consequences and evolutionary

trajectory of the symbiosis.

Burkholderia symbionts of *D. discoideum* are members of the plant beneficial

environmental group within the *Burkholderia* genus (33). Symbiont strains are

genetically diverse, belonging to three species arising from two independent lineages:

B. agricolaris, B. hayleyella, and *B. bonniea* (34). *Burkholderia* differentially impacts

host fitness according to host-symbiont genotype combinations and environmental

context (31,32,35). Symbionts generally reduce host fitness in food rich conditions but

enhance fitness in food scarce environments (31,32,35). Fitness benefits are attributed

to retention of bacteria within host spores, allowing them to reseed new environments

with bacterial food. This trait is called farming and *Burkholderia* infected hosts are thus

referred to as "farmers". *Burkholderia* symbionts themselves are poor food sources for

their hosts (31,32). However, *Burkholderia* infection appears to increase host

susceptibility to secondary bacterial infection, promoting the formation of a mini-

microbiome. It is these secondary bacteria that can serve as an amoebae food source

and thereby provide the farming benefit.

Given the importance of secondary infection in farming, we sought to explore the

underlying dynamics of this interaction. While a commonly used lab food strain, *K.*

pneumoniae, can be identified as an occasional co-infecting partner, co-infection

dynamics might vary depending on particular bacterial pairings (32). Thus, food bacteria

identity is an important environmental context that may affect outcomes. Furthermore,

the three different *Burkholderia* symbiont species have divergent evolutionary histories

of association with *D. discoideum*. While they have converged on the farming

phenotype, the effects and underlying mechanisms of infection may differ across

Burkholderia species (Haselkorn et al. in review).

Here, we reveal the density and location of secondary co-infections induced by each

Burkholderia species with a collection of secondary bacteria. Next, we clarified the

downstream benefits of *Burkholderia* infection in varied food availability contexts and

link these with host outcomes according to co-infection induction. Specifically, we

analyzed co-infection patterns and host outcomes with a variety of secondary bacteria

including: laboratory food *Klebsiella pneumoniae, Rhizobium* and *Serratia* isolates that

naturally co-occur with *D. discoideum*, and *Agrobacterium tumefaciens* and

Pseudomonas aeruginosa as pathogens that *D. discoideum* may encounter in nature.

We found that all *Burkholderia* symbionts induce some degree of secondary infection in

host sori but the density and location of secondary infections is dependent on

Burkholderia genotype and secondary bacterial identity. Contrary to previous inference,

secondary infections are predominantly extracellular with intracellular co-infections only

readily visualized in *B. agricolaris* infected spores. Overall, *B. agricolaris* induces the

highest density of combined co-infection resulting in a higher fitness benefit hosts in

food scarce environments. *B. bonniea* and *B. hayleyella* induce lower levels of

secondary co-infection but only *B. bonniea* provides significant host benefits under

specific dispersal conditions. This work illuminates the interplay between symbiont

genotypes and environmental context in mediating the expression and consequences of

novel symbiont endowed phenotypes.

Materials and Methods

<u>Bacterial Strains and Culturing</u>

Burkholderia strains were isolated in the Queller Strassmann lab from *D. discoideum*

hosts. *Klebsiella pneumoniae*, *Rhizobium* and *Serratia* originate from the Queller

Strassmann lab with *Rhizobium* and *Serratia* isolated from wild *Dictyostelium* fruiting

bodies grown directly from Virginia Mountain Lake Biological Station soil and 16s rRNA

gene sequenced with universal primers for putative identification (genbank accession

numbers MH997560 and MH997561). *Pseudomonas aeruginosa* PAO1-GFP (derived

from ATCC15692) was provided by R. Fred Inglis (36). *Agrobacterium tumefaciens*

(A136) was provided by Daniel Gage (reclassified as *Rhizobium radiobacter* 51350 in

ATCC). We used GFP labeled secondary bacteria for all experiments, with the

exception of *K. pneumoniae*, which is unlabeled when mixed with other bacteria. We

grew all bacteria on SM/5 medium (Formedium: 2g Peptone, 0.2g yeast extract, 2g

glucose, 1.9g KH_2PO_4, 1.3g $K_2HPO_4.3H_2O$, 0.49g MgO_4.anhydrous, 17g agar/liter) at

room temperature. To prepare bacteria for culturing *Dictyostelium*, we suspended

bacterial colonies from SM/5 medium into KK2 (2.2g KH2PO4 monobasic and 0.7g

K2HPO4 dibasic/liter) and set to an OD_{600nm} of 1.5. For *K. pneumoniae*/secondary

bacterial mixtures, we combined bacterial suspensions equally by volume. For

Burkholderia infections, we added 5% by volume *Burkholderia*-RFP to indicated

bacterial mixtures.

Construction of Fluorescent Bacterial Strains

We generated RFP labeled *Burkholderia* by triparental mating with *E. coli* helper strain

E1354 (pTNS3-asdEc) and *E. coli* donor strain E2072 (pmini-Tn7-gat-P1-rfp) and

confirmed identity of RFP conjugants using a *Burkholderia* specific PCR as previously

described (32,37,38). We GFP labeled *Rhizobium, Serratia*, *A. tumefaciens,* and *K.*

pneumoniae through triparental mating with *E. coli* donor WM3064 (pmini-Tn7-KS-GFP)

and *E. coli* helper E1354 (pUXBF13) as previously described (39) and confirmed

identity of GFP positive conjugants through 16s rRNA gene sequencing. *P. aeruginosa*-

GFP was described previously (36).

Dictyostelium Culture Conditions

We used *D. discoideum* clone QS864 (naturally symbiont free) for all experiments.

Cultures were initiated by plating spores on SM/5 medium with *K. pneumoniae* and

incubating under lights at room temperature until fruiting bodies developed (4-7 days).

For experiments, 10^5 spores were harvested from developed sori and plated with 200µL

of the appropriate bacterial mixtures. For all experiments (unless otherwise indicated)

we analyzed sori 5 days after plating.

For co-infection assays we plated uninfected spores on bacterial mixtures with

Burkholderia, uninfected controls were plated without *Burkholderia.* To compare spore

productivity under food variable conditions, we harvested sori from indicated co-

infection conditions and plated 10^5 spores onto SM/5 with *K. pneumoniae* at an OD_{600} of

1.5 for food-rich conditions or with heat-killed (30 min at $80^{\circ C}$) *K. pneumoniae* at an

OD_{600} of 6 for food scarce conditions.

Spore Production Assays

To harvest total spores, we flooded each plate with 5–10 mL KK2 + 0.1% Nonidet P-40

alternative and collected the entire surface contents into 15-mL Falcon tubes. We then

diluted samples in KK2 and counted spores on a hemocytometer. At least five replicates

were analyzed for each treatment.

Confocal Microscopy

We imaged spores by staining with 1% calcofluor in KK2, placing on glass bottom

culture dishes (Electron Microscopy Sciences) and overlaying with 2% agarose. We

imaged samples on an Olympus Fluoview FV1000 confocal microscope using Plan Apo

Oil 1.4NA 60X objective. Z-sections were taken every 0.5 microns at 1024 resolution.

Calcofluor was visualized with DAPI, GFP with FITC, and RFP with Cy3 then

pseudocolored grey, green, and red respectively. We imaged at least three individual

replicates and counted more than 30 spores for each.

Colony forming unit quantification

To quantify secondary bacteria, we harvested sori grown from the indicated co-culture

conditions from 6- or 14-day incubations. We suspended individual sori in KK2+0.05%

Nonidet P-40 alternative, counted spores on a hemocytometer, plated serial dilutions on

SM/5 medium and incubated plates at room temperature until colony formation (~2

days), and counted GFP colonies using a safe-light imaging system. We performed 3 or

more independent replicates for each treatment.

Streak test

Our streak test assay was initiated from the indicated co-culture conditions by touching

individual sori with sterile pipette tips and transferring them to SM/5 plates along a

~1inch streak. We incubated plates face up under lights at room temperature and

examined them 5 days (or 2 weeks) after streaking. We determined the percentage of

streaks with bacterial growth, percentage of bacterial positive streaks with fruiting

bodies, and number of fruiting bodies in positive streaks. Streaks were photographed on

a Cannon Eos7D with a macro-lens. Six sori were streaked for each replicate for at

least 4 individual replicates per condition.

Statistical analysis

We analyzed all data using R (version 3.3.1). For normally distributed data we

determined significance using a standard one-way analysis of variance (ANOVA) and a

post hoc Tukey HSD test. For non-normally distributed data we performed a Kruskall-

wallis test and *post hoc* analysis with a Dunn test using the dunnTest function in the

FSA package. We used *Burkholderia* status as fixed effects for all conditions.

Results

Burkholderia and Secondary Bacterial Combinations

To investigate the induction of secondary infection by *Burkholderia* symbionts, we

cultured an uninfected natural isolate of *D. discoideum* with different *Burkholderia*-RFP

and secondary bacteria-GFP combinations. We began with three *Burkholderia* strains:

Ba.70, Bh.11, and Bb.859, each representing one *D. discoideum* symbiont species *B.*

agricolaris, B. hayleyella, and *B. bonniea* (34). Secondary bacteria consisted of a

Klebsiella pneumoniae strain, soil isolated *Rhizobium* and *Serratia*, and lab

Agrobacterium tumefaciens and *Pseudomonas aeruginosa* strains. We chose these

representatives because: 1) *K. pneumoniae* is used widely as a lab food source for *D.*

discoideum and serves as a good starting point for experimental conditions. 2) The

Rhizobium and *Serratia* strains where co-isolated with *D. discoideum* and are thereby

ecologically relevant potential co-associates. 3) *A. tumefaciens,* in addition to its use in

plant molecular biology, is an important soil dwelling plant pathogen. As such, amoebae

may interact with *A. tumefaciens* in the environment and this could subsequently impact

the surrounding ecosystem. 4) *P. aeruginosa* is an important opportunistic human

pathogen whose association with other bacterial species in biofilms (such as pathogenic

Burkholderia cenocepacia) influences infection outcomes (40,41). Adding

Pseudomonas to the *Burkholderia-Dictyostelium* system provides a novel approach to

explore microbial interactions and virulence.

Host Outcomes Differ According to *Burkholderia* and Secondary Bacteria Conditions.

First, we examined host fitness when amoebae were co-cultured with *Burkholderia* and

secondary bacteria. We determined total spore productivity of host amoebae after one

social cycle on each labeled secondary bacterium, either alone or in a 50% mixture with

K. pneumoniae. Five percent by volume of *Burkholderia*-RFP was included to establish

infections (Figure 1). *D. discoideum* was unable to develop on any conditions where *P.*

aeruginosa was the only food source suggesting that this strain was toxic and/or

inedible for amoebae. All other conditions supported fruiting body development, but

spore productivity varied across conditions (Figure 1). In line with previous studies,

Burkholderia species differentially impact spore productivity on *K. pneumoniae*.

Typically, *B. hayleyella* was the most detrimental for host fitness with *B. agricolaris* and

B. bonniea being neutral or moderately detrimental. However, these patterns and the

degree by which symbiont altered host fitness varied across culture conditions (Figure 1

and Table 1). These results highlight the variability of fitness outcomes caused by

distinct *Burkholderia* symbionts and suggest that surrounding bacterial communities

also impact fitness outcomes.

<u>Intracellular Co-infection is Rare and Depends on *Burkholderia* and Secondary Bacterial</u>

<u>Combinations.</u>

To investigate the induction of secondary infections we imaged *D. discoideum* sori

contents after development on *Burkholderia* and secondary bacteria. We used 50/50 *K.*

*pneumoniae/*secondary bacteria-GFP conditions as they resulted in better amoebae

development than secondary bacteria-only conditions. We also imaged sori grown from

K. pneumoniae-GFP. Importantly, we do not detect any secondary bacteria in sori in the

absence of *Burkholderia* (Figure 2). Thus, these bacteria are not capable of infecting *D.*

discoideum on their own. In contrast, we can detect secondary-GFP cells in sori from

amoebae co-exposed to *Burkholderia* (Figure 3). To determine their prevalence in host

spore populations, we quantified the percent of spores intracellularly infected with

Burkholderia-RFP and with secondary bacteria-GFP. First, the percent of spores

infected with each *Burkholderia* species significantly differs (χ^2=44.02, df=2, p<0.001).

On average across conditions, *B. hayleyella* infects the most (89.2%), *B. bonniea*

infects an intermediate (68.6%), and *B. agricolaris* infects the fewest (33%) percent of

spores. However, we only readily observe intracellular secondary co-infections in *B.*

agricolaris host spores (average of 5.5% across conditions). Only rarely, if ever, do we

observe intracellular GFP in *B. hayleyella* and *B. bonniea* hosts (0.01% and 0.05%

respectively) (Figure 3). We did not observe intracellular GFP in the absence of

intracellular RFP, suggesting that secondary bacteria are only retained in *Burkholderia*

co-infected spores.

The secondary bacterium also plays a role in the prevalence of intracellular infections

for both *Burkholderia* and secondary bacteria. For instance, slightly fewer spores are

infected with *B. bonniea* when cultured with *P. aeruginosa*-GFP (29%) than with all

other bacteria (85.9-75.6%) (χ^2=10.843, df=4, p=0.028). On the flip side, significantly

more spores are infected with *B. hayleyella* when cultured with *P. aeruginosa*-GFP

(98.9%) than with all other bacteria (81.2-91%) (χ^2=10.217, df=4, p=0.036). For *B.*

agricolaris hosts, the degree of secondary co-infections significantly varied by

secondary bacteria (χ^2=15.019, df=4, p=0.004). *K. pneumoniae*-GFP is localized in only

0.2% of total spores while *Rhizobium*-GFP was localized in 14.9%. When secondary

bacterial infections are considered as a percentage of spores co-infected with *B.*

agricolaris, *Rhizobium*-GFP is co-localized in almost half of total infected spores. This

suggests that should *B. agricolaris* infection levels increase due to a higher infectious

dose or conditions that promote higher infection titers, secondary infections may

correspondingly increase.

Burkholderia Symbionts Induce Extracellular Secondary Infections

Although we found minimal intracellular co-infections in most conditions, the farming

phenotype may instead by explained by extracellular secondary infections. To get an

initial indication of extracellular co-infections, we determined the percent of confocal

images in which any extracellular GFP could be visualized (Figure 4a). We found that all

Burkholderia symbionts induced at least some level of extracellular co-infections, as we

could visualize external GFP in confocal images for each condition. Similar to our

observations for intracellular co-infections, extracellular secondary bacteria appeared

most frequently in *B. agricolaris* host sori (Figure 4a).

To quantify overall secondary co-infection, we counted GFP colony-forming units per

sori for *K. pneumoniae*-GFP$^{100\%}$ and *Rhizobium*-GFP$^{50\%}$ conditions. After six days of co-

culturing amoebae we counted the number of GFP positive colonies that developed

from individual sori (Figure 4b). No bacterial colonies were recovered when we plated

sori from fruiting bodies grown without *Burkholderia*, again indicating that these bacteria

do not by themselves infect *D. discoideum*. We found that *B. agricolaris* induces the

highest level of secondary infection, with *B. agricolaris* host sori colonized by an

average of 1.29x10^5 *K. pneumoniae*-GFP cfu's and 1.17x10^6 *Rhizobium*-GFP cfu's

(Figure 4b). We recover notable, albeit far fewer, secondary bacterial colonies from *B.*

hayleyella and *B. bonniea* hosts. *B. hayleyella* sori host an average of 206 and 169, and

B. bonniea an average of 153 and 276, *K. pneumoniae*-GFP and *Rhizobium*-GFP cfu's,

respectively.

To explore whether secondary bacteria could further amplify within fruiting bodies over

time, we also quantified R*hizobium*-GFP colony-forming units 14 days after plating. We

found that cfu's did not increase for *B. agricolaris* hosts, but dramatically increased for

B. hayleyella and *B. bonniea* hosts, which produced 2.87 x10^5 and 8.37 x10^5 GFP cfu's,

respectively. This brought the number of cfu's in all *Burkholderia* infected sori up to fairly

similar levels, perhaps representing a peak carrying capacity. However, we noticed that

the number of spores per sori for *B. hayleyella* and *B. bonniea* infected hosts appeared

to decrease over time (not shown). Thus, replication of secondary bacteria within these

sori could be damaging to spores, counter-acting potential benefits of hosting more food

bacteria.

Burkholderia Symbionts Benefit Hosts in Food Scarce Conditions in Relation to Co-

infection Induction.

Farmers have been shown to have an advantage when dispersed to food scarce

environments (31,32). This is attributed to the induction of secondary bacterial food

carriage, enabling host spores to reseed new environments with edible bacteria.

However, this benefit has previously only been measured as an average fitness

outcome across hosts infected with genotypically diverse *Burkholderia* symbionts (32).

Whether or how specific *Burkholderia* genotypes correspond with this beneficial

outcome remained unknown. Given our results demonstrating that *B. agricolaris*

induces the highest level of secondary co-infections, we speculated that *B. agricolaris*

hosts would have the highest reproductive fitness after dispersal to food scarce

environments.

To compare host fitness under different food availability conditions, we first plated

uninfected spores with *Burkholderia* and secondary bacteria (*K. pneumoniae*-GFP$^{100\%}$

and *Rhizobium*-GFP$^{50\%}$) under the same conditions employed previously. After five

days of incubation we harvested developed sori and transferred 10^5 spores with rich

(live *K. pneumoniae*-GFP) or scarce (heat killed *K. pneumoniae*) food on nutrient

medium. After five days of incubation in these conditions, we measured total spore

productivity (Figure 5a). For food rich conditions, we again found significant differences

according to amoebae infection status (F=7.41, df=3, p 0.0015 and χ^2=13.95, df=3,

p=0.0029 for *K. pneumoniae*-GFP$^{100\%}$ and *Rhizobium*-GFP$^{50\%}$ conditions respectively).

However, in this experiment only *B. hayleyella* hosts produced significantly fewer spores

than the uninfected control (Figure 5a). For food scarce conditions, spore productivity

was also significantly different according to *Burkholderia* infection status (χ^2=11.87 and

20.616, df=3, p=0.0078 and <0.001 for *K. pneumoniae*-GFP$^{100\%}$ and *Rhizobium*-GFP$^{50\%}$

conditions, respectively). Here, *B. agricolaris* hosts had the highest spore productivity

for both secondary conditions (Figure 5a). *B. bonniea* also resulted in slightly higher, but

not significantly different, spore productivity compared to the uninfected control (Figure

5a). Thus, *B. agricolaris* infections endow a benefit for their amoeba host when

dispersed to food scarce environments.

For the above assay, we evenly distributed spores on plates. If secondary bacteria are

less numerous in host sori (as for *B. hayleyella* and *B. bonniea*), they might be spread

too far from germinating spores to access and thus benefit from. Further, this assay

might not best simulate spore dispersal in nature, where spores might be deposited in

smaller denser patches by passing soil inhabitants. Therefore, we examined host fitness

in food scarce conditions using a "streak" dispersal strategy. Here, we deposited

individual sori from fruiting bodes grown on *K. pneumoniae*-GFP$^{100\%}$ or *Rhizobium*-

GFP$^{50\%}$ in small patches (~1-inch streaks) on nutrient medium. After a week of

incubation, we measured the percent of fruiting body positive streaks and the number of

fruiting bodies per each streak (Figure 5b-d).

Streaks from uninfected control sori did not produce bacterial colonies nor fruiting

bodies. Over 95% of *Burkholderia* infected sori produced streaks with bacteria,

however, the percent of streaks with fruiting bodies growing from these bacterial

colonies significantly varies across *Burkholderia* species (χ^2=13.728 and 12.127, df=2,

p=0.001 and 0.0023 for *K. pneumoniae*-GFP$^{100\%}$ and *Rhizobium*-GFP$^{50\%}$ conditions

respectively). *Burkholderia* species also significantly influences the number of fruiting

bodies per streak (χ^2=14 and 11.24, df=2, p<0.001 and 0.0036 for *K. pneumoniae*-

GFP$^{100\%}$ and *Rhizobium*-GFP$^{50\%}$ conditions respectively). *B. agricolaris* infected sori

generated significantly more fruiting bodies than *B. hayleyella* hosts from both

conditions and *B. bonniea* hosts from *K. pneumoniae*-GFP$^{100\%}$ conditions. The number

of fruiting bodies also increased the longer streak plates were left to incubate (Figure 6).

Since fruiting bodies often developed from *B. bonniea* host sori, this suggests they gain

better access to food under this dispersal strategy.

We rarely witnessed fruiting bodies from *B. hayleyella* host sori streaks, providing an

interesting counterpart to *B. bonniea*. Both symbionts induce similar densities of co-

infection (Figure 4) yet differ in downstream benefits (Figure 5). This may be explained

by the relative detriment each species exerts on its host. *B. hayleyella* reduces host

fitness compared to *B. bonniea*, whereas *B. bonniea* hosts are often indistinguishable

from uninfected counterparts (Figure 1 and 5). Thus, *B. hayleyella* toxicity may inhibit

host development despite food availability.

<u>Co-infections and Conditional Benefits are Consistent Across *Burkholderia* Species</u>

<u>Members.</u>

Our representative *Burkholderia* symbiont species significantly differed in their induction

of co-infection and host impacts. We next asked whether these phenotypes were similar

across strains of the same *Burkholderia* species. We imaged (Figure 7) and streak

tested (Figure 8) host sori for additional *Burkholderia*-RFP strains after growth on

Rhizobium-GFP$^{50\%}$. We again found noticeable, but low, levels of intracellular

Rhizobium in *B. agricolaris* infected spores, with co-infection rare or absent in *B.*

hayleyella and *B. bonniea* infected spores (Figure 5). The percent of spores infected by

Burkholderia was significantly different depending on genotype (χ^2=22.65, df=5, and

P<0.001), with *B. agricolaris* strains infecting fewer spores than *B. hayleyella* and *B.*

bonniea (Figure 5b). Despite low levels of intracellular *Rhizobium* co-infection, we

again frequently observed extracellular GFP in sori (Figure 7c).

Next, we investigated the benefits of infection by these strains in food scarce

environments (Figure 8). We found that *Burkholderia* genotype significantly influences

both the percentage of sori that generate fruiting bodies and the number of fruiting

bodies in positive streaks (χ^2=12.127, df=2, p=0.002 and χ^2=11.24, df=2, p=0.003,

respectively). In accordance with our previous pattern *B. agricolaris* infections result in

high, *B. bonniea* intermediate, and *B. hayleyella* low, levels of fruiting body production

(Figure 8). These results demonstrate that species members similarly induce co-

infections and result in similar subsequent host benefits.

Discussion:

Since elucidating the link between farming and *Burkholderia,* it has been assumed that

all *Burkholderia* symbionts allow for secondary co-infections through an intracellular co-

infection event. Intracellular co-infections of *B. agricolaris* with *K. pneumoniae* had

previously been identified (32). Co-phagocytosis with *Burkholderia* and subsequent

inhibition of phagocytic digestion was posited as a parsimonious mechanistic

explanation for the phenomenon. *Burkholderia* can be visualized within intracellular

vacuoles that appear similar to phagosomes (35) but how *Burkholderia* invades and

survive within amoebae is not currently resolved. Nonetheless, in this scenario

secondary bacteria must first be liberated from infected amoebae or spores (via

regurgitation or host cell lysis) so that surrounding amoebae may reap the benefits of

farming. Here, we found that co-intracellular infection is actually quite rare, occurring

with some frequency for only *B. agricolaris* strains, and occurring differentially across

secondary bacterial species. In contrast, all *Burkholderia* symbiont species generate

extracellular secondary bacterial infection that can be visualized for all secondary

species. The induction of overall secondary bacterial infection and its corresponding

downstream benefits is significantly different across symbionts. *B. agricolaris* strains

induce the highest levels of secondary infection (both intracellularly and overall) with *B.*

hayleyella and *B. bonniea* strains generating exclusively extracellular infections at

similarly low densities. This suggests that *Burkholderia* symbionts could induce

secondary infections by multiple mechanisms that qualitatively and quantitatively differ

between genotypes.

Overall, these results indicate that the predominant route by which different

Burkholderia symbionts induce farming leans more towards extracellular bacterial

carriage than intracellular co-infections. Susceptibility to secondary infections could be

due to *Burkholderia* symbionts compromising the primitive immune system of their

multicellular hosts. Sentinel cells serve as immune-like cells in multicellular slugs by

trapping unwanted cargo through direct phagocytosis and/or neutralization by DNA net

formation (29,30). When sentinel cells have accumulated cargo, they drop out the slug,

thereby cleansing it of potential toxic entities (29). A gene deletion that reduces sentinel

cells leads to retention of secondary bacteria through the slug stage and into the sorus

(29). *Burkholderia* host slugs have fewer sentinel cells than uninfected counterparts and

this defect goes away when hosts are cured of their *Burkholderia* symbiont via antibiotic

treatment (42). Thus, the induction of secondary infections could be an indirect

consequence of *Burkholderia* symbiosis resulting in sentinel cell reduction. This

scenario may be comparable to the phenomenon of secondary infections in mammalian

systems whereby primary infectious agents compromise the immune system of their

hosts. Despite this, we cannot rule out the possibility that extracellular secondary

infection originates from intracellular co-infection. It is possible that co-infected cells are

more susceptible to lysis, rupturing and spewing their secondary bacterial passengers

into the extracellular matrix. In either situation, secondary bacteria might then amplify

within sori.

Intracellular co-infections are most frequent with *B. agricolaris* and the soil *Rhizobium*

strain. This might reflect an ecologically relevant association between these species in

nature. *Burkholderia* and *Rhizobium* are both ubiquitous in soil and contain several

important symbiont species which have been found co-colonizing the same hosts (43–

46). Predation by amoebae in soil and aquatic systems shapes microbial community

assembly and overall food webs (47). Given the likely co-occurrence of soil amoebae

with *Burkholderia* and other soil microbes it's tempting to speculate on how these

multipartite interactions influence overall microbial communities and higher trophic

levels. Here, we show that amoebae co-disperse *Burkholderia* symbionts and

secondary bacterial hitchhikers to new environments. Thus, the impact of amoebae on

their surrounding microbial network can go well beyond predator-prey dynamics. Finally,

our observation of *Burkholderia* and *P. aeruginosa* co-infection amplifies the concern

that soil amoebae can serve as reservoirs for bacterial pathogens. These results

suggest that *Burkholderia* symbionts can increase the suite of potential pathogenic

partners hosted by amoebae.

Burkholderia-fungal associations have been well recognized for their importance in the

soil ecosystem and for their bio-restoration potential (48–50). There are compelling

parallels between *Burkholderia-Dictyostelium* and *Burkholderia-fungal* associations.

Some *Burkholderia* (notably *B. terrae*) are capable of adhering to and migrating with

growing fungal hyphae through soil (51). Similar to our system, some of these fungal

associates assist in the co-migration of other (non-migrating) bacteria (52). Several

mechanisms have been proposed to underlie these interactions, such as direct receptor

binding and indirect biofilm co-aggregation (53). *B. terrae* extracellularly colonizes

fungal hyphae but many other *Burkholderia* symbionts of diverse hots persist

intracellularly (13,54). A particularly interesting example is the *Rhizopus microsporus*

endosymbiont *B. rhizoxinica* fungi, which produces the rice seedling blight toxin (55).

Recently, secretion systems have been shown to be important for the active invasion of

B. rhizoxinica across the fungal cell wall and into the host cytoplasm (54). Secretion

systems have also been implicated in *B. psuedomallei* infections (56,57). However

many plant mutualistic *Burkholderia* species, which are closer relatives to *Burkholderia*

symbionts of *Dictyostelium,* appear to lack some of these systems (56–58). The

hypothesized portal of entry into *Dictyostelium* is via phagocytosis, which could

circumvent the need for invasion specific mechanisms. Overall, *Burkholderia* symbionts

of other hosts can help inform our understanding of the *Burkholderia*-amoebae

symbiosis and vice versa.

Biofilm formation is particularly intriguing to consider as a mechanistic explanation of

secondary infection. *Burkholderia* adherence to secondary bacteria would increase the

likelihood of co-phagocytosis or extracellular co-colonization. Different adhesive

capacities of *Burkholderia* and secondary species could explain differences in the extent

of secondary infections across bacterial combinations. Indeed, co-aggregation could be

a simple explanation for the observation of high levels of co-infection with *B. agricolaris*

and *Rhizobium*. Interestingly, recent work implicates *Dictyostelium* lectins in the farming

phenomenon, higher lectin expression was detected in farmer *D. discoideum* clones

and addition of endogenous lectins induced bacterial carriage (59). Although this work

did not consider the presence or impact of *Burkholderia*, we think *Burkholderia*

symbionts play a key role. *Burkholderia* could induce farming via induction of lectin

expression in amoebae or more simplistically, *Burkholderia* lectins may mediate

adherence of secondary bacteria. Indeed, lectin expression by *B. cenocepacia* is an

important component of biofilm formation and lectin aids in adherence of *B. cepacia* to

host tissues (40,60). Future exploration into lectin expression and adhesion

mechanisms will be helpful for clarifying these themes.

In addition to elucidating the phenomenon of secondary infections our results exemplify

the context dependency of symbiotic outcomes in this system. We found that the costs

and benefits of this symbiosis can be modified by different bacterial conditions and

spore dispersal processes. The nature and extent of farming induction by *Burkholderia*

symbionts differs across symbiont species and so do their corresponding contextual

fitness outcomes. Ultimately, further research into the mechanisms, consequences, and

ecological framework of the *Burkholderia-Dictyostelium* symbiosis will help illuminate

microbial interaction dynamics relevant to infection biology and microbial ecology.

Acknowledgements

We thank Kyle Skottke for fruitful discussions and manuscript review, Joan Strassmann

and David Queller for initial guidance in the system, and all members of the DiSalvo lab

at SIUE, particularly Jacob Miller for help with experimental discussions.

Conflict of interest We declare no conflict of interests.

Author Contributions Statement

NK and SD designed the study. NK and SD performed experiments with assistance

from ME, RN, and TH. SD and TH wrote the manuscript.

Funding

This study was supported by SIUE start-up funds from the DiSalvo lab.

Data Availability Statement

The raw data supporting the conclusions of this manuscript will be made available by

the authors, without undue reservation, upon request.

Table 1. *Burkholderia* infections significantly alter spore productivity in most (but

not all) bacterial culture conditions. Statistical Analysis of Spore Fitness from Figure

1. All DF values=3.

Condition	χ^2	p value
K. pneumoniae-GFP[100%]	23.537	>0.001
Rhizobium-GFP[100%]	20.81	>0.001
Rhizobium-GFP[50%]	18.759	>0.001
Serratia-GFP[100%]	6.411	0.09
Serratia-GFP[50%]	11.292	0.01
Agrobacterium-GFP[100%]	18.76	>0.001
Agrobacterium-GFP[50%]	16.267	>0.001
Pseudomonas-GFP[50%]	23.36	>0.001

1. Kiers ET, West SA. Evolving new organisms via symbiosis. Science. 2015 Apr
24;348(6233):392–4.

2. Margulis L, Fester R. Bellagio conference and book. Symbiosis as Source of
Evolutionary Innovation: Speciation and Morphogenesis. Conference--June 25-30,
1989, Bellagio Conference Center, Italy. Symbiosis Phila Pa. 1991;11:93–101.

3. Douglas AE. Symbiosis as a General Principle in Eukaryotic Evolution. Cold Spring
Harb Perspect Biol. 2014 Feb 1;6(2):a016113.

4. Douglas AE. Symbiotic interactions. Reprinted. Oxford: Oxford Univ. Press; 2002.
148 p. (Oxford science publications).

5.   Ochman H, Moran NA. Genes lost and genes found: evolution of bacterial
pathogenesis and symbiosis. Science. 2001 May 11;292(5519):1096–9.

6.   Dale C, Moran NA. Molecular interactions between bacterial symbionts and their
hosts. Cell. 2006 Aug 11;126(3):453–65.

7.   Wang D, Yang S, Tang F, Zhu H. Symbiosis specificity in the legume: rhizobial
mutualism. Cell Microbiol. 2012 Mar;14(3):334–42.

8.   Klironomos JN. Variation in Plant Response to Native and Exotic Arbuscular
Mycorrhizal Fungi. Ecology. 2003 Sep 1;84(9):2292–301.

9.   Chrostek E, Teixeira L. Mutualism Breakdown by Amplification of Wolbachia
Genes. Malik HS, editor. PLOS Biol. 2015 Feb 10;13(2):e1002065.

10. Álvarez-Loayza P, Jr JFW, Torres MS, Balslev H, Kristiansen T, Svenning J-C, et
al. Light Converts Endosymbiotic Fungus to Pathogen, Influencing Seedling
Survival and Niche-Space Filling of a Common Tropical Tree, Iriartea deltoidea.
PLOS ONE. 2011 Jan 31;6(1):e16386.

11. Dunbar HE, Wilson ACC, Ferguson NR, Moran NA. Aphid Thermal Tolerance Is
Governed by a Point Mutation in Bacterial Symbionts. PLOS Biol. 2007 Apr
10;5(5):e96.

12. Oliver KM, Campos J, Moran NA, Hunter MS. Population dynamics of defensive
symbionts in aphids. Proc R Soc B Biol Sci. 2008 Feb 7;275(1632):293–9.

13. Bonfante P, Anca I-A. Plants, Mycorrhizal Fungi, and Bacteria: A Network of
Interactions. Annu Rev Microbiol. 2009 Oct;63(1):363–83.

14. Bronstein JL. Conditional outcomes in mutualistic interactions. Trends Ecol Evol.
1994 Jun;9(6):214–7.

15. Hussa EA, Goodrich-Blair H. It Takes a Village: Ecological and Fitness Impacts of
Multipartite Mutualism. Annu Rev Microbiol. 2013 Sep 8;67(1):161–78.

16. Newton AC, Fitt BDL, Atkins SD, Walters DR, Daniell TJ. Pathogenesis, parasitism
and mutualism in the trophic space of microbe-plant interactions. Trends Microbiol.
2010 Aug;18(8):365–73.

17. Pérez-Brocal V, Latorre A, Moya A. Symbionts and Pathogens: What is the
Difference? In: Between Pathogenicity and Commensalism [Internet]. Springer,
Berlin, Heidelberg; 2011 [cited 2017 Dec 13]. p. 215–43. (Current Topics in
Microbiology and Immunology). Available from:
https://link.springer.com/chapter/10.1007/82_2011_190

18. Ronn R, McCaig AE, Griffiths BS, Prosser JI. Impact of Protozoan Grazing on
Bacterial Community Structure in Soil Microcosms. Appl Environ Microbiol. 2002
Dec 1;68(12):6094–105.

19. Paquet VE, Charette SJ. Amoeba-resisting bacteria found in multilamellar bodies
secreted by *Dictyostelium discoideum:* social amoebae can also package bacteria.
Kümmerli R, editor. FEMS Microbiol Ecol. 2016 Mar;92(3):fiw025.

20. Molmeret M, Horn M, Wagner M, Santic M, Abu Kwaik Y. Amoebae as training
grounds for intracellular bacterial pathogens. Appl Environ Microbiol. 2005
Jan;71(1):20–8.

21. Greub G, Raoult D. Microorganisms Resistant to Free-Living Amoebae. Clin
Microbiol Rev. 2004 Apr 1;17(2):413–33.

22. Horn M, Wagner M. Bacterial Endosymbionts of Free-living Amoebae1. J Eukaryot
Microbiol. 2004 Sep;51(5):509–14.

23. Rubeniņa I, Kirjušina M, Bērziņš A, Valciņa O, Jahundoviča I. Relationships
between Free-Living Amoeba and their Intracellular Bacteria. Proc Latv Acad Sci
Sect B Nat Exact Appl Sci. 2017 Aug 1;71(4):259–65.

24. Bozzaro S, Eichinger L. The Professional Phagocyte Dictyostelium discoideum as
a Model Host for Bacterial Pathogens. Curr Drug Targets. 2011 Jun;12(7):942–54.

25. Bozzaro S, Bucci C, Steinert M. Phagocytosis and host-pathogen interactions in
Dictyostelium with a look at macrophages. Int Rev Cell Mol Biol. 2008;271:253–
300.

26. Nasser W, Santhanam B, Miranda ER, Parikh A, Juneja K, Rot G, et al. Bacterial
Discrimination by Dictyostelid Amoebae Reveals the Complexity of Ancient
Interspecies Interactions. Curr Biol. 2013 May;23(10):862–72.

27. Skriwan C, Fajardo M, Hägele S, Horn M, Wagner M, Michel R, et al. Various
bacterial pathogens and symbionts infect the amoeba Dictyostelium discoideum.
Int J Med Microbiol. 2002 Jan;291(8):615–24.

28. smith jeff, Queller DC, Strassmann JE. Fruiting bodies of the social amoeba
Dictyostelium discoideum increase spore transport by Drosophila. BMC Evol Biol.
2014;14(1):105.

29. Chen G, Zhuchenko O, Kuspa A. Immune-like Phagocyte Activity in the Social
Amoeba. Science. 2007 Aug 3;317(5838):678–81.

30. Zhang X, Zhuchenko O, Kuspa A, Soldati T. Social amoebae trap and kill bacteria
by casting DNA nets. Nat Commun. 2016 Mar 1;7:10938.

31.   Brock DA, Douglas TE, Queller DC, Strassmann JE. Primitive agriculture in a social
amoeba. Nature. 2011 Jan 20;469(7330):393–6.

32.   DiSalvo S, Haselkorn TS, Bashir U, Jimenez D, Brock DA, Queller DC, et al.
Burkholderia bacteria infectiously induce the proto-farming symbiosis of
Dictyostelium amoebae and food bacteria. Proc Natl Acad Sci U S A. 2015 Sep
8;112(36):E5029–37.

33.   Suárez-Moreno ZR, Caballero-Mellado J, Coutinho BG, Mendonça-Previato L,
James EK, Venturi V. Common features of environmental and potentially beneficial
plant-associated Burkholderia. Microb Ecol. 2012 Feb;63(2):249–66.

34.   Brock DA, Hubert AM, Noh S, DiSalvo S, Geist KS, Haselkorn TS, et al.
Endosymbiotic adaptations in three new bacterial species associated with
Dictyostelium discoideum: Burkholderia agricolaris sp. nov., Burkholderia
hayleyella sp. nov., and Burkholderia bonniea sp. nov. 2018 Apr 19 [cited 2018
May 23]; Available from: http://biorxiv.org/lookup/doi/10.1101/304352

35.   Shu L, Brock DA, Geist KS, Miller JW, Queller DC, Strassmann JE, et al. A
complex symbiosis involving within species variation in the response of
Dictyostelium amoebae to Burkholderia bacteria. bioRxiv [Internet]. 2018 Jan 1;
Available from: http://biorxiv.org/content/early/2018/10/05/436386.abstract

36.   Inglis RF, Biernaskie JM, Gardner A, Kümmerli R. Presence of a loner strain
maintains cooperation and diversity in well-mixed bacterial communities. Proc R
Soc B Biol Sci. 2016 Jan 13;283(1822):20152682.

37.   Norris MH, Kang Y, Lu D, Wilcox BA, Hoang TT. Glyphosate Resistance as a
Novel Select-Agent-Compliant, Non-Antibiotic-Selectable Marker in Chromosomal
Mutagenesis of the Essential Genes asd and dapB of Burkholderia pseudomallei.
Appl Environ Microbiol. 2009 Oct 1;75(19):6062–75.

38.   Su S, Bangar H, Saldanha R, Pemberton A, Aronow B, Dean GE, et al.
Construction and characterization of stable, constitutively expressed, chromosomal
green and red fluorescent transcriptional fusions in the select agents, *Bacillus
anthracis* , *Yersinia pestis* , *Burkholderia mallei* , and *Burkholderia pseudomallei*.
MicrobiologyOpen. 2014 Oct;3(5):610–29.

39.   Kikuchi Y, Fukatsu T. Live imaging of symbiosis: spatiotemporal infection dynamics
of a GFP-labelled *Burkholderia* symbiont in the bean bug *Riptortus pedestris*. Mol
Ecol. 2014 Mar;23(6):1445–56.

40.   Fazli M, Almblad H, Rybtke ML, Givskov M, Eberl L, Tolker-Nielsen T. Regulation
of biofilm formation in Pseudomonas and Burkholderia species: Regulation of
biofilm formation. Environ Microbiol. 2014 Jul;16(7):1961–81.

41.   Tomlin KL, Coll OP, Ceri H. Interspecies biofilms of Pseudomonas aeruginosa and
Burkholderia cepacia. Can J Microbiol. 2001 Oct;47(10):949–54.

42. Brock DA, Callison WÉ, Strassmann JE, Queller DC. Sentinel cells, symbiotic
bacteria and toxin resistance in the social amoeba Dictyostelium discoideum. Proc
Biol Sci. 2016 27;283(1829).

43. Compant S, Nowak J, Coenye T, Clément C, Ait Barka E. Diversity and occurrence
of Burkholderia spp. in the natural environment. FEMS Microbiol Rev. 2008
Jul;32(4):607–26.

44. Estrada-De Los Santos P, Bustillos-Cristales R, Caballero-Mellado J. Burkholderia,
a Genus Rich in Plant-Associated Nitrogen Fixers with Wide Environmental and
Geographic Distribution. Appl Environ Microbiol. 2001 Jun 1;67(6):2790–8.

45. Gage DJ. Infection and Invasion of Roots by Symbiotic, Nitrogen-Fixing Rhizobia
during Nodulation of Temperate Legumes. Microbiol Mol Biol Rev. 2004 Jun
1;68(2):280–300.

46. Nguyen NH, Bruns TD. The Microbiome of Pinus muricata Ectomycorrhizae:
Community Assemblages, Fungal Species Effects, and Burkholderia as Important
Bacteria in Multipartnered Symbioses. Microb Ecol. 2015 May;69(4):914–21.

47. Edgcomb V. Marine protist associations and environmental impacts across trophic
levels in the twilight zone and below. Curr Opin Microbiol. 2016 Jun;31:169–75.

48. Banitz T, Wick LY, Fetzer I, Frank K, Harms H, Johst K. Dispersal networks for
enhancing bacterial degradation in heterogeneous environments. Environ Pollut.
2011 Oct;159(10):2781–8.

49. Ellegaard-Jensen L, Knudsen BE, Johansen A, Albers CN, Aamand J, Rosendahl
S. Fungal–bacterial consortia increase diuron degradation in water-unsaturated
systems. Sci Total Environ. 2014 Jan;466–467:699–705.

50. Rashid MI, Mujawar LH, Shahzad T, Almeelbi T, Ismail IMI, Oves M. Bacteria and
fungi can contribute to nutrients bioavailability and aggregate formation in
degraded soils. Microbiol Res. 2016 Feb;183:26–41.

51. Nazir R, Zhang M, de Boer W, van Elsas JD. The capacity to comigrate with
Lyophyllum sp. strain Karsten through different soils is spread among several
phylogenetic groups within the genus Burkholderia. Soil Biol Biochem. 2012
Jul;50:221–33.

52. Warmink JA, Nazir R, Corten B, van Elsas JD. Hitchhikers on the fungal highway:
The helper effect for bacterial migration via fungal hyphae. Soil Biol Biochem. 2011
Apr;43(4):760–5.

53. Vila T, Nazir R, Rozental S, dos Santos GMP, Calixto ROR, Barreto-Bergter E, et
al. The Role of Hydrophobicity and Surface Receptors at Hyphae of Lyophyllum sp.
Strain Karsten in the Interaction with Burkholderia terrae BS001 – Implications for

Interactions in Soil. Front Microbiol [Internet]. 2016 Oct 27 [cited 2018 Sep 7];7.
Available from: http://journal.frontiersin.org/article/10.3389/fmicb.2016.01689/full

54. Moebius N, Üzüm Z, Dijksterhuis J, Lackner G, Hertweck C. Active invasion of
bacteria into living fungal cells. eLife [Internet]. 2014 Sep 2 [cited 2018 Sep 7];3.
Available from: https://elifesciences.org/articles/03007

55. Partida-Martinez LP, Monajembashi S, Greulich K-O, Hertweck C. Endosymbiont-
Dependent Host Reproduction Maintains Bacterial-Fungal Mutualism. Curr Biol.
2007 May;17(9):773–7.

56. Gong L, Lai S-C, Treerat P, Prescott M, Adler B, Boyce JD, et al. Burkholderia
pseudomallei Type III Secretion System Cluster 3 ATPase BsaS, a
Chemotherapeutic Target for Small-Molecule ATPase Inhibitors. Blanke SR, editor.
Infect Immun. 2015 Apr;83(4):1276–85.

57. Stone JK, DeShazer D, Brett PJ, Burtnick MN. Melioidosis: molecular aspects of
pathogenesis. Expert Rev Anti Infect Ther. 2014 Dec;12(12):1487–99.

58. Angus AA, Agapakis CM, Fong S, Yerrapragada S, Estrada-de los Santos P, Yang
P, et al. Plant-Associated Symbiotic Burkholderia Species Lack Hallmark
Strategies Required in Mammalian Pathogenesis. van Schaik W, editor. PLoS
ONE. 2014 Jan 8;9(1):e83779.

59. Dinh C, Farinholt T, Hirose S, Zhuchenko O, Kuspa A. Lectins modulate the
microbiota of social amoebae. Science. 2018 Jul 27;361(6400):402–6.

60. Šulák O, Cioci G, Lameignère E, Balloy V, Round A, Gutsche I, et al. Burkholderia
cenocepacia BC2L-C Is a Super Lectin with Dual Specificity and Proinflammatory
Activity. Saper MA, editor. PLoS Pathog. 2011 Sep 1;7(9):e1002238.

FIGURES
683
Figure 1. Culture Conditions Modify the Impact of *Burkholderia* Infections on Host
Fitness. Total spore productivity for *D. discoideum* after one round of development on
the indicated GFP labeled bacterial species (Kleb= *K. pneumoniae*, Rhizo=*Rhizobium*,
Serr=*Serratia*, Agro=*A, tumefaciens,* and Pseu=*P. aeruginosa*. 50% cultures are mixed
with 50% unlabeled *K. pneumoniae* by volume). Dark grey, light grey, and medium grey
boxes indicate cultures wherein *Burkholderia* infections are initiated by inclusion of 5%
*B. agricolaris.*70-RFP, *B. hayleyella.*11-RFP, and *B. bonniea.*859-RFP, respectively.
Points within boxes indicate individual replicates. Letters indicate post-hoc significance
within panels.
693

699 **Figure 2. Secondary Bacteria do not Infect Symbiont Free Spores.** Representative
700 confocal micrographs of sori contents after growth on the indicated GFP labeled
701 bacterial cultures. Scale bar=10um.
702

Figure 3. *Burkholderia* Differentially Induce Intracellular Co-Infections. a) Representative confocal micrographs of *Burkholderia*-RFP infected sori contents after development on the indicated GFP labeled secondary bacteria. Scale bar= 10um. b) Percent of spores infected with *Burkholderia*-RFP (light grey) and co-infected with secondary bacteria-GFP (dark grey) averaged across all replicates of all secondary bacterial conditions. c) Average percent of spores infected (as in b) for each secondary culture condition. Error bars $^{\pm}$SE.

Figure 4. *Burkholderia* Induce Extracellular Co-infections. a) Percent of confocal micrographs wherein secondary bacterial-GFP cells were visualized extracellularly within sori contents. Numbers nested within bars indicate the number of images with visible GFP over the number of total images collected per condition. b) Number of GFP colony forming units (log 10) from *Burkholderia* infected sori contents when grown on *K. pneumoniae*-GFP and *Rhizobium*-GFP. Individual sori contents were harvested either 6 or 14 days after plating on the *Burkholderia*/Secondary bacterial culture conditions. These numbers may represent both extracellular and intracellularly derived secondary bacteria. Points represent the GFP-CFU count for individual replicates.

Figure 5. Farming Benefits Vary by Symbiont and Dispersal Strategy. For all
figures, top panels represent data from sori pre-grown on *K. pneumoniae*-GFP[100%] and
bottom panels represent sori pre-grown on *Rhizobium*-GFP[50%], both with (greys) and
without (white) supplementation of 5% *Burkholderia*. a) Quantification of total spores
harvested from food rich and food scarce plating conditions after transfer from *K.*
pneumoniae-GFP[100%] (top) or *Rhizobium*-GFP[50%] (bottom) plating conditions. Points
represent data from each individual replicate. b) Representative images of individual
sori streaks from fruiting bodies developed on *K. pneumoniae*-GFP[100%] (top) or
Rhizobium-GFP[50%] (bottom) plates. Top panels are magnified sections of bottom
panels. c) Percentage of bacterial positive sori streaks with observable fruiting bodies.
Error bars +SE. d) Number of fruiting bodies per fruiting body positive streaks for each
individual replicate. All letters indicated post-hoc significance within panels. For streak
tests, all sori where streaked five days after plating on *K. pneumoniae*-GFP[100%] (top
panels) or *Rhizobium*-GFP[50%] (bottom panels) and all data was collected 5 days after
streaking.
752

753
754
755
756

Figure 6. Fruiting Bodies Amplify in *Burkholderia* Infected Sori Streaks. Images of sori streaks two weeks after streak-testing. Sori were harvested from individual fruiting bodies that developed five days after plating spores on *Rhizobium*-GFP$^{50\%}$ culture conditions with 5% of the indicated *Burkholderia*-RFP strain.

Ba.70   Bh.11   Bb.859

Figure 7. Co-infection Patterns Correspond to *Burkholderia* Species. a) Representative confocal micrographs of sori after five days post plating with 5% of the indicated *Burkholderia*-RFP strain and *Rhizobium*-GFP$^{50\%}$. b) Average percent of spores visualized with intracellular *Burkholderia*-RFP (light grey) and intracellular *Rhizobium*-GFP (dark grey) for each of the indicated *Burkholderia* species. Letters indicate post-hoc significance. c) Percent of images in which external GFP was visualized out of the total number of independent images acquired. Numbers above bars represent the raw number of external GFP positive images over the number of images acquired for each *Burkholderia* infection condition.

**Figure 8. Host Benefits Correspond to *Burkholderia* Species and their Induction
of Co-infections.** a) Images of representative sori streaks from fruiting bodies grown on
Rhizobium-GFP[50%] with 5% of the indicated *Burkholderia* strain. Top panels are
magnified sections of bottom panels. b) Percentage of bacterial positive sori streaks
with observable fruiting bodies. Error bars +SE. c) Number of fruiting bodies per fruiting
body positive streaks for each individual. All letters indicated post-hoc significance
within panels. All sori where streaked five days after plating on *Rhizibium*-GFP[50%] and
all data was collected 5 days after streaking.

815
816
817
818