Effect of fire and thinning on fine-scale genetic structure and gene flow in fire-suppressed populations of sugar pine (*Pinus lambertiana* Douglas)

Brandon M. Lind[a,b], Malcolm P. North[c], Patricia E. Maloney[d], and Andrew J. Eckert[e]

October 1, 2018

[a]Forest & Conservation Sciences, University of British Columbia, Vancouver, BC V6T1Z4 Canada

[b]Integrative Life Sciences, Virginia Commonwealth University, Richmond, Virginia 23284 USA

[c]USDA Forest Service Pacific Southwest Research Station, Davis, CA 95618 USA

[d]Department of Plant Pathology and Tahoe Environmental Research Center, University of California, Davis, California 95616 USA

[e]Department of Biology, Virginia Commonwealth University, Richmond, Virginia 23284 USA

Running Title: Genetic effects of fire and thinning

Keywords: Sierra Nevada, fire, forest management, sugar pine, gene flow

Corresponding Author:

Brandon M. Lind

3041 - 2424 Main Mall

Forest Science Centre

University of British Columbia

Vancouver, British Columbia V6P 4E9

E-mail: brandon.lind@ubc.ca

Abstract

Historically, frequent, low-severity fires in dry western North American forests were a major driver of ecological patterns and processes, creating resilient ecosystems dominated by widely spaced pine species. However, a century of fire-suppression has caused overcrowding, altering forest composition to shade-tolerant species, while increasing competition and leaving trees stressed and susceptible to pathogens, insects, and high-severity fire. Exacerbating the issue, fire incidence is expected to increase with changing climate, while fire season has been observed to begin earlier and last longer than historic trends. Forest thinning and prescribed fire have been identified as important management tools to mitigate these risks. Yet little is known of how thinning, fire, or their interaction affect contemporary evolutionary processes of constituent pine species that influence fitness and play an important role in the opportunity for selection and population persistence. We assessed the impact of widely used fuel reduction treatments and prescribed fire on fine-scale gene flow on an ecologically important and historically dominant shade-intolerant pine species of the Sierra Nevada, *Pinus lambertiana* Dougl. Treatment prescription (no-thin-no-fire, thin-no-fire, and fire-and-thin) was found to differentially affect both fine-scale spatial and genetic structure as well as effective gene flow in this species. Specifically, the thin-no-fire prescription increases genetic structure (spatial autocorrelation of relatives) between adults and seedlings, while seed and pollen dispersal increase and decrease, respectively, as a function of increasing disturbance intensity. While these results may be specific to the stands at our study site, they indicate how assumptions relating to genetic effects based on spatial structure can be misleading. It is likely that these disequilibrated systems will continue to evolve on unknown evolutionary trajectories. The long-term impacts of management practices on reduced fitness from inbreeding depression should be continually monitored to ensure resilience to increasingly frequent and severe fire, drought, and pest stresses.

Introduction

Many aspects of conifer biology are affected by a tree's surrounding environment as well

as the density of hetero- and conspecifics. For instance, outcrossing rates of conifer species are

often tied to population density (Farris & Mitton 1984) and surrounding tree heights (O'Connell

et al. 2003), while removal of proximal individuals can increase pollen and gene flow distances

by reducing potential mates and removing once impeding vegetation. Thus, disturbance, *sensu*

lato, has the potential to alter contemporary demographic and reproductive dynamics through

both direct (population-level) and indirect (ecological-level) impacts (Mouillot *et al.* 2013).

Historically, natural disturbances such as fire were commonplace and equilibrated many

ecosystem functions and processes in forests of the western United States (Covington *et al.*

1994). Fire regimes in these regions had return intervals on decadal scales (10-17 years; North

et al. 2005), in contrast to wetter climates where fire return intervals were (sub)centennial (50+

years [North *et al.* 2016]). Resultantly, these ecosystems experienced frequent, low-severity

burns and were populated by fire-adapted species, creating forests dominated by resilient,

widely spaced pine trees. Yet over the past 150 years, anthropogenic influence has resulted in

forests that are now fire-suppressed and overgrown by shade-tolerant species, causing

increased competition, leaving trees stressed and susceptible to fungal and bark beetle attacks

(Bonello *et al.* 2006).

Stand densification has also increased the frequency and probability of contemporary,

high-severity fires. Between 2012 and 2014 in California alone, 14,340 fires burned 1.1 million

acres and injured or killed nearly 300 individuals (NIFC 2014). Collectively, fires across

California, the Great Basin, Southwest, and Rocky Mountain territories have burned a combined

8.8 million acres between 2014 and 2015 (NIFC 2015), while Forest Service scientists predict

future fires to reach unprecedented levels, covering over 12-15 million acres annually (USDA

Forest Service 2016a) requiring the United States Forest Service (USFS) to budget

$2,300,000,000 on wildfire management, suppression, and preparedness for the 2016 fiscal

year (USDA Forest Service 2016b). Exacerbating the issue, analyses of fire season length and

onset have shown that seasons are beginning earlier and lasting longer than historic trends

(Westerling 2006) while climate models predict extreme weather favorable to fire to become

more frequent, and ignited fires to increase in severity, size, and required suppression efforts

(Miller *et al.* 2009).

Because of these contemporaneous trends, large-scale forest thinning projects have

been implemented to simultaneously restore fire-frequent ecosystems to their pre-settlement

resilience as well as to protect urban development and human life, as fuel reduction treatments

have been shown to be an effective tool in decreasing fire severity and ignition probability (Agee

& Skinner 2005; Schwilk *et al.* 2009; Safford *et al.* 2009). For example, the Sierra Nevada

Forest Plan Amendment (USDA Forest Service 2004) mandates that 50% of initial thinning

treatments take place near urban populations, while the remaining thinning take place in natural

wildland stands. To encourage fire resiliency the USFS has implemented fuel reduction

treatments across 6.1 million acres of western, fire-suppressed forestland in 2014 (USDA Forest

Service 2016a). Further, forest and fire scientists are calling for an overhaul of management

policy to implement these thinning treatments to a far greater extent (North *et al.* 2015). While

congruent with historic forest structure, these actions will orient these already disequilibrated

systems on trajectories of unknown evolutionary consequence.

Through timber harvests, land use conversion, and fire suppression, forests have

undergone systemic shifts in composition, structure, and disturbance regimes that are

incongruous to the natural and evolutionary histories of endemic species (Collins *et al.* 2011;

Larson & Churchill 2012). Consequentially, anthropogenic forest disturbance has been at the

forefront of conservation attention for decades (Ledig 1988; 1992). The extent of human impact

on forested land has received particular attention as a result of the empirical expectations

developed from population genetic theory. Specifically, because of the reduction in individual

tree density overall, and in particular for larger trees that asymmetrically contribute gametes to

reproduction (Richardson *et al.* 2014), harvested forests are thought to be specifically subjected

to population bottlenecks, potentially altering existing mating systems or available gene pools

while decreasing genetic variability within populations and increasing differentiation from native

stands (Smouse *et al.* 2001, Cloutier *et al.* 2006; Kramer *et al.* 2008; Lowe *et al.* 2015). These

consequences can influence the fitness of affected populations, as drastic changes in gene pool

availability or mating system can alter a population's potential to adapt to local conditions where

inbreeding depression can have deleterious effects on growth and reproductive potential (e.g.,

reproductive capacity or rates of embryo abortion).

Past studies investigating the genetic effects of North American forest management

show mixed evidence of harvest influence. These studies often sub-sample populations and

primarily focus on diversity consequences across a range of molecular markers (often

microsatellites). Many management studies of North American conifers compare genotypic

diversity indices (e.g., H_E, H_O, allelic richness, etc.) between treatments to detect management

influence (Cheliak *et al.* 1988; Gömöry 1992; Buchert *et al.* 1997; Adams *et al.* 1998; Rajora *et*

al. 2000; Macdonald *et al.* 2001; Perry & Bousquet 2001; Rajora & Pluhar 2003; El-Kassaby *et*

al. 2003; Marquardt *et al.* 2007; Fageria & Rajora 2013a; b). However, the same diversity values

can manifest under completely different scenarios and tests of significance between population

values for a small number of markers may therefore be under-informative, particularly for sub-

sampled populations, as these differences can result from sampling bias or from evolutionary

processes unrelated to management. Additionally, these investigations also often employ F_{ST}

analyses to assess statistical significance between treated and untreated stands (Thomas *et al.*

1999; Perry & Bousquet 2001; Marquardt *et al.* 2007; Fageria & Rajora 2013a; b). Though when

used in this context, this test is simply signifying whether the allelic frequencies in

(sub)populations under study are likely to have been sampled from the same ancestral

population (Holsinger & Weir 2009). Very often, the treated and untreated stands are physically

adjacent (derived of a common ancestral population) and only under extreme perturbation

should significance be expected. In cases where significance is detected, and other than to

assess relative diversity between stands, such differentiation does little to inform how

management is affecting ongoing evolutionary processes affecting fitness, as such processes

may ameliorate bottlenecks due to management. It would therefore be difficult to draw such

conclusions without assessing other stand and evolutionary dynamics.

Very seldom in North American studies of forest management are evolutionary

processes influencing fitness specifically examined (but see Neale & Adams 1985). Yet when

studies are done and nonsignificant findings are found, authors generally caution interpretation

(Finkeldey & Ziehe 2004; Namroud *et al.* 2012). Very often the scale of sampling (both in terms

of numbers and spatial extent of individuals and the degree of temporal variation), as well as the

lack of investigation into evolutionary dynamics have been offered as inadequate, and that

further investigation into evolutionary consequences of natural and anthropogenic disturbance

could give valuable insight to forest managers and fill a vital knowledge gap in this regard

(Namroud *et al.* 2012). Indeed, incongruence between theoretical predictions and empirical

results from studies evaluating genetic consequences of forest disturbance has created a

paradox within the literature (Kramer *et al.* 2008). Yet as Lowe et al. (2015) point out, we may

have been looking in the wrong place. They argue that instead of simply assaying mature

cohorts to understand the genetic consequences of disturbance, future attention should include

progeny arrays as well as the relative regenerative success across a wide range of influences.

Additionally, they contend that the type and magnitude of the genetic response itself may be

better understood through the variation in mating and breeding systems of studied species. Of

particular importance, Lowe *et al.* (2015) advise scientists that the most fruitful research

endeavors will incorporate quantitative approaches to understanding evolutionary mechanisms,

specifically those connecting changes in pollination to mating systems and evolutionary fitness,

and that these efforts will likely generate critical knowledge regarding the mechanisms driving

the dynamics we observe.

Interactions between fire and forest thinning management are certain. To ensure forests

are resilient to frequent fire and disturbance, and provide habitat for public recreation and native

wildlife, the interactive impact of management and fire must be understood in an evolutionary

framework. Here, we investigated the evolutionary impact of forest management on fire-

suppressed populations of the historically dominant and ecologically important sugar pine

(*Pinus lambertiana* Dougl.) within Teakettle Experimental Forest (TEF), a USFS site located in

the central Sierra Nevada of California. Using microsatellite markers, we employ parentage

analysis and assess impact upon various processes known to affect fitness such as mating

patterns, effective dispersal distances, and fine-scale (<300m) genetic structure. Our results

show that thinning alone increases fine-scale genetic structure of constituent trees (i.e., spatial

autocorrelation of relatives), and that the vast majority of pollen and seed dispersal events take

place at this same scale. Although the genetic structure of adults is due to an interaction

between the evolutionary history of the stand and the applied treatment, mating patterns and

seedling ingrowth will determine long-term impacts of management. While effects of such

treatments will vary by location, our results show that the degree of thinning and the choice of

leave-trees should be tailored to a given stand, and that spatial structure (arrangement of

individuals across the landscape) should not be conflated with spatial genetic structure

(arrangement of relatives across the landscape). By avoiding treatments that exacerbate

genetic structure, managers may be able to decrease seed abortion due to inbreeding and thus

increase effective seed rain of species with management importance.

Methods

Study area, sampling, and focal species

Teakettle Experimental Forest (TEF) is a fire-suppressed, old-growth forest watershed in

the central Sierra Nevada mountains of California. The 1300-ha forested watershed ranges from

1900–2600m in elevation and consists of five conifer species representative of fire-suppressed

forests of the Sierra range: white fir (*Abies concolor* [Gordon] Lindley ex Hildebrand), red fir (*A.*

magnifica A. Murray), incense cedar (*Calocedrus decurrens* [Torr] Florin), Jeffrey pine (*Pinus*

jeffreyi Balf.), and sugar pine (*P. lambertiana*). Historically, fire burned the area every 11-17

years, but has been suppressed for 135 years (North *et al.* 2005) while logging had been

completely

180
181
Figure 1 *Teakettle Experimental Forest, California (Latitude: 36.9606, Longitude: -119.0258). (A) Topographic map*
and spatial arrangement of treatments (BC = burned understory thin; BN = burned no-thin; BS = burned shelterwood
thin; UC = unburned understory thin; UN = unburned no-thin; US = unburned shelterwood thin). Replicates for each
treatment are numbered one through three from south to north. (B) Mapped coordinates (Universal Transverse
Mercator) and elevation (meters) of pre-treatment adults   5cm diameter at breast height. (green: P. lambertiana,
red: P. jeffreyi, gray: A. concolor, blue: A. magnifica, orange: C. decurrens, black: Quercus, Salix, and remaining
species.).

absent within the watershed (North 2002). Six treatments were applied to neighboring 4-ha plots

at TEF (each 200m x 200m, Figure 1a) by crossing two levels of burn (no-fire and fire) with

three levels of thinning (no-thinning, overstory-thinning, and understory-thinning). The

understory thinning prescription followed guidelines in the California spotted owl (CASPO)

report (Verner *et al.* 1992), which is now widely used for fuel management in California (SNFPA

2004). Each treatment was replicated three times for a total of 18 plots covering 72ha.

Understory-thinning removed all trees with a diameter at breast height (DBH) ≤76cm and

≤≥25cm, while overstory-thinning removes all trees >25cm DBH except 18-22 of the largest

trees per hectare. Treatments were applied to the watershed over 2000 and 2001. Plot

inventories of pre-treatment (1999), and post-treatment (2001, 2004, and 2011) conditions

mapped individual trees on a 3D coordinate system that was then translated into Universal

Transverse Mercator (colored dots, Figure 1b). Only standing boles ≥5cm DBH were included in

plot inventories, which recorded species, DBH, spatial coordinates, and decay class and forest

health metrics (e.g., presence/absence of insects and pathogens). Post-treatment inventories

updated DBH, decay class, forest health metrics, and added individuals to the dataset once they

reached 5cm DBH. Here, seedling and saplings are all pine stems <5cm DBH. For these, basal

diameter and spatial coordinates were recorded over the summers of 2012 and 2013 by BML

while collecting needle tissue samples from the full census of all live *P. lambertiana* ($N = 3,135$).

Pinus lambertiana is a historically dominant member of mixed-conifer forests of the Sierra

Nevada, and continues to play important ecological roles. This species is shade-intolerant and

is an important focus of restoration in the Sierra Nevada range.

Analysis of tree spatial structure

Using plot-level *P. lambertiana* individuals, we estimated spatial structure of seedlings

and adults across 10-meter distance classes, *r*, separately using univariate inhomogeneous pair

correlation functions ($g_{inhom}(r)$) from the `spatstat` library in R (Baddeley *et al.* 2015) with an

isotropic edge correction. This statistic was chosen over Ripley's K, or its linearized version (*L*)

because of advocacy for $g_{inhom}(r)$ over these statistics (see `spatstat` manual). This analysis

tests the null hypothesis that the 2D spatial arrangement of points (adults or seedlings) is not

significantly different from complete spatial randomness (CSR; i.e., a Poisson distribution of

inter-point distances with inhomogeneous intensities of points), where support for the alternative

hypothesis is indicative of ecological factors driving spatial patterning. We calculated null

confidence envelopes for each test using 199 null simulations of CSR using the same intensity

of the pattern of individuals analyzed (equivalent to an alpha value of 0.01; see `spatstat`

manual). For trees that coincide with the null model of CSR $g_{inhom}(r) = 1$, with spatial

aggregation resulting in $g_{inhom}(r) > 1$, and with spatial inhibition resulting in $g_{inhom}(r) < 1$

(Baddeley *et al.* 2015); significance was judged using the null confidence envelopes described

above. We also repeated this analysis for shade-tolerant individuals (i.e., *A. concolor* and *A.*

magnifica grouped together). Further, we extended the univariate inhomogeneous pair

correlation function to its bivariate equivalent, $g_{inhom,i,j}(r)$, to test for spatial affinity between two

groups *i* and *j*, using similar methods as above for edge correction and null confidence

envelopes. We calculated $g_{inhom,i,j}(r)$ between unique combinations of *P. lambertiana* adults,

P. lambertiana seedlings, and shade-tolerant individuals. Hypothesis testing and interpretation

of bivariate $g_{inhom,i,j}(r)$ was carried out as with univariate $g_{inhom}(r)$. Results from these

analyses will allow us to compare standing spatial structure of trees against spatial genetic

autocorrelation (see below) and make inferences about the ecology of these species as well as

how treatments at TEF are affecting ongoing evolutionary dynamics.

DNA extraction, microsatellite amplification

Total genomic DNA was extracted according to manufacturer's protocol using the

DNeasy 96 Plant Kit (Qiagen, Germantown, MD) from finely ground *P. lambertiana* samples

within a subset of the factorial treatments at TEF: unburned-no-thin control plots (hereafter UN),

understory-thin (CASPO) plots without burn application (hereafter UC), and burned understory-

thin plots (hereafter BC) for a total of 1,348 individuals. Herein, we often refer to patterns across

these treatments in terms of increasing disturbance intensity (i.e., from UN to UC to BC). For

each individual, three chloroplast (paternally inherited, Wofford *et al.* 2014: pt71936, pt87268,

pc10) and four nuclear (biparental inheritance, Echt *et al.* 1996: rps50, rps02, rps12, rps39)

microsatellite markers were amplified (using fluorescent dyes NED, PET, VIC, and FAM) per the

original publications with minor modifications using BIO-RAD iProof high fidelity DNA

polymerase (see Supplemental Information). The chloroplast markers were chosen for their

primer conservation across *Pinus*, *Trifoliae*, *Parrya*, and *Quinquifolia* subsections of the *Pinus*

genus (Wofford *et al.* 2014) while the chosen nuclear markers have been amplified in eastern

white pine (*P. strobus* L., Echt *et al.* 1996) and both sets successfully amplified on a subset of

individuals at TEF as a proof-of-concept judged by gel electrophoresis. Multiplexed individuals

(one fluorescent dye per well) were analyzed using the Applied Biosystems 3730xl fragment

analyzer at Cornell University (http://www.biotech.cornell.edu/brc/genomics-facility) and

genotypes were called using GeneMaker v2.6.7 (see Supplemental Info;

http://www.softgenetics.com/GeneMarker.php).

Genetic diversity measures

Treatment-specific diversity measures (total number of alleles, A_T; mean number of

alleles per locus, A; effective number of alleles per locus, A_e; observed and expected

heterozygosity for nuclear markers, respectfully H_o, H_e; average number of private alleles, A_P;

and overall means for each category) were calculated for each treatment and averaged across

loci in order to compare dynamics at TEF to published studies. For estimates of H_o and H_e, only

nuclear markers were used. To quantify variation in these measures we also report standard

deviation. We also calculated hierarchical multi-locus F_ST according to Weir & Cockerham

(1984) for nuclear markers using the `hierfstat` package in R (Goudet & Jombart 2015) and

calculated treatment-specific F_ST in a similar manner in order to compare fixation indices across

treatments. Further, single- and multi-locus exclusion probabilities for parentage analysis (see

below) were calculated using python scripts modified from `gstudio` (v1.5.0; Dyer 2016).

Analysis of spatial genetic structure

To quantify spatial genetic autocorrelation at a distance class (lag) *h* (hereafter r_g^h), we

used multi-locus genetic distances (Smouse & Peakall 1999) and Euclidean geographic

distances among spatial coordinates of individuals to calculate r_g^h across distances classes *h*

corresponding to approximately 10-meter bins for *P. lambertiana* seedlings, *P. lambertiana*

adults, as well as a bivariate approximation for the clustering of *P. lambertiana* adult genotypes

to those of seedlings. For a given distance class, *h*, spatial patterning of multi-locus genotypes

are unrelated to (i.e., random relative to) the spatial patterns of individuals if $r_g^h = 0$, aggregated

if $r_g^h > 0$, and dispersed if $r_g^h < 0$. We estimated null confidence intervals by taking the 2.5[th] and

97.5[th] quantiles of $M = 1000$ estimates of $r_{g,m}^h$, where 999 of these estimates were computed by

randomly permuting individual genotypes across empirical spatial coordinates, with the M^{th}

permutation being the empirical estimate of r_g^h itself (Smouse & Peakall 1999). Using the

`PopGenReport` package in R (Adamack & Gruber 2014) we created correlograms for nuclear

and chloroplast markers both in isolation and in combination, but present only those using full

genotypes as correlograms by marker type showed similar patterns as full genotypes. We used

these correlograms to quantify spatial aggregation of genotypes so that conclusions based on

treatment effects could be compared and contextualized with ongoing evolutionary dynamics at

TEF such as that of fine-scale gene flow.

Parentage analysis

To quantify fine-scale gene flow at TEF, we conducted parentage analysis using our

genetic markers and spatial coordinates of individuals. Joint estimation of parentage and

dispersal parameters (i.e., mean dispersal distances of seed and pollen) were achieved by

expanding methods of Moran & Clark (2011). This method simultaneously estimates parentage

and dispersal kernel parameters for seed and pollen within a Bayesian framework, taking into

account genotyping error and variation in individual fecundity while treating dispersal processes

inside and outside of the mapped areas in a coherent manner, which is critical if the dispersal

kernel is to reflect both long- and short-distance movement. Here, all sampled adults are

characterized by a multi-locus genotype and a mapped coordinate. Additionally, there exists a

sample of seedlings, each of which has not only a genotype and location, but an estimated

pedigree as well, which can consider any adult as either mother or father, or of a selfing event

(though we excluded possible selfing events from analyses). The probability of the pedigree

considering two in-plot parents, before incorporating information regarding genotype, is

estimated from the probability of pollen to mother movement over the given distance and of

seed movement over the distance between mother and seedling, as well as the parental prior

distribution for fecundity and pollen production. For the study here, pollen production was

considered proportional to fecundity (as in Moran & Clark 2011) and was estimated by fitting a

2[nd]-order power polynomial regression to data from Figure 6 in Fowells & Schubert (1956)

where Cone Count = 0.0098(dbh^2) - 0.4811(dbh) + 10.651. After calculating cone counts using

this regression, we set fecundity for all adults <25cm DBH to zero given observed cone counts

from Fowells & Schubert (1956). For dispersal priors, we set the seed dispersal kernel shape

parameter, u_s, to 253.31, corresponding to a mean dispersal distance of 25m (Millar *et al.* 1992;

Fowells & Schubert 1956) while the pollen dispersal kernel shape parameter prior, u_p, was set

to 2279.72, corresponding to a mean pollen distance of 75m (Wright 1976; Neale 1983; Millar *et*

al. 1992). For priors to the standard deviation of mean dispersal we set seed (pollen) to 1013.21

(9118.90) corresponding to standard deviations of 50m (75m).

Given that either parent could have produced the offspring, the likelihood that this pair is

the true parents relative to all other possible parent pairs depends on the dispersal kernel priors

for seed and pollen, and the seed and pollen production of all trees both inside and outside of

the plot (the fraction of all possibilities; Moran & Clark 2011). To evaluate the probability of an

offspring having one parent in the plot and the other outside of the plot, a set of potential out-of-

plot parent-densities, $dp_1,\dots,dp_{20}$, each 10m progressively outside of the plot is considered

(see supplemental figure S3.1 in Moran & Clark 2011). Pollen and seed movement into the plot

is approximated by assuming first that all seed/pollen produced within each quarter-polygon, v,

originates from a tree located dp_v meters from the midpoint of each side outside of the plot. The

expected out-of-plot pollen (seeds) reaching an in-plot mother (a seedling's location) from each

quarter-polygon outside of the plot is calculated based on the average density and average

fecundities of trees outside of the plot and then multiplied by the probability of dispersal to the

point within the plot. Summing over each distance class over each side gives the total expected

out-of-plot pollen/seed dispersal to points inside of the plot. However, to calculate the probability

of an in-plot versus an out-of-plot father, the expected pollen arriving at an out-of-plot mother

from another out-of-plot father must first be calculated using the concentric polygons around the

sampled plot and the distance classes described above. The fraction of rings falling outside the

plot determines the fraction of pollen received from each distance class, dp_v, expected to come

from outside trees. Once error rates (e_1) and dropout rates (e_2) of genotyping are calculated

through regenotyping individuals (see Supplemental Information), the probability of a pedigree,

seed and dispersal parameters given the offspring genotype, distances, error rates, and

pollen/seed production can be estimated (Moran & Clark 2011). Very rarely have previous

studies investigating effects of forest management (or using parentage analysis towards such

goals) incorporated error and dropout rates into subsequent inferences.

For the current study, out-of-plot densities were extrapolated for each side of the nine

plots used at Teakettle from densities and DBH distributions (our proxy for fecundity) revealed in

pre-treatment surveys (North 2002). Due to the proximity of the treated plots, all adult trees and

seedlings across UC, BC, and UN treatments were considered simultaneously for parentage

assignment. Our methods therefore extend Moran & Clark (2011) from a single plot of sampled

individuals to multiple plots across the landscape. Additionally, instead of considering any given

pedigree as symmetrical (i.e., with no consideration for which tree was the pollen or seed donor)

we utilize genotyped markers separately to consider whether a given pedigree is for a mother-

father pair, or for a father-mother pair (i.e., we only considered nuclear markers for a potential

mother, and all markers for a potential father). The most probable pedigree for each seedling

was identified by assessing the proportion of the proposed pedigree across chains in the Gibbs

sampler (as in Moran & Clark 2011), in which we used 500,000 steps and a burn-in of 30,000.

This method was further modified to improve computational efficiency by multiprocessing

appropriate elements of the script by utilizing custom python scripts run on the VCU Center for

High Performance Computing cluster (CHiP) and the SNOW library (v0.4-2; Tierney *et al.* 2016)

in R (v3.3.3; R Core Team 2017). We replicated each run three times, and judged convergence

within and between runs in R.

Using parentage analysis to further quantify fine-scale gene flow

In addition to estimates of the mean seed and pollen dispersal from dispersal kernels

estimated during parentage analysis (see above), we used these parentage assignments to

further classify fine-scale gene flow at TEF. Using the full set of most probable pedigrees

identified from parentage analysis, we first quantified the number of in-plot vs. out-of-plot

dispersal events averaged across each replicate for a given treatment. Then, using the most

probable parentage assignment for each offspring, we quantified mean dispersal distances from

sampled mothers to seedlings, as well as between sampled fathers to sampled mothers. To

better account for uncertainty in parentage assignment (i.e., to account for fractional parentage

assignment), we also calculated mean dispersal distance by treatment by considering all

pedigrees with known individuals weighted by the probability of assignment. Specifically, for

mean seed dispersal, for each seedling we calculated the weighted average of mother-offspring

distances across pedigrees of non-zero probability that included known mothers in the dataset.

Here, each weight was the probability of assignment, $p_{seed,pedigree}$, divided by the probability of

assignment of this seedling to a known mother (1 - U_M) where U_M is the sum of the probabilities

across all non-zero pedigrees that included an unsampled mother. Treatment-level averages

were then calculated across these weighted distances. For pollen dispersal, for each seedling

we considered only pedigrees of non-zero probability where both the mother and father were

known, weighting each distance by the probability of assignment, $p_{seed,pedigree}$, divided by the

probability of assignment to known parents (1-$U_{seed,pedigree}$) where $U_{seed,pedigree}$ is the sum of

the probabilities across all non-zero pedigrees that included at least one unsampled parent.

Treatment-level averages were then calculated from these weighted distances and significance

was determined using a Kruskal-Wallis test with an alpha value of 0.05.

Scripts used in analyses described above can be found in IPython notebook format

(Pérez & Granger 2007) at https://github.com/brandonlind/teakettle.

Results

Analysis of tree spatial structure

Univariate Analysis

Across treatments, *P. lambertiana* adults generally exhibited spatial aggregation at

distance classes less than 20 meters, where this signal decreased with increasing disturbance

intensity with UN plots showing the greater magnitudes of $g_{\mathrm{inhom}}(r)$ than UC or BC plots at

these small distance classes (Figure 2 first row). For adult shade-tolerant species (*A. magnifica*

and *A. concolor* combined), the extent of spatial aggregation at large distance classes decayed

with increasing disturbance intensity (Figure 2 second row) where UC treatments generally

exhibited greater magnitudes of $g_{\mathrm{inhom}}(r)$ than BC treatments in small distance classes. For *P.*

lambertiana seedlings the spatial structure of individuals was also similar across treatments,

though UN treatments generally had significant aggregation and much larger magnitudes of

$g_{\mathrm{inhom}}(r)$ at larger distance classes than other treatments, while seedlings in BC treatments

exhibited greater magnitudes of $g_{\mathrm{inhom}}(r)$ across small distance classes than either UC or UN

plots (Figure 2 third row).

Bivariate Analysis

The spatial affinity of *P. lambertiana* seedlings to *P. lambertiana* adults,

$g_{inhom,seedling,adult}(r)$, generally ranged from randomness ($g_{inhom,seedling,adult}(r) = 1$) to spatial

inhibition ($g_{inhom,seedling,adult}(r) < 1$) with decreasing intensity of disturbance (i.e., from

undisturbed UN plots, to thin-only UC plots, to thinned-and-burned BC plots). UN plots tended to

show consistent inhibition across distance classes greater than about 15m, whereas observed

$g_{inhom,seedling,adult}(r)$ for UC plots tended to align with the lower extent of the confidence

interval with fewer instances of significant inhibition between adult and seedlings (Figure). A

similar trend for increasing spatial inhibition between *P. lambertiana* seedlings and shade-

tolerant adults ($g_{inhom,seedling,adult}(r)$), as well as for *P. lambertiana* adults and shade-tolerant

adults

UN   **UC**   **BC**

Figure 2 *Representative figures of univariate analysis of spatial structure, $g_{inhom(r)}$, by treatment replicate*
for each distance class, r. First row: adult P. lambertiana (PiLa); second row: adult shade tolerant (A.
magnifica and A. concolor = ShadeTol); third row: P. lambertiana (PiLa) seedlings. Disturbance intensity
increases by column from left to right. These figures show that with increasing disturbance intensity there
is a diminution of the degree of spatial structure within classes. Gray : null confidence envelope; Solid
black line : observed $g_{inhom}(r)$. Red dashed line : null expectation of complete spatial randomness,
$g_{inhom}(r) = 1$. Individuals are aggregated if $g_{inhom}(r) > 1$, inhibited if $g_{inhom}(r) < 1$. See Supplemental
Figures S1-S3 for all plots.

Figure 3 *Representative figures of bivariate analysis of spatial structure, $g_{inhom,i,j}(r)$, between: first row: P. lambertiana seedlings (seed) and P. lambertiana adults; second row: P. lambertiana seedlings and shade tolerant adults; third row: P. lambertiana adults to shade tolerant adults. Disturbance intensity increases by column from left to right. These figures show that the two classes compared are generally inhibited spatially by the presence of the other, and that with increasing disturbance there is a diminution of the degree of spatial inhibition between classes. Gray : null confidence envelope; Solid black line : observed $g_{inhom,i,j}(r)$. Red dashed line : null expectation of complete spatial random-ness, $g_{inhom,i,j}(r) = 1$. Individuals are aggregated if $g_{inhom,i,j}(r) > 1$, inhibited if $g_{inhom,i,j}(r) < 1$. The gray shading in the third column of the first row indicates the null confidence envelope extended beyond the limit of the y-axis, where the pattern of the confidence envelope seen in the third column of the second and third rows is caused by sample size varying among distance classes. It should be noted that the observed values for all comparisons generally fall below the y = 1 expectation except for some short distance classes. See Supplemental Figures S4-S6 for all plots.*

$(g_{inhom,PiLa-adult,shadetol}(r))$, was also observed (Figure 3) where UN treatments generally had

a greater inhibition than UC or BC plots, though BC plots also exhibited some evidence of

spatial inhibition between groups. The results from the uni- and bivariate analyses of spatial

patterns at TEF suggest that pines are generally clustered with other pines, shade-tolerant

individuals are clustered with other shade-tolerant individuals, but spatial patterns of shade-

tolerant adults generally show spatial inhibition with pine individuals of both classes.

Additionally, *P. lambertiana* seedlings showed similar clustering across all treatments,

suggesting a similar pattern of response to the environment by individual trees. Further,

together with the univariate spatial clustering of *P. lambertiana* seedlings at small distance

classes, these bivariate results suggest there may be ecological drivers influencing realized

patterns of seedlings across microenvironments at TEF (e.g., perhaps sites with decreased

competition for [or optimal levels of] water, nutrients, or light).

Diversity measures

To compare our results with measures often used across the literature to investigate

genetic effects of forest management we calculated various genetic diversity measures (see

Methods). Genetic diversity measures (Table 1) seemed to be most influenced by census size

across the various measures we estimated here. For instance, census size increased from BC

(n = 109 individuals) to UN (n = 557 individuals) to UC (n = 682 individuals) where related

diversity measures of A_{T}, A, A_e, and A_P followed this trend. Observed heterozygosity was

greatest for UN plots, followed by BC and UC plots, while expected heterozygosity decreased

from UC to BC to UN (Table 1). Thus, no trend was observed between diversity measures and

increasing disturbance treatment at TEF.

Hierarchical *F*-statistics were calculated with nuclear markers to compare the extent of

fixation within and across treatment types, with individuals nested in replicates, replicates

nested in treatments, and treatments nested within TEF. The overall multilocus F_{ST} ($F_{rep,TEF}$) at

TEF was 0.075, consistent with estimates of many *Pinus* species across various spatial scales

(Howe et

Table 1 *Genetic diversity measures (standard deviation) by treatment. N : census number of individuals*
[adults, seedlings]; A_T : total number of alleles; A : mean number of alleles per locus; A_e : effective
number of alleles (harmonic mean across loci); H_o, H_e : respectively the observed and expected
heterozygosity for nuclear markers; A_P : average number of private alleles. For A, A_e, H_o, and H_e, values
indicate averages across loci, where values for each locus were calculated across all three treatment
replicates simultaneously. H_o and H_e used only nuclear markers, whereas other genetic diversity columns

Treatment	N	A_T	A	A_e	H_o	H_e	A_P
UN	557 [236,321]	180	25.71 (6.50)	3.23 (1.58)	0.87 (0.06)	0.77 (0.06)	46
UC	682 [307,375]	210	30.00 (7.76)	6.20 (3.07)	0.57 (0.30)	0.84 (0.10)	73
BC	109 [42,67]	107	15.29 (6.80)	4.80 (2.46)	0.82 (0.08)	0.82 (0.07)	5
Mean	449 [195,254]	165.67	23.67	4.74	0.75	0.81	41.3

considered all loci.

al. 2003), suggesting that the majority of genetic variation was partitioned more so within plots

than between plots. The $F_{rep,TEF}$ for individual markers varied: rps02 ($F_{rep,TEF}$ = 0.019), rps12

($F_{rep,TEF}$ = 0.037), rps39 ($F_{rep,TEF}$ = 0.148), rps50 ($F_{rep,TEF}$ = 0.103). Considering only genotypes

across replicates of a given treatment, treatment-level estimates of $F_{rep,tx}$ also varied ($F_{rep,UN}$ =

0.011, $F_{rep,UC}$ = 0.109, $F_{rep,BC}$ = 0.035) but showed no pattern with increasing disturbance

intensity. Pairwise $F_{rep,tx}$ comparisons between treatments were calculated by considering

genotypes across two treatments simultaneously and were used to compare the extent of

fixation across disturbance intensity. Here, the three comparisons ranged from 0.050 (UC and

UN) to 0.055 (UC and BC) to 0.075 (BC and UN) indicative of increasing relative fixation with

increasing disparity for the intensity of disturbance for a given comparison.

Analysis of spatial genetic structure

Analysis of spatial genetic autocorrelation (sensu Smouse & Peakall 1999) was carried

out at TEF to better understand how treatment affects standing genetic structure (*P. lambertiana*

adults x *P. lambertiana* adults), how this standing genetic structure relates to the genetic

structure of seedlings (*P. lambertiana* seedlings x *P. lambertiana* seedlings), and the tendency

of alike genotypes to be aggregated or inhibited across the treatments as the stands continue to

develop

489   **UN**   **UC**   **BC**

491
492

493
494

Figure 4 *Analysis of spatial genetic structure (sensu Smouse & Peakall 1999) between P. lambertiana*
adults (first row), P. lambertiana seedlings (second row), and between P. lambertiana adults and
seedlings (third row) by treatment (columns) across distance classes within plots (main panel) or across
TEF (insets). Values of $r_g^h = 0$ indicate random spatial patterns of genotypes, $r_g^h > 0$ indicate clustering of
alike genotypes, and $r_g^h < 0$ indicate spatial inhibition of alike genotypes.

after treatment (*P. lambertiana* adults x *P. lambertiana* seedlings). In all comparisons, spatial

genetic structure in BC treatments did not differ significantly from a random spatial distribution of

genotypes (last column Figure), perhaps due to the relatively small census sizes (Table 1) in

distance-class bins. However, there seems to be an effect of treatment on the spatial patterning

of genotypes of adults in the UC and UN stands (first row Figure). While the natural fire-

suppressed stands (UN) exhibited small but significant spatial genetic structure for most

distance classes up to 200m, UC stands resulted in significant aggregation of adult genotypes

at a greater degree than UN up to 150m, where genotypes became spatially inhibited up to the

maximum distances in stands ($200(\sqrt{2})$m; Figure). These patterns resulted in spatial

distributions of seedling genotypes that were randomly distributed except for very short distance

classes in UN treatments, and for UC seedlings, resulted in the general pattern observed for UC

adults albeit to a higher degree of both aggregation and inhibition (second row Figure). As a

result, alike genotypes between adults and seedlings were aggregated up to 150m in UC plots,

whereas this relationship in UN treatments resulted in negative values of $r_{g,adult,seed}^h$ that

bordered the confidence envelope for spatial inhibition but were not significantly different from a

random spatial distribution of genotypes (third row of Figure 4). While the genetic structure of

adults is due to the interaction of the effect of treatment on pretreatment conditions, the long-

term dynamics of these stands will be influenced by seedling ingrowth. These results suggest

that UC treatments may, in the long term, increase the relatedness of individuals across short

spatial scales less than 150m relative to either BC or UN treatments. This will be particularly

exacerbated if gene flow occurs at similarly fine spatial scales (see below).

Quantifying fine-scale gene flow

In-plot vs. out-of-plot dispersal events

To understand how gene flow across plots is influenced by treatment, we quantified the

number of in-plot and out of plot dispersal events from pedigrees identified as

527
Figure 5 *Mother-offspring dispersal events by treatment for (A) dispersal between in-plot individuals, (B)*
dispersal into plot from an out-of-plot mother, and (C) the ratio of these values. There were no events in
which a known mother dispersed seed to another plot, therefore B is utilizing information from parentage
analysis that indicated the mother of a given seedling was not sampled. Orange letters within each plot

show significant differences between medians, as inferred from separate Kruskal-Wallis tests (see main
text of Results). Vertical lines indicate standard deviations.

most probable from our parentage analysis. To account for sample size differences, we also

calculated the ratio of these values. The number of in-plot and out-of-plot dispersal events

between mother and offspring differed by treatment (Figure A-B) but not significantly so ($p >$

0.4297). The ratio of these values differed by treatment as well (Figure C), with UC having the

greatest proportion of in-plot dispersal events but overall there were no significant differences

among treatments ($p = 0.1926$).

We next quantified the number of in-plot and out-of-plot dispersal events of pollen from

the most probable pedigrees identified from parentage analysis. In these cases, out-of-plot

pollen dispersal events were tallied as an in-plot mother receiving pollen from an unsampled or

out-of-plot father. As with mother-offspring dispersal events we also calculated a ratio of these

values. The UC treatment exhibited the most in-plot pollen dispersal events, followed by UN and

BC (Figure A), though these comparisons were not significant using a Kruskal-Wallis test ($p =$

0.5073). UN and UC treatments exhibited similar levels of out-of-plot dispersal events (Figure

B), which differed (though not significantly, $p = 0.1376$) from BC out-of-plot events. The ratio of

in-plot vs. out-of-plot dispersal events increased with increasing disturbance (Figure C) but did

not differ significantly ($p = 0.1030$).

Median dispersal distances by treatment

Considering the most probable parentage from our model, we calculated the median

seed dispersal distances between offspring and known mothers, as well as between the median

pollen dispersal between known mothers and fathers (see Methods). Median seed dispersal

varied by treatment, being greatest for UN and decreasing with increasing disturbance intensity

(Figure A). Results from a Kruskal-Wallis test indicated significant differences between groups

($p = 0.0480$), with *post hoc* tests indicating significant differences between UN and BC ($H =$

4.34, $p = 0.0372$) but not between UN and UC ($H = 2.77$, $p = 0.0959$) or between UC and BC (H

$= 2.75$, $p = 0.0970$; Figure A). Median pollen dispersal also varied by treatment, being greatest

for BC treatments, followed by UN and UC treatments, which did not differ significantly (*p* =

0.1381; Figure B).

Figure 6 *Father-mother dispersal events by treatment for (A) dispersal between in-plot individuals, (B)*
dispersal into plot from an out-of-plot mother, and (C) the ratio of these values. Plot-level tallies were
those of in-plot mothers receiving pollen from either an in-plot father (A) or an out-of-plot (sampled or
unsampled) father (B). Orange letters within each plot show significant differences between medians, as
inferred from Kruskal-Wallis tests (see main text of Results). Vertical lines indicate standard deviations.

569
Figure 7 *Dispersal distances for seed (A) and pollen (B) calculated from the most probable pedigree from*
parentage analysis, considering only pedigrees with known mothers (A) or known parents (B). Orange
letters within each plot show significant differences between medians, as inferred from Kruskal-Wallis
tests for mother-offspring and father-mother dispersal distances (see main text of Results).

These realized distances were roughly in line with, but smaller than, mean dispersal distances

estimated from dispersal kernel shape parameters in the parentage analysis: mean seed

dispersal = 65m (95% credible interval: 57-75); mean pollen dispersal = 170m (95% CI: 150-

190; Figure).

To consider uncertainty in parentage assignment, we calculated weighted average

dispersal distances for seed and pollen dispersal (see Methods). Assignments to mothers of

out-of-plot adults were less common than for assignments to in-plot fathers, as can be seen

from the blocks (replicates) within treatment of Figure . Using this set of fractional parentage, we

calculated weighted average distances for each seed and nested these distances within

treatments (see Methods). We first considered mother-offspring and father-mother dispersals

from fractional parentage where the identified adults could originate in any treatment at TEF.

Distances differed significantly by treatment (Figure A; $H = 7.91$, $p = 0.0191$) where UN and UC

were significantly different ($H = 8.11$, $p = 0.0044$) but not between any other comparison (H

range = [0.0042,0.6755], $p > 0.4111$). Father-mother distances (Figure B) also differed by

treatment ($H = 41.16$, $p = 1.15E-9$), with median dispersal distance decreasing from BC to UN to

590
Figure 8 *Fitted 2D-t dispersal kernel for seed (red) and pollen (black) using shape parameters inferred*
from parentage analysis (sensu Moran & Clark 2011). Dashed lines show the 95% credible interval. This
figure is truncated at the maximum distance within plots ($200\sqrt{2}m$) to focus on differences at short
distances.

UC, where all pairwise considerations were also significant (*H* range = [5.21,27.18], *p* range =

[1.85E-07, 0.0224]).

Because the proximity of the treatment replicates at TEF may interact with dispersal

distance estimates, we also considered dispersal distances within plot tallied within treatments

using weighted distances as described above. Median values of mother-offspring in-plot

distances decreased with increasing disturbance intensity (Figure C) and differed by treatment

(*H* = 47.10, *p* = 5.91E-11), but only between UN and UC (*H = 4.29*, *p* = 0.0382) and between

UN and BC (*H* = 5.83, *p* = 0.0253) and not between UC and BC treatments (*H* = 0.95, *p* =

0.3291). In-plot father-mother distances (Figure D) were significantly different across treatments

(*H* = 13.89, *p* = 0.0010), with BC having greater distances that either UN (*H* = 5.83, *p* = 0.0157)

or UC (*H* = 5.07, *p* = 0.0242), and UC exhibiting greater distances than UN (*H* = 5.00, *p* =

0.0253).

Figure 9 *Fractional parentage across parentage analysis cycles for (A) maternal assignment and (B)*
paternal assignment (see Methods) with adult individuals along x-axes and seedling individuals along y-
axes. Each cell represents the fraction of the cycles a particular seedling was assigned to a given adult
(black ~ 0 to red to orange to yellow to white ~1).

618   ***Figure 10*** *Dispersal distances between mothers and offspring (first column) and between fathers and*
619   *mothers (second column) using assigned adults from any location (A-B) and for only in-plot individuals (C-*
620   *D). Orange letters within each plot show significant differences between medians, as inferred from*
621   *separate Kruskal-Wallis tests (see main text of Results).*

Discussion

Frequent fires were commonplace in historic forests of the Sierra Nevada, where forests

exhibited relatively lower tree densities and a higher proportion of pine species (North *et al*.

2005; Knapp *et al.* 2013). Yet post-settlement fire suppression has led to forest densification

that has caused instability in these systems and has increased the chances of uncharacteristic

high-severity wildfire. As a result, thinning prescriptions are used to increase the resilience of

constituent stands (SNFPA 2004; Agee & Skinner 2005; Schwilk *et al.* 2009; Safford *et al.*

2009). While these prescriptions can mimic the density-reducing effects of fire, and reduce fire

severity, it is currently unknown how thinning, in isolation or through its interaction with

managed fire, will alter evolutionary dynamics of ecologically important species such as *P.*

lambertiana (SNEP 1996). We characterized spatial and genetic structure of a fire-suppressed

forest treated with a common thinning prescription in the Sierra Nevada and compared this with

its interaction with prescribed fire and with a no-thin-no-fire control treatment. Our results

suggest that spatial structure of constituent species is a result of the interaction between

treatment and ecology where pines are often clustered with other pines, shade-tolerant trees

are often clustered with other shade-tolerant trees, and pine seedlings often are inhibited by

both adult pine and shade-tolerant individuals. While genetic diversity statistics are informative

of stand-level diversity, they are less informative regarding ongoing evolutionary dynamics as a

result of treatment as they do little to predict inbreeding of future generations nor the scale at

which mating events are to occur. Used in isolation, diversity indices leave researchers to

speculate about ongoing processes and future outcomes, while monitoring of processes that

affect fitness provides more meaningful inferences which can be directly used by land

managers.

From the analysis of spatial genetic structure (*sensu* Smouse & Peakall 1999), and

despite spatial inhibition between adults and seedlings across treatments, our results suggest

that unburned thinned stands (UC treatments) result in the increase of fine-scale similarity of

adult to seedling genotypes relative to control (UN treatments) or thinned-and-burned stands

(BC treatments). Parentage analysis offered additional quantification of fine-scale gene flow and

suggested that effective seed and pollen dispersal within plots generally decreased and

increased, respectively, with the increasing intensity of disturbance, perhaps due to an increase

in microsite suitability for *P. lambertiana*, or for adults, the availability of potential mates. Our

results were measured from individuals remaining or regenerating 13 years post-treatment, very

near the historical fire return interval for this area. Thus, ongoing dynamics should be monitored,

and will likely change through time, as stands with different treatments continue to develop and

respond to subsequent disturbances such as fire.

The genetic effects of forest management

With some exceptions, studies investigating the genetic consequences of forest

management have centered around the impact on genetic diversity indices (see Table 1 in

Ratnam *et al*. 2014). This focus is likely due to the fact that highly outcrossing tree species often

suffer from elevated inbreeding depression, where survival and reproduction of subsequent

generations may be impacted. In such cases, genetic diversity has been used as an index for

evolutionary potential, likely attributable to the consequences of the relative contribution of

additive genetic variance to phenotypic variance (i.e., narrow-sense heritability) in the breeder's

equation (Lynch & Walsh 1998), but the use of heritability itself as a measure of evolvability

comes with important caveats (e.g., see Hansen *et al*. 2011). Further, such diversity indices

have been used to assess the relative reduction of alleles due to harvest intensity, where the

removal of individuals from stands will likely reduce the diversity of alleles present. Here,

management resulting in population bottlenecks is of concern. While these premises are

important to investigate, the use of genetic diversity indices as the sole method for inference of

management impact are limiting with regard to evolutionary outcomes. If the focus is to be on

management impact on evolutionary potential, processes that influence evolutionary fitness

should be investigated instead (e.g., mating systems, effective dispersal, fecundity, spatial

genetic structure, pollen pool heterogeneity, juvenile survival; Lowe *et al.* 2015). Many traits with

fitness consequences in trees are of a polygenic basis (Lind *et al.* 2018), where any given

underlying positive-effect locus has minimal influence on the trait. In such cases, fixation (as

measured by a handful of putatively neutral markers) at some of the underlying causative loci

can be ameliorated by selection for combinations of alleles at other loci. Therefore, while alleles

with little to no effect on fitness are informative for demographic processes, these should not be

conflated with loci under selection, particularly loci under strong negative selection with

important implications for inbreeding depression. Such neutral markers could be better utilized

in assessing consequences *within* processes that directly affect fitness. However, in cases

where spatial genetic relatedness is increased as a result of management, or individuals

become increasingly sparse, wasted reproductive effort (e.g., embryo abortion, or high juvenile

mortality) due to increased instances of consanguineous or self-mating events may play an

important role in ongoing population dynamics (Kärkkäinen *et al.* 1999; Sorensen 2001),

particularly when seed rain of heterospecifics exceeds effective reproductive output of historical

or ecologically important species (e.g., as for *P. lambertiana* at TEF, Zald *et al.* 2008).

Dispersal dynamics of tree species

692   The analysis of spatial genetic structure and gene flow within and across populations of

trees can elucidate ongoing evolutionary dynamics, as this spatial structure is a result of

selective and neutral processes acting across temporal and spatial scales (Hardy & Vekemans

1999; Oddou-Muratorio *et al.* 2004; Robledo-Arnuncio *et al.* 2004; Oddou-Muratorio *et al.* 2011).

Thus, quantifying dispersal and mating system is an important component in understanding

such patterns. There are multiple biological and ecological factors that shape dispersal

dynamics and resulting mating systems, such as population density, degree of fragmentation,

manner of pollination (e.g., anemophily, entomophily, or zoophily), relative reproductive output,

phenotype (such as crown shape or height), interannual climatic variation, as well as stochastic

variables such as wind direction and strength (Burczyk *et al*. 1996; Dow & Ashley 1998;

Robledo-Arnuncio *et al*. 2004, Burczyk *et al*. 2004; O'Connell *et al*. 2004). Compared with

herbaceous and annual plants, trees have more extensive gene flow (Hamrick *et al*. 1992),

though such distances are idiosyncratic to a given population, species, and system. For

instance, estimates of pollen dispersal for *Pinus sylvestris* varied from between 17-29m based

on paternity assignment (Robledo-Arnuncio *et al*. 2004) to 136m (Robledo-Arnuncio & Gil 2005)

using the TwoGener method (Smouse *et al*. 2001) where 4.3% of mating events came from

pollen dispersed over 30km (Petit & Hampe 2006; Savolainen *et al*. 2007). Seed dispersal

distances can also vary idiosyncratically, particularly for winged seeds or those that are also

dispersed by animals, such as with *P. lambertiana*.

Spatial genetic structure will be a function of these dispersal consequences as well as

their ecological interaction with the environment. While much of the quantification of such

structure in trees has been carried out at regional or continental scales, examples exist for

investigations at fine spatial scales below a few hundred meters. For instance, Marquardt *et al*.

(2007) assessed spatial genetic structure of eastern white pine (*Pinus strobus* L.) as a function

of management influence at Menominee Indian Reservation in northeastern Wisconsin. While

spatial genetic structure within 100m differed by population, the strongest autocorrelation

occurred at the least disturbed site (Marquardt *et al*. 2007). However, while they sampled both

adults and natural regeneration they did not distinguish these two groups when inferring spatial

genetic structure. Conversely, in Norway spruce (*Picea abies* L. Karst.) populations of northern

Italy, Scotti *et al*. (2008) assessed spatial genetic structure of mitochondrial (maternally

inherited) and chloroplast (paternally inherited) loci across both adults and saplings. While

chloroplast haplotypes were uncorrelated across most distance classes up to 90m for both

classes, the maternally inherited mitochondrial markers showed strong affinity below 30m,

where this affinity was greater for saplings than for adults. This pattern was seen for *P.*

lambertiana individuals at TEF as well, where both adults and seedlings were genetically

structured at small distance classes in UC treatments, though seedling genotypes were

clustered to a higher degree than adults (Figure). To our knowledge however, few instances in

the literature compare both spatial structure of trees with spatial genetic structure of tree

genotypes. At TEF, seedlings were clustered at fine spatial scales across all treatments likely

due to microsite suitability (as most cached seeds will likely persist only in suitable sites), but

were only clustered genetically in UC treatments. As such, without genotypic data, investigators

may be lead to spurious conclusions where it may be assumed that clustering of individuals also

indicates clustering of genotypes. Further, ingrowth of *P. lambertiana* in UC treatments will likely

be more related to nearby individuals, which may cause inbreeding and embryo abortion to a

greater degree in subsequent generations than in other stands at TEF.

Management implications

Our results suggest that management is affecting dispersal through the availability of

suitable microsites for seedling establishment, as well as through the availability of mates. As

disturbance intensity increased at TEF, mean effective seed dispersal generally decreased

while effective pollen dispersal generally increased (Figure A-B), likely due to the proximity of

suitable (e.g., unshaded) microsites and the availability of potential mates, respectfully. Using

the inferred dispersal kernels (Figure), the vast majority of dispersal occurs across small

distance classes, with the estimated probability of dispersal of pollen below 150m accounting for

more than 90.2% of pollen dispersal events, while dispersal of seed below 50m (150m)

accounts for 87.3% (99.2%) of dispersal events across TEF. Such a dispersal tendency will

drive spatial genetic structure and will interact with environment (including management) to

ultimately determine the patterns we observe across the landscape. Because UC treatments

generally resulted in an increased spatial affinity of alike genotypes between adults and

seedlings (Figure), short-term dynamics (decadal scales) may be dominated by mating events

between related individuals. However, long-term dynamics will likely affect this structure as well.

The strong levels of spatial genetic structure observed in seedlings have been shown to

decrease in adult stages because of self-thinning processes in other tree species (Hamrick *et al.*

1993; Epperson & Alvarez-Buylla 1997; Chung *et al.* 2003; Oddou-Muratorio *et al.* 2004), and

may well occur at TEF as well. Even so, such consequences are dependent upon initial

structure that may vary to differing degrees in undisturbed stands, or across the landscape.

Long-term dynamics should be monitored as these stands continue to develop and respond to

contemporaneous ecological pressures.

Conclusion

Understanding how thinning and fire prescriptions intended to decrease fire severity and

restore ecosystem resilience influence evolutionary dynamics of historically dominant and

ecologically important pine species is of paramount significance. We found that treatment of fire-

suppressed populations of *P. lambertiana* differentially affects fine-scale spatial and genetic

structure, and that seed and pollen dispersal increase and decrease, respectively, with

disturbance intensity. Such dynamics are likely to remain unequilibrated in the short term, and

therefore management would benefit from further monitoring of evolutionary dynamics that

affect fitness in these forests (e.g., reproductive output, survival of seedlings). Further

monitoring across broader spatial scales would also inform how these management

prescriptions affect dynamics across a greater extent of environmental heterogeneity and how

these evolutionary dynamics vary by locality. Such information will allow management to

prescribe   treatments   in   a   regionally-   and   site-specific   manner.

Acknowledgments

The authors wish to thank Britta Austin, Casey Harless, Erin Hobson, Nate Stearrett, and

Alexandrea Stylianou for help with field and lab work (and for being great people), VCU's Center

for High Performance Computing, and to VCU for start-up funds awarded to AJE.

References

Adamack AT, Gruber B (2014) PopGenReport: simplifying basic population genetic analyses in
R. Methods in Ecology & Evolution 5:384-387.

Adams WT, Zuo J, Shimizu JY, Tappeiner JC (1998) Impact of alternative regeneration
methods on genetic diversity in coastal Douglas-fir. Forest Science 44:390–396.

Agee JK, Skinner CN (2005) Basic principles of forest fuel reduction treatments. Forest Ecology
and Management 211:83–96.

Baddeley A, Rubak E, Turner R (2015) Spatial point patterns: Methodology and applications
with R. London: Chapman and Hall/CRC press, 2015. URL:
http://www.crcpress.com/Spatial-Point-Patterns-Methodology-and-Applications-with-
R/Baddeley-Rubak-Turner/9781482210200/

Bonello P, Gordon TR, Herms DA, Wood DL, Erbilgin N (2006) Nature and ecological
implications of pathogen-induced systemic resistance in conifers: A novel hypothesis.
Physiological and Molecular Plant Pathology 68: 95–104.

Buchert GP, Rajora OP, Hood JV, Dancik BP (1997) Effects of Harvesting on Genetic Diversity
in Old-Growth Eastern White Pine in Ontario, Canada. Conservation Biology 11:747–758.

Burczyk J, Adams WT, Shimizu JY (1996) Mating patterns and pollen dispersal in a natural
knobcone pine (*Pinus attenuata* Lemmon.) stand. Heredity 77:251–260.

Burczyk J, DiFazio SP, Adams WT (2004) Gene flow in forest trees: how far do genes really
travel? Forest Genetics 11:179.

Cheliak WM, Murray G, Pitel JA (1988) Genetic effects of phenotypic selection in white spruce.
Forest Ecology and Management 24:139–149.

Chung MY, Epperson BK, Chung MG (2003) Genetic structure of age class in *Camellia japonica*
(Theaceae). Evolution 57:62–73.

Cloutier D, Kanashiro M, Ciampi AY, Schoen DJ (2006) Impact of selective logging on
inbreeding and gene dispersal in an Amazonian tree population of *Carapa guianensis* Aubl.
Molecular Ecology 16:797–809.

Collins BM, Everett RG, Stephens SL (2011) Impacts of fire exclusion and recent managed fire
on forest structure in old growth Sierra Nevada mixed-conifer forests. Ecosphere 2:1–14.

Covington WW, Everett RL, Steele R et al. (1994) Historical and anticipated changes in forest
ecosystems of the inland west of the United States. Journal of Sustainable Forestry 2:13–
63.

Dyer (2016) gstudio: Tools related to the spatial analysis of genetic marker data. R package
version 1.5.0. https://dyerlab.ces.vcu.edu/2015/01/16/gstudio/

Echt CS, MayMarquardt P, Hseih M, Zahorchak R (1996) Characterization of microsatellite
markers in eastern white pine. Genome 39:1102–1108.

El-Kassaby YA, Dunsworth BG, Krakowski J (2003) Genetic evaluation of alternative silvicultural
systems in coastal montane forests: western hemlock and amabilis fir. Theoretical &
Applied Genetics 107:598–610.

Epperson BK, Alvarez-Buylla ER (1997) Limited seed dispersal and genetic structure in life
stages of *Cecropia obtusifolia*. Evolution 51:275–282.

Fageria MS, Rajora OP (2013a) Effects of harvesting of increasing intensities on genetic
diversity and population structure of white spruce. Evolutionary Applications 6:778–794.

Fageria MS, Rajora OP (2013b) Effects of silvicultural practices on genetic diversity and
population structure of white spruce in Saskatchewan. Tree Genetics & Genomes 10:287–
296.

Farris MA, Mitton JB (1984) Population density, outcrossing rate, and heterozygote superiority
in ponderosa pine. Evolution 8:1151–1154.

Finkeldey R, Ziehe M (2004) Genetic implications of silvicultural regimes. Forest Ecology and
Management 197:231–244.

Fowells HA, Schubert GH (1956) Seed crops of forest trees in the pine region of California.
USDA Forest Service, Technical Bulletin 150, Washington, D.C.

Gömöry D (1992) Effect of stand origin on the genetic diversity of Norway spruce (*Picea abies*
Karst.) populations. Forest Ecology and Management 54:215–223.

Hamrick JL, Godt MJW, Sherman-Broyles SL (1992) Factors influencing levels of genetic
diversity in woody plant species. New Forests. 6:95–124.

Hamrick JL, Murawsky DA, Nason JD (1993) The influence of seed dispersal mechanisms on
the genetic structure of tropical tree populations. Vegetatio 107-108:281–297.

Hansen TF, Pelabon C, Houle D (2011) Heritability is not evolvability. Evolutionary Biology
38:258-277

Hardy OJ, Vekemans X (1998) Isolation by distance in a continuous population: reconciliation
between spatial autocorrelation analysis and population genetics models. Heredity 83:145-
154.

Holsinger KE, Weir BS (2009) Genetics in geographically structured populations: defining,
estimating and interpreting *F*. Nature Reviews Genetics 10:639–650.

Jerome Goudet and Thibaut Jombart (2015). hierfstat: Estimation and Tests of Hierarchical F-
Statistics. R package version 0.04-22. https://CRAN.R-project.org/package=hierfstat

Kärkkäinen K, Savolainen O, Koski V (1999) Why do plants abort so many developing seeds:
bad offspring or bad maternal genotypes? Evolutionary Ecology 13:305-317.

Kramer AT, Ison JL, Ashley MV, Howe HF (2008) The Paradox of Forest Fragmentation
Genetics. Conservation Biology 22:878–885.

Knapp EE, Skinner CN, North MP, Estes BL (2013) Long-term overstory and understory change
following logging and fire exclusion in a Sierra Nevada mixed-conifer forest. Forest Ecology
and Management 310:903–914.

Larson AJ, Churchill D (2012) Tree spatial patterns in fire-frequent forests of western North
America, including mechanisms of pattern formation and implications for designing fuel
reduction and restoration treatments. Forest Ecology and Management 267:74–92.

Ledig FT (1988) The conservation of diversity in forest trees: why and how should genes be
conserved? BioScience 38:471–479.

Ledig FT (1992) Human impacts on genetic diversity in forest ecosystems. Oikos 63:87–108.

Lind BM, Menon M, Bolte CE, Faske TM, Eckert AJ (2018) The genomics of local adaptation in
trees: are we out of the woods yet? Tree Genetics & Genomes 14:29.

Lowe AJ, Cavers S, Boshier D, Breed MF, Hollingsworth PM (2015) The resilience of forest
fragmentation genetics - no longer a paradox - we were just looking in the wrong place.
Heredity 115:97–99.

Lynch M, Walsh B (1998) Genetics and analysis of quantitative traits. Sinauer Sunderland, MA

Macdonald SE, Thomas BR, Cherniawsky DM, Purdy BG (2001) Managing genetic resources of
lodgepole pine in west-central Alberta: patterns of isozyme variation in natural populations
and effects of forest management. Forest Ecology and Management 152:45–58.

Marquardt PE, Echt CS, Epperson BK, Pubanz DM (2007) Genetic structure, diversity, and
inbreeding of eastern white pine under different management conditions. Canadian Journal
of Forest Research 37:2652–2662.

Millar C, Graumlich LJ, Delany DL, Westfall RD, King JC (2002) Response of high-elevation
conifers. In: Proceedings of the Eighteenth Annual Pacific Climate Workshop (p. 57).

Miller JD, Safford HD, Crimmins M, Thode AE (2009) Quantitative Evidence for Increasing
Forest Fire Severity in the Sierra Nevada and Southern Cascade Mountains, California and
Nevada, USA. Ecosystems 12:16–32.

Moran EV, Clark JS (2011) Estimating seed and pollen movement in a monoecious plant: a
hierarchical Bayesian approach integrating genetic and ecological data. Molecular Ecology
20:1248–1262.

Mouillot D, Graham NAJ, Villéger S, Mason NWH, Bellwood DR (2013) A functional approach
reveals community responses to disturbances. Trends in Ecology & Evolution 28:167–177.

Namroud M-C, Bousquet J, Doerksen T, Beaulieu J (2012) Scanning SNPs from a large set of
expressed genes to assess the impact of artificial selection on the undomesticated genetic
diversity of white spruce. Evolutionary Applications 5:641–656.

Neale DB (1983) Population genetic structure in Douglas-fir shelterwood regeneration system in
southwest Oregon. PhD dissertation. Oregon State University.

Neale DB, Adams WT (1985) The mating system in natural and shelterwood stands of Douglas-
fir. Theoretical & Applied Genetics 71:201–207.

NIFC (2014) National Interagency Coordination Center Incident Management Situation Report -
October 2014. National Interagency Fire Center, 1–7.

NIFC (2015) National Interagency Coordination Center Incident Management Situation Report -
September 2015. National Interagency Fire Center, 1–9.

North MP (2002) The Teakettle Experiment. Proceedings of a Symposium on the Kings River
Sustainable Forest Ecosystems Project: progress and current status: January 26, 1998,
Clovis, California, 47.

North, M, Collins, B, Safford, H, and Stephenson, N (2016) Montane Forests. Pages 553-578
in H. Mooney and E. Zavelta, (eds.) Ecosystems of California, U.C. Press, Berkeley, CA.
984 pp.

North MP, Hurteau MD, Fiegener R, Barbour M (2005) Influence of fire and El Nino on tree
recruitment varies by species in Sierran mixed conifer. Forest Science 51:187–197.

North MP, Stephens SL, Collins BM et al. (2015) Reform forest fire management. Science
349:1280–1281.

O'Connell LM, Russell J, Ritland K (2004) Fine-scale estimation of outcrossing in western
redcedar with microsatellite assay of bulked DNA. Heredity 93:443–449.

Oddou-Muratorio S, Demesure-Musch B, Pelissier R (2004) Impacts of gene flow and logging
history on the local genetic structure of a scattered tree species, *Sorbus torminalis* L.
Crantz. Crantz. Molecular Ecology 132), 3689–3702.

Oddou-Muratorio S, Klein EK, Vendramin GG, Fady B (2011) Spatial vs. temporal effects on
demographic and genetic structures: the roles of dispersal, masting and differential
mortality on patterns of recruitment in *Fagus sylvatica*. Molecular Ecology 20:1997–2010.

Pérez F, Granger BE (2007) IPython: A system for interactive scientific computing. Computing
in Science and Engineering 9: 21-29.

Perry DJ, Bousquet J (2001) Genetic diversity and mating system of post-fire and post-harvest
black spruce: an investigation using codominant sequence-tagged-site (STS) markers.
Canadian Journal of Forest Research 31:32–40.

Petit RJ, Hampe A (2006) Some evolutionary consequences of being a tree. Annual Review of
Ecology 37:187–214.

R Core Team (2017) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rajora OP, Pluhar SA (2003) Genetic diversity impacts of forest fires, forest harvesting, and
alternative reforestation practices in black spruce (*Picea mariana*). Theoretical and Applied
Genetics 106:1203–1212.

Rajora OP, Rahman MH, Buchert GP, Dancik BP (2000) Microsatellite DNA analysis of genetic
effects of harvesting in old-growth eastern white pine (*Pinus strobus*) in Ontario, Canada.
Molecular Ecology 9:339–348.

Ratnam W, Rajora OP, Finkeldey R, Aravanopoulos F, Bouvet J-M, Vaillancourt RE, Kanashiro
M, Fady B, Tomita M, Vinson C (2014) Forest Ecology and Management. Forest Ecology
and Management 333:52–65.

Richardson JL, Urban MC, Bolnick DI, Skelly DK (2014) Microgeographic adaptation and the
spatial scale of evolution. Trends in Ecology & Evolution 29:165–176.

Robledo-Arnuncio JJ, Alía R, Gil L (2004) Increased selfing and correlated paternity in a small
population of a predominantly outcrossing conifer, *Pinus sylvestris*. Molecular Ecology
13:2567–2577.

Robledo-Arnuncio JJ, Gil L (2004) Patterns of pollen dispersal in a small population of *Pinus
sylvestris* L. revealed by total-exclusion paternity analysis. Heredity 94:13–22.

Safford HD, Schmidt DA, Carlson CH (2009) Effects of fuel treatments on fire severity in an area
of wildland–urban interface, Angora Fire, Lake Tahoe Basin, California. Forest Ecology and
Management 258:773–787.

Savolainen O, Pyhäjärvi T, Knürr T (2007) Gene Flow and Local Adaptation in Trees. Annual
Review of Ecology, Evolution, and Systematics 38:595–619.

Schwilk DW, Keeley JE, Knapp EE et al. (2009) The National Fire and Fire Surrogate study:
effects of fuel reduction methods on forest vegetation structure and fuels. Ecological
applications 19:285–304.

Scotti I, Gugerli F, Pastorelli R, Sebastiani F, Vendramin GG (2008) Maternally and paternally
inherited molecular markers elucidate population patterns and inferred dispersal processes

on a small scale within a subalpine stand of Norway spruce (*Picea abies* [L.] Karst.). Forest
ecology and management 255:3806-3812.

Sierra Nevada Forest Plan Amendment – SNFPA (2004) Final Environmental Impact
Statement: vols.1–6. USDA Forest Service, Pacific Southwest Region, Vallejo, CA.

Sierra Nevada Ecosystem Project - SNEP (1996) Final Report to Congress: vols.1–3. University
of California, Centers for Water and Wildland Resources, Davis, CA.

Smouse PE, Dyer RJ, Westfall RD, Sork VL (2001) Two-generation analysis of pollen flow
across a landscape. I. Male gamete heterogeneity among females. Evolution 55:260–271.

Smouse PE, Peakall R (1999) Spatial autocorrelation analysis of individual multiallele and
multilocus genetic structure. Heredity 82:561–573.

Sorenson FC (1983) Geographic variation in seedling Douglas-fir (*Pseudotsuga menziesii*) from
the western Siskiyou Mountains of Oregon. Ecology 64:696-702.

Thomas BR, Macdonald SE, Hicks M, Adams DL, Hodgetts RB (1999) Effects of reforestation
methods on genetic diversity of lodgepole pine: an assessment using microsatellite and
randomly amplified polymorphic DNA markers. Theoretical and Applied Genetics 98:793–
801.

Tierney L, Rossini AJ, Li N, Sevcikova H (2016) snow: Simple Network of Workstations. R
package version 0.4-2. https://CRAN.R-project.org/package=snow

USDA Forest Service (2004) Sierra Nevada Forest Plan Amendment. Final Supplemental
Environmental Impact Statement, Record of Decision. Pacific Southwest Region, R5-MB-
046, 1–72.

USDA Forest Service (2016a) Fiscal Year 2016 Budget Overview. 1–57.

USDA Forest Service (2016b) Fiscal Year 2016 Budget Justification. 1–553.

Verner J, McKelvey KS, Noon BR, Gutierrez RJ, Gould Jr., GI, Beck TW (1992) The California
spotted owl: a technical assessment of its current status. General Technical Report PSW-
133. USDA Forest Service, Pacific Southwest Research Station, Albany, CA, 285 pp.

Weir BC, Cockerham CC (1984) Estimating *F*-statistics for the analysis of population structure.
Evolution, 1358-1370.

Westerling AL (2006) Warming and Earlier Spring Increase Western U.S. Forest Wildfire
Activity. Science 313:940–943.

Wofford AM, Finch K, Bigott A, Willyard A (2014) A Set of Plastid Loci for Use in Multiplex
Fragment Length Genotyping for Intraspecific Variation in *Pinus* (Pinaceae). Applications in
Plant Sciences 2:1400002.

Wright J (1976) Introduction to forest genetics. Academic Press, New York.

Zald HSJ, Gray AN, North MP, Kern RA (2008) Initial tree regeneration responses to fire and
thinning treatments in a Sierra Nevada mixed-conifer forest, USA. Forest Ecology and
Management 256:168–179.