Title: Reorganization of thalamocortical connections in congenitally blind humans

Running title: Thalamic plasticity in congenital blindness

Authors: Theo F. Marins[a,b], Maite Russo[c], Erika C. Rodrigues[a], Marina Monteiro[a], Jorge Moll[a], Daniel Felix[a], Julia Bouzas[a], Helena Arcanjo[d]; Claudia D. Vargas[c], Fernanda Tovar-Moll[a,b,1]

[a] D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil
[b]Post-Graduation Program in Morphological Sciences (PCM) of the Institute of Biomedical Sciences (ICB), Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil; [c]Institute of Biophysics Carlos Chagas Filho (IBCCF), Federal University of Rio de Janeiro (UFRJ); [d]Centro de Oftalmologia Especializada

[1]Corresponding Author: fernanda.tovarmoll@idor.org

ABSTRACT

Cross-modal plasticity in blind individuals has been reported over the past decades showing that non-visual information is carried and processed by 'visual' brain structures. However, the structural underpinnings of cross-modal plasticity in congenitally blind individuals remain unclear despite multiple efforts. We mapped thalamocortical connectivity and assessed cortical thickness and integrity of white matter of ten congenitally blind individuals and ten sighted controls. We hypothesized an aberrant thalamocortical pattern of connectivity taking place in the absence of visual stimuli from birth as a potential mechanism of cross-modal plasticity. In addition to the increased cortical thickness of the primary visual cortex and reduced integrity of visual white matter bundles, we observed structural connectivity changes between the thalamus and occipital and temporal cortices. Specifically, the thalamic territory dedicated to connections with the occipital cortex was smaller and displayed weaker connectivity in congenitally blind individuals. In contrast, those connecting with the temporal cortex showed greater volume and increased connectivity compared to sighted controls. The abnormal pattern of thalamocortical connectivity included the lateral and medial geniculate nuclei and the pulvinar nucleus. For the first time in humans, a remapping of structural thalamocortical connections involving both unimodal and multimodal thalamic nuclei has been demonstrated, shedding light on the possible mechanisms of cross-modal plasticity in humans. The present findings may help understand the functional adaptations commonly observed in congenitally blind individuals.

Keywords: cross-modal plasticity, congenital blindness, diffusion tensor imaging, thalamus, cortical thickness

Abbreviations: WM = White Matter; DTI = Diffusion Tensor Imaging; LGN = Lateral Geniculate Nucleus; MGN = Medial Geniculate Nucleus; Congenitally blind = CB; Sighted controls = SC.

INTRODUCTION

Blindness and vision impairment affects at least 2.2 billion people around the

world[1]. A significant part of these patients has congenital blindness, which can be caused

by congenital anomalies and infantile glaucoma, among other causes[2]. Congenital

blindness represents an intriguing model for understanding how the brain builds and

maintains its fundamental principles of organization and hierarchy of processing in the

absence of one of the major sources of input. The impact of congenital blindness on

spared sensory systems has also been a focus of study over the past years. A better

characterization of the basis of blindness-related brain alterations, including the well-

described phenomenon of cross-modal plasticity, can pave the way for developing and

optimizing new inclusive devices and brain-machine interfaces.

Numerous brain alterations can occur in response to the lack of visual input,

including in white matter (WM) and gray matter (GM) of both cortical and subcortical

brain structures[3–5]. Congenitally/early blind individuals often exhibit atrophy of optic

chiasm and reduced microstructural integrity of optic radiations and geniculocalcarine

tract[6–11], as investigated by Diffusion Tensor Imaging (DTI). Also, splenium of the corpus

callosum, inferior longitudinal fasciculus, and inferior fronto-occipital fasciculus, major

WM bundles connecting different cortical areas involved in visual processing and

perception, are impacted by the absence of visual input[10,12–16]. The way congenital and

early blindness affect the primary visual cortex structure is a matter of debate. Whereas

increased cortical thickness and volume have been reported[17–19], the opposite effects have

also been observed[6,8,20,21].

Functional neuroimaging studies have shown that visual cortical structures in the

blind brain often participate in the processing of non-visual information, which is called

cross-modal plasticity. In blind individuals, this reorganization is commonly observed as

a visual takeover by the auditory and tactile systems[22–27]. Still, it has also been reported

involving language[28,29], olfaction, and gustation[30,31] very often leading to increased

abilities and behavioral improvement in the remaining senses. In addition, recruitment of

'visual' brain areas in blindness has been reported during auditory[18,32–35], tactile[23], and

linguistic [29] tasks, and correlation with behavioral gains has also been shown[33,36–39].

Moreover, virtual lesions targeted to the 'visual' cortex induced by Transcranial Magnetic

Stimulation (TMS) can temporarily impair performance on auditory[12] and tactile[40] tasks

in blind individuals. Altogether, these findings suggest that the brain undergoes critical

changes in the absence of visual input, which pave the way for cross-modal plasticity.

While evidence corroborates cross-modal plasticity in blindness, neuroanatomical

correlates underlying this phenomenon are controversial and may involve a wide range

of brain structural changes, including the development of new connections and

unmasking /rewiring of the existing ones[41]. Indeed, it is unclear how and which pathways

convey non-visual information to the visual cortex in the blind brain. Most findings on

the possible anatomical underpinnings of cross-modal plasticity and its mechanisms come

from animal models of blindness studies. They point to the emergence of direct

connections between deprived visual, auditory, and somatosensory cortices, indicating

that non-visual information would reach the visual cortex by direct corticocortical

connectivity, a transient connection that becomes stabilized during development by the

lack of appropriate visual stimuli[42–48]. On the other hand, the preservation of

geniculocalcarine pathways in blind individuals[8,49,50] has been reported, which supports

the idea that adaptive plasticity may rely upon the conservation of thalamocortical

pathways. In the naturally blind mole rat, the remnant visual pathways carry auditory

information to the visual cortex as their visual nuclei of the thalamus receive input from

the inferior colliculus, an important auditory center[51,52] Moreover, in addition to the

preserved thalamocortical connections, visual cortical areas receive input from auditory

and somatosensory nuclei of the thalamus in blind mice and opossum[46,53]. These findings

suggest that robust changes in the thalamocortical connectivity may occur in response to

the lack of visual input and could also explain the cross-modal plasticity observed in

congenitally blind individuals.

Thus, despite remarkable changes that have been reported in the congenitally

blind brain, the structural underpinnings of cross-modal plasticity are still uncertain in

humans. In the present study, we combined neuroimaging techniques based on diffusion-

weighted imaging, probabilistic tractography, and brain morphometry to interrogate the

thalamic (re)mapping of cortical connections of congenitally blind individuals.

Specifically, we tested the hypothesis that structural changes of thalamocortical

projections involved in visual and multimodal sensory processing would occur in

response to the absence of appropriate visual input from birth. In addition, we

hypothesized that congenital blindness impairs the appropriate maturation of 'visual'

areas, reflected in WM and GM changes. Our results point to a remapping of the thalamic

connections with both temporal and occipital cortices in congenitally blind individuals

compared to a matched sample of sighted controls and an increased cortical thickness in

the primary visual cortex (V1) and impaired WM microstructure of visual structures.

METHODS

Subjects

This study was conducted in accordance with the ethical standards-compliant of

the Declaration of Helsinki and has been approved by the IDOR / Copa D'Or Ethics and

Scientific Committee. Ten congenitally blind (CB; mean age: 31.8, standard deviation:

8.7, 6 males) individuals and ten sex- and age-matched sighted controls (SC; mean age:

32.2, standard deviation: 6.66, 6 males) were included in the study. All participants were

right-handed, had no history of neurologic or psychiatric diseases, and were not taking

brain-active medication. For the inclusion of the CB individuals, their medical records

and additional clinical exams (such as the visual evoked potentials) have been reviewed.

Afterward, they were examined by an experienced ophthalmologist, confirming that they

were totally blind or had only minimal residual light sensitivity. All blind participants

were Braille readers. The cohort characteristics are summarized in Table 1.

Table 1. Cohort characteristics

Patient ID	Age	Gender	Cause of congenital blindness	Handedness	Braille
CB01	35	F	Anophthalmia	Right	Yes
CB02	36	M	Macular dystrophy	Right	Yes
CB03	32	F	Glaucoma	Right	Yes
CB04	18	M	Glaucoma	Right	Yes
CB05	40	M	Glaucoma	Right	Yes
CB06	27	M	Glaucoma	Right	Yes
CB07	22	M	Retinal detachment	Right	Yes
CB08	44	M	Unknown	Right	Yes
CB09	24	F	Retinopathy of prematurity	Right	Yes
CB10	40	F	Unknown	Right	Yes

M: male. F: female

Data Acquisition

Brain imaging was performed at D'Or Institute for Research and Education

(IDOR) in a 3T Achieva scanner (Philips Medical Systems, the Netherlands) using an

eight-channel SENSE head coil. Imaging consisted in a high-resolution 3D T1-weighted

image ($1mm^3$ isotropic, TR/TE (ms) = 7.2 / 3.4, FOV= 240 x 240, matrix = 240, slice

thickness = 1mm) and diffusion-weighted (2 x 2 x $2mm^3$ isotropic, no gap, TR/TE (ms)=

10150/60, FOV = 224 x 224, matrix = 112 x 112) with diffusion sensitization gradients

applied in 64 noncollinear directions, with a b factor of 1000 s/mm^2, one volume without

diffusion weighting.

Data Analysis

Before data preprocessing, images were visually inspected for excessive

movements or artifacts. Diffusion-weighted images were preprocessed and analyzed

using FSL[54] toolboxes. In each subject, original data were corrected for the effects of head

movement and eddy currents using eddy correct, and a brain mask was created by running

BET[55] on the B=0 (no diffusion weighting) image. We created FA images using FDT

(with DTIFIT algorithm), part of FSL[56]. All subjects' data were aligned into standard

space (Montreal Neurological Institute, MNI) using the nonlinear registration tool

FNIRT, and the mean FA image was created and thinned to create a mean FA skeleton.

The FA map of each subject was projected onto skeletonized FA, and the resulting data

was fed into voxel-wise cross-subject statistics using TBSS[57]. We tested for group

differences with an unpaired t-test using Randomise[58] for permutation-based (5000

permutations) non-parametric testing of whole skeleton FA. Also, analysis restricted to

the whole thalamus was separately conducted.

Connectivity-based segmentation of the thalamus was also performed using FDT,

as described previously[59], in each hemisphere separately. Six cortical masks were

predefined using the Harvard-Oxford atlas: Prefrontal, Precentral, Postcentral, Posterior

Parietal (parietal cortex, except for the postcentral cortex), Occipital, and Temporal. Since

the Oxford Thalamic Connectivity Probability Atlas does not include the lateral and

medial geniculate nuclei (LGN and MGN, respectively), both right and left thalamic

masks were obtained from an atlas on the Colin-27 Average Brain[60]. All masks were

registered into the subjects' space using nonlinear registration. After BEDPOSTX[56],

probabilistic tracking was conducted with the PROBTRACKX[56] tool, in which the

unilateral thalamus was defined as the seed region and the six ipsilateral cortical masks

as classification targets. Following, we calculated the number of samples reaching each
target mask as a proportion of the total number of samples reaching any target mask.
Then, the hard segmentation of each thalamus based on its connectivity to each of the six
ipsilateral cortical areas was performed. The volume of each resultant segment of the
thalamus was normalized by the volume of the ipsilateral thalamus **at** the subject level.
Group analysis of the normalized volume was performed with SPSS 20.0 (IBM
Corporation, New York) using a repeated-measures analysis of variance (ANOVA;
within-subjects factors: 'segment' and 'side'; between-subjects factor: 'group'). We used
Bonferroni to adjust for multiple comparisons.

To perform a voxel-wise analysis of the thalamocortical connectivity and compare
them between groups, the resultant six segments at the subject level were transformed
back to the MNI space. Each segment was overlapped to create a single mask, thus
consisting in a sum of all individual segment masks from both groups. The voxel-wise
analysis was conducted within this mask separately for each thalamic segment. Group
differences were investigated with an unpaired t-test using Randomise[58] for permutation-
based (5000 permutations) non-parametric testing of the connectivity maps for each
segment.

Cortical thickness analysis
Cortical thickness analysis was performed based on high-resolution 3D T1-
weighted gradient-echo images using Freesurfer software
(http://surfer.nmr.mgh.harvard.edu/). Preprocessing steps included removal of non-brain
tissue, segmentation of the subcortical white matter and deep gray matter volumetric
structures, intensity normalization, tessellation of the gray matter/white matter boundary,
automated topology correction, and surface deformation following intensity gradients.

To investigate group differences, we applied the command-line Freesurfer group
analysis stream, including correction for multiple comparisons. First, the data from every
subject was assembled by resampling each subject's data onto a common space, spatially
smoothing it, and concatenating all the subject's resampled and smoothed data into a
single file. We defined the design matrix and design contrast comparing both groups,
resulting in a t-test analysis of the whole brain. The statistical results were then submitted
to a cluster-wise correction for multiple comparisons ($p<0.05$).

For simplicity, group differences are referred to as "increase" and "decrease"
throughout the manuscript taking the SC group as a reference. The coordinates in the

figures are given according to MNI space, and results are plotted on the MNI standard
brain.
158
159 **Data availability**
160   Data that support the present findings are available from the corresponding author
161 upon reasonable request.
162
RESULTS
Reorganization of thalamocortical connections
The thalamus segmentation based on the structural connectivity patterns to six
predefined cortical areas (Prefrontal, Precentral, Postcentral, Posterior Parietal,
Temporal, and Occipital) successfully resulted in six thalamic segments on each side (Fig.
1) in all participants.
The analysis of thalamic volumes revealed an interaction effect between
'segment*group' (F(2.32, 41.80)=4.76, p=0.01). *Posthoc* analysis with Bonferroni
correction for multiple comparisons on the interaction 'segment*group' indicated that the
SC and CB groups significantly differed in the volume of the Occipital segment
(p=0.013), being reduced in the CB group; and Temporal segment (p=0.002) being
increased in the CB group (Fig. 1; Table 2).
In addition, there was a main effect of 'segment' (F(2.32, 41.80)=635,92,
p<0.005) and an interaction between 'segment*side' (F(2.56 46.01)=9.77, p<0.005).
Pairwise comparisons were run to investigate the main effect of 'segment' and interaction
'segment*side' adjusting for Bonferroni (Table S1 and S2).
179
180 Figure 1 around here.

Table 2. Normalized volume comparisons of thalamic segments

Segment	Side	SC		CB	
		Mean Volume	SD	Mean Volume	SD
Prefrontal p=0.064	R	0.561224	0.0372437	0.515953	0.0578148
	L	0.537903	0.0424005	0.505637	0.0457513
Precentral p=0.14	R	0.055635	0.0125891	0.060761	0.0150288
	L	0.067851	0.0290276	0.078154	0.0259177
Occipital* p=0.013	R	0.11052	0.0286398	0.071793	0.0451293
	L	0.138718	0.0443775	0.088397	0.0174996
Posterior Parietal p=0.97	R	0.065566	0.0267344	0.064867	0.0174419
	L	0.049694	0.026557	0.05505	0.0229696
Postcentral p=0.07	R	0.044252	0.0156764	0.040539	0.0148974
	L	0.060081	0.0209732	0.071497	0.0407492
Temporal* p=0.002	R	0.162803	0.0458977	0.246086	0.0842971
	L	0.145753	0.0363323	0.201264	0.0570562

187 *Segments that showed significant (p<0.05) between-group differences after Bonferroni
188 correction for multiple comparisons.
189 L= left thalamus; R= right thalamus. Mean volume refers to the normalized mean volume
190 in arbitrary units.
191

To investigate whether congenital blindness alters the anatomical connectivity
pattern between the thalamus and the cortex, we conducted a group comparison using
threshold-free cluster enhancement (TFCE, FWE-corrected, $p<0.05$) of the probability
maps of connectivity of each thalamic segment. This analysis revealed that the
connectivity of the thalamic segment connecting with the occipital and the temporal

cortices significantly differed between groups, with the thalamo-temporal connectivity
being increased and the thalamo-occipital connectivity being reduced in the CB group.
Specific regions showing increased thalamo-temporal connectivity in the CB group were
observed in both thalami, specifically in the bilateral LGN, bilateral pulvinar, bilateral
MGN, left medial dorsal nucleus, left anterior nucleus, left ventral anterior nucleus and
left lateral posterior nucleus (Fig. 2). On the other hand, reduced thalamo-occipital
anatomical connectivity in the CB group was restricted to the territory of the pulvinar and
lateral posterior nucleus on the left thalamus (Fig. 2). Further analysis revealed that both
group differences in thalamocortical connectivity (increased thalamo-temporal
connectivity and decreased thalamo-occipital connectivity) partially shared a common
neural territory on the left pulvinar (Fig. 2). Thalamic projections to the Prefrontal,
Precentral, Postcentral, and Posterior Parietal segments did not show statistically
significant differences between groups.

Figure 2 around here.

White Matter Microstructure Changes in Blind Subjects
We investigated differences in WM microstructure using TBSS for fractional
anisotropy (FA). Whole-brain (threshold-free cluster enhancement (TFCE), corrected
$p<0.05$) analysis showed decreased FA in the major WM structures connecting the
occipital cortex, such as the splenium of the corpus callosum, bilateral inferior
longitudinal fasciculi, bilateral inferior fronto-occipital fasciculi, and bilateral superior
longitudinal fasciculi (Fig. 3) in the CB group. Additional analysis restricted to the whole
thalamus revealed a diffuse decrease in FA in the CB group, including left and right
pulvinar, medial dorsal nucleus, right ventral lateral nucleus, right lateral posterior, and
left lateral nucleus (Fig. 3).

Figure 3 around here.

Increased Cortical Thickness in Blind Subjects

Whole-brain comparison of cortical thickness (whole-brain analysis, cluster-
corrected for multiple comparisons, $p < 0.05$; Fig. 4A) revealed greater thickness of the
left primary visual cortex (V1, Brodmann area 17 (BA17), Size= 268.24 mm^2, X=-5.2,
Y=-86.4, Z=10.3, MNI) in the CB group (CB group > SC group). This single cluster was
then masked to further investigate thickness in each participant (Fig. 4B). No other
cortical region showed group differences in thickness.

Figure 4 around here.

DISCUSSION

The present findings indicate that the absence of appropriate visual input from
birth in humans leads to a wide range of structural brain changes. For the first time, we
described a structural remapping of thalamocortical connectivity, which may help
understand the functional adaptations commonly observed in congenitally blind
individuals.

The analysis of a sample of right-handed CB individuals, Braille readers, brought
new evidence of reduced thalamic areas projecting to/from the occipital cortex compared
to SC. On the other hand, a greater volume of the thalamic territory dedicated to
connections with the temporal cortex was observed in CB compared to SC. Additionally,
the voxel-wise analysis revealed that thalamic nuclei such as bilateral LGN, MGN, and
pulvinar were more connected to the temporal cortex in CB. In contrast, left pulvinar and
lateral posterior nuclei displayed weaker projections to the left occipital cortex in this
group. Interestingly, we found strengthened connectivity of the left pulvinar to the
temporal cortex over the occipital cortex, suggesting a possible rerouting of the pulvinar
connections in CB. Moreover, we confirmed previous findings by showing that visual
cortical areas exhibit greater thickness, besides a diffuse impairment in the microstructure
of major visual WM bundles.

Our findings bring new evidence in line with previous studies showing
impairment of thalamic nuclei involved in visual and multimodal processing, such as
LGN and pulvinar. Using voxel-based morphometry (VBM), previous studies have
shown volume reductions in thalamic visual centers in CB individuals, including dorsal
LGN and pulvinar[13,61], identified with the aid of brain atlases. A similar approach has

been used to describe volume reduction in WM close to the LGN in six bilaterally anophthalmic subjects compared with sighted subjects[8]. However, it is worth noting that the present study was not limited to comparing the volume of thalamic structures between groups. Instead, we went beyond by mapping the thalamocortical projections and exploring a possible remapping of thalamic territories and their connectivity pattern to/from the cortex. Thus, one key difference between the present and previous studies is that visual-related thalamic nuclei investigated here were identified using subject-level tractography. Interestingly, the thalamic territory occupied by projections to/from the temporal cortex showed greater volume, whereas those to/from the occipital cortex showed reduced volume in CB individuals compared to the SC, which brings new insights into the thalamic remodeling in congenital blindness. The comparison of thalamic volumes obtained from different approaches, such as probabilistic tractography and VBM, must be explored in future studies considering their technical aspects and biological underpinnings. For example, we employed a probabilistic tractography algorithm to map the connectivity distributions from individual voxels within the thalamus in each participant, while VBM relies on the boundaries between thalamic nuclei[62].

Even though interesting, volume comparisons of thalamic structures have low intrinsic sensitivity, as differences in connectivity may not be reflected as volume changes. This would be even more tricky since thalamocortical remapping may be presented as both shrinkage and expansions of nuclei territories in a non-homogeneous and distributed pattern. For the first time, we disentangled this by comparing the voxel-wise differences in thalamocortical connectivity between groups, revealing the pulvinar's central role in remapping thalamocortical connectivity. WM connections between pulvinar/lateral posterior nuclei and occipital cortex were decreased in the CB group. On the other hand, increased connectivity to/from the temporal cortex was observed in the pulvinar. These findings suggest a thalamic shift of cortical connections in blind individuals, prioritizing the temporal projections over the occipital ones.

The pulvinar is the largest, higher-order multimodal thalamic nucleus, involved mainly in visual attention[63]. It displays a wide range of cortical connectivity beyond the visual cortex, roughly to the gray matter of all lobes and subcortical structures such as the amygdala and superior colliculus[63–65]. Evidence suggests that cortico-cortical integration via pulvinar is an important pathway for transferring visual information between cortical

areas[65,66]. Thus, it is reasonable to hypothesize that the pulvinar may become a central

player in the dynamics of brain plasticity, possibly by integrating different sensory

modalities at associative levels and paving multimodal plasticity in blind individuals.

However, the pulvinar typically connects to higher-order areas of the temporal cortex

such as inferior, ventral, and lateral temporal areas, temporoparietal junction, and mesial

temporal cortex[64,67,68]. Therefore, it is impossible to conclude whether the aberrant

connectivity pattern of the pulvinar observed here relies on new thalamocortical

projections to the temporal cortex or a mechanism of remodeling the preexisting ones.

Hence, the role of pulvinar in cross-modal plasticity needs to be considered and further

explored in future studies.

It is worth noting that our findings add new insights to the still-developing

literature on thalamocortical connectivity changes in blind humans. Even though Reislev

and colleagues[69] have investigated possible differences in thalamocortical connectivity

between CB and SC groups, evidence of thalamic remapping has not been found. For the

first time using a voxel-wise approach, we reported converging evidence that thalamic

projections to/from the temporal cortex are increased as opposed to those to/from the

occipital cortex. First, comparisons of the thalamic territory connecting to the temporal

and occipital cortices showed increased and decreased volume in the CB group,

respectively. Second, the voxel-wise analysis pointed to an aberrant pattern of

thalamocortical connectivity involving pulvinar/lateral posterior nucleus, LGN, and

MGN, among others. Thus, despite similarities, we used original analysis to describe the

impact of congenital blindness on the thalamocortical connectivity in humans.

In the past years, several authors have described increased V1 thickness in the

blind brain[18,19,39,70], which can be explained by the lack of correct pruning during

ontogenesis due to insufficient input[19,71,72]. On the other hand, the absence of appropriate

input may also result in an impaired structure of V1[6,8,15,21]. In the present study, the CB

group displayed greater thickness of the left but not the right V1 (BA 17) than SC.

Although the changes of cortical thickness in blind individuals have been reported

bilaterally in previous studies, the most prominent results are often reported in the left

occipital cortex in right-handers[18,70], indicating lateralization effects of plasticity in these

cases. Indeed, the visual cortex is highly lateralized for a wide range of stimuli[73–76]. Non-

visual tasks, such as spatial and pitch discrimination of sounds[34], voice perception[77] and

recognition[35], sound localization[36], syntactic[29,] and semantic[28] processing of sentences,

elicit lateralized visual cortex activation in blind individuals. In addition, the lateralization
of functional connectivity to associative brain areas has also been reported[34,78]. Due to
the nature of hemispheric dominance of the visual cortex that differently affects both
hemispheres depending on the task, it is hard to define precisely what aspects of brain
plasticity underpin the left lateralization of increased cortical thickness in CB. One
possible mechanism affecting the left V1 is that input to V1 seems to be a critical part of
cortical maturation and pruning[19,71,72]. Indeed, we found evidence of left-lateralized
impairment effects on thalamo-occipital connections in CB. Thus, the involvement of
thalamocortical connectivity in this phenomenon should be addressed in future studies.

We did not find evidence of thalamic remapping of connections to the
somatosensory and motor cortices, for example, in CB. However, careful interpretation
of the results must consider two key points. First, the connectivity-based segmentation of
the thalamus used in the present study is a 'winner-takes-all' approach in which the most
probable connections are considered while the least probable ones are discarded. Hence,
plasticity involving somatosensory and motor remodeling of thalamic connections may
not have been strong enough to be detected by our methods. As a result, the 'normal'
pattern of thalamic segmentation would mask meaningful but less structured connections.
For example, even though weaker, aberrant somatosensory, auditory and motor thalamic
nuclei are connected with V1 in blind opossum[46]. In addition, V1 activation following
tactile stimulation may be driven by aberrant connections between the
ventroposteriorlateral nucleus and LGN in CB individuals[79]. Second, although we did not
intend to assess corticocortical connections between visual and the remaining sensory
modalities, this kind of direct connectivity has been reported in animal models of
blindness and must be considered a possible mechanism of cross-modal plasticity in
humans. In fact, the intermodal connections during ontogenesis seem to be altered by the
lack of visual input in animals[48,53,80–82]. In humans, indirect measurement based on
functional connectivity pointed to direct connections between the occipital and auditory
cortex[83] and primary somatosensory cortex[84,85]. Moreover, the visual cortex of enucleated
animals also displays both abnormal corticocortical and thalamocortical connections with
somatosensory, auditory, and motor areas[46], suggesting that both mechanisms may
coexist to support cross-modal plasticity in the blind brain.
In accordance with previous studies that showed decreased FA of WM in CB[10,13–
15], we observed focal impairment of WM integrity in CB individuals impacting most

occipital connections to the associative cortical areas. Structures showing decreased FA were splenium of the corpus callosum, bilateral inferior longitudinal fasciculi, bilateral inferior fronto-occipital fasciculi, and bilateral superior longitudinal fasciculus, which are known for their role in processing and integration of visual information. Lower FA values refer to WM integrity and myelination impairments and are commonly pointed to as a biomarker of developmental changes, axonal degeneration, and plasticity. Indeed, visual input is crucial for maturation and refinement of early crude connectivity of the visual system,[71,82,86] suggesting that decreased myelination of visual WM tracts in CB individuals might reflect the lack of maturation of the system rather than axonal degeneration per se. Moreover, the thalamus also displayed reduced integrity in the CB group, indicating that microstructural impairment is not restricted to corticocortical bundles. FA analysis of GM structures may represent a measure of cytoarchitecture[87] and is in line with previous studies[69]. In addition to the reduced FA, previous studies suggest that the lack of visual input impacts WM volume compared with sighted controls[7,8,16].

The present findings raise intriguing questions that should be addressed in future studies. For example, the CB group showed increased structural connectivity between the ventral anterior nuclei and the temporal cortex. Given that most studies focused on exploring abnormal thalamocortical connectivity involving V1, the possible involvement of a motor thalamic nucleus, such as the ventral anterior, deserves further investigation. In addition, our sample of blind individuals includes only CB. Despite similar approaches have targeted both CB and late blind individuals[69], future studies should employ voxel-wise investigation of thalamocortical connectivity in people with acquired blindness to elucidate to what extent the present thalamocortical connectivity changes depend on visual experience.

In summary, the present study corroborates previous findings pointing to the blind brain as a model of a wide range of plastic phenomena. For the first time in humans, the remapping of thalamocortical connections involving both unimodal and multimodal thalamic nuclei is described, which may represent a mechanism of how non-visual stimuli are relayed to the 'visual' cortex. Future studies should employ neurophysiologic approaches to correlate these changes with functional plasticity often observed in the absence of visual stimuli from birth.

ACKNOWLEDGEMENTS

392   We thank Ivanei Bramati and Débora Lima for their support during data analysis

393 and writing. The authors declare that the research was conducted in the absence of any

394 commercial or financial relationships that could be construed as a potential conflict of

395 interest.

FUNDING

This work was funded by the Foundation for Research Support in the State of Rio

de Janeiro (FAPERJ), the National Council of Scientific and Technological Development

(CNPq), the Capes Foundation, and intramural grants of the D'Or Institute for Research

and Education (IDOR).

COMPETING INTERESTS

The authors report no competing interests.

REFERENCES

1.   WHO. *World Report on Vision*. Vol 214.; 2019.

2.   Gogate P, Gilbert C, Zin A. Severe visual Impairment and blindness in infants:
Causes and opportunities for control. *Middle East African Journal of*
Ophthalmology. 2011;18(2):109-114. doi:10.4103/0974-9233.80698

3.   Anurova I, Carlson S, Rauschecker JP. Overlapping Anatomical Networks
Convey Cross-Modal Suppression in the Sighted and Coactivation of "visual"
and Auditory Cortex in the Blind. *Cerebral Cortex*. 2019;29(11):4863-4876.
doi:10.1093/cercor/bhz021

4.   Aguirre GK, Datta R, Benson NC, et al. Patterns of individual variation in visual
pathway structure and function in the sighted and blind. *PLoS ONE*.
2016;11(11):1-24. doi:10.1371/journal.pone.0164677

5.   Lao Y, Kang Y, Collignon O, et al. A study of brain white matter plasticity in
early blinds using tract-based spatial statistics and tract statistical analysis.
NeuroReport. 2015;26(18):1151-1154. doi:10.1097/WNR.0000000000000488

6.   Noppeney U, Friston KJ, Ashburner J, Frackowiak R, Price CJ. Early visual
deprivation induces structural plasticity in gray and white matter [1]. *Current*
Biology. 2005;15(13):488-490. doi:10.1016/j.cub.2005.06.053

7.   Park HJ, Jeong SO, Kim EY, et al. Reorganization of neural circuits in the blind
on diffusion direction analysis. *NeuroReport*. 2007;18(17):1757-1760.
doi:10.1097/WNR.0b013e3282f13e66

8.   Bridge H, Cowey A, Ragge N, Watkins K. Imaging studies in congenital
anophthalmia reveal preservation of brain architecture in "visual" cortex. *Brain*.
2009;132(12):3467-3480. doi:10.1093/brain/awp279

9.   Shu N, Liu Y, Li J, Li Y, Yu C, Jiang T. Altered anatomical network in early
blindness revealed by diffusion tensor tractography. *PLoS ONE*. 2009;4(9):1-13.
doi:10.1371/journal.pone.0007228

10.   Reislev NL, Dyrby TB, Siebner HR, Kupers R, Ptito M. Simultaneous
434   Assessment of White Matter Changes in Microstructure and Connectedness in
435   the Blind Brain. *Neural Plasticity*. 2016;2016. doi:10.1155/2016/6029241

436   11.   Ptito M, Paré S, Dricot L, Cavaliere C, Tomaiuolo F, Kupers R. A quantitative
437   analysis of the retinofugal projections in congenital and late-onset blindness.
438   *NeuroImage: Clinical*. 2021;32. doi:10.1016/J.NICL.2021.102809

439   12.   Collignon O, Lassonde M, Lepore F, Bastien D, Veraart C. Functional cerebral
440   reorganization for auditory spatial processing and auditory substitution of vision
441   in early blind subjects. *Cerebral Cortex*. 2007;17(2):457-465.
442   doi:10.1093/cercor/bhj162

443   13.   Ptito M, Schneider FCG, Paulson OB, Kupers R. Alterations of the visual
444   pathways in congenital blindness. *Experimental Brain Research*. 2008;187(1):41-
445   49. doi:10.1007/s00221-008-1273-4

446   14.   Shu N, Li J, Li K, Yu C, Jiang T. Abnormal diffusion of cerebral white matter in
447   early blindness. *Human Brain Mapping*. 2009;30(1):220-227.
448   doi:10.1002/hbm.20507

449   15.   Leporé N, Voss P, Lepore F, et al. Brain structure changes visualized in early-
450   and late-onset blind subjects. *Neuroimage*. 2010;49(1):134-140.
451   doi:10.1016/j.neuroimage.2009.07.048

452   16.   Dietrich S, Hertrich I, Kumar V, Ackermann H. Experience-related structural
453   changes of degenerated occipital white matter in late-blind Humans-A diffusion
454   tensor imaging study. *PLoS ONE*. 2015;10(4):1-19.
455   doi:10.1371/journal.pone.0122863

456   17.   Qin W, Liu Y, Jiang T, Yu C. The Development of Visual Areas Depends
457   Differently on Visual Experience. *PLoS ONE*. 2013;8(1):1-10.
458   doi:10.1371/journal.pone.0053784

459   18.   Anurova I, Renier LA, de Volder AG, Carlson S, Rauschecker JP. Relationship
460   Between Cortical Thickness and Functional Activation in the Early Blind.
461   *Cerebral Cortex*. 2015;25(8):2035-2048. doi:10.1093/cercor/bhu009

462   19.   Jiang J, Zhu W, Shi F, et al. Thick visual cortex in the early blind. *Journal of
463   Neuroscience*. 2009;29(7):2205-2211. doi:10.1523/JNEUROSCI.5451-08.2009

464   20.   Voss P, Pike BG, Zatorre RJ. Evidence for both compensatory plastic and disuse
465   atrophy-related neuroanatomical changes in the blind. *Brain*. 2014;137(4):1224-
466   1240. doi:10.1093/brain/awu030

467   21.   Pan WJ, Wu G, Li CX, Lin F, Sun J, Lei H. Progressive atrophy in the optic
468   pathway and visual cortex of early blind Chinese adults: A voxel-based
469   morphometry magnetic resonance imaging study. *Neuroimage*. 2007;37(1):212-
470   220. doi:10.1016/j.neuroimage.2007.05.014

471   22.   Sadato N, Pascual-Leone A, Grafman J, et al. Activation of the primary visual
472   cortex by Braille reading in blind subjects. *Nature*. 1996;380(6574):526-528.
473   doi:10.1038/380526a0

474   23.   Büchel C, Price C, Frackowiak RSJ, Friston K. Different activation patterns in
475   the visual cortex of late and congenitally blind subjects. *Brain*. 1998;121(3):409-
476   419. doi:10.1093/brain/121.3.409

477   24.   Röder B, Teder-Sälejärvi W, Sterr A, Rösler F, Hillyard SA, Neville HJ.
478   Improved auditory spatial tuning in blind humans. *Nature*. 1999;400(6740):162-
479   166. doi:10.1038/22106

480   25.   Weeks R, Horwitz B, Aziz-Sultan A, et al. A positron emission tomographic
481   study of auditory localization in the congenitally blind. *Journal of Neuroscience*.
482   2000;20(7):2664-2672. doi:10.1523/jneurosci.20-07-02664.2000

483   26.   Burton H, Snyder AZ, Diamond JB, Raichle ME. Adaptive changes in early and
484   late blind: A fMRI study of verb generation to heard nouns. *Journal of*
485   *Neurophysiology*. 2002;88(6):3359-3371. doi:10.1152/jn.00129.2002

486   27.   Wan CY, Wood AG, Reutens DC, Wilson SJ. Early but not late-blindness leads
487   to enhanced auditory perception. *Neuropsychologia*. 2010;48(1):344-348.
488   doi:10.1016/j.neuropsychologia.2009.08.016

489   28.   Röder B, Stock O, Bien S, Neville H, Rösler F. Speech processing activates
490   visual cortex in congenitally blind humans. *European Journal of Neuroscience*.
491   2002;16(5):930-936. doi:10.1046/j.1460-9568.2002.02147.x

492   29.   Lane C, Kanjlia S, Omaki A, Bedny M. "Visual" cortex of congenitally blind
493   adults responds to syntactic movement. *Journal of Neuroscience*.
494   2015;35(37):12859-12868. doi:10.1523/JNEUROSCI.1256-15.2015

495   30.   Rosenbluth R, Grossman ES, Kaitz M. Performance of early-blind and sighted
496   children on olfactory tasks. *Perception*. 2000;29(1):101-110. doi:10.1068/p3001

497   31.   Cuevas I, Plaza P, Rombaux P, de Volder AG, Renier L. Odour discrimination
498   and identification are improved in early blindness. *Neuropsychologia*.
499   2009;47(14):3079-3083. doi:10.1016/j.neuropsychologia.2009.07.004

500   32.   de Volder AG, Catalan-Ahumada M, Robert A, et al. Changes in occipital cortex
501   activity in early blind humans using a sensory substitution device. *Brain*
502   *Research*. 1999;826(1):128-134. doi:10.1016/S0006-8993(99)01275-5

503   33.   Gizewski ER, Gasser T, de Greiff A, Boehm A, Forsting M. Cross-modal
504   plasticity for sensory and motor activation patterns in blind subjects.
505   *Neuroimage*. 2003;19(3):968-975. doi:10.1016/S1053-8119(03)00114-9

506   34.   Collignon O, Vandewalle G, Voss P, et al. Functional specialization for auditory-
507   spatial processing in the occipital cortex of congenitally blind humans. *Proc Natl*
508   *Acad Sci U S A*. 2011;108(11):4435-4440. doi:10.1073/pnas.1013928108

509   35.   Hölig C, Föcker J, Best A, Röder B, Büchel C. Crossmodal plasticity in the
510   fusiform gyrus of late blind individuals during voice recognition. *Neuroimage*.
511   2014;103:374-382. doi:10.1016/j.neuroimage.2014.09.050

512   36.   Gougoux F, Zatorre RJ, Lassonde M, Voss P, Lepore F. A functional
513   neuroimaging study of sound localization: Visual cortex activity predicts
514   performance in early-blind individuals. *PLoS Biology*. 2005;3(2):0324-0333.
515   doi:10.1371/journal.pbio.0030027

516   37.   Stevens AA, Snodgrass M, Schwartz D, Weaver K. Preparatory activity in
517   occipital cortex in early blind humans predicts auditory perceptual performance.
518   *Journal of Neuroscience*. 2007;27(40):10734-10741.
519   doi:10.1523/JNEUROSCI.1669-07.2007

520   38.   Voss P, Gougoux F, Zatorre RJ, Lassonde M, Lepore F. Differential occipital
521   responses in early- and late-blind individuals during a sound-source
522   discrimination task. *Neuroimage*. 2008;40(2):746-758.
523   doi:10.1016/j.neuroimage.2007.12.020

524   39.   Voss P, Zatorre RJ. Occipital cortical thickness predicts performance on pitch
525   and musical tasks in blind individuals. *Cerebral Cortex*. 2012;22(11):2455-2465.
526   doi:10.1093/cercor/bhr311

527   40.   Cohen LG, Celnik P, Pascual-Leone A, et al. Functional relevance of cross-
528   modal plasticity in blind humans. *Nature*. 1997;389(6647):180-183.
529   doi:10.1038/38278

530   41.   Bavelier D, Neville HJ. Cross-modal plasticity: Where and how? *Nature Reviews*
531   *Neuroscience*. 2002;3(6):443-452. doi:10.1038/nrn848

42. Falchier A, Clavagnier S, Barone P, Kennedy H. Anatomical evidence of
multimodal integration in primate striate cortex. *Journal of Neuroscience*.
2002;22(13):5749-5759. doi:10.1523/jneurosci.22-13-05749.2002
43. Kingsbury MA, Lettman NA, Finlay BL. Reduction of early thalamic input alters
adult corticocortical connectivity. *Developmental Brain Research*.
2002;138(1):35-43. doi:10.1016/S0165-3806(02)00456-X
44. Rockland KS, Ojima H. Multisensory convergence in calcarine visual areas in
macaque monkey. *International Journal of Psychophysiology*. 2003;50(1-2):19-
26. doi:10.1016/S0167-8760(03)00121-1
45. Clavagnier S, Falchier A, Kennedy H. Long-distance feedback projections to area
V1: Implications for multisensory integration, spatial awareness, and visual
consciousness. *Cognitive, Affective and Behavioral Neuroscience*.
2004;4(2):117-126. doi:10.3758/CABN.4.2.117
46. Karlen SJ, Kahn DM, Krubitzer L. Early blindness results in abnormal
corticocortical and thalamocortical connections. *Neuroscience*. 2006;142(3):843-
858. doi:10.1016/j.neuroscience.2006.06.055
47. Hall AJ, Lomber SG. Auditory cortex projections target the peripheral field
representation of primary visual cortex. *Experimental Brain Research*.
2008;190(4):413-430. doi:10.1007/s00221-008-1485-7
48. Charbonneau V, Laramée ME, Boucher V, Bronchti G, Boire D. Cortical and
subcortical projections to primary visual cortex in anophthalmic, enucleated and
sighted mice. *European Journal of Neuroscience*. 2012;36(7):2949-2963.
doi:10.1111/j.1460-9568.2012.08215.x
49. Pallas SL, Littman T, Moore DR. Cross-modal reorganization of callosal
connectivity without altering thalamocortical projections. *Proc Natl Acad Sci U S
A*. 1999;96(15):8751-8756. doi:10.1073/pnas.96.15.8751
50. Zhang Y, Wan S, Ge J, Zhang X. Diffusion tensor imaging reveals normal
geniculocalcarine-tract integrity in acquired blindness. *Brain Research*.
2012;1458:34-39. doi:10.1016/j.brainres.2012.04.012
51. Bronchti G, Heil P, Scheich H, Wollberg Z. Auditory pathway and auditory
activation of primary visual targets in the blind mole rat (Spalax ehrenbergi): I. 2-
deoxyglucose study of subcortical centers. *Journal of Comparative Neurology*.
1989;284(2):253-274. doi:10.1002/cne.902840209
52. Doron N, Wollberg Z. Cross-modal neuroplasticity in the blind mole rat spalax
ehrenbergi: A WGA-HRP tracing study. *NeuroReport*. 1994;5(18):2697-2701.
doi:10.1097/00001756-199412000-00072
53. Laemle LK, Strominger NL, Carpenter DO. Cross-modal innervation of primary
visual cortex by auditory fibers in congenitally anophthalmic mice. *Neuroscience
Letters*. 2006;396(2):108-112. doi:10.1016/j.neulet.2005.11.020
54. Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and
structural MR image analysis and implementation as FSL. *Neuroimage*.
2004;23(SUPPL. 1):S208-S219. doi:10.1016/J.NEUROIMAGE.2004.07.051
55. Smith SM. Fast robust automated brain extraction. *Human Brain Mapping*.
2002;17(3):143-155. doi:10.1002/HBM.10062
56. Behrens TEJ, Woolrich MW, Jenkinson M, et al. Characterization and
propagation of uncertainty in diffusion-weighted MR imaging. *Magnetic
Resonance in Medicine*. 2003;50(5):1077-1088. doi:10.1002/MRM.10609
57. Smith SM, Jenkinson M, Johansen-Berg H, et al. Tract-based spatial statistics:
Voxelwise analysis of multi-subject diffusion data. *Neuroimage*.
2006;31(4):1487-1505. doi:10.1016/j.neuroimage.2006.02.024

58. Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation
inference for the general linear model. *Neuroimage*. 2014;92:381-397.
doi:10.1016/J.NEUROIMAGE.2014.01.060
59. Behrens TEJ, Woolrich MW, Jenkinson M, et al. Characterization and
Propagation of Uncertainty in Diffusion-Weighted MR Imaging. *Magnetic
Resonance in Medicine*. 2003;50(5):1077-1088. doi:10.1002/mrm.10609
60. Chakravarty MM, Bertrand G, Hodge CP, Sadikot AF, Collins DL. The creation
of a brain atlas for image guided neurosurgery using serial histological data.
Neuroimage. 2006;30(2):359-376. doi:10.1016/j.neuroimage.2005.09.041
61. Cecchetti L, Ricciardi E, Handjaras G, Kupers R, Ptito M, Pietrini P. Congenital
blindness affects diencephalic but not mesencephalic structures in the human
brain. *Brain Structure and Function*. 2016;221(3):1465-1480.
doi:10.1007/s00429-014-0984-5
62. Iglesias JE, Insausti R, Lerma-Usabiaga G, et al. A probabilistic atlas of the
human thalamic nuclei combining ex vivo MRI and histology. *Neuroimage*.
2018;183:314-326. doi:10.1016/J.NEUROIMAGE.2018.08.012
63. Benarroch EE. Pulvinar: Associative role in cortical function and clinical
correlations. *Neurology*. 2015;84(7):738-747.
doi:10.1212/WNL.0000000000001276
64. Leh SE, Chakravarty MM, Ptito A. The connectivity of the human pulvinar: A
diffusion tensor imaging tractography study. *International Journal of Biomedical
Imaging*. 2008;2008(1). doi:10.1155/2008/789539
65. Saalmann YB, Pinsk MA, Wang L, Li X, Kastner S. Pulvinar regulates
information transmission between cortical areas based on attention demands.
Science (1979). 2012;337(6095):753-756. doi:10.1016/S0140-6736(02)11602-
3.Association
66. Theyel BB, Llano DA, Sherman SM. The corticothalamocortical circuit drives
higher-order cortex in the mouse. *Nature Neuroscience*. 2010;13(1):84-88.
doi:10.1038/nn.2449
67. Arcaro MJ, Pinsk MA, Kastner S. The anatomical and functional organization of
the human visual pulvinar. *Journal of Neuroscience*. 2015;35(27):9848-9871.
doi:10.1523/JNEUROSCI.1575-14.2015
68. Rosenberg DS, Mauguière F, Catenoix H, Faillenot I, Magnin M. Reciprocal
thalamocortical connectivity of the medial pulvinar: A depth stimulation and
evoked potential study in human brain. *Cerebral Cortex*. 2009;19(6):1462-1473.
doi:10.1093/cercor/bhn185
69. Reislev NH, Dyrby TB, Siebner HR, Lundell H, Ptito M, Kupers R.
Thalamocortical Connectivity and Microstructural Changes in Congenital and
Late Blindness. *Neural Plasticity*. 2017;2017(November).
doi:10.1155/2017/9807512
70. Park HJ, Lee JD, Kim EY, et al. Morphological alterations in the congenital blind
based on the analysis of cortical thickness and surface area. *Neuroimage*.
2009;47(1):98-106. doi:10.1016/j.neuroimage.2009.03.076
71. Bourgeois JP, Jastreboff PJ, Rakic P. Synaptogenesis in visual cortex of normal
and preterm monkeys: Evidence for intrinsic regulation of synaptic
overproduction. *Proc Natl Acad Sci U S A*. 1989;86(11):4297-4301.
doi:10.1073/pnas.86.11.4297
72. Luo L, O'Leary DDM. Axon retraction and degeneration in development and
disease. *Annual Review of Neuroscience*. 2005;28:127-156.
doi:10.1146/annurev.neuro.28.061604.135632

73. Corballis PM, Funnell MG, Gazzaniga MS. Hemispheric asymmetries for simple
visual judgments in the split brain. *Neuropsychologia*. 2002;40(4):401-410.
doi:10.1016/S0028-3932(01)00100-2
74. Rossion B, Joyce CA, Cottrell GW, Tarr MJ. Early lateralization and orientation
tuning for face, word, and object processing in the visual cortex. *Neuroimage*.
2003;20(3):1609-1624. doi:10.1016/j.neuroimage.2003.07.010
75. Hougaard A, Jensen BH, Amin FM, Rostrup E, Hoffmann MB, Ashina M.
Cerebral asymmetry of fMRI-BOLD responses to visual stimulation. *PLoS ONE*.
2015;10(5):1-16. doi:10.1371/journal.pone.0126477
76. Davies-Thompson J, Johnston S, Tashakkor Y, Pancaroglu R, Barton JJS. The
relationship between visual word and face processing lateralization in the
fusiform gyri: A cross-sectional study. *Brain Research*. 2016;1644:88-97.
doi:10.1016/j.brainres.2016.05.009
77. Gougoux F, Belin P, Voss P, Lepore F, Lassonde M, Zatorre RJ. Voice
perception in blind persons: A functional magnetic resonance imaging study.
Neuropsychologia. 2009;47(13):2967-2974.
doi:10.1016/j.neuropsychologia.2009.06.027
78. Sabbah N, Authié CN, Sanda N, et al. Increased functional connectivity between
language and visually deprived areas in late and partial blindness. *Neuroimage*.
2016;136:162-173. doi:10.1016/j.neuroimage.2016.04.056
79. Müller F, Niso G, Samiee S, Ptito M, Baillet S, Kupers R. A thalamocortical
pathway for fast rerouting of tactile information to occipital cortex in congenital
blindness. *Nature Communications*. 2019;10(1):1-9. doi:10.1038/s41467-019-
13173-7
80. Berman NEJ. Alterations of visual cortical connections in cats following early
removal of retinal input. *Developmental Brain Research*. 1991;63(1-2):163-180.
doi:10.1016/0165-3806(91)90076-U
81. Bock AS, Kroenke CD, Taber EN, Olavarria JF. Retinal input influences the size
and corticocortical connectivity of visual cortex during postnatal development in
the ferret. *Journal of Comparative Neurology*. 2012;520(5):914-932.
doi:10.1002/cne.22738
82. Laramée ME, Bronchti G, Boire D. Primary visual cortex projections to
extrastriate cortices in enucleated and anophthalmic mice. *Brain Structure and
Function*. 2014;219(6):2051-2070. doi:10.1007/s00429-013-0623-6
83. Klinge C, Eippert F, Röder B, Büchel C. Corticocortical connections mediate
primary visual cortex responses to auditory stimulation in the blind. *Journal of
Neuroscience*. 2010;30(38):12798-12805. doi:10.1523/JNEUROSCI.2384-
10.2010
84. Wittenberg GF, Werhahn KJ, Wassermann EM, Herscovitch P, Cohen LG.
Functional connectivity between somatosensory and visual cortex in early blind
humans. *European Journal of Neuroscience*. 2004;20(7):1923-1927.
doi:10.1111/j.1460-9568.2004.03630.x
85. Ioannides AA, Liu L, Poghosyan V, et al. MEG reveals a fast pathway from
somatosensory cortex to occipital areas via posterior parietal cortex in a blind
subject. *Frontiers in Human Neuroscience*. 2013;7(JUL):1-12.
doi:10.3389/fnhum.2013.00429
86. Sengpiel F, Kind PC. The role of activity in development of the visual system.
Current Biology. 2002;12(23):818-826. doi:10.1016/S0960-9822(02)01318-0
87. Assaf Y. Imaging laminar structures in the gray matter with diffusion MRI.
Neuroimage. 2019;197:677-688. doi:10.1016/J.NEUROIMAGE.2017.12.096

88. Chakravarty MM, Bertrand G, Hodge CP, Sadikot AF, Collins DL. The creation
of a brain atlas for image guided neurosurgery using serial histological data.
Neuroimage. 2006;30(2):359-376. doi:10.1016/j.neuroimage.2005.09.041

Fig. 1 Thalamocortical Connectivity in Sighted Controls and Congenitally Blind

Individuals. (A) Six cortical masks were predefined using the Harvard-Oxford atlas:

Prefrontal, Precentral, Postcentral, Posterior Parietal (parietal cortex, except for the

postcentral cortex), Occipital, and Temporal. Resultant thalamic segments showing

statistically significant connectivity to each cortical area in the (B) SC (sighted controls,

top) and (C) CB (congenitally blind, middle) groups. The color bar represents the p-

value after FWE correction. The normalized volume of the Occipital and Temporal

segments showed group differences (p<0.001, Bonferroni correction for multiple

comparisons).

Figure 2. Thalamocortical Connectivity Changes in Congenitally Blind
Individuals. The yellow box (left) depicts thalamic areas that exhibited increased
connectivity with the temporal cortex, including MGN, LGN and pulvinar bilaterally. The
blue box (right) depicts thalamic territory that exhibited decreased connectivity with the
occipital cortex in congenitally blind individuals, namely the left pulvinar/lateral
posterior nucleus. The white box (middle) shows thalamic territories obtained from an
atlas based on the Colin-27 Average Brain[88] and depicts the location of LGN (green),
MGN (dark pink), pulvinar (red), medial dorsal (yellow), ventral anterior (orange),
anterior (purple), and lateral posterior (light pink) nuclei. A graphical overlay (dark blue)
of thalamic areas that exhibited both increased connectivity to the temporal cortex and
decreased connectivity to the occipital cortex (p<0.05, FWE-corrected) in CB individuals
is shown (white box, bottom). L = left; R = right; A = anterior; P = posterior. The
coordinates are given according to the MNI space and plotted on the MNI standard
brain. Color bars represent the t-value.

Fig.3 FA comparison (CB group < SC group). White matter tracts (whole brain) showed reduced FA (fractional anisotropy) in the CB (congenitally blind) group, as compared to SC (sighted controls) group FWE-corrected, p< 0.05). FA difference in the thalamus, resulting from a separate analysis, is also displayed. The color bar represents the t value corrected for multiple comparisons. ILF = inferior longitudinal fasciculus; IFOF = inferior fronto-occipital fasciculus; CC = corpus callosum; SLF = superior longitudinal fasciculus.

Figure 4. Increased Cortical Thickness in the Congenitally Blind Group. (A) Increased cortical thickness in the left V1 (BA17) in the CB (congenitally blind) group compared to the SC (sighted controls) group, whole-brain analysis. (B) The mean cortical

729   thickness (in mm) of each group is displayed for visualization of the clustered values and

730   not for inference (error bar represents standard deviation).

734   **Supplementary material**

736   **Table S1. Investigation of the effect of 'segment'.**

Segment (pairwise comparison)		p-value (adjusted)
Frontal	Motor	$4.55 \times 10^{-130}*$
mean = 0.539 (sd = 0.049)	Posterior Parietal	$3.2 \times 10^{-132}*$
	Somatosensory	$3.01 \times 10^{-133}*$
	Occipital	$1.49 \times 10^{-122}*$
Motor	Occipital	$7.36 \times 10^{-4}*$
mean = 0.064 (sd = 0.024)	Posterior Parietal	1
	Somatosensory	1
	Temporal	$2.2 \times 10^{-32}*$
Occipital	Posterior Parietal	$4.3 \times 10^{-4}*$
mean = 0.1 (sd = 0.039)	Somatosensory	$1.28 \times 10^{-5}*$
	Temporal	$6.62 \times 10^{-18}*$
Posterior Parietal	Somatosensory	1
mean = 0.058 (sd = 0.024)	Temporal	$1.39 \times 10^{-33}*$
Somatosensory	Temporal	$1.65 \times 10^{-37}*$
mean = 0.047 (sd = 0.021)		
Temporal	Frontal	$4.04 \times 10^{-101}*$
mean = 0.193 (sd = 0.063)		

737   Mean and standard deviation (sd) refer to the normalized mean volume in arbitrary units.

738   P values were adjusted for Bonferroni correction. Asterisks indicate statistically

739   significant differences.

743   **Table S2. Investigation of the interaction 'segment*side'.**

Segment	Volume (mean(sd))		p-value (adjusted)
	Left side	Right side	
Frontal	0.536 (0.053)	0.541 (0.046)	0.766
Motor	0.074 (0.027)	0.053 (0.014)	0.00308*
Occipital	0.11 (0.035)	0.09 (0.041)	0.112
Posterior Parietal	0.051 (0.025)	0.064 (0.021)	0.0812
Somatosensory	0.059 (0.021)	0.036 (0.012)	0.000101*
Temporal	0.169 (0.056)	0.216 (0.063)	0.0171*

744 Volume and standard deviation (sd) refer to the normalized mean volume in arbitrary

745 units. P values were adjusted for Bonferroni correction. Asterisks indicate statistically

746 significant differences.

749