1

Transcriptional dynamics following freezing stress reveal selection for mechanisms of
freeze tolerance at the poleward range margin in the cold water intertidal barnacle
Semibalanus balanoides

Running title: Transcription following freezing in *S. balanoides*

Katie E. Marshall[a], Eddy J. Dowle[bc], Alexandra Petrunina[d], Gregory Kolbasov[d] and Benny
K. K. Chan[e*]

[a]Department of Zoology, University of British Columbia, Vancouver, Canada

[b]Department of Integrative Biology, University of Colorado, Denver, USA

[c]Department of Anatomy, University of Otago, Dunedin, New Zealand

[d]Moskovskij gosudarstvennyj universitet imeni M V Lomonosova Biologiceskij fakul'tet,
Department of Invertebrate Zoology

[e]Academia Sinica, Biodiversity Research Center, Taipei, Taiwan

*Corresponding author: Benny K.K. Chan

chankk@gate.sinica.edu.tw

Abstract

The ability to survive freezing has repeatedly evolved across multiple phyla. This suggests that

the mechanisms of freeze tolerance must be readily evolvable from basal physiological traits.

While several biochemical correlates to freeze tolerance have been described, the mechanism that

confers freeze tolerance is still not well understood. To understand both the basic biochemical

mechanisms of freeze tolerance as well as their role in local adaptation at the poleward range

edge, we conducted a transcriptomic study on two populations (one from the poleward range

margin in the White Sea, Russia, and one from the central coast of British Columbia, Canada) of

the cold water acorn barnacle *Semibalanus balanoides* on a time series following a freezing

event. We found that the British Columbia population (at the equatorward range margin) was

significantly less freeze tolerant than the White Sea population (at the poleward range margin).

After assembling and annotating a *de novo* transcriptome for *S. balanoides*, we found that the

patterns of differential transcript expression following freezing were almost entirely non-

overlapping between the two populations, with the White Sea population expressing a series of

heat shock proteins in response to freezing stress as well as several aquaporins, while the British

Columbia population expressed a series of proteases instead, indicating severe protein damage.

We found strong evidence of purifying selection on the significantly upregulated transcripts in

the White Sea population, suggesting local adaptation to freezing threat. Taken together, this

shows the importance of freeze tolerance to population survival at the poleward range margin,

and highlights the central roles of aquaporins and heat shock proteins to the trait of freeze

tolerance across taxa.

Keywords: freeze tolerance, transcriptomics, heat shock proteins, barnacles, local adaptation

Introduction

The occurrence of species range edges is one of the central problems in biogeography

(Sexton, Mcintyre, Angert, & Rice, 2009). Broadly speaking, it is believed that equatorward

range edges are due to biotic interactions, while poleward range edges are due to abiotic effects

(Louthan, Doak, & Angert, 2015). Given that minimum temperatures have the steepest and most

consistent latitudinal gradient (Sunday, Bates, & Dulvy, 2012), this suggests that low temperature

tolerance is central for understanding the presence of poleward range edges. Therefore,

understanding the mechanisms and adaptive potential of low temperature tolerance is central for

delineating the role of local adaptation in the maintenance of poleward range edges.

Low temperature tolerance includes responses to two overlapping, but distinct,

physiological stressors: the direct effects of low temperature on reaction rates, and managing the

potential and spread of internal ice. In ectothermic organisms, biochemical reaction rates

decrease due to Arrhenius effects as well as by decreased membrane fluidity at low temperatures,

which causes a general decrease in rates of physiological functioning (reviewed in R. E. Lee,

2010). At body temperatures below 0 °C, organisms also risk internal ice formation. Surviving

these two stressors, while linked by temperature, appear to involve different physiological

mechanisms. Some organisms like the eastern spruce budworm (*Choristoneura fumiferana*

Clemens) are able to suppress their freezing point well below -30 °C by the production of low

molecular weight cryoprotectants and antifreeze proteins, and can survive low temperatures as

long as they do not freeze (Han & Bauce, 1995; Katie E. Marshall & Sinclair, 2015; Qin, Doucet,

Tyshenko, & Walker, 2007). Others like the woolly bear caterpillar (*Pyrrharctia isabella* Smith)

induce and survive freezing at relatively high temperatures (c.a. -10 °C), but are unable to survive

temperatures below -16 °C (Boardman, Terblanche, & Sinclair, 2011; Layne & Kuharsky, 2000).

Being able to survive freezing is therefore a physiological trait that is likely mechanistically

distinct from being able to survive low temperature (reviewed in Toxopeus & Sinclair, 2018).

The ability to survive internal ice formation is a paraphyletic trait broadly distributed

across the animal kingdom, with freeze-tolerant species from multiple phyla including

arthropods, molluscs, and vertebrates (Sinclair, Addo-Bediako, & Chown, 2003; Storey &

Storey, 1996; Toxopeus & Sinclair, 2018). Freeze tolerance appears to have evolved

independently on at least 20 occasions, and this is almost certainly a large underestimate (Sinclair

et al., 2003). The repeated emergence of this trait suggests that the underlying mechanisms must

be readily evolvable and that freeze tolerance is possible in a variety of underlying physiological

conditions. Despite the ubiquity of the trait, and over 200 years of research, the physiological

mechanisms underlying animal freeze tolerance remain poorly understood (Lee, 2010; Sømme,

2000; Teets & Denlinger, 2014; Toxopeus & Sinclair, 2018).

Surviving internal ice formation is a unique challenge for animals. They need to be able

to manage potential mechanical damage from ice crystal formation, cellular dehydration as water

leaves the cell to join the growing extracellular ice lattice, and sustained periods of anoxia since

oxygen cannot perfuse easily to frozen tissue (Lee, 2010). Many biochemical correlates to

survival of these challenges have been described; most commonly this involves the mass

biosynthesis of low molecular weight polyols and sugars that act as osmoprotectants and

cryoprotectants that can also decrease the melting (and therefore freezing) point of cells for

avoiding intracellular ice formation (Koštál, Korbelová, Poupardin, Moos, & Šimek, 2016;

Toxopeus & Sinclair, 2018). However there is no correlation between quantity of low molecular

weight cryoprotectants and degree of freeze tolerance—for instance, woolly bear caterpillars *P.*

isabella accumulate circa 2.5 M glycerol as a cryoprotectant but have a lower lethal temperatures

of only circa -16 °C (Boardman et al., 2011; K. E. Marshall & Sinclair, 2011), while the freeze

tolerant goldenrod gall fly (*Eurosta solidaginis* Fitch) has glycerol content of circa 0.3 M but a

lower lethal temperature below -50 °C (Lee, Dommell, Joplin, & Denlinger, 1995). Freeze

tolerant intertidal barnacles and mussels appear to not accumulate polyols or sugars at all

(Johnston & Clarke, 1990). It therefore appears that while these biochemical correlates might be

associated with enhanced survival of freezing, they do not explain the physiological mechanisms

of the ability to tolerate internal ice.

More recent work has focused on the role of proteins in freeze tolerance. Antifreeze

proteins may protect against intracellular freezing, which is almost always lethal even in very

freeze tolerant animals (Davies, 2014; but see review in Sinclair & Renault, 2010). In the

extracellular spaces, they may inhibit recrystallization, keeping ice crystals small enough to avoid

mechanical damage. However they have been described from only a few freeze tolerant species,

and their role in freeze tolerance remains controversial (Duman, Bennett, Sformo, Hochstrasser,

& Barnes, 2004; Toxopeus & Sinclair, 2018). Aquaporins facilitate the movement of water out

of the cell during freezing and glycerol transporters to facilitate the movement of sugars into the

cell have been reasonably well-explored in plant freeze tolerance (Peng, Arora, Li, Wang, &

Fessehaie, 2008; Philip, Yi, Elnitsky, & Lee, 2008; Storey & Storey, 2013; Yi et al., 2011). In

animals, aquaporins are implicated in the freeze tolerance in *E. solidaginis* and the wood frog

(*Lithobates sylvatica* LeConte), but how taxonomically broadly important they are for freeze

tolerance is not currently well understood (Philip, Kiss, & Lee, 2011; Philip et al., 2008; Storey

& Storey, 2013; Yi et al., 2011). Finally, heat shock proteins are constitutively upregulated in

overwintering insects and have been implicated in plant freeze tolerance, but are not generally

thought to be involved in responses to acute freezing stress in animals (King & MacRae, 2015;

Rinehart et al., 2006). However, the Antarctic midge (*Belgica antarctica* Jacobs) upregulates

Hsp70 in response to repeated freezing episodes, and freeze tolerant drosophilid fly (*Chymomyza*

costata Zetterstedt) upregulate small heat shock proteins following freezing suggest, suggesting

that heat shock proteins may be more important to freeze tolerance than previously thought

(Teets, Kawarasaki, Lee, & Denlinger, 2011).

While the biochemical correlates of freeze tolerance have been explored in broad array of

terrestrial organisms, very little is known about the biochemical mechanisms of freeze tolerance

in intertidal organisms (Dennis, Loomis, & Hellberg, 2014; Murphy, 1983). In temperate and

polar regions, intertidal organisms including barnacles, mussels, and snails are almost without

exception freeze tolerant (Murphy, 1983; Waller, Worland, Convey, & Barnes, 2006; but see

Sinclair, Marshall, Singh, & Chown, 2004). This is necessary because of the unique nature of the

intertidal—at low tide in the winter, the wetness of the habitat and the year-round presence of

food in the digestive tract makes freezing a constant threat when animals are emersed. Some

efforts have been made to determine the underlying biochemical mechanisms, but they remain

elusive (Ansart & Vernon, 2003; Dennis et al., 2014; Murphy, 1983; Storey & Storey, 2013). It

has been hypothesized that anaerobic byproducts such as strombine and taurine may be important

in freeze tolerance of *Mytilus sp.* mussels (Loomis, Carpenter, & Crowe, 1988), but much

remains unknown about how intertidal species survive freezing.

Here we present the first (to our knowledge) transcriptomic study following freezing

stress in an intertidal animal. We explore potential mechanisms of freeze tolerance in the acorn

barnacle (*Semibalanus balanoides* Linnaeus) from two populations that differ widely in the

intensity of freezing stress, one from the central coast of British Columbia, Canada which

experience relatively mild winters, and one from the White Sea in Russia which experiences

significantly harsher winters. We hypothesize that barnacles from British Columbia will be less

freeze tolerant (will have lower survival following freezing) than barnacles from the White Sea.

If this is true, we will infer that transcriptional responses to freezing in the White Sea population

likely enhance freeze tolerance. In addition, if transcripts upregulated in this population are

important for survival in poleward range edge habitats, we predict that we will see evidence of

selective sweeps on transcripts identities that are associated with biochemical mechanisms of

freeze tolerance and are upregulated following freezing stress. We find strong interpopulation

differences in response to freezing stress, and that this includes wholescale transcription of heat

shock proteins following freezing stress, as well as the transcription of aquaporins (which show

significant evidence of selective sweeps) and potentially new classes of antifreeze proteins. This

suggests that these protein-based mechanisms of freeze tolerance may be broadly universal across

freeze tolerant phyla, and that they may have selective advantages in intertidal environments.

Methods

Study system and collection

The barnacle *S. balanoides* is a common species in rocky intertidal zones through the

Arctic Circle, including northwestern Europe and both coasts of North America.   In western

North America, it is found as far south as the central coast of British Columbia (Flight & Rand,

2012; Schmidt, Bertness, & Rand, 2000; Wethey, 1983). Because of its polar distribution as well

as its intertidal habitat, *S. balanoides* regularly experiences freezing events throughout the winter

(Cook & Gabott, 1970; Cook & Lewis, 1971; Dennis J. Crisp & Ritz, 1967).   Late-stage larvae

(cyprids) of *S. balanoides* have even been found embedded in ice during winter, and successfully

settle and metamorphose after spring thaw (Pineda, DiBacco, & Starczak, 2005). Barnacles do

not appear to use polyol cryoprotectants for their freeze tolerance, but little is known about the

mechanisms of their freeze tolerance (Cook & Gabott, 1970; D.J. Crisp, Davenport, & Gabbott,

1977).

We collected *S. balanoides* from mid intertidal habitats by gathering small stones covered

with large groups of barnacles at the White Sea Biological Station (Moscow State University,

Russia; 66.5551° N, 34.1976° E) as well as from Calvert Island, British Columbia, Canada

(51.5559° N, 128.0365° W) during the spring of 2015. The climate on Calvert Island is

moderated by the Alaska Current which brings warm water from the south, and thus winter

temperatures rarely drop below -10 °C, while in the White Sea overwinter temperatures regularly

drop below -20 °C (Figure S1-S2), thus we expected barnacles from the White Sea be more

tolerant of freezing. We have installed robobarnacles (Chan, Lima, Williams, Seabra, & Wang,

2016) in the White Sea to record the estimated body temperature of similarly-shaped *Tetraclita*

spp. barnacles during the winter months (Figure S1). Barnacles for transcriptomic experiments

were collected in June in both habitats, wrapped in wet newspaper, then were placed in a

common seawater aquarium maintained at 10 °C and aerated by an airstone for two weeks before

experimentation.

Experimental design and freezing exposures

After acclimation, we measured barnacle test diameter with digital calipers along the

longest edge to the nearest mm. To simulate an intertidal freezing event, we cooled barnacles by

placing the rocks they had settled on into a Styrofoam box, then placed the box in a freezer set to

-20 °C. We continuously monitored barnacle temperature by placing a 36 AWG gauge Type T

(copper-constantan) thermocouple through the plates into the barnacle body. The thermocouple

was attached to a Picotech TC-08 interface (Pico Technology, Cambridge UK) which relayed

temperature data to an attached computer every 0.5 s in real time. We maintained barnacle

temperature to within 0.5 °C of the desired temperature by opening and closing the freezer door.

Barnacle supercooling points (SCPs) were recorded as the lowest temperature attained before

sudden release of heat due to the latent heat of crystallization (Sinclair, Coello Alvarado, &

Ferguson, 2015). With this procedure, we achieved cooling rates of circa 1.2 °C/min, which is

within the range found in their natural habitat during a low tide (Figure S1). To measure

differences in freeze tolerance, we exposed barnacles from both populations to -10 or -6 °C for 4

h, then warmed them by placing them immediately into seawater at 10 °C to mimic inundation

during high tide conditions. Survival was monitored for two days following and scored based on

the ability to close the top plates when removed from seawater, then compared using generalized

linear models with a binomial distribution in R (Team, 2016), while SCPs were compared using

ANOVA.

To measure transcriptional changes following freezing, we exposed 48 barnacles from

each population to -6 °C for 4 h using the above procedure. We sampled groups from each

population at four different time points: before the freezing event took place ("control"), after the

freezing event but before thawing in seawater ("before thawing"), 4 h after thawing ("4h

recovery") and 28 h after thawing ("28h recovery"; experimental design presented in Figure 1).

For each population × timepoint, we rapidly dissected out the soft tissue of 12 barnacles from

each population and flash-froze tissue in groups of three individuals in liquid nitrogen for four

total biological replicates/population/timepoint.

Transcriptomics

Total RNA was extracted from 32 pools of barnacles using a TRIzol protocol. Sequencing

libraries were prepared and sequenced by the Academia Sinica Bioinformatics Center. Libraries

were prepared using Illumina's stranded mRNA TruSeq RNA sample prep v2 kit as per

manufacturer's instructions. Cleaned libraries were sequenced on 2 lanes of 181 bp paired-end

Illumina HiSeq 2000 at the Academia Sinica Sequencing Center.

Our general approach for data analysis was informed by best practices for RNA-seq

(Conesa et al., 2016). The preliminary data quality checks and analysis were conducted in

Galaxy (Afgan et al., 2016). We clipped Illumina adapters from reads and removed low quality

and short read sequences using Trimmomatic's default settings (version 0.36.3; Bolger, Lohse, &

Usadel, 2014). We then took the trimmed reads and assembled a transcriptome using Trinity

(default settings for paired end reads, version 2.4.0, Grabherr et al., 2011), then clustered

redundant sequences using CD-HIT-EST with a sequence identity cut-off of 0.9 (Fu, Niu, Zhu,

Wu, & Li, 2012). We then aligned these sequences to the NCBI NR database using Diamond

(Buchfink, Xie, & Huson, 2014), then filtered out all reads that did not map to metazoans

(viruses, algae, bacteria, etc.) using MEGAN6 (Huson et al., 2016) to produce a *de novo*

transcriptome for *S. balanoides*. To annotate the transcriptome, we used Blast2GO (default

settings, version 5.1.13, Gotz et al., 2008) to blastx the transcripts to the NCBI arthropod nr

database, then mapped to Interpro IDs and GO terms to produce a final, annotated transcriptome.

To quantify gene expression following a freezing event, we then mapped our trimmed

reads to this transcriptome using HISAT2 (D. Kim, Langmead, & Salzberg, 2015), then

assembled and quantified transcripts using StringTie (Pertea et al., 2015), gffCompare (Trapnell

et al., 2010), and FeatureCounts (Liao, Smyth, & Shi, 2014). Following quantification, we tested

for differences in transcript abundance among time points and populations using DESeq2 (Love,

Huber, & Anders, 2014) to identify differentially expressed transcripts (adjusted p-value < 0.05)

within each population as a function of timepoint, while controlling for multiple comparisons

using the Benjamini-Hochberg multiple comparisons procedure.

We then used Blast2GO to test for functional enrichment of the significantly differentially

regulated transcripts within each population. We simplified the significantly enriched GO list for

plotting using REVIGO to eliminate semantically redundant terms, using the "tiny" setting

(Supek, Bošnjak, Škunca, & Šmuc, 2011). We also annotated each sequence with the nearest

KEGG orthology term using the KEGG Automatic Annotation Server (Moriya, Itoh, Okuda,

Yoshizawa, & Kanehisa, 2007), then tested for KEGG pathway enrichment using the Gage and

Pathview (Luo & Brouwer, 2013) packages in R.

Population differentiation

To quantify genetic differences between the two geographic sites, single nucleotide

polymorphisms (SNPs) were called using ANGSD 0.916 (Korneliussen, Albrechtsen, & Nielsen,

2014). The 'before thawing' samples were removed from the analysis as their transcript

expression profiles and therefore individual transcripts read coverage levels, differed markedly

from the other samples. Mapped reads were first filtered to remove PCR duplicates and

secondary mapping. Within ANGSD missing values were assigned for base calls of a phred score

<30. Loci were limited to those that had a depth of at least 10 reads per sample and an across all

samples Minor Allele Frequency (MAF) estimate of >0.05. As each sample is comprised of three

individuals (pooled) genotypes could not be called, instead sample MAF were estimated using

the allele counts method in ANGSD. Sample MAF estimates were used to conduct a Principle

Components Analysis (PCA) within R (prcomp function; PCA presented in Figure S3).

247   We tested whether there was evidence for selection on the differentially expressed

transcripts at either of the two geographic sites. As with the MAF estimates, the 'before thawing'

samples, PCR duplicates, and secondary mapping were removed. Samples were grouped into two

groups by geographic site, with each of the two groups were comprised of 36 individuals total

(12 samples of three individuals each). A pileup file was generated and Tajima's D (Tajima,

1989) estimates were generated for both locations using PoPoolation (Robert Kofler et al., 2011).

Within PoPoolation missing values were assigned for base calls of a phred score <20 and a

corrected Tajima's D was calculated for both geographic sites. Tajima's D estimates were

compared between the two sites within R (via a T-test). To further examine genetic

differentiation between the two geographic sites. PoPoolation2 (R. Kofler, Pandey, & Schlotterer,

2011) was also used to estimate Fisher Exact Tests (per site) and F_{ST} values (per site and sliding

window).

Results

Assessing freeze tolerance

Supercooling point (mean = -3.2 ± 0.51 °C) did not differ between barnacle populations, nor did

SCP differ with basal diameter of test (p > 0.4 for both terms, n = 8 – 15 per population).

Barnacles from the White Sea had significantly higher survival at both exposure temperatures (81

– 95% vs. 27 – 78%; Figure 2; deviance explained = 48.4, df = 1,18, p < 0.001), and exposure to

-10 °C significantly decreased survival in both populations relative to exposure at -6 °C (deviance

explained = 74.8, df = 1, 17, p < 0.001; n = 92 -152 for each combination).

Sequencing and assembly

A total of 776.69 million 181 bp reads were produced from two lanes of Illumina Hiseq 2000,

with an average of 24.75 million/library for the White Sea population and 23.79 million/library

for the British Columbia population. Of these, an average of 14.2 million reads/library in the

White Sea population and 13.1 million reads/library in the British Columbia population passed

quality control by Trimmomatic. These reads were then used to assemble a *de novo*

transcriptome using Trinity, which was then reduced to 61,881 sequences using CD-HIT-EST.

Of these, a total of 61,779 were matched to taxonomic hits by Diamond. We then filtered non-

metazoan sequences (RNA viruses, algae, and other microorganisms) out using MEGAN, and

were left with a total of 46,535 sequences that were an average of 1277 base pairs long (N_{50} =

1084 bp). An average of 4.84 million reads/library in the White Sea population and 3.63 million

reads/library in the British Columbia population were mapped to the final transcriptome.

Pattern of transcript expression

We found the patterns of significantly differentially expressed transcripts after a freezing event

differed strongly between the two populations (Figure 3, Figure 4). In the more freeze tolerant

population (White Sea), no transcripts were differentially expressed during freezing, while after

four hours of recovery, 191 transcripts were differentially expressed (almost evenly split between

up and downregulated). By 28 hours after freezing, this had decreased to 185 transcripts, with

slightly more upregulated than down. This pattern was dramatically different in the less freeze

tolerant population (British Columbia), which had a total of 11,050 transcripts differentially

regulated during the freezing event, the majority of which (9432) were significantly upregulated.

By four hours of recovery, this number had decreased to 51 (again dominated by 46 upregulated

transcripts), and by 28 hours of recovery this had increased to 1739 transcripts (again dominated

by 977 significantly upregulated transcripts).

Identities of differentially expressed transcripts

The identities of the top differentially expressed transcripts following freezing were

functionally very different between the two populations. The few transcripts that both

populations upregulated immediately following freezing were putative transcription factors

(annotated by Blast2Go as "proto-oncogenes"; Table 1). In the British Columbia population, the

most noticeable pattern is the massive upregulation of transcripts during the freezing event

(>9000 transcripts, Figures 3 and 4). This included several ribosomal RNAs, mitochondrially-

encoded electron transport chain subunits, and histone methyltransferases and deacetylases

(Supplementary Data 1). Among the downregulated transcripts includes several that Blast2Go

annotated as "macrophage mannose receptors" which are functionally assigned the "carbohydrate

binding" GO term (Supplementary Data 1). At four hours following freezing, this pattern has

largely disappeared, with only a total of 46 significantly upregulated transcripts (Figure 3 and 4).

These include several transcripts that Blast2Go identified as serine proteases, while the

downregulated transcripts include one identified as "Peter Pan", a transcript annotated as

"macrophage mannose receptor", and an INO80 complex subunit (Supplementary Data 1). By 28

hours following freezing, differential expression has increased again (Figure 3 and 4), with a total

of 50 upregulated transcripts that BLAST to likely proteases and 13 that BLAST to ubiquitin

activity, suggesting significant protein scavenging. By contrast, in the White Sea population, a

large number of transcripts associated with heat shock proteins were the most highly upregulated

transcripts following freezing (Table 2). Two transcripts that were identified as E3 ubiquitin

ligases were also upregulated at both 4 and 28 hours following freezing (Supplementary Data 1).

Four and 28 hours following freezing, the down-regulated transcripts included a variety of

transcripts that perhaps most notably included several mitochondrially-encoded electron transport

chain subunits (Supplementary Data 1) including NADH dehydrogenase subunits 1, 2, 4, 5, and

6, cytochrome b, ATP synthase F_0 subunit 6, as well as cytochrome c oxidase subunits 1, 2, and 3

(Supplementary Data 1). No nuclear-encoded electron transport chain subunits were

downregulated in our dataset.

Functional enrichment of molecular functions of differentially expressed transcripts

The functions of the transcripts similarly differed significantly between the two

populations (Tables 2, 3, Figure 5). In the British Columbia population, we found significant

enrichment of 134 GO terms after REVIGO filtering in the downregulated and 80 in the

upregulated transcripts during freezing, including several electron transport chain-related terms in

both sets (Figure 5). By four hours following freezing there was no significant enrichment of GO

terms, and by 28 hours following freezing there was significant enrichment of 56 GO terms in the

downregulated transcripts with only 15 in the upregulated transcripts (Figure 5). The most

notable of these enriched terms included "UDP-glucose-6-dehydrogenase activity" and

"scavenger receptor activity". By contrast, in the White Sea population, as we had no significant

differential expression of transcripts during freezing, we did not find any significantly enriched

GO terms at this timepoint. While we did not find any significant enrichment of GO terms

among upregulated transcripts in the White Sea population four hours following recovery, we did

find a series of terms associated with the electron transport chain including "oxidoreductase

activity", "heme-copper terminal oxidase activity", and "hydrogen ion transmembrane activity"

that were significantly enriched among transcripts that were significantly downregulated at this

timepoint (Figure 5). Interestingly, all of these downregulated terms were associated with

transcripts that encoded only mitochondrially-encoded electron chain subunits. By 28 hours of

recovery, the pattern had changed significantly with no enriched GO terms among the

downregulated transcripts, but among the significantly upregulated transcripts the terms "glycerol

transport", "cellular water homeostasis", and "water transport" were significantly enriched

(which all were annotated as aquaporins).

Using pathview and Gage to examine significantly enriched gene sets based on KEGG

orthology we found additional functional enrichment. In the British Columbia population, before

thawing we found significant enrichment of the "spliceosome" term and underrepresentation of

the "epithelial cell signaling in *Helicobacter pylori*" term. Four hours after thawing, we observed

significant overrepresentation of the terms "ribosome biogenesis in eukaryotes" and

"spliceosome", and underrepresentation of "Parkinson's disease", "Alzheimer's disease" and

"oxidative phosphorylation", although we note upon examination that the first two terms all

indicated the same underrepresentation of electron transport chain subunits. There was no

functional enrichment 28 hours following freezing in the British Columbia population. Finally,

in the White Sea population we observed significant overrepresentation of terms including

"ribosome biogenesis in eukaryotes", "spliceosome", and "bacterial invasion of epithelial cells"

at 28 hours following freezing, but no terms were overrepresented at any other timepoints.

Population differentiation

The estimated F_{ST} value (0.45; 9,164 loci passing filter) suggests little gene flow occurs between

the two populations. Across all differentially expressed transcripts Tajima D estimates averaged

-1.2 and -1.4 for White Sea and British Columbia respectively, whereas non-differentially

expressed transcripts averaged -0.6 (T-test p-value <0.001) and -0.8 (T-test p-value <0.001) for

White Sea and British Columbia respectively, suggesting selective sweeps on these differentially-

expressed transcripts in each population (Table 4). Tajima's D estimates for aquaporins in the

White Sea population averaged -2.6 compared to the average across differently expressed

transcripts of -1.42, indicative of a potential selective sweep in these transcripts (Table 4).

Discussion

Intertidal organisms constantly risk freezing through the winter. We find, consistent with

our hypothesis, that the northern White Sea population of *S. balanoides* is significantly more

freeze tolerant than the British Columbia population. To our knowledge, there is currently only

one genome-sequenced freeze tolerant animal (the Antarctic midge *Belgica antarctica*; Kelley et

al., 2014) and no genome-sequenced barnacles, so we assembled a *de novo* transcriptome to study

transcriptional responses to freezing stress. In addition, the differing timelines of transcription

suggest two very different sets of processes taking place in each population following a freezing

event, which adds to our limited understanding of the molecular mechanisms of freeze tolerance.

Finally, the low Tajima's D values on these differentially expressed transcripts suggests that local

adaptation to freezing has taken place in the White Sea population. Here we have shown the first

study of transcriptional responses following a freezing stress in a non-insect, and demonstrated

evidence of selection on transcripts associated with freeze tolerance in a natural population.

One of the most puzzling results we found is the massive upregulation of transcripts

during the freezing exposure in the British Columbia population (Figures 3 and 4). There are

three possibilities to explain this pattern: this represents genuine upregulation of transcription,

degradation of mRNA causing greater numbers of smaller transcripts, or damage to translational

machinery causing mRNA to accumulate. The first hypothesis of increased rates of *de novo*

mRNA transcription is possible since the cytoplasm of cells generally remains unfrozen during a

freezing exposure. However this is difficult to reconcile with the generally-accepted rate of

transcription in invertebrates of about 2kb/minute (Ardehali & Lis, 2009), particularly at a

subzero temperature where biochemical events would be expected to be proceed significantly

more slowly. This upregulation is also strikingly absent in the White Sea population, suggesting

increased transcription rates during freezing is not a general nor adaptive response to freezing

stress. The second hypothesis of degradation of mRNA due to osmotic shock or direct freeze-

induced damage is certainly possible, which could increase the total number of detected mRNA

transcripts due to cleavage of larger molecules into several smaller ones. However, we note that

the average transcript length of these upregulated transcripts is significantly larger (1761 bp vs.

1398 bp for the whole transcriptome), which is inconsistent with the possibility of mRNA

degradation. We also note that, at least in *Arabidopsis thaliana*, low temperature tends to

increase mRNA half-life (Chiba et al., 2013). The third hypothesis, cold-induced damage to

ribosomal structure or translation initiation machinery, is possible since translation is associated

with mRNA degradation (Bicknell & Ricci, 2017). At least in bacteria, the ribosome is

particularly sensitive to cold shock and several cellular mechanisms for surviving cold shock are

focused on ribosomal proteins (Phadtare, 2004). In addition, we note several ribosomal mRNAs

are among the top-upregulated transcripts in this experimental group, as well as several

transcripts associated with eukaryotic translation initiation machinery (Supplementary Data 1).

Finally, we also note that at 4 hours following freezing, we find that the KEGG term "ribosome

biogenesis" is significantly enriched among upregulated transcripts in the British Columbia

population, suggesting repair or rebuilding of ribosomal subunits (Table S1). If the ribosomes are

damaged and unable to process mRNA, we might expect a buildup of mRNA (Bicknell & Ricci,

2017). If this is true, we would expect that the White Sea population produces ribosomes that are

more cold shock resistant. Further functional studies, in addition to further transcriptomic work

in freeze tolerant animals, is clearly necessary to differentiate among these potential explanations.

The role(s) of heat shock proteins in freeze tolerance is not well understood, but

upregulation of Hsp70 has been documented in response to repeated freezing in *B. antarctica*

(Teets et al., 2011). While cold-induced denaturation of proteins has been well studied

(Graziano, 2010; Todgham, Hoaglund, & Hofmann, 2007), during survivable freezing significant

desiccation of cells is believed to occur (Storey, 1997; Teets & Denlinger, 2014). It is possible

that this desiccation leads directly to protein denaturation through the disruption of the hydration

shell around proteins during desiccating periods. In the more freeze tolerant White Sea

population, we observed significant upregulation of several shock proteins following freezing—it

is unclear whether these are involved with assisting in refolding of damaged proteins or are

preparatory for a second stress event. By contrast, the less freeze tolerant population upregulated

proteolysis-related transcripts (with significant enrichment of GO terms such as "serine

endopeptidase activity" and "scavenger receptor") following freezing, suggesting significant

damage to protein structure has taken place and the only possible solution is to degrade these

proteins. Our study points to the importance of maintenance of protein stability at low

temperature in an ecological setting.

In addition to direct effects on protein structure, we also saw very different patterns of

differential regulation of aquaporins between the two populations following freezing. In the

White Sea population, we saw significant enrichment of GO categories associated with

aquaporins, which we did not observe in the British Columbia population. This suggests that

only the White Sea population is generally upregulating aquaporin transcription following

freezing. It is believed that aquaporins aid in water transport out of the cell and that

aquaglyceroporins may assist in transport of glycerol into the cell during freezing events (Philip

et al., 2008; Storey & Storey, 2013; Yi et al., 2011). While significant glycerol accumulation is

not associated with freeze tolerance in marine invertebrates (Storey & Storey, 2013), it is likely

that managing water content in the cell is essential for freezing survival. Thus far aquaporins

have been associated with freeze tolerance in the wood frog *Rana sylvatica* (Philip et al., 2008)

and the goldenrod gall fly *Eurosta solidaginis* (Yi et al., 2011)*,* and with this study we

demonstrate their importance in a new class of freeze tolerant organisms, suggesting that this may

be a generalized mechanism for freeze tolerance. In addition, we found very low Tajima's D

values in aquaporins that were upregulated following freezing (Table 4), suggesting that selective

sweeps had taken place on the sequence of these putative aquaporins. Follow-up with functional

studies, particularly employing the frog oocyte expression system, would provide interesting

evidence and further structural information in this poorly understood system (Lind et al., 2017).

The role of antifreeze proteins in freeze tolerance is unclear and little is generally known

about invertebrate antifreeze proteins (Davies, 2014; H. J. Kim et al., 2017). We found many

transcripts (>200) that were annotated with the term "macrophage mannose receptors" that were

upregulated following freezing in both populations. While we cannot definitively assign a

function to these based solely on a transcriptomic study, we do note that these are C-type lectins,

which are believed to be paralogous to Type II antifreeze proteins in fish (Graham, Lougheed,

Ewart, & Davies, 2008; Ng & Hew, 1992). Indeed, when we examined the BLAST results, many

of these sequences had high similarity to Type II antifreeze or ice structuring proteins in fish.

Several freeze tolerant species are thought to produce antifreeze proteins to inhibit

recrystallization of ice at low temperatures (Davies, 2014; Storey & Storey, 2013), although

functional studies are necessary for confirming ice recrystallization inhibition activity.

Antifreeze proteins are found broadly scattered across the domains of life, and appear to readily

evolve (Davies, 2014). In addition, they appear to frequently have been transferred across

species through horizontal gene transfer (Davies, 2014; Kiko, 2010). While this study was not

designed to differentiate between the hypotheses of similar structure due to convergent evolution

or by horizontal gene transfer, further sequencing studies may be able to address this point.

One of the most consistent results we found in both populations is downregulation of

transcripts associated with the electron transport chain. There are two potential explanations for

this downregulation—either transcription of mitochondrially-encoded transcripts was inhibited,

or the number of mitochondria themselves were downregulated following a freezing stress. This

is likely an adaptive response since a degraded or malfunctioning electron transport chain can

cause significant oxidative stress, as can a sudden reduction in ATP demand as we might expect

at low temperatures (Somero, Lockwood, & Tomanek, 2017). Reducing mitochondrial numbers

as a way to reduce metabolic rate is a seasonal adaptation in the overwintering larvae of the moth

Gynaephora groenlandica (Kukal, Duman, & Serianni, 1989), and this may be a similar

phenomena. While we cannot distinguish between these explanations in this study, we

hypothesize that reducing mitochondrial activity facilitates freeze tolerance in our barnacles and

suggest studies isolating mitochondria to measure mitochondrial respiration and histology to

determine mitochondrial numbers following freezing stress.

Conclusions

Here we have presented the first transcriptional time series following freezing in a freeze

tolerant barnacle. By comparing two populations that differ strongly in their freeze tolerance, we

can control for genetic background and thereby infer the transcripts that are associated with

greater freeze tolerance. We find transcriptional evidence for protein damage during a freezing

event, and suggest that heat shock protein upregulation appears to be important for recovery from

freezing. We also document aquaporin expression associated with freezing in a third freeze

tolerant species, and describe a series of new putative antifreeze proteins. These transcripts were

associated with selective sweeps in the White Sea population of barnacles, suggesting local

adaptation to freezing conditions has occurred. Taken together, we shed new light on the

mechanisms of adaptation to the poleward range margin in animals.

Acknowledgements

The authors would like to acknowledge the Company of Biologists for travel funding to KEM, a
University of Oklahoma startup grant and Nigel Fish for computing resources for KEM, as well
as Taiwan-Russian collaboration funds from MOST, Taiwan (106-2923-B-001 -002 -MY3) and
RFBR (grants17-54-52006 MNT_a, 18-04-00624 A, 15-29-02447 ofi_m) and Academia Sinica
Senior Investigator Award (AS-IA-105-L03) to BKKC for funding for sequencing. We would
like to thank the Hakai Institute and Mary O'Connor for collecting barnacles in British Columbia,
as well as for access to temperature data. We would also like to acknowledge Yao-Fong Tsao
and Vanessa Pei-Chen Tsai for laboratory assistance in sample preparation, and Dr. Mei-Yeh Lu
for her invaluable advice for library preparation. Finally, we would like to thank Jantina
Toxopeus and two anonymous referees for comments that have greatly improved the manuscript.

References

Afgan, E., Baker, D., van den Beek, M., Blankenberg, D., Bouvier, D., Čech, M., … Goecks, J. (2016). The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2016 update. *Nucleic Acids Research*, *44*(W1), W3–W10. doi:10.1093/nar/gkw343

Ansart, A., & Vernon, P. (2003). Cold hardiness in molluscs. *Acta Oecologica*, *24*(2), 95–102. doi:10.1016/S1146-609X(03)00045-6

Ardehali, M. B., & Lis, J. T. (2009). Tracking rates of transcription and splicing in vivo. *Nature Structural & Molecular Biology*, *16*(11), 1123–1124. doi:10.1038/nsmb1109-1123

Bicknell, A. A., & Ricci, E. P. (2017). When mRNA translation meets decay. *Biochemical Society Transactions*, *45*(2), 339–351. doi:10.1042/BST20160243

Boardman, L., Terblanche, J. S., & Sinclair, B. J. (2011). Transmembrane ion distribution during recovery from freezing in the woolly bear caterpillar *Pyrrharctia isabella* (Lepidoptera: Arctiidae). *Journal of Insect Physiology*, *57*(8), 1154–1162. doi:10.1016/j.jinsphys.2011.04.022

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)*, *30*(15), 2114–20. doi:10.1093/bioinformatics/btu170

Buchfink, B., Xie, C., & Huson, D. H. (2014). Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, *12*(1), 59–60. doi:10.1038/nmeth.3176

Chan, B. K. K., Lima, F. P., Williams, G. A., Seabra, R., & Wang, H. Y. (2016). A simplified biomimetic temperature logger for recording intertidal barnacle body temperatures. *Limnology and Oceanography: Methods*, *14*(7), 448–455. doi:10.1002/lom3.10103

Chiba, Y., Mineta, K., Hirai, M. Y., Suzuki, Y., Kanaya, S., Takahashi, H., … Naito, S. (2013). Changes in mRNA Stability Associated with Cold Stress in Arabidopsis Cells. *Plant and Cell Physiology*, *54*(2), 180–194. doi:10.1093/pcp/pcs164

Conesa, A., Madrigal, P., Tarazona, S., Gomez-Cabrero, D., Cervera, A., McPherson, A., …
Mortazavi, A. (2016). A survey of best practices for RNA-seq data analysis. *Genome*
Biology, *17*(1), 13. doi:10.1186/s13059-016-0881-8

Cook, P. A., & Gabott, P. A. (1970). Seasonal changes in the free glycerol level in the body parts
of the adult barnacle *Balanus balanoides*. *Marine Biology*, *7*(1), 11–13.
doi:10.1007/BF00346803

Cook, P. A., & Lewis, A. H. (1971). Acquisition and loss of cold-tolerance in adult barnacles
(*Balanus balanoides*) kept under laboratory conditions. *Marine Biology*, *9*(1), 26–30.
doi:10.1007/BF00348814

Crisp, D. J., Davenport, J., & Gabbott, P. A. (1977). Freezing tolerance in *Balanus balanoides*.
Comparative Biochemistry and Physiology Part A: Physiology, *57*(3), 359–361.
doi:10.1016/0300-9629(77)90206-7

Crisp, D. J., & Ritz, D. A. (1967). Changes in temperature tolerance of *Balanus balanoides*
during its life-cycle. *Helgoländer Wissenschaftliche Meeresuntersuchungen*, *15*(1–4), 98–
115. doi:10.1007/BF01618612

Davies, P. L. (2014). Ice-binding proteins: A remarkable diversity of structures for stopping and
starting ice growth. *Trends in Biochemical Sciences*, *39*(11), 548–555.
doi:10.1016/j.tibs.2014.09.005

Dennis, A. B., Loomis, S. H., & Hellberg, M. E. (2014). Latitudinal Variation of Freeze
Tolerance in Intertidal Marine Snails of the Genus *Melampus* (Gastropoda: Ellobiidae).
Physiological and Biochemical Zoology, *87*(4), 517–526. doi:10.1086/676138

Duman, J. G., Bennett, V., Sformo, T., Hochstrasser, R., & Barnes, B. M. (2004). Antifreeze
proteins in Alaskan insects and spiders. *Journal of Insect Physiology*, *50*(4), 259–266.
doi:10.1016/J.JINSPHYS.2003.12.003

Flight, P. A., & Rand, D. M. (2012). Genetic variation in the acorn barnacle from allozymes to
population genomics. *Integrative and Comparative Biology*, *52*(3), 418–429.
doi:10.1093/icb/ics099

Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: accelerated for clustering the next-
547   generation sequencing data. *Bioinformatics*, *28*(23), 3150–3152.
548   doi:10.1093/bioinformatics/bts565

549   Gotz, S., Garcia-Gomez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J., …
550   Conesa, A. (2008). High-throughput functional annotation and data mining with the
551   Blast2GO suite. *Nucleic Acids Research*, *36*(10), 3420–3435. doi:10.1093/nar/gkn176

552   Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., … Regev, A.
553   (2011). Full-length transcriptome assembly from RNA-Seq data without a reference
554   genome. *Nature Biotechnology*, *29*(7), 644–52. doi:10.1038/nbt.1883

555   Graham, L. A., Lougheed, S. C., Ewart, K. V., & Davies, P. L. (2008). Lateral transfer of a
556   lectin-like antifreeze protein gene in fishes. *PLoS ONE*, *3*(7), e2616.
557   doi:10.1371/journal.pone.0002616

558   Graziano, G. (2010). On the molecular origin of cold denaturation of globular proteins. *Physical*
559   *Chemistry Chemical Physics*, *12*(42), 14245. doi:10.1039/c0cp00945h

560   Han, E. N., & Bauce, E. (1995). Non-Freeze Survival of Spruce Budworm Larvae,
561   *Choristoneura-fumiferana*, at Subzero Temperatures during Diapause. *Entomologia*
562   *Experimentalis Et Applicata*, *75*(1), 67–74.

563   Huson, D. H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., … Tappu, R. (2016).
564   MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale
565   Microbiome Sequencing Data. *PLOS Computational Biology*, *12*(6), e1004957.
566   doi:10.1371/journal.pcbi.1004957

567   Johnston, I. A., & Clarke, A. (1990). Cold Adaptation in Marine Organisms [and Discussion].
568   *Philosophical Transactions of the Royal Society B: Biological Sciences*, *326*(1237), 655–
569   667. doi:10.1098/rstb.1990.0037

570   Kelley, J. L., Peyton, J. T., Fiston-Lavier, A.-S., Teets, N. M., Yee, M.-C., Johnston, J. S., …
571   Denlinger, D. L. (2014). Compact genome of the Antarctic midge is likely an adaptation to
572   an extreme environment. *Nature Communications*, *5*(1), 4611. doi:10.1038/ncomms5611

Kiko, R. (2010). Acquisition of freeze protection in a sea-ice crustacean through horizontal gene
transfer? *Polar Biology*, *33*(4), 543–556. doi:10.1007/s00300-009-0732-0

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low
memory requirements. *Nature Methods*, *12*(4), 357–60. doi:10.1038/nmeth.3317

Kim, H. J., Lee, J. H., Hur, Y. B., Lee, C. W., Park, S.-H., & Koo, B.-W. (2017). Marine
Antifreeze Proteins: Structure, Function, and Application to Cryopreservation as a Potential
Cryoprotectant. *Marine Drugs*, *15*(2). doi:10.3390/md15020027

King, A. M., & MacRae, T. H. (2015). Insect Heat Shock Proteins During Stress and Diapause.
Annual Review of Entomology, *60*(1), 59–75. doi:10.1146/annurev-ento-011613-162107

Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik, A., …
Schlötterer, C. (2011). PoPoolation: A Toolbox for Population Genetic Analysis of Next
Generation Sequencing Data from Pooled Individuals. *PLoS ONE*, *6*(1), e15925.
doi:10.1371/journal.pone.0015925

Kofler, R., Pandey, R. V., & Schlotterer, C. (2011). PoPoolation2: identifying differentiation
between populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics*,
27(24), 3435–3436. doi:10.1093/bioinformatics/btr589

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next
Generation Sequencing Data. *BMC Bioinformatics*, *15*(1), 356. doi:10.1186/s12859-014-
0356-4

Koštál, V., Korbelová, J., Poupardin, R., Moos, M., & Šimek, P. (2016). Arginine and proline
applied as food additives stimulate high freeze tolerance in larvae of *Drosophila*
melanogaster. *The Journal of Experimental Biology*, *219*(15), 2358–2367.
doi:10.1242/jeb.142158

Kukal, O., Duman, J. G., & Serianni, A. S. (1989). Cold-induced mitochondrial degradation and
cryoprotectant synthesis in freeze-tolerant arctic caterpillars. *Journal of Comparative*
Physiology B, *158*(6), 661–671. doi:10.1007/BF00693004

Layne, J. R., & Kuharsky, D. K. (2000). Triggering of cryoprotectant synthesis in the woolly bear
caterpillar (*Pyrrharctia isabella* Lepidoptera: Arctiidae). *Journal of Experimental Zoology*,
286(4), 367–371.

Lee, R. E. (2010). A primer on insect cold tolerance. In D. L. Denlinger & R. E. Lee (Eds.), *Low*
Temperature Biology of Insects (pp. 3–34). Cambridge, UK: Cambridge University Press.

Lee, R. E., Dommell, R. A., Joplin, K. H., & Denlinger, D. L. (1995). Cryobiology of the freeze-
tolerant gall fly *Eurosta solidaginis*: overwintering energetics and heat shock proteins.
Climate Research, *5*, 61–67. doi:10.3354/cr005061

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930.
doi:10.1093/bioinformatics/btt656

Lind, U., Järvå, M., Rosenblad, M. A., Pingitore, P., Karlsson, E., Wrange, A. L., … Blomberg,
A. (2017). Analysis of aquaporins from the euryhaline barnacle *Balanus improvisus* reveals
differential expression in response to changes in salinity. *PLoS ONE*, *12*(7), e0181192.
doi:10.1371/journal.pone.0181192

Loomis, S. H., Carpenter, J. F., & Crowe, J. H. (1988). Identification of strombine and taurine as
cryoprotectants in the intertidal bivalve *Mytilus edulis*. *Biochimica et Biophysica Acta*,
943(2), 113–118. doi:10.1016/0005-2736(88)90542-1

Louthan, A. M., Doak, D. F., & Angert, A. L. (2015). Where and When do Species Interactions
Set Range Limits? doi:10.1016/j.tree.2015.09.011

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
doi:10.1186/s13059-014-0550-8

Luo, W., & Brouwer, C. (2013). Pathview: an R/Bioconductor package for pathway-based data
integration and visualization. *Bioinformatics*, *29*(14), 1830–1831.
doi:10.1093/bioinformatics/btt285

Marshall, K. E., & Sinclair, B. J. (2011). The sub-lethal effects of repeated freezing in the woolly
bear caterpillar *Pyrrharctia isabella*. *Journal of Experimental Biology*, *214*(7), 1205–1212.
doi:10.1242/jeb.054569

Marshall, K. E., & Sinclair, B. J. (2015). The relative importance of number, duration and
intensity of cold stress events in determining survival and energetics of an overwintering
insect. *Functional Ecology*, *29*(3), 357–366. doi:10.1111/1365-2435.12328

Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C., & Kanehisa, M. (2007). KAAS: An
automatic genome annotation and pathway reconstruction server. *Nucleic Acids Research*,
35(SUPPL.2), W182–W185. doi:10.1093/nar/gkm321

Murphy, D. J. (1983). Freezing resistance in intertidal invertebrates. *Annual Review of*
Physiology, *45*, 289–299. doi:10.1146/annurev.ph.45.030183.001445

Ng, N. F. L., & Hew, C. L. (1992). Structure of an antifreeze polypeptide from the sea raven:
Disulfide bonds and similarity to lectin-binding proteins. *Journal of Biological Chemistry*,
267(23), 16069–16075. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/1644794

Peng, Y., Arora, R., Li, G., Wang, X., & Fessehaie, A. (2008). Rhododendron catawbiense
plasma membrane intrinsic proteins are aquaporins, and their over-expression compromises
constitutive freezing tolerance and cold acclimation ability of transgenic *Arabidopsis* plants.
Plant, Cell and Environment, *31*(9), 1275–1289. doi:10.1111/j.1365-3040.2008.01840.x

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., & Salzberg, S. L.
(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.
Nature Biotechnology, *33*(3), 290–5. doi:10.1038/nbt.3122

Phadtare, S. (2004). Recent developments in bacterial cold-shock response. *Current Issues in*
Molecular Biology, *6*, 125–136. Retrieved from http://www.caister.com/cimb/v/v6/125.pdf

Philip, B. N., Kiss, A. J., & Lee, R. E. (2011). The protective role of aquaporins in the freeze-
tolerant insect *Eurosta solidaginis*: functional characterization and tissue abundance of
EsAQP1. *The Journal of Experimental Biology*, *214*(Pt 5), 848–857.
doi:10.1242/jeb.051276

Philip, B. N., Yi, S.-X., Elnitsky, M. a, & Lee, R. E. (2008). Aquaporins play a role in

desiccation and freeze tolerance in larvae of the goldenrod gall fly, *Eurosta solidaginis*. *The*

Journal of Experimental Biology, *211*(Pt 7), 1114–1119. doi:10.1242/jeb.016758

Pineda, J., DiBacco, C., & Starczak, V. (2005). Barnacle larvae in ice: Survival, reproduction,

and time to postsettlement metamorphosis. *Limnology and Oceanography*, *50*(5), 1520–

1528. doi:10.4319/lo.2005.50.5.1520

Qin, W., Doucet, D., Tyshenko, M. G., & Walker, V. K. (2007). Transcription of antifreeze

protein genes in *Choristoneura fumiferana*. *Insect Molecular Biology*, *16*(4), 423–434.

doi:10.1111/j.1365-2583.2007.00743.x

Rinehart, J. P., Hayward, S. A. L., Elnitsky, M. A., Sandro, L. H., Lee, R. E., & Denlinger, D. L.

(2006). Continuous up-regulation of heat shock proteins in larvae, but not adults, of a polar

insect. *Proceedings of the National Academy of Sciences*, *103*(38), 14223–14227.

doi:10.1073/pnas.0606840103

Schmidt, P. S., Bertness, M. D., & Rand, D. M. (2000). Environmental heterogeneity and

balancing selection in the acorn barnacle *Semibalanus balanoides*. *Proceedings of the Royal*

Society B, *267*(1441), 379–384. doi:10.1098/rspb.2000.1012

Sexton, J. P., Mcintyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and Ecology of

Species Range Limits. *Annu. Rev. Ecol. Evol. Syst*, *40*, 415–36.

doi:10.1146/annurev.ecolsys.110308.120317

Sinclair, B. J., Addo-Bediako, A., & Chown, S. L. (2003). Climatic variability and the evolution

of insect freeze tolerance. *Biological Reviews*, *78*(2), 181–195.

doi:10.1017/s1464793102006024

Sinclair, B. J., Coello Alvarado, L. E., & Ferguson, L. V. (2015). An invitation to measure insect

cold tolerance: Methods, approaches, and workflow. *Journal of Thermal Biology*.

doi:10.1016/j.jtherbio.2015.11.003

Sinclair, B. J., & Renault, D. (2010). Intracellular ice formation in insects: Unresolved after 50

years? *Comparative Biochemistry and Physiology - A Molecular and Integrative*

Physiology, *155*(1), 14–18. doi:10.1016/j.cbpa.2009.10.026

Somero, G. N., Lockwood, B. L., & Tomanek, L. (2017). Oxygen and Metabolism. In *Biochemical Adaptation* (pp. 29–138).

Sømme, L. (2000). The history of cold hardiness research in terrestrial arthropods. *Cryo-Letters*, *21*(5), 289–296.

Storey, K. B. (1997). Organic Solutes in Freezing Tolerance. *Comparative Biochemistry and Physiology Part A: Physiology*, *117*(3), 319–326. doi:10.1016/S0300-9629(96)00270-8

Storey, K. B., & Storey, J. M. (1996). Natural freezing survival in animals. *Annual Review of Ecology and Systematics*, *27*, 365–386. doi:10.1146/annurev.ecolsys.27.1.365

Storey, K. B., & Storey, J. M. (2013). Molecular biology of freezing tolerance. *Comprehensive Physiology*, *3*(3), 1283–1308. doi:10.1002/cphy.c130007

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global redistribution of animals. *Nature Climate Change*, *2*(9), 686–690. doi:10.1038/nclimate1539

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. *PLoS ONE*, *6*(7), e21800. doi:10.1371/journal.pone.0021800

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, *123*(3), 585–95. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/2513255

Team, R. C. (2016). R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from https://www.r-project.org/

Teets, N. M., & Denlinger, D. L. (2014). Surviving in a frozen desert: environmental stress physiology of terrestrial Antarctic arthropods. *The Journal of Experimental Biology*, *217*(Pt 1), 84–93. doi:10.1242/jeb.089490

Teets, N. M., Kawarasaki, Y., Lee, R. E., & Denlinger, D. L. (2011). Survival and energetic costs of repeated cold exposure in the Antarctic midge, *Belgica antarctica*: a comparison between

frozen and supercooled larvae. *The Journal of Experimental Biology*, *214*(Pt 5), 806–814.

doi:10.1242/jeb.060624

Todgham, A. E., Hoaglund, E. A., & Hofmann, G. E. (2007). Is cold the new hot? Elevated

ubiquitin-conjugated protein levels in tissues of Antarctic fish as evidence for cold-

denaturation of proteins in vivo. *Journal of Comparative Physiology B*, *177*(8), 857–866.

doi:10.1007/s00360-007-0183-2

Toxopeus, J., & Sinclair, B. J. (2018). Mechanisms underlying insect freeze tolerance. *Biological*

Reviews. doi:10.1111/brv.12425

Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., … Pachter,

L. (2010). Transcript assembly and quantification by RNA-Seq reveals unannotated

transcripts and isoform switching during cell differentiation. *Nature Biotechnology*, *28*(5),

511–5. doi:10.1038/nbt.1621

Waller, C. L., Worland, M. R., Convey, P., & Barnes, D. K. a. (2006). Ecophysiological

strategies of Antarctic intertidal invertebrates faced with freezing stress. *Polar Biology*,

29(12), 1077–1083. doi:10.1007/s00300-006-0152-3

Wethey, D. S. (1983). Geographic limits and local zonation: the barnacles *Semibalanus* (*Balanus*)

and *Chthamalus* in New England. *The Biological Bulletin*, *165*(1), 330–341.

doi:10.2307/1541373

Yi, S. X., Benoit, J. B., Elnitsky, M. A., Kaufmann, N., Brodsky, J. L., Zeidel, M. L., … Lee, R.

E. (2011). Function and immuno-localization of aquaporins in the Antarctic midge *Belgica*

antarctica. *Journal of Insect Physiology*, *57*(8), 1096–1105.

doi:10.1016/j.jinsphys.2011.02.006

Data accessibility

The original sequencing reads have been archived at the NCBI Short Read Archive under Bioproject number PRJNA496531 and will be made accessible on publication. The assembled and annotated *Semibalanus balanoides* transcriptome has deposited as a Transcriptome Shotgun Assembly at the NCBI database under the same number. Other data (read counts) have been deposited on the Open Science Framework at XXXXXX and will be made accessible upon publication.

Author Contributions

KEM, AP, GK, and BKKC designed the research

BKKC, AP and GK collected the barnacles from the White Sea

KEM, AP, GK, and BKKC performed the laboratory research

KEM and ED analyzed the data

KEM wrote the first draft of the paper, AP, GK, BKKC, and ED contributed to writing

Figure captions and Table captions

Table 1. Transcripts that are significantly differentially regulated in both British Columbia and White Sea populations. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Table 2. Transcripts that are significantly differentially regulated at both 4 and 28 hours following freezing in the White Sea population. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Table 3. Transcripts that are significantly differentially regulated at both 4 and 28 hours following freezing in the British Columbia population. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Table 4. Tajima D estimates among transcripts. Estimates are averaged across transcripts that either contain or do not contain the search terms. No transcripts annotated as "aquaporins" had sufficient coverage to estimate Tajima D in the British Columbia population. The total number of transcripts passing filter for each test is in brackets. Numbers marked with an asterisk indicate a statistically-significant (t-test, $P < 0.05$) difference between the Tajima D estimates of transcripts containing the search term and those not. Numbers marked with a "^" indicates terms that had insufficient numbers of transcripts to run a t-test.

Figure 1. Outline of experimental design, showing timepoints that barnacles from each population were sampled.

Figure 2. Survival of barnacles from the White Sea and British Columbia populations following a 4 h exposure to either -10 or -6 °C. Asterisk indicates a significant difference in survival between populations in a generalized linear model with binomial error, n = 92 – 151 per population/temperature.

Figure 3. Timecourse of the number of differentially expressed transcripts in barnacles from the White Sea (WS) and British Columbia (BC) with expression calculated relative to control (0 h) for each population using DESeq2.

Figure 4. Venn diagrams illustrating the number of differentially expressed (DE) transcripts following a four hour freezing exposure at -6 °C relative to unfrozen control for each population. Up arrows indicate significantly upregulated transcripts, down arrows indicate significantly downregulated transcripts, and a tilde indicates contraregulated transcripts which are upregulated in one group, but downregulated in another. Areas indicate rank order of number of DE transcripts. A) Number of DE transcripts four hours following freezing between the White Sea and British Columbia populations. B) Number of DE transcripts four hours following freezing between the White Sea and British Columbia populations. C) Number of DE transcripts in the White Sea population (there were no DE transcripts before thawing). D) Number of DE transcripts in the British Columbia population.

Figure 5. Significantly enriched GO terms at each timepoint in each population following a four
hour freezing exposure at -6 °C relative to unfrozen control for each population as determined by
Fisher's exact test and REVIGO. Blue indicates terms enriched in the White Sea population,
while red indicates terms enriched in the British Columbia population. Asterisks indicate terms
with long names that have been shortened for space. All full terms are available in
Supplementary Data 2.

Table 1. Transcripts that are significantly differentially regulated in both British Columbia
and White Sea populations. GO terms are listed only for molecular function. Only the top
5 up and down-regulated (shaded) transcripts are given (supplementary data 1 for all).

Timepoint	Description	Log$_2$ Fold Change in White Sea	Log$_2$ Fold Change in British Columbia	GO Terms
4 hours following freezing	Proto-oncogene c-partial	2.79	6.01	DNA binding; transcription factor activity, sequence-specific DNA binding; sequence-specific DNA binding
	Proto-oncogene c-partial	1.86	2.32	None assigned
	Aromatic-L-amino-acid decarboxylase	1.94	4.12	carboxy-lyase activity
28 hours following freezing	Williams-Beuren syndrome chromosomal region 27	-1.82	-1.17	methyltransferase activity; S-adenosylmethionine-dependent methyltransferase activity
	CAD	-0.92	-1.59	aspartate carbamoyltransferase activity; dihydroorotase activity; carbamoyl-phosphate synthase (ammonia) activity; carbamoyl-phosphate synthase (glutamine-hydrolyzing) activity; ATP

			binding; metal ion binding; amino acid binding
Plasma alpha-L-fucosidase	-1.09	-1.40	alpha-L-fucosidase activity
alpha- partial	1.47	2.44	GTP binding; GTPase activity; structural constituent of cytoskeleton; microtubule-based process
pleckstrin homology domain-containing family F member 2	1.07	0.61	metal ion binding
dystroglycan-like isoform X1	1.04	0.92	calcium ion binding
dystroglycan isoform X1	0.89	0.95	calcium ion binding
Villin quail	0.87	1.79	None assigned

Table 2. Transcripts that are significantly differentially regulated at both 4 and 28 hours following freezing in the White Sea population. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Description	Log$_2$ Fold Change at 4 hours	Log$_2$ Fold Change at 28 hours	GO Names
heat shock partial	2.79	6.01	ATP binding
heat shock 68-like	1.86	2.32	ATP binding
heat shock 70 A1-partial	1.94	4.12	ATP binding
heat shock 70 partial	1.82	1.17	ATP binding
heat shock partial	0.92	1.59	ATP binding
gibberellin 20 oxidase	-1.09	-1.40	oxidoreductase activity
sodium-dependent nutrient amino acid transporter	-1.47	-2.44	symporter activity
BAI1-associated 3	-1.07	-0.61	None assigned
Williams-Beuren syndrome chromosomal region 27	-1.04	-0.92	methyltransferase activity; S-adenosylmethionine-dependent methyltransferase activity
None assigned	-0.89	-0.95	None assigned

802 **Table 3. Transcripts that are significantly differentially regulated at both 4 and 28 hours**
803 **following freezing in the British Columbia population. GO terms are listed only for**
804 **molecular function. Only the top 5 up and down-regulated (shaded) transcripts are given**
805 **(supplementary data 1 for all).**

Description	Log$_2$ Fold Change at 4 hours	Log$_2$ Fold Change at 28 hours	GO Names
Cytochrome partial	6.61	5.62	iron ion binding
Proto-oncogene c- partial	6.31	2.52	transcription factor activity, sequence-specific DNA binding
Proto-oncogene c- partial	6.01	2.28	transcription factor activity, sequence-specific DNA binding
techylectin-5A-like isoform X1	3.71	6.85	carbohydrate binding
Proto-oncogene c- partial	2.32	1.77	None assigned
None assigned	-1.02	-1.01	None assigned
INO80 complex subunit	-0.74	-0.59	(Process: DNA repair)
Peter pan	-0.70	-0.49	rRNA binding

Table 4. Tajima D estimates among transcripts. Estimates are averaged across transcripts
that either contain or do not contain the search terms. No transcripts annotated as
"aquaporins" had sufficient coverage to estimate Tajima D in the British Columbia
population. The total number of transcripts passing filter for each test is in brackets.
Numbers marked with an asterisk indicate a statistically-significant (t-test, P < 0.05)
difference between the Tajima D estimates of transcripts containing the search term and
those not. Numbers marked with a "^" indicates terms that had insufficient numbers of
transcripts to run a t-test.

Search term	British Columbia transcripts not containing search term	British Columbia transcripts containing search term	White Sea transcripts not containing search term	White Sea transcripts containing search term
All Differentially Expressed (DE) Transcripts	-0.64 (13129)	-1.18* (1672)	-0.81 (14765)	-1.42* (875)
Transcripts DE in White Sea	-0.70 (14759)	-1.19* (42)	-0.83 (15393)	-1.44* (247)
"Transcription Factor"	-0.70 (14800)	-1.16$^{\wedge}$ (1)	-0.84 (15631)	-2.48* (9)
"Aquaporin"	NA	NA	-0.84 (15637)	-2.60$^{\wedge}$ (3)
"Macrophage Mannose Receptor"	-0.70 (14799)	-1.11$^{\wedge}$ (2)	-0.84 (15636)	-1.32^ (4)

Before thawing

4h after freezing

28h after freezing

NADH dehydrogenase (ubiquinone) activity
oxidoreductase activity, acting on the aldehyde or oxo group of donors*
oxidoreductase activity, oxidizing metal ions
oligosaccharyl transferase activity
potassium ion binding
drug binding
FK506 binding
ATPase coupled ion transmembrane transporter activity
threonine–type endopeptidase activity
electron carrier activity
RNA polymerase II transcription cofactor activity
peptidyl–prolyl cis–trans isomerase activity
sulfur compound binding
oxidoreductase activity, acting on NAD(P)H
oxidoreductase activity, acting on the aldehyde or oxo group of donors
magnesium ion binding
oxidoreductase activity, acting on the CH–OH group of donors*
signal transducer activity, downstream of receptor
structural constituent of ribosome
translation factor activity, RNA binding
isomerase activity
N–acetyltransferase activity
enzyme binding
protein serine/threonine kinase activity
transcription factor activity, protein binding
coenzyme binding
GTP binding
guanyl nucleotide binding
ubiquitin–like protein transferase activity
ligase activity
transferase activity, transferring glycosyl groups
cofactor binding
oxidoreductase activity
nucleoside phosphate binding
hydrolase activity, acting on acid anhydrides
small molecule binding
anion binding
carbohydrate derivative binding
structural molecule activity
catalytic activity
transferase activity
binding
oxidoreductase activity, acting on diphenols*
oxidoreductase activity, acting on NAD(P)H*
oxidoreductase activity, acting on a heme group of donors
heme–copper terminal oxidase activity
electron carrier activity
oxidoreductase activity, acting on NAD(P)H
hydrogen ion transmembrane transporter activity
oxidoreductase activity
transporter activity
adenylylsulfate kinase activity
sulfate adenylyltransferase (ATP) activity
alpha–L–fucosidase activity
galactosylceramide sulfotransferase activity
nucleobase–containing compound transmembrane transporter activity
hydrolase activity, acting on glycosyl bonds

UDP–glucose 6–dehydrogenase activity
purine–nucleoside phosphorylase activity
protein transmembrane transporter activity
superoxide–generating NADPH oxidase activity
wide pore channel activity
calcium–dependent phospholipid binding
scavenger receptor activity
serine–type endopeptidase activity
aminoacyl–tRNA ligase activity
serine hydrolase activity
chitin binding
actin binding
macromolecular complex binding
calcium ion binding
peptidase activity
hydrolase activity

polyol transmembrane transporter activity
water channel activity
water transmembrane transporter activity
organic hydroxy compound transmembrane transporter activity

N–acetylmuramoyl–L–alanine amidase activity
adenosylhomocysteinase activity
sulfate adenylyltransferase (ATP) activity
hydrolase activity, acting on ether bonds
hydrogen ion transmembrane transporter activity
NADH dehydrogenase activity
electron carrier activity
oxidoreductase activity, acting on NAD(P)H
structural constituent of ribosome
monooxygenase activity
iron ion binding
oxidoreductase activity, acting on paired donors*
hydrolase activity, acting on glycosyl bonds
carbohydrate binding
translation factor activity, RNA binding
coenzyme binding
hydrolase activity, hydrolyzing O–glycosyl compounds
GTP binding
oxidoreductase activity
cofactor binding
guanyl nucleotide binding
actin binding
structural molecule activity
anion binding
transporter activity
carbohydrate derivative binding
nucleoside phosphate binding
small molecule binding
catalytic activity

White Sea
British Columbia

0.0 0.2 0.4 0.6 0.8 1.0 1.0 0.8 0.6 0.4 0.2 0.0
0.0 0.2 0.4 0.6 0.8 1.0 1.0 0.8 0.6 0.4 0.2 0.0
0.0 0.2 0.4 0.6 0.8 1.0 1.0 0.8 0.6 0.4 0.2 0.0

Downregulated   Upregulated
Downregulated   Upregulated
Downregulated   Upregulated

$$\frac{\text{Differentially expressed transcripts}}{\text{Total transcripts in category}}$$